

Crystal structure of tris(1,10-phenanthroline- $\kappa^2 N,N'$)iron(II) bis[bis(trifluoromethylsulfonyl)imide] monohydrate

Kazunori Teramoto,^a Takeshi Kawasaki,^b Toshikazu Nishide^a and Yasuhisa Ikeda^{c*}

^aDepartment of Chemical Biology and Applied Chemistry, College of Engineering, Nihon University, 1 Nakagawara Tokusada Tamura, Koriyama 963-8642, Japan,
^bDepartment of Chemistry, Faculty of Science, Toho University, 2-2-1 Miyama, Funabashi, Chiba 274-8510, Japan, and ^cResearch Laboratory for Nuclear Reactors, Tokyo Institute of Technology, 2-12-1-N1-34 Ookayama, Meguro-ku, Tokyo 152-8550, Japan. *Correspondence e-mail: yikeda@nr.titech.ac.jp

Received 3 December 2014; accepted 9 December 2014

Edited by M. Weil, Vienna University of Technology, Austria

The crystal structure of the title complex, $[Fe(C_{12}H_8N_2)_3][(CF_3SO_2)_2N]_2 \cdot H_2O$, is constructed by one octahedral $[Fe(\text{phen})_3]^{2+}$ (phen = 1,10-phenanthroline) cation (point group symmetry 2), two Tf_2N^- [bis(trifluoromethylsulfonyl)imide] anions, and one water molecule of crystallization (point group 2). The Fe–N bond lengths are indicative of a d^6 low-spin state for the Fe^{II} ion in the complex. The dihedral angle between the phen ligands in the cation is $87.64(6)^\circ$. The Tf_2N^- counter-anion is non-coordinating, with the $-CF_3$ groups arranged in a *trans* fashion with respect to each other, leading to an *anti,anti* conformation of the $-CF_3$ groups and $-SO_2N^-$ moieties relative to the S–C bonds. The water molecule of crystallization connects two O atoms of the Tf_2N^- anions through weak hydrogen bonds. C–H \cdots O hydrogen-bonding interactions are also observed, consolidating the packing of the molecules into a three-dimensional network structure.


Keywords: crystal structure; 1,10-phenanthroline; iron(II) complex; complex salt; bis(trifluoromethylsulfonyl)imide; low-spin d^6 Fe^{II} ions; hydrogen bonding.

CCDC reference: 1038289

1. Related literature

For the synthesis of the anhydrous title complex, see: Teramoto *et al.* (2014). For typical Fe–N bond lengths of low-spin d^6 Fe^{II} ions, see: Deng *et al.* (2001); Setifi *et al.* (2013). Crystal structures of complexes with the $[Fe(\text{phen})_3]^{2+}$ cation were reported by Koh (1994), Potočnák *et al.* (2014) and Zhong

(2012). In the crystal structure of the ionic liquid choline bis(trifluoromethylsulfonyl)imide (Nockemann *et al.*, 2009), the free Tf_2N^- anion has a similar conformation to that in the title compound.



2. Experimental

2.1. Crystal data

$[Fe(C_{12}H_8N_2)_3](C_2F_6NO_4S_2)_2 \cdot H_2O$
 $M_r = 1174.78$
Monoclinic, $C2/c$
 $a = 20.7745(15)$ Å
 $b = 16.0107(12)$ Å
 $c = 13.3084(10)$ Å
 $\beta = 91.657(1)^\circ$

$$V = 4424.7(6) \text{ \AA}^3$$

$$Z = 4$$

Mo $K\alpha$ radiation

$$\mu = 0.65 \text{ mm}^{-1}$$

$$T = 100 \text{ K}$$

$$0.42 \times 0.11 \times 0.10 \text{ mm}$$

2.2. Data collection

Bruker APEXII CCD area-detector diffractometer
Absorption correction: multi-scan (*SADABS*; Bruker, 2007)
 $T_{\min} = 0.773$, $T_{\max} = 0.938$

13659 measured reflections
4910 independent reflections
3247 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.065$

2.3. Refinement

$R[F^2 > 2\sigma(F^2)] = 0.048$
 $wR(F^2) = 0.122$
 $S = 1.02$
4910 reflections
338 parameters
2 restraints

H atoms treated by a mixture of independent and constrained refinement
 $\Delta\rho_{\max} = 0.71 \text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.90 \text{ e \AA}^{-3}$

Table 1
Selected bond lengths (Å).

Fe1–N1	1.977 (3)	Fe1–N3	1.966 (3)
Fe1–N2	1.974 (3)		

Table 2
Hydrogen-bond geometry (Å, °).

D–H \cdots A	D–H	H \cdots A	D \cdots A	D–H \cdots A
O5–H5W \cdots O4 ⁱ	0.88 (1)	2.23 (6)	2.963 (4)	141 (7)
C2–H2 \cdots O3 ⁱⁱ	0.95	2.50	3.433 (4)	166
C14–H14 \cdots O2	0.95	2.53	3.481 (5)	174

Symmetry codes: (i) $x - \frac{1}{2}$, $y - \frac{1}{2}$, z ; (ii) $-x + 1$, $-y + 2$, $-z + 1$.

Data collection: *APEX2* (Bruker, 2007); cell refinement: *SAINT* (Bruker, 2007); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL* and *PLATON* (Spek, 2009).

Acknowledgements

We thank Professor Takafumi Kitazawa of Toho University for his useful comments.

Supporting information for this paper is available from the IUCr electronic archives (Reference: WM5100).

References

- Bruker (2007). *APEX2, SAINT* and *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Deng, R. M. K., Simon, S., Dillon, K. B. & Goeta, A. E. (2001). *Acta Cryst. C* **57**, 4–6.
- Koh, L. L., Xu, Y., Hsieh, A. K., Song, B., Wu, F. & Ji, L. (1994). *Acta Cryst. C* **50**, 884–886.
- Nockemann, P., Binnemans, K., Thijss, B., Parac-Vogt, T. N., Merz, K., Mudring, A.-V., Menon, P. C., Rajesh, R. N., Cordoyannis, G., Thoen, J., Leys, J. & Glorieux, C. (2009). *J. Phys. Chem. B*, **113**, 1429–1437.
- Potočnák, I., Váhovská, L. & Herich, P. (2014). *Acta Cryst. C* **70**, 432–436.
- Setifi, Z., Setifi, F., Ng, S. W., Oudahmane, A., El-Ghazzi, M. & Avignant, D. (2013). *Acta Cryst. E* **69**, m12–m13.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Spek, A. L. (2009). *Acta Cryst. D* **65**, 148–155.
- Teramoto, K., Nishide, T., Okumura, S., Takao, K. & Ikeda, Y. (2014). *Electrochemistry*, **82**, 566–572.
- Zhong, K.-L. (2012). *Acta Cryst. C* **68**, m259–m264.

supporting information

Acta Cryst. (2015). E71, m8–m9 [doi:10.1107/S2056989014026966]

Crystal structure of tris(1,10-phenanthroline- κ^2N,N')iron(II) bis[bis(trifluoromethylsulfonyl)imide] monohydrate

Kazunori Teramoto, Takeshi Kawasaki, Toshikazu Nishide and Yasuhisa Ikeda

S1. Experimental

Red powders of $[Fe(\text{phen})_3](\text{Tf}_2\text{N})_2$ were synthesized as described in the literature by Teramoto *et al.* (2014). The title complex was crystallized by cooling a hot concentrated aqueous solution of $[Fe(\text{phen})_3](\text{Tf}_2\text{N})_2$.

S2. Refinement

The H atom of the water molecule was located in a difference map and was refined by applying a restraint for the O—H bond length (0.85 (1) Å). The remaining H atoms were positioned geometrically with C—H = 0.95 Å. All H atoms were constrained to ride on their parent atoms with $U_{\text{iso}}(\text{H}) = 1.2 U_{\text{eq}}(\text{C})$ or $U_{\text{iso}}(\text{H}) = 1.5 U_{\text{eq}}(\text{O})$.





Figure 1


View of the $[Fe(\text{phen})_3]^{2+}$, Tf_2N^- and H_2O molecular units. Displacement ellipsoids are represented at the 30% probability level. Hydrogen atoms were omitted for clarity. [Symmetry code: i) $-x + 1, y, -z + 1/2$.]

**Figure 2**

Hydrogen bonds between the H₂O molecule and Tf₂N⁻ anions. [Symmetry codes: ii) $x - 1/2, y - 1/2, z$, iii) $-x + 3/2, y - 1/2, -z + 3/2$.]

Tris(1,10-phenanthroline- κ^2N,N')iron(II) bis[bis(trifluoromethylsulfonyl)imide] monohydrate

Crystal data



$$M_r = 1174.78$$

Monoclinic, C2/c

Hall symbol: -C 2yc

$$a = 20.7745 (15) \text{ \AA}$$

$$b = 16.0107 (12) \text{ \AA}$$

$$c = 13.3084 (10) \text{ \AA}$$

$$\beta = 91.657 (1)^\circ$$

$$V = 4424.7 (6) \text{ \AA}^3$$

$$Z = 4$$

$$F(000) = 2368$$

$$D_x = 1.764 \text{ Mg m}^{-3}$$

Mo K α radiation, $\lambda = 0.71073 \text{ \AA}$

Cell parameters from 1623 reflections

$$\theta = 2.2\text{--}23.5^\circ$$

$$\mu = 0.65 \text{ mm}^{-1}$$

$$T = 100 \text{ K}$$

Block, red

$$0.42 \times 0.11 \times 0.10 \text{ mm}$$

Data collection

Bruker APEXII CCD area-detector
diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

Detector resolution: 8.333 pixels mm⁻¹

phi and ω scans

Absorption correction: multi-scan
(SADABS; Bruker, 2007)

$$T_{\min} = 0.773, T_{\max} = 0.938$$

$$13659 \text{ measured reflections}$$

$$4910 \text{ independent reflections}$$

$$3247 \text{ reflections with } I > 2\sigma(I)$$

$$R_{\text{int}} = 0.065$$

$$\theta_{\max} = 27.2^\circ, \theta_{\min} = 1.6^\circ$$

$$h = -26 \rightarrow 24$$

$$k = -17 \rightarrow 20$$

$$l = -17 \rightarrow 16$$

Refinement

Refinement on F^2

Least-squares matrix: full

$$R[F^2 > 2\sigma(F^2)] = 0.048$$

$$wR(F^2) = 0.122$$

$$S = 1.02$$

$$4910 \text{ reflections}$$

$$338 \text{ parameters}$$

$$2 \text{ restraints}$$

Primary atom site location: structure-invariant
direct methods

Secondary atom site location: difference Fourier
map

Hydrogen site location: inferred from
neighbouring sites

H atoms treated by a mixture of independent
and constrained refinement

$$w = 1/[\sigma^2(F_o^2) + (0.0533P)^2 + 0.4369P]$$

where $P = (F_o^2 + 2F_c^2)/3$

$$(\Delta/\sigma)_{\max} = 0.001$$

$$\Delta\rho_{\max} = 0.71 \text{ e \AA}^{-3}$$

$$\Delta\rho_{\min} = -0.90 \text{ e \AA}^{-3}$$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Fe1	0.5000	0.81145 (4)	0.2500	0.01247 (17)
S1	0.70284 (4)	0.87060 (6)	0.73999 (6)	0.0194 (2)
S2	0.81435 (4)	0.96272 (6)	0.78549 (6)	0.0190 (2)
N1	0.41901 (12)	0.80991 (17)	0.32441 (19)	0.0128 (6)
N2	0.52464 (12)	0.72317 (17)	0.34722 (19)	0.0136 (6)
N3	0.53087 (12)	0.90330 (18)	0.33681 (19)	0.0141 (6)
N4	0.77157 (14)	0.88188 (19)	0.7924 (2)	0.0231 (7)
O1	0.66576 (12)	0.81717 (18)	0.80049 (19)	0.0320 (7)
O2	0.67338 (11)	0.94387 (16)	0.69810 (18)	0.0243 (6)
O3	0.80627 (11)	1.01191 (16)	0.69622 (17)	0.0235 (6)
O4	0.87849 (11)	0.94170 (17)	0.81960 (19)	0.0264 (6)
O5	0.5000	0.5160 (4)	0.7500	0.102 (2)
H5W	0.4670 (6)	0.4824 (12)	0.742 (7)	0.153*
F1	0.75397 (11)	0.73840 (14)	0.65849 (17)	0.0369 (6)
F2	0.75628 (11)	0.84768 (16)	0.56672 (17)	0.0409 (6)
F3	0.66724 (10)	0.78257 (14)	0.58392 (17)	0.0343 (6)
F4	0.78834 (11)	0.98981 (14)	0.97358 (15)	0.0318 (6)
F5	0.82027 (10)	1.09733 (13)	0.89196 (16)	0.0281 (5)
F6	0.72349 (9)	1.05100 (14)	0.86804 (15)	0.0283 (5)
C1	0.36704 (15)	0.8593 (2)	0.3148 (2)	0.0150 (7)
H1	0.3646	0.8974	0.2601	0.018*
C2	0.31626 (16)	0.8573 (2)	0.3816 (2)	0.0175 (8)
H2	0.2807	0.8941	0.3725	0.021*
C3	0.31802 (15)	0.8018 (2)	0.4603 (3)	0.0177 (8)
H3	0.2841	0.8008	0.5067	0.021*
C4	0.37042 (16)	0.7465 (2)	0.4717 (2)	0.0160 (7)
C5	0.42054 (15)	0.7547 (2)	0.4026 (2)	0.0143 (7)
C6	0.37791 (16)	0.6843 (2)	0.5492 (3)	0.0190 (8)
H6	0.3445	0.6769	0.5956	0.023*
C7	0.43081 (16)	0.6365 (2)	0.5575 (2)	0.0197 (8)
H7	0.4339	0.5955	0.6090	0.024*
C8	0.47740 (15)	0.7066 (2)	0.4134 (2)	0.0153 (7)
C9	0.48305 (16)	0.6464 (2)	0.4898 (2)	0.0164 (7)
C10	0.54083 (16)	0.5999 (2)	0.4945 (3)	0.0206 (8)
H10	0.5475	0.5585	0.5449	0.025*

C11	0.58709 (17)	0.6152 (2)	0.4258 (2)	0.0193 (8)
H11	0.6255	0.5830	0.4268	0.023*
C12	0.57795 (16)	0.6777 (2)	0.3545 (2)	0.0173 (8)
H12	0.6114	0.6884	0.3090	0.021*
C13	0.56412 (15)	0.9013 (2)	0.4249 (2)	0.0177 (8)
H13	0.5741	0.8486	0.4542	0.021*
C14	0.58467 (16)	0.9738 (2)	0.4751 (3)	0.0199 (8)
H14	0.6086	0.9698	0.5368	0.024*
C15	0.57025 (16)	1.0499 (2)	0.4352 (3)	0.0209 (8)
H15	0.5840	1.0993	0.4692	0.025*
C16	0.53488 (16)	1.0558 (2)	0.3435 (3)	0.0178 (8)
C17	0.51693 (15)	0.9803 (2)	0.2973 (2)	0.0153 (7)
C18	0.51659 (16)	1.1325 (2)	0.2942 (3)	0.0214 (8)
H18	0.5282	1.1843	0.3246	0.026*
C19	0.72140 (18)	0.8058 (2)	0.6315 (3)	0.0236 (8)
C20	0.78431 (17)	1.0287 (2)	0.8856 (3)	0.0213 (8)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Fe1	0.0125 (3)	0.0142 (4)	0.0107 (3)	0.000	0.0000 (3)	0.000
S1	0.0223 (5)	0.0212 (5)	0.0148 (4)	0.0016 (4)	-0.0001 (4)	0.0002 (4)
S2	0.0201 (5)	0.0208 (5)	0.0161 (4)	0.0048 (4)	-0.0013 (4)	-0.0022 (4)
N1	0.0138 (14)	0.0118 (15)	0.0125 (14)	-0.0030 (12)	-0.0016 (11)	-0.0011 (11)
N2	0.0119 (14)	0.0160 (16)	0.0128 (14)	0.0010 (12)	-0.0006 (11)	-0.0017 (12)
N3	0.0112 (13)	0.0188 (16)	0.0124 (14)	0.0001 (12)	0.0017 (11)	-0.0010 (12)
N4	0.0255 (16)	0.0209 (18)	0.0223 (16)	0.0033 (14)	-0.0104 (14)	-0.0022 (13)
O1	0.0329 (15)	0.0405 (19)	0.0229 (14)	-0.0055 (13)	0.0053 (12)	0.0070 (13)
O2	0.0242 (13)	0.0233 (16)	0.0249 (14)	0.0049 (11)	-0.0071 (11)	-0.0013 (11)
O3	0.0247 (13)	0.0297 (16)	0.0162 (13)	0.0038 (11)	0.0001 (11)	0.0037 (11)
O4	0.0198 (13)	0.0316 (17)	0.0276 (14)	0.0076 (12)	-0.0043 (11)	-0.0029 (12)
O5	0.088 (5)	0.098 (6)	0.120 (6)	0.000	-0.005 (5)	0.000
F1	0.0399 (14)	0.0241 (14)	0.0461 (15)	0.0071 (11)	-0.0105 (12)	-0.0124 (11)
F2	0.0504 (15)	0.0476 (17)	0.0257 (12)	-0.0160 (12)	0.0163 (12)	-0.0074 (11)
F3	0.0352 (13)	0.0325 (14)	0.0343 (13)	-0.0027 (11)	-0.0124 (11)	-0.0101 (11)
F4	0.0465 (14)	0.0333 (14)	0.0153 (11)	0.0019 (11)	-0.0021 (10)	0.0005 (10)
F5	0.0301 (12)	0.0205 (13)	0.0334 (13)	-0.0013 (10)	-0.0050 (10)	-0.0065 (10)
F6	0.0210 (11)	0.0354 (14)	0.0283 (12)	0.0071 (10)	-0.0019 (9)	-0.0127 (10)
C1	0.0140 (16)	0.0155 (19)	0.0152 (17)	0.0005 (14)	-0.0026 (14)	-0.0021 (14)
C2	0.0137 (17)	0.018 (2)	0.0201 (18)	0.0011 (14)	-0.0022 (14)	-0.0013 (15)
C3	0.0131 (16)	0.020 (2)	0.0195 (18)	-0.0058 (14)	0.0033 (14)	-0.0044 (15)
C4	0.0156 (17)	0.017 (2)	0.0156 (17)	-0.0053 (14)	-0.0012 (14)	-0.0030 (14)
C5	0.0174 (17)	0.0144 (19)	0.0111 (16)	-0.0036 (14)	-0.0003 (14)	-0.0037 (13)
C6	0.0212 (18)	0.0158 (19)	0.0201 (18)	-0.0044 (15)	0.0036 (15)	-0.0002 (15)
C7	0.0262 (19)	0.020 (2)	0.0136 (17)	-0.0033 (16)	0.0047 (15)	0.0024 (15)
C8	0.0170 (17)	0.0142 (19)	0.0145 (17)	-0.0051 (14)	-0.0006 (14)	-0.0030 (14)
C9	0.0202 (18)	0.016 (2)	0.0125 (17)	-0.0021 (15)	-0.0025 (14)	-0.0031 (14)
C10	0.0284 (19)	0.019 (2)	0.0143 (17)	0.0016 (16)	-0.0038 (15)	0.0010 (15)

C11	0.0193 (17)	0.021 (2)	0.0172 (18)	0.0057 (15)	-0.0021 (14)	0.0007 (15)
C12	0.0168 (17)	0.019 (2)	0.0157 (17)	0.0019 (15)	0.0002 (14)	-0.0026 (14)
C13	0.0139 (17)	0.025 (2)	0.0135 (17)	0.0044 (15)	-0.0012 (14)	0.0009 (15)
C14	0.0147 (17)	0.029 (2)	0.0163 (18)	0.0011 (16)	-0.0010 (14)	-0.0068 (15)
C15	0.0174 (18)	0.022 (2)	0.0231 (19)	-0.0041 (15)	0.0027 (15)	-0.0117 (16)
C16	0.0153 (17)	0.019 (2)	0.0195 (18)	-0.0003 (15)	0.0032 (14)	-0.0038 (15)
C17	0.0127 (17)	0.019 (2)	0.0148 (17)	0.0025 (14)	0.0038 (14)	-0.0014 (14)
C18	0.0191 (19)	0.015 (2)	0.030 (2)	0.0010 (14)	0.0018 (15)	-0.0055 (16)
C19	0.028 (2)	0.019 (2)	0.023 (2)	-0.0039 (17)	-0.0014 (16)	-0.0033 (16)
C20	0.0235 (19)	0.025 (2)	0.0156 (18)	0.0052 (16)	-0.0017 (15)	-0.0019 (15)

Geometric parameters (\AA , $^{\circ}$)

Fe1—N1	1.977 (3)	C2—C3	1.372 (5)
Fe1—N1 ⁱ	1.977 (3)	C2—H2	0.9500
Fe1—N2	1.974 (3)	C3—C4	1.408 (5)
Fe1—N2 ⁱ	1.974 (3)	C3—H3	0.9500
Fe1—N3	1.966 (3)	C4—C5	1.415 (4)
Fe1—N3 ⁱ	1.966 (3)	C4—C6	1.439 (5)
S1—O1	1.417 (3)	C5—C8	1.414 (5)
S1—O2	1.429 (3)	C6—C7	1.341 (5)
S1—N4	1.581 (3)	C6—H6	0.9500
S1—C19	1.827 (4)	C7—C9	1.439 (5)
S2—O3	1.431 (2)	C7—H7	0.9500
S2—O4	1.435 (2)	C8—C9	1.403 (5)
S2—N4	1.574 (3)	C9—C10	1.413 (5)
S2—C20	1.824 (4)	C10—C11	1.368 (5)
N1—C1	1.341 (4)	C10—H10	0.9500
N1—C5	1.366 (4)	C11—C12	1.388 (5)
N2—C8	1.364 (4)	C11—H11	0.9500
N2—C12	1.326 (4)	C12—H12	0.9500
N3—C13	1.343 (4)	C13—C14	1.400 (5)
N3—C17	1.368 (4)	C13—H13	0.9500
O5—H5W	0.875 (10)	C14—C15	1.360 (5)
F1—C19	1.318 (4)	C14—H14	0.9500
F2—C19	1.324 (4)	C15—C16	1.410 (5)
F3—C19	1.329 (4)	C15—H15	0.9500
F4—C20	1.327 (4)	C16—C17	1.401 (5)
F5—C20	1.330 (4)	C16—C18	1.439 (5)
F6—C20	1.327 (4)	C17—C17 ⁱ	1.425 (6)
C1—C2	1.400 (4)	C18—C18 ⁱ	1.345 (7)
C1—H1	0.9500	C18—H18	0.9500
N3—Fe1—N3 ⁱ	83.17 (16)	C7—C6—C4	121.7 (3)
N3—Fe1—N2 ⁱ	174.17 (11)	C7—C6—H6	119.1
N3 ⁱ —Fe1—N2 ⁱ	94.38 (11)	C4—C6—H6	119.1
N3—Fe1—N2	94.38 (11)	C6—C7—C9	121.1 (3)
N3 ⁱ —Fe1—N2	174.17 (11)	C6—C7—H7	119.5

N2 ⁱ —Fe1—N2	88.54 (16)	C9—C7—H7	119.5
N3—Fe1—N1 ⁱ	92.05 (11)	N2—C8—C9	123.7 (3)
N3 ⁱ —Fe1—N1 ⁱ	89.01 (11)	N2—C8—C5	116.3 (3)
N2 ⁱ —Fe1—N1 ⁱ	82.60 (11)	C9—C8—C5	120.0 (3)
N2—Fe1—N1 ⁱ	96.38 (11)	C8—C9—C10	116.6 (3)
N3—Fe1—N1	89.01 (11)	C8—C9—C7	118.7 (3)
N3 ⁱ —Fe1—N1	92.05 (11)	C10—C9—C7	124.7 (3)
N2 ⁱ —Fe1—N1	96.38 (11)	C11—C10—C9	119.2 (3)
N2—Fe1—N1	82.60 (11)	C11—C10—H10	120.4
N1 ⁱ —Fe1—N1	178.58 (17)	C9—C10—H10	120.4
O1—S1—O2	118.91 (16)	C10—C11—C12	120.1 (3)
O1—S1—N4	108.49 (16)	C10—C11—H11	120.0
O2—S1—N4	116.70 (16)	C12—C11—H11	120.0
O1—S1—C19	103.70 (17)	N2—C12—C11	122.9 (3)
O2—S1—C19	104.80 (16)	N2—C12—H12	118.6
N4—S1—C19	101.81 (17)	C11—C12—H12	118.6
O3—S2—O4	118.49 (16)	N3—C13—C14	122.6 (3)
O3—S2—N4	116.61 (16)	N3—C13—H13	118.7
O4—S2—N4	108.02 (16)	C14—C13—H13	118.7
O3—S2—C20	104.65 (16)	C15—C14—C13	119.8 (3)
O4—S2—C20	103.83 (15)	C15—C14—H14	120.1
N4—S2—C20	103.19 (17)	C13—C14—H14	120.1
C1—N1—C5	117.1 (3)	C14—C15—C16	120.1 (3)
C1—N1—Fe1	129.7 (2)	C14—C15—H15	120.0
C5—N1—Fe1	112.9 (2)	C16—C15—H15	120.0
C12—N2—C8	117.4 (3)	C17—C16—C15	116.6 (3)
C12—N2—Fe1	129.9 (2)	C17—C16—C18	118.2 (3)
C8—N2—Fe1	112.6 (2)	C15—C16—C18	125.2 (3)
C13—N3—C17	117.1 (3)	N3—C17—C16	123.9 (3)
C13—N3—Fe1	130.2 (2)	N3—C17—C17 ⁱ	115.67 (18)
C17—N3—Fe1	112.7 (2)	C16—C17—C17 ⁱ	120.4 (2)
S1—N4—S2	124.92 (19)	C18 ⁱ —C18—C16	121.4 (2)
N1—C1—C2	123.0 (3)	C18 ⁱ —C18—H18	119.3
N1—C1—H1	118.5	C16—C18—H18	119.3
C2—C1—H1	118.5	F1—C19—F2	107.8 (3)
C3—C2—C1	119.7 (3)	F1—C19—F3	108.7 (3)
C3—C2—H2	120.1	F2—C19—F3	107.6 (3)
C1—C2—H2	120.1	F1—C19—S1	111.6 (3)
C2—C3—C4	119.5 (3)	F2—C19—S1	111.1 (3)
C2—C3—H3	120.2	F3—C19—S1	109.9 (3)
C4—C3—H3	120.2	F4—C20—F5	108.1 (3)
C3—C4—C5	116.9 (3)	F4—C20—F6	108.5 (3)
C3—C4—C6	125.4 (3)	F5—C20—F6	108.6 (3)
C5—C4—C6	117.6 (3)	F4—C20—S2	111.0 (3)
N1—C5—C8	115.5 (3)	F5—C20—S2	108.8 (2)

N1—C5—C4	123.6 (3)	F6—C20—S2	111.8 (2)
C8—C5—C4	120.8 (3)		

Symmetry code: (i) $-x+1, y, -z+1/2$.

Hydrogen-bond geometry (\AA , $^\circ$)

$D\text{—H}\cdots A$	$D\text{—H}$	$H\cdots A$	$D\cdots A$	$D\text{—H}\cdots A$
O5—H5W \cdots O4 ⁱⁱ	0.88 (1)	2.23 (6)	2.963 (4)	141 (7)
C2—H2 \cdots O3 ⁱⁱⁱ	0.95	2.50	3.433 (4)	166
C14—H14 \cdots O2	0.95	2.53	3.481 (5)	174

Symmetry codes: (ii) $x-1/2, y-1/2, z$; (iii) $-x+1, -y+2, -z+1$.