

5,6-Dihydroxy-7,8-dimethoxyflavone

Lin-Lin Jing, Xiao-Fei Fan, Peng-Cheng Fan, Lei He and Zheng-Ping Jia*

Department of Pharmacy, Lanzhou General Hospital of PLA, Key laboratory of the prevention and cure for the plateau environment damage PLA, 730050, Lanzhou Gansu, People's Republic of China

Correspondence e-mail: zhengping_jia@yahoo.cn


Received 12 April 2013; accepted 25 May 2013

Key indicators: single-crystal X-ray study; $T = 295\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.004\text{ \AA}$; R factor = 0.070; wR factor = 0.236; data-to-parameter ratio = 15.9.

The title compound (systematic name: 5,6-dihydroxy-7,8-dimethoxy-2-phenylchromen-4-one), $C_{17}\text{H}_{14}\text{O}_6$, is a flavone that was isolated from the petroleum ether-soluble fraction of the rare traditional Chinese medicinal herb *Saussurea involucrata*. The flavone molecule is almost planar, with a dihedral angle between the planes of the benzopyran-4-one group and the attached phenyl group of $1.89(6)^\circ$. The 5-hydroxy group forms a strong intramolecular hydrogen bond with the carbonyl group, resulting in a six-membered hydrogen-bonded ring. The 6-hydroxy group also forms an intramolecular O—H···O contact. In the crystal, the molecules are linked by O—H···O and C—H···O hydrogen bonds and π – π interactions [$3.37(2)$ – $3.39(2)\text{ \AA}$], which build up a three-dimensional network.

Related literature

For biological activity of *Saussurea involucrata*, see: Zheng *et al.* (1993); Gao *et al.* (2005); Tao *et al.* (2010); Ma *et al.* (2011); Jia *et al.* (2005); Liu *et al.* (1985). For related structures, see: Xiong *et al.* (2009); Vijayalakshmi *et al.* (1986); Paula *et al.* (2002).



Experimental

Crystal data

$C_{17}\text{H}_{14}\text{O}_6$
 $M_r = 314.28$
Triclinic, $P\bar{1}$

Data collection

Bruker APEXII CCD diffractometer
5252 measured reflections

3368 independent reflections
1786 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.036$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.070$
 $wR(F^2) = 0.236$
 $S = 1.03$
3368 reflections

212 parameters
H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 0.32\text{ e \AA}^{-3}$
 $\Delta\rho_{\text{min}} = -0.32\text{ e \AA}^{-3}$

Table 1
Hydrogen-bond geometry (\AA , $^\circ$).

| $D-\text{H} \cdots A$ | $D-\text{H}$ | $\text{H} \cdots A$ | $D \cdots A$ | $D-\text{H} \cdots A$ |
|----------------------------|--------------|---------------------|--------------|-----------------------|
| O4—H4···O3 ⁱ | 0.82 | 2.05 | 2.764 (4) | 146 |
| C12—H12···O6 ⁱⁱ | 0.93 | 2.58 | 3.234 (4) | 128 |
| O3—H3···O2 | 0.82 | 1.84 | 2.564 (3) | 146 |
| O4—H4···O3 | 0.82 | 2.34 | 2.767 (3) | 113 |

Symmetry codes: (i) $-x, -y + 1, -z$; (ii) $-x + 2, -y + 1, -z + 1$.

Data collection: *APEX2* (Bruker, 2007); cell refinement: *SAINT* (Bruker, 2007); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3* (Farrugia, 2012); software used to prepare material for publication: *SHELXTL* (Sheldrick, 2008) and *PLATON* (Spek, 2009).

We thank the Natural Science Foundation of China (grant No. 81202458) and China Postdoctoral Science Foundation (grant No. 2012M521926) for financial support.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: RK2401).

References

- Bruker (2007). *APEX2* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Farrugia, L. J. (2012). *J. Appl. Cryst.* **45**, 849–854.
- Gao, X., Zhang, Z. M., Xu, A. X. & Lei, X. Y. (2005). *Zhongguo Yao Xue Za Zhi*, **40**, 1062–1065.
- Jia, J. M., Wu, C. F. & Liu, W. (2005). *Biol. Pharm. Bull.* **9**, 1612–1614.
- Liu, L. S., Xiao, X. H. & Zhang, L. D. J. (1985). *Lanzhou Univ. Nat. Sci.*, **4**, 80–83.
- Ma, H. P., Fan, P. C., Jing, L. L., Yao, J., He, X. Y., Yang, Y., Chen, K. M. & Jia, Z. P. (2011). *J. Ethnopharmacol.* **137**, 1510–1515.
- Paula, V. F., Barbosa, L. C. A., Errington, W., Howarth, O. W. & Cruz, M. P. (2002). *J. Braz. Chem. Soc.* **13**, 276–280.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Spek, A. L. (2009). *Acta Cryst. D* **65**, 148–155.
- Tao, Y., Zhong, Z. Z., Zhi, L. Y. & Chen, H. B. (2010). *J. Ethnopharmacol.* **128**, 405–411.
- Vijayalakshmi, J., Rajan, S. S., Srinivasan, R. & Ramachandran Nair, A. G. (1986). *Acta Cryst. C* **42**, 1752–1754.
- Xiong, H.-P., Wu, Z.-J., Chen, F.-T. & Chen, W.-S. (2009). *Acta Cryst. E* **65**, o3276–o3277.
- Zheng, R. L., Liu, G. S., Xing, G. X., Jia, Z. J., Du, M. & Tan, L. Q. (1993). *Acta Pharmacol. Sin.* **14**, S47–S49.

supplementary materials

Acta Cryst. (2013). E69, o1096 [doi:10.1107/S1600536813014451]

5,6-Dihydroxy-7,8-dimethoxyflavone

Lin-Lin Jing, Xiao-Fei Fan, Peng-Cheng Fan, Lei He and Zheng-Ping Jia

Comment

Saussurea involucrata (Kar. et Kir) Sch.-Bip is one of the precious Tibetan herbs that have been used for a long period of time. Modern pharmacological studies have reported that the herb exhibit a wide range of bioactivities, including antioxidation, anti-inflammatory, anti-fatigue, anti-hypoxia, anti-cancer and analgesic effects. (Zheng *et al.*, 1993; Gao *et al.*, 2005; Tao *et al.*, 2010; Ma *et al.*, 2011; Jia *et al.*, 2005; Liu *et al.*, 1985; Paula *et al.*, 2002).

Our chemical investigation of this herbs for components with anti-hypoxia activity resulted in the isolation of the title compound and crystal growth one, suitable for X-ray diffraction.

The molecular structure of title compound is almost planar; the dihedral angle between the benzopyran-4-one group and the attached phenyl group is 1.89 (6) $^{\circ}$ (Fig. 1).

In the crystal, the 5-hydroxy group forms a strong intramolecular hydrogen bond with the carbonyl group, resulting in a six-membered ring (Fig. 1). The centrosymmetrical dimers of title compound is linked by intermolecular O4—H4···O3ⁱ hydrogen bonds. Non-classical C—H···O hydrogen bonds (C2—H2···O2ⁱⁱ, C15—H15···O2ⁱⁱ and C14—H14···O4ⁱⁱⁱ) and π – π interactions between molecule pairs (C3—C4—C5···C5ⁱ—C4ⁱ—C3ⁱ = 3.37 (2)–3.39 (2) \AA are found in the crystal structure. All of these interactions build up a three-dimensional network. Symmetry codes: (i) -x, -y+1, -z; (ii) 1-x, -y, -z; (iii) 1+x, 1+y, z.

Experimental

The air-dried whole plants (10 kg) of *S. involucrata* were milled and extracted with 70% ethanol (150 L \times 3) for 2 h each time at 351 K. The resulting extract was concentrated to give ethanol extract (2.4 kg). The ethanol extract was suspended in water and extracted successively with equal volumes petroleum ether, *EtOAc* and *n-BuOH*. The petroleum ether extract (1.2 kg) was subjected to a silica gel column eluted with petroleum ether–*EtOAc* (10:0 to 1:2, *v/v*) to afford five fractions. Fraction 5 was purified repeatedly over a Sephadex LH-20 column with a mixture of CHCl₃–MeOH (1:1, *v/v*) to give title compound (500 mg). Single crystals suitable for X-ray diffraction analysis were obtained by slow evaporation from MeOH at room temperature.


¹H-NMR (600 MHz, *DMS-d6*), δ (p.p.m.): 5.53 (1*H*, s, —OH), 12.27 (1*H*, s, —OH), 7.94 (2*H*, m), 7.57 (3*H*, m), 6.70 (1*H*, s), 4.16 (3*H*, s), 4.12 (3*H*, s). ¹³C-NMR (100 MHz, *DMS-d6*), δ (p.p.m.): 183.0, 164.2, 146.8, 143.1, 142.1, 133.3, 133.1, 132.0, 131.4, 129.1, 126.3, 106.8, 105.1, 62.2, 61.5.

Refinement

In the structure the H atoms were positioned geometrically and refined with using a riding model: C—H = 0.96 \AA for methyl H; C—H = 0.97 \AA for methylene H and C—H = 0.93 \AA for aryl H with $U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}(\text{C})$ for methyl H and $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$ for other. Hydroxy H atoms were positioned with O—H = 0.82 \AA and $U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}(\text{O})$.

Computing details

Data collection: *APEX2* (Bruker, 2007); cell refinement: *SAINT* (Bruker, 2007); data reduction: *SAINT* (Bruker, 2007); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3* (Farrugia, 2012); software used to prepare material for publication: *SHELXTL* (Sheldrick, 2008) and *PLATON* (Spek, 2009).

**Figure 1**

Molecular structure of the title compound with the atom numbering scheme. Displacement ellipsoids are drawn at the 30% probability level. H atoms are presented as a small spheres of arbitrary radius. Hydrogen bond is shown as dashed lines.

5,6-dihydroxy-7,8-dimethoxy-2-phenylchromen-4-one*Crystal data*

$C_{17}H_{14}O_6$
 $M_r = 314.28$
Triclinic, $P\bar{1}$
 $a = 7.953 (6)$ Å
 $b = 8.548 (6)$ Å
 $c = 10.951 (8)$ Å
 $\alpha = 96.602 (8)^\circ$
 $\beta = 92.282 (8)^\circ$
 $\gamma = 100.279 (7)^\circ$
 $V = 726.3 (9)$ Å³

$Z = 2$
 $F(000) = 328$
 $D_x = 1.437 \text{ Mg m}^{-3}$
Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å
Cell parameters from 1607 reflections
 $\theta = 2.4\text{--}28.0^\circ$
 $\mu = 0.11 \text{ mm}^{-1}$
 $T = 295 \text{ K}$
Block, orange
 $0.21 \times 0.16 \times 0.09$ mm

Data collection

Bruker APEXII CCD
diffractometer
Radiation source: fine-focus sealed tube
Graphite monochromator
 φ and ω scans

5252 measured reflections
3368 independent reflections
1786 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.036$
 $\theta_{\text{max}} = 28.3^\circ$, $\theta_{\text{min}} = 2.4^\circ$

$h = -9 \rightarrow 10$
 $k = -10 \rightarrow 11$

$l = -14 \rightarrow 14$

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.070$

$wR(F^2) = 0.236$

$S = 1.03$

3368 reflections

212 parameters

0 restraints

Primary atom site location: structure-invariant
direct methods

Secondary atom site location: difference Fourier
map

Hydrogen site location: inferred from
neighbouring sites

H-atom parameters constrained

$w = 1/[\sigma^2(F_o^2) + (0.1286P)^2]$

where $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{\text{max}} < 0.001$

$\Delta\rho_{\text{max}} = 0.32 \text{ e } \text{\AA}^{-3}$

$\Delta\rho_{\text{min}} = -0.32 \text{ e } \text{\AA}^{-3}$

Special details

Geometry. All s.u.'s (except the s.u. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell s.u.'s are taken into account individually in the estimation of s.u.'s in distances, angles and torsion angles; correlations between s.u.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell s.u.'s is used for estimating s.u.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

| | x | y | z | $U_{\text{iso}}^*/U_{\text{eq}}$ |
|-----|------------|------------|---------------|----------------------------------|
| O1 | 0.6956 (2) | 0.4357 (2) | 0.24860 (16) | 0.0386 (5) |
| O2 | 0.3542 (3) | 0.1892 (2) | -0.02333 (18) | 0.0489 (6) |
| O3 | 0.1717 (2) | 0.4078 (2) | 0.00746 (18) | 0.0450 (5) |
| H3 | 0.1947 | 0.3212 | -0.0196 | 0.067* |
| O4 | 0.1217 (3) | 0.6852 (2) | 0.14665 (19) | 0.0512 (6) |
| H4 | 0.0594 | 0.6291 | 0.0909 | 0.077* |
| O5 | 0.3574 (3) | 0.8349 (2) | 0.32644 (18) | 0.0528 (6) |
| O6 | 0.6457 (2) | 0.7095 (2) | 0.38048 (16) | 0.0450 (5) |
| C1 | 0.7247 (3) | 0.2981 (3) | 0.1857 (2) | 0.0354 (6) |
| C2 | 0.6135 (4) | 0.2132 (3) | 0.0965 (3) | 0.0405 (7) |
| H2 | 0.6368 | 0.1175 | 0.0576 | 0.049* |
| C3 | 0.4604 (3) | 0.2653 (3) | 0.0593 (2) | 0.0359 (6) |
| C4 | 0.4325 (3) | 0.4143 (3) | 0.1259 (2) | 0.0303 (6) |
| C5 | 0.2887 (3) | 0.4797 (3) | 0.1000 (2) | 0.0334 (6) |
| C6 | 0.2615 (3) | 0.6188 (3) | 0.1678 (2) | 0.0367 (6) |
| C7 | 0.3817 (4) | 0.6939 (3) | 0.2618 (2) | 0.0377 (7) |
| C8 | 0.5273 (3) | 0.6333 (3) | 0.2885 (2) | 0.0360 (6) |
| C9 | 0.5500 (3) | 0.4924 (3) | 0.2197 (2) | 0.0338 (6) |
| C10 | 0.8873 (3) | 0.2593 (3) | 0.2294 (2) | 0.0362 (6) |
| C11 | 0.9879 (4) | 0.3551 (4) | 0.3266 (3) | 0.0457 (7) |
| H11 | 0.9524 | 0.4461 | 0.3637 | 0.055* |
| C12 | 1.1388 (4) | 0.3168 (4) | 0.3683 (3) | 0.0555 (9) |
| H12 | 1.2037 | 0.3809 | 0.4340 | 0.067* |

| | | | | |
|------|------------|------------|------------|-------------|
| C13 | 1.1942 (4) | 0.1838 (4) | 0.3130 (3) | 0.0558 (9) |
| H13 | 1.2969 | 0.1584 | 0.3407 | 0.067* |
| C14 | 1.0974 (4) | 0.0895 (4) | 0.2172 (3) | 0.0569 (9) |
| H14 | 1.1342 | -0.0010 | 0.1805 | 0.068* |
| C15 | 0.9459 (4) | 0.1264 (4) | 0.1742 (3) | 0.0492 (8) |
| H15 | 0.8826 | 0.0620 | 0.1079 | 0.059* |
| C16 | 0.2964 (6) | 0.8185 (5) | 0.4437 (4) | 0.0927 (14) |
| H16A | 0.1833 | 0.7547 | 0.4353 | 0.139* |
| H16B | 0.2929 | 0.9224 | 0.4862 | 0.139* |
| H16C | 0.3712 | 0.7671 | 0.4897 | 0.139* |
| C17 | 0.7790 (5) | 0.8185 (4) | 0.3370 (3) | 0.0655 (10) |
| H17A | 0.8386 | 0.7615 | 0.2774 | 0.098* |
| H17B | 0.8575 | 0.8694 | 0.4048 | 0.098* |
| H17C | 0.7313 | 0.8982 | 0.2995 | 0.098* |

Atomic displacement parameters (\AA^2)

| | U^{11} | U^{22} | U^{33} | U^{12} | U^{13} | U^{23} |
|-----|-------------|-------------|-------------|--------------|--------------|--------------|
| O1 | 0.0383 (11) | 0.0392 (11) | 0.0415 (10) | 0.0216 (9) | -0.0029 (8) | -0.0015 (8) |
| O2 | 0.0437 (12) | 0.0463 (12) | 0.0544 (12) | 0.0171 (9) | -0.0080 (9) | -0.0127 (9) |
| O3 | 0.0383 (12) | 0.0452 (12) | 0.0511 (12) | 0.0182 (9) | -0.0083 (9) | -0.0080 (9) |
| O4 | 0.0429 (13) | 0.0525 (13) | 0.0612 (14) | 0.0296 (10) | -0.0074 (10) | -0.0096 (10) |
| O5 | 0.0668 (15) | 0.0453 (12) | 0.0497 (12) | 0.0305 (11) | 0.0005 (10) | -0.0101 (9) |
| O6 | 0.0451 (12) | 0.0519 (12) | 0.0370 (11) | 0.0143 (10) | -0.0040 (9) | -0.0044 (9) |
| C1 | 0.0384 (16) | 0.0329 (14) | 0.0393 (15) | 0.0161 (12) | 0.0063 (12) | 0.0063 (11) |
| C2 | 0.0404 (17) | 0.0367 (15) | 0.0476 (16) | 0.0201 (13) | 0.0026 (13) | -0.0020 (12) |
| C3 | 0.0344 (15) | 0.0381 (15) | 0.0365 (14) | 0.0121 (12) | 0.0019 (11) | 0.0009 (11) |
| C4 | 0.0321 (14) | 0.0321 (14) | 0.0290 (13) | 0.0118 (11) | 0.0047 (10) | 0.0031 (10) |
| C5 | 0.0310 (15) | 0.0388 (15) | 0.0315 (13) | 0.0112 (12) | 0.0005 (11) | 0.0027 (10) |
| C6 | 0.0340 (15) | 0.0395 (15) | 0.0406 (15) | 0.0186 (12) | 0.0018 (11) | 0.0035 (11) |
| C7 | 0.0423 (17) | 0.0371 (15) | 0.0376 (14) | 0.0198 (13) | 0.0069 (12) | -0.0001 (11) |
| C8 | 0.0402 (16) | 0.0409 (15) | 0.0277 (13) | 0.0140 (12) | 0.0011 (11) | -0.0017 (10) |
| C9 | 0.0320 (15) | 0.0403 (15) | 0.0329 (14) | 0.0166 (12) | 0.0016 (11) | 0.0052 (11) |
| C10 | 0.0338 (15) | 0.0364 (15) | 0.0433 (15) | 0.0141 (12) | 0.0069 (12) | 0.0121 (11) |
| C11 | 0.0458 (18) | 0.0488 (18) | 0.0464 (16) | 0.0202 (14) | 0.0006 (13) | 0.0051 (13) |
| C12 | 0.046 (2) | 0.068 (2) | 0.0549 (19) | 0.0154 (17) | -0.0082 (15) | 0.0124 (16) |
| C13 | 0.0407 (19) | 0.065 (2) | 0.071 (2) | 0.0237 (17) | 0.0030 (16) | 0.0252 (18) |
| C14 | 0.048 (2) | 0.0475 (19) | 0.082 (2) | 0.0252 (16) | 0.0080 (17) | 0.0113 (17) |
| C15 | 0.0446 (18) | 0.0439 (17) | 0.0628 (19) | 0.0200 (14) | 0.0016 (14) | 0.0043 (14) |
| C16 | 0.134 (4) | 0.090 (3) | 0.066 (2) | 0.055 (3) | 0.041 (2) | -0.006 (2) |
| C17 | 0.064 (2) | 0.063 (2) | 0.060 (2) | -0.0004 (18) | 0.0005 (17) | -0.0104 (16) |

Geometric parameters (\AA , ^\circ)

| | | | |
|-------|-----------|---------|-----------|
| O1—C1 | 1.355 (3) | C7—C8 | 1.384 (4) |
| O1—C9 | 1.372 (3) | C8—C9 | 1.389 (4) |
| O2—C3 | 1.250 (3) | C10—C11 | 1.393 (4) |
| O3—H3 | 0.8200 | C10—C15 | 1.388 (4) |
| O3—C5 | 1.360 (3) | C11—H11 | 0.9300 |
| O4—H4 | 0.8200 | C11—C12 | 1.374 (4) |

| | | | |
|---------------|-----------|---------------|------------|
| O4—C6 | 1.359 (3) | C12—H12 | 0.9300 |
| O5—C7 | 1.375 (3) | C12—C13 | 1.377 (5) |
| O5—C16 | 1.404 (4) | C13—H13 | 0.9300 |
| O6—C8 | 1.372 (3) | C13—C14 | 1.365 (4) |
| O6—C17 | 1.417 (4) | C14—H14 | 0.9300 |
| C1—C2 | 1.343 (4) | C14—C15 | 1.378 (4) |
| C1—C10 | 1.467 (4) | C15—H15 | 0.9300 |
| C2—H2 | 0.9300 | C16—H16A | 0.9600 |
| C2—C3 | 1.429 (4) | C16—H16B | 0.9600 |
| C3—C4 | 1.450 (4) | C16—H16C | 0.9600 |
| C4—C5 | 1.393 (4) | C17—H17A | 0.9600 |
| C4—C9 | 1.385 (4) | C17—H17B | 0.9600 |
| C5—C6 | 1.384 (4) | C17—H17C | 0.9600 |
| C6—C7 | 1.391 (4) | | |
| | | | |
| C1—O1—C9 | 119.6 (2) | C4—C9—C8 | 121.7 (2) |
| C5—O3—H3 | 109.5 | C11—C10—C1 | 121.1 (2) |
| C6—O4—H4 | 109.5 | C15—C10—C11 | 120.8 (3) |
| C7—O5—C16 | 114.1 (3) | C15—C10—C11 | 118.1 (3) |
| C8—O6—C17 | 112.7 (2) | C10—C11—H11 | 119.6 |
| O1—C1—C10 | 111.1 (2) | C12—C11—C10 | 120.8 (3) |
| C2—C1—O1 | 121.9 (2) | C12—C11—H11 | 119.6 |
| C2—C1—C10 | 127.0 (2) | C11—C12—H12 | 119.9 |
| C1—C2—H2 | 119.0 | C11—C12—C13 | 120.2 (3) |
| C1—C2—C3 | 122.1 (2) | C13—C12—H12 | 119.9 |
| C3—C2—H2 | 119.0 | C12—C13—H13 | 120.2 |
| O2—C3—C2 | 123.7 (2) | C14—C13—C12 | 119.5 (3) |
| O2—C3—C4 | 120.9 (2) | C14—C13—H13 | 120.2 |
| C2—C3—C4 | 115.4 (2) | C13—C14—H14 | 119.5 |
| C5—C4—C3 | 121.9 (2) | C13—C14—C15 | 121.0 (3) |
| C9—C4—C3 | 119.3 (2) | C15—C14—H14 | 119.5 |
| C9—C4—C5 | 118.8 (2) | C10—C15—H15 | 119.8 |
| O3—C5—C4 | 120.6 (2) | C14—C15—C10 | 120.3 (3) |
| O3—C5—C6 | 118.5 (2) | C14—C15—H15 | 119.8 |
| C6—C5—C4 | 120.9 (2) | O5—C16—H16A | 109.5 |
| O4—C6—C5 | 122.7 (2) | O5—C16—H16B | 109.5 |
| O4—C6—C7 | 118.5 (2) | O5—C16—H16C | 109.5 |
| C5—C6—C7 | 118.8 (2) | H16A—C16—H16B | 109.5 |
| O5—C7—C6 | 118.7 (2) | H16A—C16—H16C | 109.5 |
| O5—C7—C8 | 119.5 (2) | H16B—C16—H16C | 109.5 |
| C8—C7—C6 | 121.7 (2) | O6—C17—H17A | 109.5 |
| O6—C8—C7 | 121.1 (2) | O6—C17—H17B | 109.5 |
| O6—C8—C9 | 120.8 (2) | O6—C17—H17C | 109.5 |
| C7—C8—C9 | 118.1 (2) | H17A—C17—H17B | 109.5 |
| O1—C9—C4 | 121.7 (2) | H17A—C17—H17C | 109.5 |
| O1—C9—C8 | 116.6 (2) | H17B—C17—H17C | 109.5 |
| | | | |
| O1—C1—C2—C3 | 2.1 (4) | C3—C4—C9—C8 | -178.3 (2) |
| O1—C1—C10—C11 | 2.1 (4) | C4—C5—C6—O4 | -179.0 (2) |

| | | | |
|----------------|------------|-----------------|------------|
| O1—C1—C10—C15 | −177.4 (3) | C4—C5—C6—C7 | 0.8 (4) |
| O2—C3—C4—C5 | −0.7 (4) | C5—C4—C9—O1 | −179.0 (2) |
| O2—C3—C4—C9 | 177.8 (2) | C5—C4—C9—C8 | 0.3 (4) |
| O3—C5—C6—O4 | 0.9 (4) | C5—C6—C7—O5 | 177.9 (2) |
| O3—C5—C6—C7 | −179.3 (2) | C5—C6—C7—C8 | 0.3 (4) |
| O4—C6—C7—O5 | −2.3 (4) | C6—C7—C8—O6 | 179.2 (2) |
| O4—C6—C7—C8 | −179.9 (3) | C6—C7—C8—C9 | −1.0 (4) |
| O5—C7—C8—O6 | 1.6 (4) | C7—C8—C9—O1 | −180.0 (2) |
| O5—C7—C8—C9 | −178.6 (2) | C7—C8—C9—C4 | 0.7 (4) |
| O6—C8—C9—O1 | −0.2 (4) | C9—O1—C1—C2 | −1.1 (4) |
| O6—C8—C9—C4 | −179.5 (2) | C9—O1—C1—C10 | 179.3 (2) |
| C1—O1—C9—C4 | −1.3 (4) | C9—C4—C5—O3 | 179.1 (2) |
| C1—O1—C9—C8 | 179.4 (2) | C9—C4—C5—C6 | −1.1 (4) |
| C1—C2—C3—O2 | 180.0 (3) | C10—C1—C2—C3 | −178.3 (2) |
| C1—C2—C3—C4 | −0.9 (4) | C10—C11—C12—C13 | 1.0 (5) |
| C1—C10—C11—C12 | 179.0 (3) | C11—C10—C15—C14 | 1.5 (5) |
| C1—C10—C15—C14 | −178.9 (3) | C11—C12—C13—C14 | −0.6 (5) |
| C2—C1—C10—C11 | −177.5 (3) | C12—C13—C14—C15 | 0.7 (5) |
| C2—C1—C10—C15 | 3.0 (4) | C13—C14—C15—C10 | −1.2 (5) |
| C2—C3—C4—C5 | −179.9 (2) | C15—C10—C11—C12 | −1.5 (5) |
| C2—C3—C4—C9 | −1.3 (4) | C16—O5—C7—C6 | 103.2 (4) |
| C3—C4—C5—O3 | −2.4 (4) | C16—O5—C7—C8 | −79.2 (4) |
| C3—C4—C5—C6 | 177.5 (2) | C17—O6—C8—C7 | −92.6 (3) |
| C3—C4—C9—O1 | 2.4 (4) | C17—O6—C8—C9 | 87.6 (3) |

Hydrogen-bond geometry (Å, °)

| D—H···A | D—H | H···A | D···A | D—H···A |
|----------------------------|------|-------|-----------|---------|
| O4—H4···O3 ⁱ | 0.82 | 2.05 | 2.764 (4) | 146 |
| C12—H12···O6 ⁱⁱ | 0.93 | 2.58 | 3.234 (4) | 128 |
| O3—H3···O2 | 0.82 | 1.84 | 2.564 (3) | 146 |
| O4—H4···O3 | 0.82 | 2.34 | 2.767 (3) | 113 |
| C11—H11···O1 | 0.93 | 2.34 | 2.675 (4) | 101 |

Symmetry codes: (i) $-x, -y+1, -z$; (ii) $-x+2, -y+1, -z+1$.