

Research Article

Network Pharmacology-Based and Molecular Docking-Based Analysis of Suanzaoren Decoction for the Treatment of Parkinson's Disease with Sleep Disorder

Yan-yun Liu ¹, Li-hua Yu,² Juan Zhang ³, Dao-jun Xie ³, Xin-xiang Zhang,¹ and Jia-ming Yu¹

¹Graduate School, Anhui University of Traditional Chinese Medicine, Hefei, Anhui 230038, China

²College of Integrated Chinese and Western Medicine, Anhui University of Traditional Chinese Medicine, Hefei, Anhui 230038, China

³Center of Neurology, Anhui Provincial Hospital of Traditional Chinese Medicine, Hefei, Anhui 230031, China

Correspondence should be addressed to Juan Zhang; 1477210980@qq.com

Received 17 April 2021; Revised 18 August 2021; Accepted 3 September 2021; Published 8 October 2021

Academic Editor: Bilal Alatas

Copyright © 2021 Yan-yun Liu et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

This study is aimed at exploring the possible mechanism of action of the Suanzaoren decoction (SZRD) in the treatment of Parkinson's disease with sleep disorder (PDSD) based on network pharmacology and molecular docking. Traditional Chinese Medicine Systems Pharmacology (TCMSP) was used to screen the bioactive components and targets of SZRD, and their targets were standardized using the UniProt platform. The disease targets of "Parkinson's disease (PD)" and "Sleep disorder (SD)" were collected by OMIM, GeneCards, and DisGeNET databases. Thereafter, the protein-protein interaction (PPI) network was constructed using the STRING platform and visualized by Cytoscape (3.7.2) software. Then, the DAVID platform was used to analyze the Gene Ontology (GO) enrichment analysis and Kyoto Encyclopedia of Genes and Genomes (KEGG) pathway. Cytoscape (3.7.2) software was also used to construct the network of the "herb-component-target-pathway." The core active ingredients and core action targets of the drug were verified by molecular docking using AutoDock software. A total of 135 Chinese herbal components and 41 corresponding targets were predicted for the treatment of PDSD using SZRD. Fifteen important signaling pathways were screened, such as the cancer pathway, TNF signaling pathway, PI3K-AKT signaling pathway, HIF-1 signaling pathway, and Toll-like receptor signaling pathway. The results of molecular docking showed that the main active compounds could bind to the representative targets and exhibit good affinity. This study revealed that SZRD has the characteristics and advantages of "multicomponent, multitarget, and multipathway" in the treatment of PDSD; among these, the combination of the main active components of quercetin and kaempferol with the key targets of *AKT1*, *IL6*, *MAPK1*, *TP53*, and *VEGFA* may be one of the important mechanisms. This study provides a theoretical basis for further study of the material basis and molecular mechanism of SZRD in the treatment of PDSD.

1. Introduction

Parkinson's disease (PD) is a common neurodegenerative disease in middle-aged and elderly individuals. Clinically, symptoms include classic motor symptoms—tremor, myotonia, bradykinesia, and postural imbalance—as well as nonmotor symptoms—sleep disorders, smell disorders, autonomic nervous dysfunction, and cognitive and mental

disorders [1]. Sleep disorders (SD), the most common non-motor symptoms of PD, seen in 60–90% of these patients, are one of the common nocturnal symptoms [2]. Treatment modalities of PD with sleep disorder (PDSD), used in clinical practice, include levodopa, dopamine receptor agonists, benzodiazepines, and melatonin [3–5]. However, long-term use of such drugs may enhance restless leg syndrome, periodic limb movements, and rapid eye movement (REM) sleep

TABLE 1: SZRD shares chemical composition information table.

Number	Mol ID	Molecule name	OB%	DL	Drug
A	MOL000359	Sitosterol	36.91	0.75	<i>Glycyrrhiza glabra</i> , <i>Rhizoma chuanxiong</i>
B	MOL000211	Mairin	55.38	0.78	<i>Glycyrrhiza glabra</i> , <i>Semen ziziphi spinosae</i>
C	MOL000422	Kaempferol	41.88	0.24	<i>Glycyrrhiza glabra</i> , <i>Rhizoma anemarrhenae</i>

TABLE 2: Chemical information sheet of major active ingredients.

Number	Mol ID	Molecule name	OB%	DL	Degree
GC84	MOL000098	Quercetin	46.43	0.28	127
C	MOL000422	Kaempferol	41.88	0.24	104
GC74	MOL000500	Vestitol	74.66	0.21	33
GC8	MOL003896	7-Methoxy-2-methyl isoflavone	42.56	0.20	30
GC11	MOL004328	Naringenin	58.29	0.21	29
ZM2	MOL004373	Anhydroaricarin	45.41	0.44	27
GC9	MOL000392	Formononetin	69.67	0.21	26
ZM6	MOL000449	Stigmasterol	43.48	0.76	24
GC5	MOL002565	Medicarpin	49.22	0.34	24
GC63	MOL000497	Licochalcone a	40.79	0.29	24
GC41	MOL004891	Shinpterocarpin	80.3	0.73	23
GC65	MOL004978	2-[(3R)-8,8-Dimethyl-3,4-dihydro-2H-pyrano[6,5-f] chromen-3-yl]-5-methoxyphenol	36.21	0.52	23
SZR1	MOL001522	(S)-Coclaurine	42.35	0.24	22
GC6	MOL004980	Isorhamnetin	39.71	0.33	22
GC76	MOL005003	Licoagrocarpin	58.81	0.58	21
GC25	MOL004835	Glypallichalcone	61.60	0.19	20
GC64	MOL004974	3'-Methoxyglabridin	46.16	0.57	20

GC: *Glycyrrhiza glabra*; C: common components of *Glycyrrhiza glabra* and *Rhizoma anemarrhenae*; ZM: *Rhizoma anemarrhenae*; SZR: *Semen ziziphi spinosae*.

behavior disorder symptoms [6]. Furthermore, melatonin had little effect on objective sleep parameters [7]. Suanzaoren decoction (SZRD), derived from JinKuiYaoLue, is composed of five traditional Chinese medicines: *Semen ziziphi spinosae*, *Glycyrrhiza glabra*, *Rhizoma anemarrhenae*, *Poria cocos*, and *Rhizoma chuanxiong* [8]. It has the effect of nourishing the blood, calming the mind, clearing away heat, and eliminating annoyance, mainly treating liver and blood deficiency and insomnia caused by heat deficiency [9–12]. PD belongs to the category of “fibrillation disease” in traditional Chinese medicine, and it is more common in the elderly [13]. Clinical evidence shows that SZRD has a remarkable curative effect on insomnia [14–16]. However, the mechanism of action of fibrillation disease merging with wakefulness is not clear. Therefore, with the systematic research methods of network pharmacology and molecular docking, the overall analysis of the “herb-component-target-pathway” was conducted in this study to explore the possible mechanism of SZRD in the treatment of PDS and to provide new theoretical support for the clinical treatment of PDS.

2. Materials and Methods

2.1. Screening and Target Prediction of Active Components of SZRD. Application analysis platform and database system pharmacology of Chinese medicine (TCMSP, [https://](https://tcmssp.com/tcmssp.php)

tcmssp.com/tcmssp.php) [17] were used to retrieve active ingredients of SZRD and predict the targets of active ingredients. We use pharmacodynamics to select active ingredients satisfying both oral bioavailability (OB) $\geq 30\%$ and drug – likeness (DL) ≥ 0.18 [18, 19]. TCMSP was used for the prediction of targets of active ingredients. At the same time, the UniProt database (<https://www.uniprot.org/>) [20], which was set for human species, was used to standardize the drug target of each active ingredient.

2.2. Screening of Disease-Related Targets. The targets related to the PD and SD were obtained through retrieving the OMIM (<https://omim.org/search/advanced/>) [21], GeneCards (<https://www.genecards.org/>) [22], and DisGeNET database search (<https://www.disgenet.org/>) [23] using the keyword “Parkinson’s diseases” or “Sleep disorder.”

2.3. Screening of Common Targets of Diseases and Drugs. The online Venny 2.1 mapping platform (<http://bioinfo.g.cnbcsc.es/tools/venny/index.html>) was used to map “Parkinson’s diseases,” “Sleep disorder,” and “SZRD” targets and to get the intersection targets.

2.4. Common Target PPI Network Construction. The intersection targets were imported into the STRING database (<https://string-db.org/cgi/input>) [24]. A confidence, ≥ 0.4 ,




FIGURE 1: Composition-target network of SZRD. Circles are for traditional Chinese medicine; octagons are compound component; diamonds are target. GC: *Glycyrrhiza glabra*; SZR: *Semen ziziphi spinosae*; FL: *Poria cocos*; ZM: *Rhizoma anemarrhenae*; CX: *Rhizoma chuanxiong*; A: common components of *Glycyrrhiza glabra* and *Rhizoma chuanxiong*; B: common components of *Glycyrrhiza glabra* and *Semen ziziphi spinosae*; C: common components of *Glycyrrhiza glabra* and *Rhizoma anemarrhenae*.




FIGURE 2: Disease-drug target Venn diagram.

was taken, and the free nodes were hidden to construct the protein-protein interaction (PPI) network. This PPI network was further processed by Cytoscape 3.7.2 software [25] to realize visualization and screen out the core targets.

2.5. *GO and KEGG Enrichment Analysis.* For the screened core targets, the DAVID data platform (<https://david.ncifcrf.gov/tools.jsp>) [26] was used for Gene Ontology (GO) functional annotations and Kyoto Encyclopedia of Genes and Genomes (KEGG) pathway enrichment analysis. Select “Homo species” on the DAVID platform; further analyze the SZRD for PDSD-related biological processes (BP), cellular component (CC), molecular function (MF), and signal pathway; and use the bioinformatic online platform (<https://www.bioinformatics.com.cn/>) [27] to visualize the result analysis.

2.6. *Constructing the “H-C-T-P” Network.* Cytoscape 3.7.2 software was used to construct the network of “herb-component-target-pathway” (H-C-T-P). This “H-C-T-P” network together with the screened main active ingredients, core targets, and concentrated main signal pathways was used to systematically analyze the possible mechanism of SZRD in the treatment of PDSD.

TABLE 3: Core target information table.

Target	Degree	Betweenness centrality (BC)	Closeness centrality (CC)
AKT1	123	0.064225	0.731518
IL6	110	0.033377	0.693727
MAPK3	109	0.052416	0.693727
TP53	105	0.027917	0.676259
VEGFA	100	0.016309	0.664311
TNF	98	0.018693	0.661972
CASP3	97	0.019316	0.661972
JUN	97	0.021842	0.664311
EGF	95	0.020902	0.657343
MAPK8	95	0.017033	0.657343
PTGS2	91	0.032141	0.646048
MAPK1	90	0.012068	0.639456
EGFR	89	0.016470	0.641638
MYC	89	0.013782	0.639456
STAT3	88	0.011530	0.635135
FOS	87	0.033525	0.639456
CXCL8	85	0.016357	0.626667
MMP9	85	0.027723	0.632997
IL1B	82	0.013500	0.626667
CAT	81	0.038221	0.624585
ESR1	79	0.010994	0.616393
CCND1	77	0.006779	0.610390
CCL2	76	0.005212	0.608414
NOS3	74	0.020902	0.614379
MTOR	71	0.006984	0.598726
MAPK14	71	0.005773	0.600639
IL10	70	0.003254	0.593060
MMP2	69	0.003821	0.596825
PPARG	68	0.008861	0.594937
ICAM1	67	0.002937	0.591195
RELA	67	0.014775	0.585670
ERBB2	66	0.009451	0.587500
AR	64	0.017042	0.589342
HMOX1	63	0.006109	0.580247
STAT1	63	0.006329	0.574924
SERPINE1	62	0.004200	0.580247
IL4	61	0.002350	0.576687
IL2	60	0.002819	0.569697
VCAM1	60	0.002099	0.574924
BCL2L1	59	0.002173	0.567976
CASP8	58	0.002424	0.564565

2.7. Docking and Verification of Potential Active Ingredients with Core Target Molecules. The 2D structures of the potential active ingredients in SZRD were downloaded from the TCMSP database while the 3D structure of the PDS D docking targets (top 5 of degree in PPI network) treated by SZRD was downloaded from the Worldwide Protein Data Bank (PDB) database (<https://www.rcsb.org/>) [28]. They were imported

to AutoDockTools [29] for hydrogenation, dehydration, and other pretreatments. Then, molecular docking of the receptor and ligand was conducted to analyze its binding activity. The docking results were visualized using the PyMol software [30].

3. Results

3.1. The Active Components and Effective Targets of SZRD. By searching the TCMSP database, 135 different active ingredients of SZRD were screened, including 9 in *Semen ziziphi spinosae*, 15 in *Poria cocos*, 15 in *Rhizoma anemarrhenae*, 7 in *Rhizoma chuanxiong*, and 92 in *Glycyrrhiza glabra*. The results showed that these different drugs contained common active ingredients, namely A, B, and C as shown in Table 1.

Cytoscape 3.7.2 software was used to screen the active ingredients with a degree ≥ 20 in SZRD. As shown in Table 2, 17 major components, which include quercetin, kaempferol, vestitol, 7-methoxy-2-methyl isoflavone, naringenin, anhydroicaritin, formononetin, stigmaterol, licochalcone A, and isorhamnetin, were obtained using this software.

A total of 204 different drug targets were screened using the TCMSP database and were standardized by the UniProt database, which included the 26 in *Semen ziziphi spinosae*, 16 in *Poria cocos*, 90 in *Rhizoma anemarrhenae*, 23 in *Rhizoma chuanxiong*, and 193 in *Glycyrrhiza glabra*. The data of potential active ingredients and potential targets of SZRD in the treatment of PDS D were imported into Cytoscape 3.7.2 software to obtain a diagram of the traditional Chinese medicine component-target network (Figure 1).

3.2. Related Targets for Disease. After combining the three databases and deleting repeated targets, a total of 9777 PD-related targets and 10748 SD-related targets were obtained from the OMIM, DisGeNET, and GeneCards databases.

3.3. Common Targets for Diseases and Drugs. A total of 9777 PD-related targets, 10748 SD-related targets, and 204 SZRD drug prediction targets were imported using the Venny online mapping platform. After mapping, 189 intersection targets of SZRD and PDS D were obtained (Figure 2).

3.4. PPI Network Construction. A total of 189 targets were imported into the STRING platform to construct a PPI network. Then, 189 nodes and 3426 edges were also obtained using this platform. The double median of "Degree," that is, "Degree ≥ 58 ," was used to screen the intersection targets. Thus, 41 nodes, 765 edges, and a total of 41 core targets for SZRD treatment of PDS D were obtained (Table 3). Import the PPI network information obtained from the STRING platform into Cytoscape (3.7.2) software for visualization (Figure 3).

3.5. GO and KEGG Enrichment Analysis. The GO function enrichment analysis of the 41 core targets was performed on the DAVID platform, and a total of 1423 GO items were obtained, including 1325 BP, 36 CC, and 62 MF. The first 15, 8, and 14 items were selected based on the *P* value for visual analysis (Figure 4). Results showed that the treatment of PDS D by SZRD mainly involves BP such as cell migration,




FIGURE 3: Core target PPI network. As the figure shows, the larger area of the circle could be considered as more important in this network.




FIGURE 4: GO function enrichment results of SZRD in the treatment of PDSD.

angiogenesis, leukocyte differentiation, cell proliferation, stress, cell aging, cell adhesion, and cell regeneration. These targets pass through cytokine receptor binding, transcription factor binding specificity, phosphatase binding, DNA binding domain specificity, protein structure combining, G protein coupled receptor molecules, and other functions, and

they play a role in the cell membrane, RNA polymerase II transcription factor complex, nucleus and organelle outer membrane and plasma membrane protein complexes, and extracellular matrix components.

One hundred and nine signal pathways were enriched by KEGG pathway analysis of the core targets using the DAVID

TABLE 4: KEGG pathway enrichment results.

Term	%	Count	P value	Related genes
hsa05200: pathways in cancer	65.8536	27	2.30E-23	CXCL8, PTGS2, RELA, EGFR, MAPK8, CASP8, CCND1, MYC, CASP3, ERBB2, AKT1, MAPK1, MAPK3, JUN, EGF, STAT1, MMP2, STAT3, FOS, MMP9, MTOR, VEGFA, AR, IL6, PPARG, TP53, BCL2L1
hsa04668: TNF signaling pathway	43.9024	18	3.19E-21	JUN, VCAM1, FOS, PTGS2, MAPK14, TNF, MMP9, RELA, ICAM1, IL6, MAPK8, CASP8, IL1B, CASP3, CCL2, AKT1, MAPK1, MAPK3
hsa05161: hepatitis B	43.9024	18	7.08E-19	JUN, CXCL8, STAT1, STAT3, FOS, TNF, MMP9, RELA, IL6, MAPK8, CASP8, CCND1, MYC, CASP3, AKT1, MAPK1, TP53, MAPK3
hsa05142: Chagas disease	41.4634	17	1.04E-19	IL10, JUN, CXCL8, SERPINE1, FOS, MAPK14, TNF, IL2, RELA, IL6, MAPK8, CASP8, IL1B, CCL2, AKT1, MAPK1, MAPK3
hsa05205: proteoglycans in cancer	41.4634	17	4.91E-15	MMP2, STAT3, MAPK14, ESR1, TNF, MMP9, EGFR, MTOR, VEGFA, CCND1, MYC, CASP3, ERBB2, AKT1, MAPK1, TP53, MAPK3
hsa04151: PI3K-Akt signaling pathway	39.0243	16	3.03E-10	NOS3, EGF, EGFR, IL2, MTOR, RELA, VEGFA, IL4, IL6, CCND1, MYC, AKT1, MAPK1, TP53, BCL2L1, MAPK3
hsa04010: MAPK signaling pathway	36.5853	15	5.72E-11	JUN, EGF, FOS, MAPK14, TNF, EGFR, RELA, MAPK8, MYC, IL1B, CASP3, AKT1, MAPK1, TP53, MAPK3
hsa05212: pancreatic cancer	34.1463	14	1.31E-17	STAT1, EGF, STAT3, EGFR, RELA, VEGFA, MAPK8, CCND1, ERBB2, AKT1, MAPK1, TP53, BCL2L1, MAPK3
hsa04066: HIF-1 signaling pathway	34.1463	14	2.96E-15	NOS3, EGF, STAT3, SERPINE1, EGFR, MTOR, RELA, VEGFA, IL6, ERBB2, AKT1, MAPK1, HMOX1, MAPK3
hsa04620: Toll-like receptor signaling pathway	34.1463	14	1.11E-14	JUN, CXCL8, STAT1, FOS, MAPK14, TNF, RELA, IL6, MAPK8, CASP8, IL1B, AKT1, MAPK1, MAPK3
hsa05164: influenza A	34.1463	14	7.36E-12	JUN, CXCL8, STAT1, MAPK14, TNF, RELA, ICAM1, IL6, MAPK8, IL1B, CCL2, AKT1, MAPK1, MAPK3
hsa05145: toxoplasmosis	31.7073	13	5.84E-13	IL10, STAT1, STAT3, MAPK14, TNF, RELA, MAPK8, CASP8, CASP3, AKT1, MAPK1, BCL2L1, MAPK3
hsa05160: hepatitis C	31.7073	13	5.83E-12	CXCL8, STAT1, EGF, STAT3, MAPK14, TNF, EGFR, RELA, MAPK8, AKT1, MAPK1, TP53, MAPK3
hsa05152: tuberculosis	31.7073	13	1.73E-10	IL10, STAT1, MAPK14, TNF, RELA, IL6, MAPK8, CASP8, IL1B, CASP3, AKT1, MAPK1, MAPK3
hsa04068: FoxO signaling pathway	29.2682	12	1.45E-10	IL10, IL6, MAPK8, CCND1, EGF, STAT3, CAT, MAPK1, AKT1, MAPK14, EGFR, MAPK3

platform. According to the P value < 0.05 and the number of genes ≥ 12 , 15 signal pathways with high probability were screened out for visual analysis as shown in Table 4 and Figure 5. Moreover, Figure 5 shows that SZRD treatment of PDS may be mainly related to TNF, PI3K-Akt, MAPK, HIF-1, Toll-like receptor, FoxO, and other signaling pathways.

3.6. Construction of “H-C-T-P” Network. The “H-C-T-P” network was constructed using the 17 major components,

41 core targets, and 15 signal pathways of SZRD in the treatment of PDS (Figure 6).

3.7. Molecular Docking Results and Analysis. According to Table 3, the top 5 targets of degree are AKT1, IL6, MAPK1, TP53, and VEGFA. The docking targets with the two active components with the highest degree of quercetin (degree = 127) and kaempferol (degree = 104) in SZRD were docking. As shown in Table 5, the binding energy of




FIGURE 5: KEGG enrichment bubble diagram.

quercetin, kaempferol with AKT1, IL6, MAPK1, TP53, and VEGFA was all less than $-5.0 \text{ kcal}\cdot\text{mol}^{-1}$, showing good binding force. The binding of AKT1 to quercetin and kaempferol and TP53 to kaempferol is shown in Figure 7.

4. Discussion

4.1. Understanding of PDS in Traditional Chinese Medicine and Western Medicine. In Western medicine, PDS is believed to be associated with a variety of factors which may also be related to the increase or decrease of serum vitamin D, melatonin, serum cystatin (Cys) C, homocysteine (Hcy), and dopamine levels in the striatum caused by PD itself [31–34].

There is no related record of PDS in traditional Chinese medicine classes, but according to its related clinical symptoms, it can be classified as a combination of “fibrillation” and “insomnia.” SZRD is a classic prescription, with tranquilizing properties, mainly treating “deficiency of liver and blood, deficiency of heat, and internal disturbance.” This prescription uses a lot of *Semen ziziphi spinosae* that nourishes the blood and liver and gives peace of mind. *Rhizoma anemarrhenae* nourishes Yin and clears the heat, and when the evil heat is gone, the healthy Qi is

restored. “*Poria cocos* calms the heart and tranquilizes the mind.” Moreover, the theory of the properties of drugs says it is “good at calming the mind.” *Rhizoma anemarrhenae* and *Poria cocos* are used to help *Semen ziziphi spinosae* calm the mind from vexation. The nature of *Rhizoma chuanxiong* is scattered, “Qi medicine in the blood,” and is introduced into the liver meridian. Furthermore, this regulates Qi activity to help the liver recover its ability to dredge and release Qi. Rihua Zi medicine said that it “cures all wind, all Qi, all strain, all blood, and replenish five kinds of weakness.” *Rhizoma chuanxiong* is used in combination with King medicine, which has a magical effect to help sanguification and liver recuperation. Then, *Glycyrrhiza glabra* is used to harmonize the various drugs. All the drugs work together, to calm the mind in addition to the effect of vexation.

4.2. The Mechanism Prediction of SZRD for PDS. Based on the network pharmacological research method, a total of 17 major active components of SZRD, which include quercetin, kaempferol, visistin, 7-methoxy-2-methyl isoflavone, naringenin, anhydroicaritin, formononetin, stigmasterol, licochalcone A, and isorhamnetin, in the treatment of PDS were obtained.




FIGURE 6: H-C-T-P network diagram. Triangle is SZRD; octagons are compound component; diamonds are target; V is pathway.

TABLE 5: Docking results of target protein and active compound.

Core target	PDB ID	Binding energy/(kcal Mol ⁻¹)	
		Quercetin	Kaempferol
AKT1	1UNQ	-6.18	-6.51
IL-6	1ALU	-5.72	-5.90
MAPK3	4QTB	-5.45	-5.33
TP53	1YC5	-5.92	-6.08
VEGFA	4QAF	-5.61	-5.67

Quercetin, kaempferol, and isorhamnetin are flavonoids which can scavenge reactive oxygen free radicals and achieve antioxidant and neuroprotective effects [35–38]. In the H₂O₂-induced PD model PC-12 cells, quercetin treatment increased cell viability, reduced mutagenesis of antioxidant enzymes, and reduced apoptosis of cells and hippocampal neurons [39]. In addition, quercetin can significantly reduce inflammation and oxidative damage in the striatum of PD model rats induced by 6-hydroxy-dopamine (6-OHDA) or 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine (MPTP) [40]. Filomeni et al. found that kaempferol can significantly protect neuroblastoma cells (SH-SY5Y nerve cells) and major neurons from rotenone damage [41]. Moreover, it can reduce protease lysis and nuclear apoptosis and significantly reduce oxidative stress levels and mitochondrial hydroxyl compound content [41]. We also found that kaempferol may play a neuroprotective role by inhibiting cysteine proteinase-3 and by reducing brain cell apoptosis [42]. At the same time, quercetin and kaempferol have sed-

ative and hypnotic effects, and both can also improve the effects of sleep disorders [43–46].

According to the GO and KEGG pathway enrichment analysis results, we speculated that the mechanism of SZRD in the treatment of PDSD may be mainly related to TNF, PI3K-Akt, MAPK, HIF-1, Toll-like receptors, and FoxO-associated signaling pathways. Such as involving *AKT1*, *IL6*, *MAPK3*, *TP53*, and *VEGFA* multiple key targets, cell aging, adhesion, regeneration, and angiogenesis, leukocyte differentiation is closely related to the process of cell metabolism. Recent studies have shown that PI3K-Akt, MAPK, and FOXO are all involved in the process of human cell apoptosis [47–49]. *AKT1*, an important member of the AKT (protein kinase B, PKB) family, is an intracellular serine/threonine involved in a variety of cellular BP, and its activation is mainly dependent on the PI3K signaling pathway [50]. *AKT1* is activated by phosphorylation, which in turn activates downstream signaling molecules [51]. The PI3K/Akt signal transduction pathway is involved in a variety of cytokines, and it has been found that by reducing the phosphorylation levels of PI3K, AKT, and mTOR, the transduction of this pathway can play a role in the treatment of MPTP-induced PD mice [52]. Studies have shown that cytokines are involved in regulating sleep, among which interleukin-1 (IL-1), interleukin-6 (IL-6), and tumor necrosis factor (TNF- α and TNF- β) play a role in promoting sleep [53–55]. Elevation of interleukin-1 β (IL-1 β) and TNF- α can directly act on central sleep-wake neurons, causing increased nonrapid eye movement (NREM) sleep and regulating the pathologic circadian rhythm [56, 57]. IL-1 β and TNF- α , as sleep-regulating cytokines, can induce the production of




FIGURE 7: Molecular docking diagram of chemical composition to target: (a) 1UNQ-quercetin; (b) 1UNQ-kaempferol; (c) 1YC5-kaempferol.

each other [58]. After the activation of the PI3K-Akt signaling pathway, the activity of inflammatory mediator genes was upregulated, which promoted the production of a large number of cytokines and the increase in the levels of TNF- α , IL-6, and IL-1 β , thus playing a role in regulating sleep [59]. Mitogen-activated protein kinase (MAPK) is a kind of serine/threonine protein kinases in cells. The MAPK-mediated signal transduction pathway is mainly related to the inflammatory response and plays a role in the phosphorylation of protein c-Jun [60]. Studies have shown that SP600125, a specific inhibitor of the JNK signaling pathway, significantly reduced the expression of phosphorylated c-Jun

in the substantia nigra of the midbrain in PD model mice [61]. It was found that MAPK, AKT1, and IL-6 in the serum of patients with anxiety were all increased, and there was a significant positive correlation between anxiety severity and sleep quality. With the improvement of anxiety state and sleep state, the expression of MAPK decreases gradually [62–64]. The FOXO signaling pathway is interrelated with the PI3K-Akt and MAPK signaling pathways [65, 66]. FoxO3a protein is an important subtype of the FoxO family and is an important cytokine in PI3K-Akt signal transduction [67]. After phosphorylation and modification of FoxO3a, specific downstream target genes can be activated

to induce autophagy and apoptosis of cells [68]. For example, when the MAPK pathway is activated or the PI3K-Akt pathway is inhibited, FoxO3a is dephosphorylated, thereby regulating different downstream factors and promoting cell apoptosis [68]. HIF-1 and its target gene *VEGF* can play a protective role in PD through mechanisms such as antioxidant stress, and the overexpression of *VEGF* can promote the proliferation and differentiation of neurons and reduce MPTP-induced substantia nigra cell injury [69]. The expression of angiogenic factor *VEGF* increased with an increase in the severity of sleep disturbance [70].

5. Conclusions

In this study, we used the network pharmacology research method to predict the chemical composition, target, and signal pathways at multiple levels. The prediction results were verified by molecular docking technology. The results show that SZRD plays an important role in the treatment of sleep disorders associated with PD through “multicomponent, multitarget, and multipathway,” which provides a new theoretical basis for further experimental research and clinical treatment.

Data Availability

The data used to support the findings of this study are available from the corresponding author upon request.

Conflicts of Interest

The authors declare that they have no conflicts of interest.

Authors' Contributions

Yan-yun Liu and Li-hua Yu contributed equally to this work.

Acknowledgments

The authors acknowledge the web database platform and software for data analysis. I would like to thank my tutor Professor Juan Zhang for her guidance on the ideas and writing of the article. Thanks are due to Professor Yu and Professor Xie for their guidance on the content of my article. Thanks are due to my classmates for their help in data collation and analysis. This article could not have been finished without their help. This study was totally supported by the National Natural Foundation of China (No. 81774299) and the Scientific Research Project of the Chinese National Medical Association (No. 2020ZY070-410033).

References

- [1] A. Berardelli, G. K. Wenning, A. Antonini et al., “EFNS/MDS-ES recommendations for the diagnosis of Parkinson’s disease,” *European Journal of Neurology*, vol. 20, no. 1, pp. 16–34, 2013.
- [2] “Guidelines for the treatment of Parkinson’s disease in China (fourth edition),” *Chinese Journal of Neurology*, vol. 53, no. 12, pp. 973–986, 2020.
- [3] Y. C. Pan and F. L. Wang, “Evaluation of the clinical effect of carbidopa-levodopa controlled-release tablet in the treatment of Parkinson’s disease with sleep disorder,” *World Journal of Sleep Medicine*, vol. 6, no. 5, pp. 544–546, 2019.
- [4] Z. Y. Liu and J. Wang, “Role of dopaminergic agonists in the treatment of non-motor symptoms of Parkinson’s disease,” *Shanghai Med Pharm*, vol. 36, no. 3, pp. 9–17, 2015.
- [5] H. J. Wei, M. Du, and H. Y. Bai, “Correlations of melatonin and glutathione levels with oxidative stress mechanism in Parkinson’s disease,” *Acta Academiae Medicinae Sinicae*, vol. 41, no. 2, pp. 183–187, 2019.
- [6] C. Juri, P. Chaná, J. Tapia, C. Kunstmann, and T. Parrao, “Quetiapine for insomnia in Parkinson Disease,” *Clinical Neuropharmacology*, vol. 28, no. 4, pp. 185–187, 2005.
- [7] A. P. Deng, M. Q. Wu, P. Q. Yuan, and L. Yanag, “Research progress of sleep disorders in patients with Parkinson’s disease,” *Journal of nursing*, vol. 26, no. 18, pp. 27–31, 2019.
- [8] Q. H. Zhou, H. L. Wang, X. L. Zhou et al., “Efficacy and safety of Suanzaoren decoction for chronic insomnia disorder in adults: study protocol for randomised, double-blind, double-dummy, placebo-controlled trial,” *BMJ Open*, vol. 7, no. 4, article e014280, 2017.
- [9] Y. Wang, X. Yang, P. F. Xia et al., “Review of chemical constituents, pharmacological effects and clinical applications of Suanzaoren decoction and prediction and analysis of its Q-markers,” *China journal of Chinese materia medica*, vol. 45, no. 12, pp. 2765–2771, 2020.
- [10] X. Y. Niu, B. S. He, Y. du et al., “The investigation of immunoprotective and sedative hypnotic effect of total polysaccharide from Suanzaoren decoction by serum metabolomics approach,” *Journal of Chromatography B*, vol. 1086, pp. 29–37, 2018.
- [11] L. H. Zhan, Y. J. Dong, K. Yang et al., “Soporific effect of modified Suanzaoren decoction and its effects on the expression of CCK-8 and orexin-A,” *Evidence-based Complementary and Alternative Medicine*, vol. 2020, Article ID 6984087, 17 pages, 2020.
- [12] B. Yang, A. H. Zhang, H. Sun et al., “Metabolomic study of insomnia and intervention effects of Suanzaoren decoction using ultra-performance liquid-chromatography/electrospray-ionization synapt high-definition mass spectrometry,” *Journal of Pharmaceutical and Biomedical Analysis*, vol. 58, pp. 113–124, 2012.
- [13] W. M. Yang, Y. C. Bao, H. Wang et al., “Diagnosis and treatment of fibrillation (Parkinson’s disease),” *Clinical Journal of Traditional Chinese Medicine*, vol. 24, no. 11, pp. 1125–1126, 2012.
- [14] X. Liu, P. Wang, L. Ding, S. B. Zhang, K. Shi, and Q. Y. You, “Effects of Suanzaoren decoction on the expression of nuclear receptor PPAR γ and its co-activator PGC-1 α in elderly chronic sleep deprivation rats,” *Modernization of Traditional Chinese Medicine and Materia Medica-World Science and Technology*, vol. 23, no. 5, pp. 1339–1346, 2021.
- [15] Y. X. Gong, “Clinical effect evaluation of Suanzaoren decoction on insomnia,” *Inner Mongolia Journal of Traditional Chinese Medicine*, vol. 40, no. 6, pp. 12–14, 2021.
- [16] K. T. Zhou and Y. X. Yu, “Clinical efficacy and safety of modified Suanzaoren decoction combined with escitalopram tablets in the treatment of insomnia,” *Chinese Journal of Clinical Rational Drug Use*, vol. 14, no. 15, pp. 134–136, 2021.

- [17] J. Ru, P. Li, J. Wang et al., "TCMSP: a database of systems pharmacology for drug discovery from herbal medicines," *Journal of Cheminformatics*, vol. 6, article 24735618, pp. 1–6, 2014.
- [18] S. Wang, H. Wang, and Y. Lu, "Tianfoshen oral liquid: a CFDA approved clinical traditional Chinese medicine, normalizes major cellular pathways disordered during colorectal carcinogenesis," *Oncotarget*, vol. 8, no. 9, pp. 14549–14569, 2017.
- [19] Y. Y. Wang, W. D. Yu, C. L. Shi et al., "Network pharmacology of Yougui pill combined with Buzhong Yiqi decoction for the treatment of sexual dysfunction," *Evidence-Based Complementary and Alternative Medicine*, vol. 2019, Article ID 1243743, 10 pages, 2019.
- [20] The UniProt Consortium, "UniProt: the universal protein knowledgebase," *Nucleic Acids Research*, vol. 45, no. D1, pp. D158–D169, 2017.
- [21] A. Hamosh, A. F. Scott, J. S. Amberger, C. A. Bocchini, and V. McKusick, "Online Mendelian Inheritance in Man (OMIM), a knowledgebase of human genes and genetic disorders," *Nucleic acids research*, vol. 33, no. Database issue, article 15608251, pp. D514–D517, 2005.
- [22] M. Safran, I. Dalah, J. Alexander et al., "GeneCards version 3: the human gene integrator," *Database*, vol. 2010, article 20689021, 2010.
- [23] J. Piñero, J. M. Ramírez-Anguita, J. Saüch-Pitarch et al., "The DisGeNET knowledge platform for disease genomics: 2019 update," *Nucleic Acids Research*, vol. 48, no. D1, pp. D845–D855, 2020.
- [24] D. Szklarczyk, A. L. Gable, D. Lyon et al., "STRING V11: protein-protein association networks with increased coverage, supporting functional discovery in genome-wide experimental datasets," *Nucleic Acids Research*, vol. 47, no. D1, pp. D607–D613, 2019.
- [25] P. Shannon, A. Markiel, O. Ozier et al., "Cytoscape: a software environment for integrated models of biomolecular interaction networks," *Genome Research*, vol. 13, no. 11, pp. 2498–2504, 2003.
- [26] D. W. Huang, B. T. Sherman, and R. A. Lempicki, "Systematic and integrative analysis of large gene lists using DAVID bioinformatics resources," *Nature protocols*, vol. 4, no. 1, pp. 2498–2504, 2009.
- [27] "Heatmap was plotted by <https://www.bioinformatics.com.cn/>, a free online platform for data analysis and visualization".
- [28] D. S. Goodsell, C. Zardecki, L. di Costanzo et al., "RCSB protein data bank: enabling biomedical research and drug discovery," *Protein Science*, vol. 29, no. 1, pp. 52–65, 2020.
- [29] N. El-Hachem, B. Haibe-Kains, A. Khalil, F. H. Kobeissy, and G. Nemer, "AutoDock and AutoDockTools for protein-ligand docking: beta-site amyloid precursor protein cleaving enzyme 1 (BACE1) as a case study," *Methods in Molecular Biology*, vol. 1598, article 28508374, pp. 391–403, 2017.
- [30] E. H. Baugh, S. Lyskov, B. D. Weitzner, and J. J. Gray, "Real-time PyMOL visualization for Rosetta and PyRosetta," *PLOS ONE*, vol. 6, no. 8, article e21931, 2011.
- [31] Z. W. Ma, P. C. He, and T. Ma, "Correlation analysis of sleep disturbance and serum vitamin D level in patients with Parkinson's disease," *Journal of Applied Clinical Medicine*, vol. 24, no. 24, pp. 53–56, 2020.
- [32] L. Y. Li, *Correlation between plasma melatonin levels and non-motor symptoms in patients with Parkinson's disease*, Zhengzhou university, 2020.
- [33] L. Wang, M. L. Huang, and R. Fu, "Correlation between serum Cysc, Hcy levels and sleep disorders in patients with early Parkinson's disease," *Chinese Journal of Gerontology*, vol. 40, no. 3, pp. 571–574, 2020.
- [34] X. F. Luo, Y. Li, J. Hu, and C. H. Zhang, "Regulating effect of gardenoside on sleep disturbance in experimental Parkinson's disease rats and its mechanism," *Journal of Jilin University (Medical Edition)*, vol. 46, no. 6, pp. 1177–1181, 2020.
- [35] Y. H. Tang, Y. Y. Li, H. Y. Yu et al., "Quercetin attenuates chronic ethanol hepatotoxicity: Implication of "free" iron uptake and release," *Food and Chemical Toxicology*, vol. 67, article 24569067, pp. 131–138, 2014.
- [36] P. Soucek, E. Kondrova, J. Hermanek et al., "New model system for testing effects of flavonoids on doxorubicin-related formation of hydroxyl radicals," *Anti-Cancer Drugs*, vol. 22, no. 2, pp. 176–184, 2011.
- [37] P. Rajendran, T. Rengarajan, N. Nandakumar, R. Palaniswami, Y. Nishigaki, and I. Nishigaki, "Kaempferol, a potential cytostatic and cure for inflammatory disorders," *European journal of medicinal chemistry*, vol. 86, article 25147152, pp. 103–112, 2014.
- [38] Y. H. Choi, "The cytoprotective effect of isorhamnetin against oxidative stress is mediated by the upregulation of the Nrf2-dependent HO-1 expression in C2C12 myoblasts through scavenging reactive oxygen species and ERK inactivation," *General Physiology and Biophysics*, vol. 35, no. 2, pp. 145–154, 2016.
- [39] D. K. Bao, J. K. Wang, X. B. Pang, and H. L. Liu, "Protective effect of quercetin against oxidative stress-induced cytotoxicity in rat pheochromocytoma (PC-12) cells," *Molecules*, vol. 22, no. 7, p. 1122, 2017.
- [40] S. Shamsheer and K. Puneet, "Piperine in combination with quercetin halt 6-OHDA induced neurodegeneration in experimental rats: biochemical and neurochemical evidences," *Neuroscience research*, vol. 133, article 29056550, pp. 38–47, 2018.
- [41] G. Filomeni, I. Graziani, D. de Zio et al., "Neuroprotection of kaempferol by autophagy in models of rotenone-mediated acute toxicity: possible implications for Parkinson's disease," *Neurobiology of Aging*, vol. 33, no. 4, pp. 767–785, 2012.
- [42] Y. Z. He, M. P. Ding, Y. K. Wang et al., "Protective effect of kaempferol on cerebral ischemia-reperfusion injury in rats," *Chinese Journal of Traditional Chinese Medicine*, vol. 27, no. 8, pp. 1673–1675, 2009.
- [43] Q. J. Chen and X. Gan, "Research progress on extraction of quercetin from *Sophora japonica* and its effect on central nervous system," *Chemical and biological engineering*, vol. 24, no. 3, pp. 11–13, 2007.
- [44] Y. H. Tao, *Study on enzyme inhibitors of γ -aminobutyric acid (GABA) metabolism*, Huazhong University of Science and Technology, Wuhan, 2006.
- [45] W. J. Chen, *Experimental study on anticonvulsant and sedative and hypnotic effects of bupleurin A*, Southern Medical University, Guangzhou, 2013.
- [46] J. W. Zhong, L. X. Chen, and B. C. Cai, "Effect of Chinese herb oyster on hypnotic effect of pentobarbital sodium," *Chinese Archives of Traditional Chinese Medicine*, vol. 27, no. 3, pp. 499–501, 2009.
- [47] Y. B. L. Rademacher, K. A. Matkowskyj, E. D. LaCount, and E. H. Carchman, "Topical application of a dual PI3K/mTOR inhibitor prevents anal carcinogenesis in a human papillomavirus mouse model of anal cancer," *European Journal of Cancer Prevention*, vol. 28, no. 6, pp. 483–491, 2019.

- [48] Z. Jiang, J. L. Zhang, J. T. Qiu, and S. Cui, "The PD-1/PD-L1 binding inhibitor BMS-202 suppresses the synthesis and secretion of gonadotropins and enhances apoptosis via p38MAPK signaling pathway," *Drug Development Research*, vol. 34309063, pp. 1–8, 2021.
- [49] Y. Y. Shi, X. T. Meng, Y. N. Xu, and X. J. Tian, "Role of FOXO protein's abnormal activation through PI3K/AKT pathway in platinum resistance of ovarian cancer," *Journal of Obstetrics and Gynaecology Research*, vol. 47, no. 6, pp. 1946–1957, 2021.
- [50] X. T. Guo, J. F. Liang, and X. B. Zhang, "Research progress on the relationship between PI3K/Akt/mTOR signaling pathway and Parkinson's disease," *Journal of Modern Medicine & Health*, vol. 31, no. 7, pp. 1019–1021, 2015.
- [51] R. L. Guo, J. Ya, X. J. Li, W. L. Wang, Z. F. Mo et al., "Expression of PI3K/Akt signaling pathway in tumors and its role in proliferation and apoptosis," *Hebei Medical Journal*, vol. 34, no. 12, pp. 1863–1864, 2012.
- [52] R. X. Lv, L. L. Du, F. H. Zhou, L. X. Zhang, and X. Y. Liu, "Study on the mechanism of rosmarinic acid inhibiting PI3K/Akt/mTOR signaling pathway and promoting autophagy to alleviate Parkinson's disease," *Journal of Ningxia Medical University*, vol. 41, no. 12, pp. 1189–1194, 2019.
- [53] W. Shearer, J. Reuben, J. Mullington et al., "Soluble TNF- α receptor 1 and IL-6 plasma levels in humans subjected to the sleep deprivation model of spaceflight," *The Journal of Allergy and Clinical Immunology*, vol. 107, no. 1, pp. 165–170, 2001.
- [54] P. Entzian, K. Linnemann, M. Schlaak, and P. Zabel, "Obstructive sleep apnea syndrome and circadian rhythms of hormones and cytokines," *American Journal of Respiratory and Critical Care Medicine*, vol. 153, no. 3, pp. 1080–1086, 1996.
- [55] L. Zhang, R. H. Zhang, Y. Shen, S. Z. Qiao, Z. L. Hui, and J. Chen, "Shimian granules improve sleep, mood and performance of shift nurses in association changes in melatonin and cytokine biomarkers: a randomized, double-blind, placebo-controlled pilot study," *Chronobiology International*, vol. 37, no. 4, pp. 592–605, 2020.
- [56] Y. Huang, L. Xia, and G. H. Chen, "Study on changes of tumor necrosis factor- α in patients with primary insomnia," *Chinese Journal of Clinicians (electronic version)*, vol. 7, no. 7, pp. 2877–2879, 2013.
- [57] K. Wright, A. Drake, D. Frey et al., "Influence of sleep deprivation and circadian misalignment on cortisol, inflammatory markers, and cytokine balance," *Brain, Behavior, and Immunity*, vol. 47, article 25640463, pp. 24–34, 2015.
- [58] A. S. Guo, C. Liu, B. S. Li, Z. J. Zhu, W. G. Chen, and A. H. Li, "Effects of acupoint embedding thread on monoamine transmitters, IL-1 β and TNF- α contents in hypothalamus of rats with insomnia," *Shanghai Journal of Acupuncture and Moxibustion*, vol. 33, no. 7, pp. 672–675, 2014.
- [59] J. Luyendyk, G. Schabbauer, M. Tencati, T. Holscher, R. Pawlinski, and N. Mackman, "Genetic analysis of the role of the PI3K-Akt pathway in lipopolysaccharide-induced cytokine and tissue factor gene expression in monocytes/macrophages," *Journal of Immunology*, vol. 180, no. 6, pp. 4218–4226, 2008.
- [60] S. Y. Liu, P. Xu, X. L. Luo, J. F. Hu, and X. H. Liu, "(7R,8S)-Dehydrodiconiferyl alcohol suppresses lipopolysaccharide-induced inflammatory responses in BV2 microglia by inhibiting MAPK signaling," *Neurochemical Research*, vol. 41, no. 7, pp. 1570–1577, 2016.
- [61] Y. S. Wang, Z. F. Wei, Q. Y. Tian, Z. F. Zhang, H. X. Zhou, and Y. X. Zhang, "Regulation of JNK pathway in the process of inflammation and apoptosis of substantia nigra cells in subacute Parkinson's disease model mice," *Journal of Clinical and Experimental Pathology*, vol. 5, pp. 601–605, 2007.
- [62] N. Yoshihiko, A. Taku, Y. Hikaru, S. Norio, T. Minoru, and T. Hiroshi, "Yokukansan enhances the proliferation of B65 neuroblastoma," *Journal of Traditional and Complementary Medicine*, vol. 7, no. 1, pp. 34–44, 2017.
- [63] C. X. Yang, N. Sun, Y. Ren et al., "Association between AKT1 gene polymorphisms and depressive symptoms in the Chinese Han population with major depressive disorder," *Neural Regeneration Research*, vol. 7, no. 3, pp. 235–239, 2012.
- [64] A. Niraula, K. G. Witcher, J. F. Sheridan, and J. P. Godbout, "Interleukin-6 induced by social stress promotes a unique transcriptional signature in the monocytes that facilitate anxiety," *Biological Psychiatry*, vol. 85, no. 8, pp. 679–689, 2019.
- [65] H. C. Wu, J. W. Zhang, Z. G. Sun, S. Xiang, Y. Qiao, and F. Lian, "Effects of electroacupuncture on expression of PI3K/Akt/Foxo3a in granulosa cells from women with Shen (kidney) deficiency syndrome undergoing in vitro fertilization-embryo transfer," *Chinese Journal of Integrative Medicine*, vol. 25, no. 4, pp. 252–258, 2019.
- [66] F. Matsuda, N. Inoue, A. Maeda et al., "Expression and function of apoptosis initiator FOXO3 in granulosa cells during follicular atresia in pig ovaries," *Journal of Reproduction and Development*, vol. 57, no. 1, pp. 151–158, 2011.
- [67] D. M. Guo, L. L. Xiao, H. J. Hu, M. H. Liu, L. Yang, and X. Lin, "FGF21 protects human umbilical vein endothelial cells against high glucose-induced apoptosis via PI3K/Akt/Fox3a signaling pathway," *Journal of Diabetes and its Complications*, vol. 32, no. 8, pp. 729–736, 2018.
- [68] A. Tseng, L. H. Wu, S. S. Shieh, and D. L. Wang, "SIRT3 interactions with FOXO3 acetylation, phosphorylation and ubiquitination mediate endothelial cell responses to hypoxia," *The Biochemical Journal*, vol. 464, no. 1, pp. 157–168, 2014.
- [69] Z. Zhang, J. Yan, Y. Chang, S. ShiDu Yan, and H. Shi, "Hypoxia inducible factor-1 as a target for neurodegenerative diseases," *Current Medicinal Chemistry*, vol. 18, no. 28, pp. 4335–4343, 2011.
- [70] Y. B. Dong, G. Y. Wu, T. Zhu et al., "VEGF promotes cartilage angiogenesis by phospho-ERK1/2 activation of Dll4 signaling in temporomandibular joint osteoarthritis caused by chronic sleep disturbance in Wistar rats," *Oncotarget*, vol. 8, no. 11, pp. 17849–17861, 2017.