

Long-Term Efficacy and Low Adverse Events of Methylprednisolone Pulses Combined to Low-Dose Glucocorticoids for Systemic Sclerosis: A Retrospective Clinical Study of 10 Years' Follow-Up [Letter]

Mingrui Xie

Medway NHS Foundation Trust, Gillingham, Kent, UK

Correspondence: Mingrui Xie, Medway NHS Foundation Trust, Windmill Road, Gillingham, Kent, ME7 5NY, UK, Tel +447857121148, Email mingrui.xie@nhs.net

Dear editor

I read with great interest the research article by Cheng et al¹ on long-term low dose glucocorticoid treatment for systemic sclerosis (SSc). Rheumatic diseases can be debilitating, often only controlled by high dose steroids which comes with its own toxicity profile. It is encouraging to see research into limiting detrimental side effects of treatment for these patients, hence I would like to share my thoughts on this.

It is impressive that the authors thought of alternative steroid regimens in order to bring maximal benefits to patients. However, even low dose corticosteroids can be detrimental and an article the authors cited also reported the same, where fractures have occurred quite early on even in patients on low doses.² Thus, it is highly commendable that although the authors' main focus is on steroids, they have taken other steroid-sparing agents into consideration. Unfortunately, as they mentioned, none of the conventional immunosuppressive therapies for SSc have shown to be effective for skin fibrosis, but nintedanib could be helpful for SSc with pulmonary involvement such as interstitial lung disease. This brings to mind the GiACTA trial³ which showed tocilizumab as a successful agent to maintain remission for giant cell arteritis, another rheumatic disease, leading to it being licensed for use in several countries. Similarly, there are early trials showing benefits of tocilizumab for SSc.⁴ Another drug worth considering is rituximab, where small trials have shown benefits⁵ and even resulted in regulatory approval for SSc in Japan. These are all promising and I hope there can be more investigations into them in future.

I appreciate the authors noting the limits of the study, one of which is the lack of blinding, and I am curious as to why it was not done. Although most of the clinical evaluations are quite objective, the modified Rodnan skin score could still be subjective despite efforts to standardize it, and not having blinding could definitely bias the assessor.

Even though this study only reports the result of 46 patients, which seems like a small sample size, I note that the authors started off with more and 128 patients were lost to follow up during the 10 years. It would be helpful to know the reasons for that as it would give a clearer picture on how patients tolerated steroids long term.

I also note the authors excluded SSc patients with any other organ involvement other than skin and lungs. I hope they can elaborate on the reason as SSc can affect the heart, kidneys, bowel, and muscles, in order to help readers understand the population of the trial better.

Therefore, it is good that alternative steroid regimens that seemingly have a lower rate of adverse effects are being studied. However, because of the well-documented toxicity of steroids and having seen many examples of that in my own

clinical practice, I eagerly await further studies of a larger sample size and also potentially other steroid-sparing anti-inflammatory agents.

Disclosure

The author reports no conflicts of interest in this communication.

References

1. Cheng H, Yu Z, Yan CL, Yang HD, Gao C, Wen HY. Long-term efficacy and low adverse events of methylprednisolone pulses combined to low-dose glucocorticoids for systemic sclerosis: a retrospective clinical study of 10 years' follow-up. *J Inflamm Res*. 2022;15:4421–4433. doi:10.2147/JIR.S373387
2. Adami G, Saag KG. Glucocorticoid-induced osteoporosis: 2019 concise clinical review. *Osteoporos Int*. 2019;30(6):1145–1156. doi:10.1007/s00198-019-04906-x
3. Stone JH, Tuckwell K, Dimonaco S, et al. Trial of tocilizumab in giant-cell arteritis. *N Engl J Med*. 2017;377(4):317–328. doi:10.1056/NEJMoa1613849
4. Panopoulos ST, Tektonidou MG, Bournia VK, Arida A, Sfikakis PP. Anti-interleukin 6 therapy effect for refractory joint and skin involvement in systemic sclerosis: a real-world, single-center experience. *J Rheumatol*. 2022;49(1):68–73. doi:10.3899/jrheum.210273
5. Ebata S, Yoshizaki A, Oba K, et al. Safety and efficacy of rituximab in systemic sclerosis (DESIREs): open-label extension of a double-blind, investigators-initiated, randomised, placebo-controlled trial. *Lancet Rheumatol*. 2022;4(8):e546–e555. doi:10.1016/S2665-9913(22)00131-X

Dove Medical Press encourages responsible, free and frank academic debate. The content of the Journal of Inflammation Research 'letters to the editor' section does not necessarily represent the views of Dove Medical Press, its officers, agents, employees, related entities or the Journal of Inflammation Research editors. While all reasonable steps have been taken to confirm the content of each letter, Dove Medical Press accepts no liability in respect of the content of any letter, nor is it responsible for the content and accuracy of any letter to the editor.

Journal of Inflammation Research

Dovepress

Publish your work in this journal

The Journal of Inflammation Research is an international, peer-reviewed open-access journal that welcomes laboratory and clinical findings on the molecular basis, cell biology and pharmacology of inflammation including original research, reviews, symposium reports, hypothesis formation and commentaries on: acute/chronic inflammation; mediators of inflammation; cellular processes; molecular mechanisms; pharmacology and novel anti-inflammatory drugs; clinical conditions involving inflammation. The manuscript management system is completely online and includes a very quick and fair peer-review system. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/journal-of-inflammation-research-journal>

<https://doi.org/10.2147/JIR.S387159>