

Association Between Nitrate-Reducing Oral Bacteria and Cardiometabolic Outcomes: Results From ORIGINS

Charlene E. Goh, BDS, DrPH; Pauline Trinh, MPH; Paolo C. Colombo, MD; Jeanine M. Genkinger, PhD; Barun Mathema, PhD; Anne-Catrin Uhlemann, MD, PhD; Charles LeDuc, PhD; Rudolph Leibel, MD; Michael Rosenbaum, MD; Bruce J. Paster, PhD; Moïse Desvarieux, MD, PhD; Panos N. Papapanou, DDS, PhD; David R. Jacobs Jr, PhD; Ryan T. Demmer, PhD

Background—The enterosalivary nitrate-nitrite-nitric oxide pathway is an alternative pathway of nitric oxide generation, potentially linking the oral microbiome to insulin resistance and blood pressure (BP). We hypothesized that increased abundance of nitrate-reducing oral bacteria would be associated with lower levels of cardiometabolic risk cross-sectionally.

Methods and Results—ORIGINS (Oral Infections, Glucose Intolerance, and Insulin Resistance Study) enrolled 300 diabetes mellitus-free adults aged 20 to 55 years (mean=34±10 years) (78% women). Microbial DNA was extracted from subgingival dental plaque (n=281) and V3–V4 regions of the 16S rRNA gene were sequenced to measure the relative abundances of 20 a priori-selected taxa with nitrate-reducing capacity. Standardized scores of each taxon's relative abundance were summed, producing a nitrate-reducing taxa summary score (NO₃TSS) for each participant. Natural log-transformed homeostatic model assessment of insulin resistance, plasma glucose, systolic BP, and diastolic BP were regressed on NO₃TSS in multivariable linear regressions; prediabetes mellitus and hypertension prevalence were regressed on NO₃TSS using modified Poisson regression models. Nitrate-reducing bacterial species represented 20±16% of all measured taxa. After multivariable adjustment, a 1-SD increase in NO₃TSS, was associated with a −0.09 (95% CI, −0.15 to −0.03) and −1.03 mg/dL (95% CI, −1.903 to −0.16) lower natural log-transformed homeostatic model assessment of insulin resistance and plasma glucose, respectively. NO₃TSS was associated with systolic BP only among patients without hypertension; 1-SD increase in NO₃TSS was associated with −1.53 (95% CI, −2.82 to −0.24) mm Hg lower mean systolic BP. No associations were observed with prediabetes mellitus and hypertension.

Conclusions—A higher relative abundance of oral nitrate-reducing bacteria was associated with lower insulin resistance and plasma glucose in the full cohort and with mean systolic BP in participants with normotension. (*J Am Heart Assoc.* 2019;8:e013324. DOI: 10.1161/JAHA.119.013324.)

Key Words: epidemiology • high blood pressure • insulin resistance • nitrate • oral microbiome

Increasing evidence suggests that the digestive tract microbiome (ie, bacteria colonizing the oral cavity and the gastrointestinal tract) may contribute to the development of insulin resistance,^{1,2} type 2 diabetes mellitus,^{3,4} and hypertension.^{5,6} These associations between the oral microbiome and

increased cardiometabolic risk are most commonly hypothesized to result from a chronic inflammatory response to a dysbiotic subgingival microbiome.⁷ However, a possible alternative mechanism is via the production of the physiologically important gaseous transmitter, nitric oxide (NO).

From the Faculty of Dentistry, National University of Singapore, Singapore (C.E.G.); Department of Epidemiology, Columbia University, Mailman School of Public Health (P.T., J.M.G., B.M., M.D.), Division of Cardiology, Department of Medicine, Columbia University (P.C.C.), Herbert Irving Comprehensive Cancer Center, Columbia University Irving Medical Center (J.M.G.), Division of Infectious Diseases and Microbiome and Pathogen Genomics Core, Department of Medicine, Columbia University Irving Medical Center (A.-C.U.), Division of Molecular Genetics, Departments of Pediatrics and Medicine (C.L., R.L., M.R.), and Division of Periodontics, Section of Oral and Diagnostic Sciences, College of Dental Medicine (P.N.P.), Columbia University, New York, NY; The Forsyth Institute, Cambridge, MA (B.J.P.); Department of Oral Medicine, Infection, and Immunity, Harvard School of Dental Medicine, Boston, MA (B.J.P.); INSERM UMR 1153, Centre de Recherche Epidemiologie et Statistique Paris Sorbonne Cité (CRESS), METHODS Core, Paris, France (M.D.); Division of Epidemiology and Community Health, School of Public Health, University of Minnesota, Minneapolis, MN (D.R.J., R.T.D.).

Accompanying Data S1, Tables S1 through S7, and Figure S1 are available at <https://www.ahajournals.org/doi/suppl/10.1161/JAHA.119.013324>

Correspondence to: Ryan T. Demmer, PhD, Division of Epidemiology and Community Health, School of Public Health, University of Minnesota, 300 West Bank Office Building, 1300 S 2nd Street, Minneapolis, MN 55454. E-mail: demm0009@umn.edu

Received May 18, 2019; accepted September 18, 2019.

© 2019 The Authors. Published on behalf of the American Heart Association, Inc., by Wiley. This is an open access article under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

Clinical Perspective

What Is New?

- The oral microbiome plays an important role in the enterosalivary nitrate-nitrite-nitric oxide pathway.
- To the best of our knowledge, this is the first study to directly examine the relationship between specific nitrate-reducing oral microbiota and cardiometabolic outcomes in a population setting.
- Our results support the hypothesis that oral nitrate-reducing bacteria play a beneficial role in blood pressure regulation and insulin resistance.

What Are the Clinical Implications?

- If this relationship proves to be causal, oral microbial risk factors for cardiometabolic outcomes may be identified, and further research could yield useful treatments that manipulate the oral microbiome to improve cardiometabolic health.

NO is an important signaling molecule involved in many physiological processes, including endothelial function, vasodilation, immune function, glucose metabolism, and blood pressure (BP) control.⁸ A loss of NO production and bioavailability has been implicated in the pathogenesis of insulin resistance and hypertension.^{9,10} NO production was originally thought to occur solely through the endogenous conversion of L-arginine and oxygen into NO and L-citrulline by NO synthases found in the endothelium and other tissues. However, it has recently been discovered that NO production can also occur via the reduction of salivary nitrates by nitrate-reducing oral bacteria to form nitrites, which are then swallowed and made systemically bioavailable for further reduction into NO in the blood vessels and tissues.⁸ This so-called enterosalivary nitrate-nitrite-NO pathway presents a novel and biologically plausible mechanism by which oral bacteria might influence the systemic bioavailability of NO and the development of related clinical cardiometabolic outcomes in humans.

The enterosalivary pathway is thought to underlie the strong evidence from experimental studies suggesting that increased dietary nitrate intake has beneficial systemic cardiometabolic effects. A systematic review of 13 trials lasting 1 to 6 weeks found an \approx 4.1 and 2.0 mm Hg reduction in systolic BP (SBP) and diastolic BP (DBP), respectively, following daily nitrate supplementation.¹¹ Studies in mice have shown that dietary nitrate can improve insulin signaling and reverse features of metabolic syndrome.^{12,13} Reduced plasma glucose and improved insulin sensitivity following nitrate supplementation have also been observed in some human studies, although the results are less conclusive.^{14–16}

The direct role of nitrate-reducing oral bacteria in the enterosalivary pathway of NO production is supported by several small experimental studies that use antibacterial mouthwash to reduce the overall oral bacteria. Antibacterial mouthwash use significantly blunts the BP and plasma glucose reductions observed following experimental nitrate supplementation.^{14,17,18} Notably, even in the absence of exogenous nitrate supplementation, a decrease in salivary and plasma nitrite, and an \approx 3 mm Hg increase in SBP and DBP was observed after antibacterial mouthwash use.¹⁹ This finding suggests that oral microbiota play a continuous role in BP regulation through the nitrate-nitrite-NO pathway.

To our knowledge, no study has directly investigated the relationship between specific oral microbiota with known nitrate-reducing capacity and cardiometabolic outcomes in a population setting. Only a few clinical trials have directly correlated abundance of nitrate-reducing oral bacteria with cardiometabolic outcomes,^{20,21} and associations for only a few species of nitrate-reducing bacteria were reported. Furthermore, the population distribution of oral nitrate-reducing bacteria remains unexplored and it is unknown whether nuanced variation, rather than the extreme differences in nitrate-reducing taxa created by mouthwash use, is beneficially related to cardiometabolic parameters. The purpose of this study is to examine the cross-sectional relationship between subgingival nitrate-reducing bacteria and cardiometabolic outcomes in diabetes mellitus-free adults enrolled in ORIGINS (Oral Infections, Glucose Intolerance, and Insulin Resistance Study).⁴ We hypothesize that higher relative abundance of nitrate-reducing oral bacteria will be associated with lower levels of insulin resistance and BP, as well as a lower prevalence of prediabetes mellitus and hypertension.

Methods

The data that support the findings of this study are available from the corresponding author upon reasonable request.

Description of ORIGINS

ORIGINS is a cohort study investigating the relationship between subgingival microbial community composition and impaired glucose metabolism.⁴ The cross-sectional data used in this study are from the baseline wave 1 (n=300 participants) enrolled from February 2011 to 2013, with the following inclusion criteria: (1) aged 20 to 55 years; (2) no diabetes mellitus (type 1 or type 2) based on participant self-report, glycated hemoglobin values <6.5%, and fasting plasma glucose <126 mg/dL; and (3) no self-reported history of myocardial infarction, congestive heart failure, stroke, or chronic inflammatory conditions. All participants reported not

taking antibiotics in the past 30 days. Participants underwent oral examinations (including periodontal measurements), collection of oral bacteria specimens, blood draw after an overnight fast, and in-person anthropometric assessments at the same visit. Columbia University's institutional review board approved the study protocol. All participants provided written informed consent.

Of the 300 participants, the present analyses include only the 281 participants without missing 16S rRNA data or important baseline cardiometabolic risk factors.

Bacterial Assessment and Identification

Subgingival plaque samples (n=281) were collected from prespecified sites. The mesiobuccal site of the second-most posterior tooth in the lower left quadrant (excluding third molars) of each participant was sampled using sterile curettes after removal of the supragingival plaque.⁴ The samples were suspended in 300 μ L of TE buffer (50 mmol/L Tris, 1 mmol/L EDTA; pH 7.6), and microbial DNA was extracted using the MasterPure Gram Positive DNA Purification Kit (Epicentre).²²

Next-generation sequencing of the 16S rRNA gene was performed at the Forsyth Institute. The Human Oral Microbiome Identification using Next Generation Sequencing (HOMINGS) methodology^{22,23} is designed specifically for oral taxa, generating species-level information with high precision. Briefly, 50 ng of DNA was used to amplify the V3–V4 region of the 16S rRNA gene (using 341F/806R universal primers) and PCR products were purified using AMPure beads. Amplicons were then sequenced on the MiSeq (Illumina) platform. Paired-end reads were joined using QIIME join fastq; minimum overlap was set to 70 bp, and the percent max difference was 25%. Nonbarcoded sequences and sequences with a Phred quality score <25 were excluded. Samples with <5000 reads were excluded from the analyses. Overall, 18 531 931 sequences were generated for final analysis (median of 75 977 sequences per sample).

HOMINGS follows an in silico hybridization process. A BLAST program, called “ProbeSeq for HOMINGS,” uses specially designed in silico species-specific 16S rRNA-based oligonucleotide probes to identify species taxa and frequency.²⁴ An array is created using the raw sequence files and the program loops through examining one sequence at a time, looking for a “string” that fully matches one of the probes. The total number of matches, or unique in silico-hybridization events, are then counted with each match representing the conceptual identification of 1 bacterial cell. ProbeSeq is an iterative process, and sequences not detected by a species-level probe are then processed against genus-level probes. The final HOMINGS data output for each individual are expressed as the relative abundance of each target taxa (by dividing the respective HOMINGS hits for that

taxa by the sum of all taxa hits within the individual, ie, percent proportions of each target taxa). Overall, each sample had an average of 22% (SD=12%) unmapped reads that matched to neither species nor genus probe. Using HOMINGS, 668 different taxa were identified in ORIGINS, with an average of 182 (SD=50) taxa identified in each participant sample.

Operationalization of Nitrate-Reducing Oral Bacteria Exposure

Exposure to nitrate-reducing oral bacteria was defined by creating a summary score comprising oral bacteria species previously identified in the literature as being associated with nitrate-reduction capacity.^{25,26} From the list of 28 putative nitrate-reducing oral species (Table S1), 20 taxa were identified by HOMINGS.²⁷ These 20 nitrate-reducing bacteria overall showed low correlations with each other (Figure S1). To address the skewed distributional properties of using proportional data, a variance-stabilizing arcsin-square root transformation commonly used in microbiome analyses was first applied to the relative abundance of each taxa.²⁸ The arcsin-square root transformed relative abundance of each taxa was then standardized by dividing by its SD, as per an a priori approach described elsewhere.⁴ Standardized values for each of the nitrate-reducing taxa were then summed, creating a summary score representing the total nitrate-reducing microbiota community exposure in the sample. The standardization gives equal weight to each taxa, and, without complete knowledge of their nitrate-reducing capacity, prevents a summary score from being dominated by the most abundant taxa (Table S2).

Outcomes

Insulin resistance and plasma glucose

Plasma glucose and insulin levels were measured from blood collected following an overnight fast. Insulin resistance was measured using homeostatic model assessment of insulin resistance (HOMA-IR) values calculated from fasting insulin and glucose levels.²⁹

Prediabetes Mellitus

Prediabetes mellitus (yes/no) was defined in accordance with the American Diabetes Association criteria as follows: (1) fasting plasma glucose \geq 100 mg/dL and <126 mg/dL; or (2) glycated hemoglobin \geq 5.7% and <6.5%.³⁰

SBP and DBP

Seated resting SBP and DBP were measured in triplicate and the last 2 measurements averaged to obtain our continuous measures of mean SBP and DBP (mm Hg).

Hypertension

Hypertension (yes/no) was defined in accordance with the most recent 2017 American Heart Association criteria as follows: (1) an SBP recording of ≥ 130 mm Hg; or (2) a DBP recording ≥ 80 mm Hg.³¹ Participants were also classified as having hypertension if a diagnosis of hypertension was self-reported.

Risk Factor Assessment

Cardiometabolic risk factors were measured by trained research assistants as previously described.^{4,32} Participant body mass index was calculated as weight in kilograms/height in meters.² Questionnaires were administered to obtain information on: age, sex, race/ethnicity (non-Hispanic black, non-Hispanic white, Hispanic, other), educational level (high school completion, college/vocational training, advanced degrees), cigarette smoking (current, former, or never smoking, and duration/intensity of smoking). Overall dietary pattern was assessed using the Alternative Healthy Eating Index 2010 (AHEI 2010) that was created based on foods and nutrients predictive of chronic disease risk.³³ The index consists of several components: vegetables, fruits, whole grains, nuts and vegetable protein, red/processed meat, sugar-sweetened beverages and fruit juice, trans fats, polyunsaturated fats, long-chain fatty acids, sodium, and alcohol consumption. Each food group has a range of 0 to 10 points, which are then summed to create the overall score. The AHEI 2010 score ranges from 0 to 110, with higher AHEI scores associated with a lower risk of coronary heart disease and diabetes mellitus.³³ Leisure-time physical activity was assessed and converted to metabolic equivalents, and participants were categorized into 4 leisure-time physical activity categories as previously described.³² Measures of periodontitis were obtained from the clinical periodontal examinations as previously published,⁴ and periodontal status was measured by the percentage of periodontal sites with attachment loss ≥ 3 mm. (see Data S1 for additional information on risk factor operationalization.)

Statistical Analysis

To address the skewed distribution of HOMA-IR values, insulin resistance was operationalized as natural log-transformed HOMA-IR (lnHOMA-IR) in the analyses. Geometric means are presented after back-transforming predicted means obtained in regression analyses described below. Multivariable models regressed continuous measures of lnHOMA-IR, plasma glucose levels, SBP, and DBP (dependent variables) on the continuous summary score for nitrate-reducing bacteria (NO₃TSS) in separate regressions for each outcome. Results were also presented visually in categories of increasing

intervals of SD. Because the outcomes of prediabetes mellitus and hypertension were common in our study population, relative risk regression models using a modified Poisson regression with robust error variance were used to calculate the prevalence ratios instead of odds ratios.³⁴ To avoid the possibility of behavioral modification and medications (after a hypertension diagnosis) from masking the associations with bacteria, sensitivity analyses for SBP and DBP outcomes were conducted using only the 187 participants with normotension. Exploratory analyses of the relationship between the 20 individual bacteria taxa and the cardiometabolic outcomes of interest were also conducted.

All multivariable regressions were adjusted for the potential confounders of age, sex, race, and smoking status a priori based on previous studies, with education, body mass index, percentage of probing sites with attachment loss ≥ 3 mm, and dietary pattern additionally included as they were associated with the exposure and outcomes at an $\alpha=0.20$ level of significance (Table S3). Additional sensitivity analysis was also conducted in which alcohol use and physical activity were added to the regression model.

Results

Sample Characteristics

The demographic characteristics of the ORIGINS cohort (n=281) are presented in Table S4. The mean age of our study population was 34 years (SD=10 years), and the majority were women (78%), college educated (67%), and never smokers (79%). A total of 42% of participants had none or mild periodontitis as defined per the Centers for Disease Control and Prevention/American Academy of Periodontology guidelines.³⁵ A total of 95% of the a priori–selected sites from which subgingival plaque was sampled had a probing depth ≤ 3 mm, and the remaining 5% (11 sites) had a probing depth of 4 mm. The prevalence of prediabetes mellitus and hypertension in this population was 18% (n=50) and 33% (n=93), respectively. The characteristics of the participants with normotension were comparable to the whole sample (Table S4).

Prevalence and Relative Abundance of Individual Nitrate-Reducing Bacteria

The mean relative abundances and prevalence of the 20 individual nitrate-reducing bacterial taxa are presented in Figure 1. *Rothia dentocariosa* had the highest mean relative abundance (7.9%) and was detected in all participants, whereas *Propionibacterium acnes* had the lowest relative abundance (0.0002%) and was detected in only 6% of participants. However, it should be noted that at such low relative abundance ($<0.0005\%$) the reliability of this taxa




Figure 1. The prevalence (%) and mean relative abundance (%) of the 20 nitrate-reducing taxa measured in subgingival plaque samples among the 281 participants in ORIGINS (Oral Infections, Glucose Intolerance, and Insulin Resistance Study).

distribution is poor. Participants had a mean total relative abundance of nitrate-reducing taxa of 20% (SD=16%; range: 0.09%–86%), and many of the nitrate-reducing bacteria species were present in most participants. The mean nitrate-reducing bacterial summary score NO₃TSS was ≈0 (SD=5.42).

Association of Nitrate-Reducing Bacterial Summary Score With Insulin Resistance, Plasma Glucose, and Prediabetes Mellitus

The mean (IQR) HOMA-IR and mean (SD) plasma glucose values in this population were 1.75 (1.45) and 85 mg/dL (7.6 mg/dL),

respectively. A higher NO₃TSS was associated with lower insulin resistance. Every 1-SD higher NO₃TSS, was associated with a −0.09 (95% CI, −0.15 to −0.03) lower lnHOMA-IR, controlling for age, sex, race, education, body mass index, smoking status, percentage of probing sites with attachment loss ≥3 mm, and dietary pattern (Table 1). The geometric means of the HOMA-IR values across increasing SD intervals of NO₃TSS were 1.85 (95% CI, 1.55–2.22), 1.89 (95% CI, 1.66–2.16), 1.59 (95% CI, 1.38–1.83), and 1.46 (95% CI, 1.22–1.73) (linear trend $P=0.003$). Mean values of lnHOMA-IR are presented in Figure 2.

Inverse associations were also observed between NO₃TSS and baseline plasma glucose (Table 1). In multivariable

Table 1. Mean Difference in Natural Log-Transformed Homeostasis Model Assessment for Insulin Resistance (lnHOMA-IR), Plasma Glucose Levels (mg/dL), and Prevalence Ratio of Prediabetes for Every 1 Standard Deviation (STD) Increase in Nitrate-Reducing Taxa Summary Score (NO₃TSS)

Model	Insulin Resistance (lnHOMA-IR)	Glucose, mg/dL	Prediabetes Mellitus (Prevalence Ratio)
	NO ₃ TSS (1 SD)	NO ₃ TSS (1 SD)	NO ₃ TSS (1 SD)
1	−0.10 (−0.16 to −0.03)	−1.05 (−1.93 to −0.16)	0.90 (0.66–1.21)
2	−0.08 (−0.14 to −0.02)	−0.85 (−1.67 to −0.04)	0.95 (0.74–1.21)
3	−0.08 (−0.13 to −0.02)	−0.81 (−1.61 to −0.02)	0.93 (0.73–1.19)
4	−0.08 (−0.14 to −0.02)	−0.84 (−1.64 to −0.03)	0.94 (0.73–1.21)
5	−0.09 (−0.15 to −0.03)	−1.03 (−1.90 to −0.16)	0.79 (0.61–1.03)
6	−0.09 (−0.15 to −0.02)	−1.43 (−2.30 to −0.58)	0.83 (0.63–1.09)

Model 1: crude (n=281). Model 2: adjusted for age, sex, race, and education (n=281). Model 3: adjusted for age, sex, race, education, body mass index (BMI), and smoking (n=281). Model 4: adjusted for age, sex, race, education, BMI, smoking, and percentage of periodontal sites with attachment loss ≥3 mm (n=280). Model 5: adjusted for age, sex, race, education, BMI, smoking, percentage of periodontal sites with attachment loss ≥3 mm, and dietary pattern (n=253). Model 6: adjusted for age, sex, race, education, BMI, smoking, percentage of periodontal sites with attachment loss ≥3 mm, dietary pattern, alcohol use, and physical activity (n=226). lnHOMA-IR indicates natural log-transformed homeostatic model assessment of insulin resistance; NO₃TSS, nitrate-reducing taxa summary score.




Figure 2. Natural log-transformed homeostatic model assessment of insulin resistance (HOMA-IR) values, plasma glucose, systolic blood pressure (SBP), and diastolic blood pressure (DBP) (95% CI) across increasing SD intervals of nitrate-reducing taxa summary score (NO_3TSS). Adjusted for age, sex, race, education, body mass index (BMI), smoking, percentage of probing sites with attachment loss ≥ 3 mm, and dietary pattern. Note: <-1 SD ($n=43$); ≥ -1 to 0 SD ($n=99$); ≥ 0 to 1 SD ($n=96$); and ≥ 1 SD ($n=43$).

analyses, every 1-SD higher NO_3TSS was associated with a -1.03 (95% CI, -1.90 to -0.16) lower plasma glucose level (mg/dL). Prediabetes mellitus prevalence tended to decrease as NO_3TSS levels increased (0.79; 95% CI, 0.61–1.03) (Table 1).

Association of Nitrate-Reducing Bacterial Summary Score With BP and Hypertension

The mean (SD) SBP and DBP was 117 mm Hg (12 mm Hg) and 75 mm Hg (10 mm Hg), respectively, in the whole sample; 112 mm Hg (9 mm Hg) and 70 mm Hg (6 mm Hg), respectively, for participants with normotension ($n=187$) and 128 mm Hg (11 mm Hg) and 84 mm Hg (9 mm Hg), respectively, for participants with hypertension ($n=93$).

When examining the association between NO_3TSS and BP outcomes in the full sample, NO_3TSS was not significantly associated with SBP or DBP (Table 2). However, in sensitivity analyses including only participants with normotension, the effect estimates were similarly inverse but larger for SBP. A SD higher NO_3TSS was associated with a -1.53 mm Hg (95% CI, -2.82 to -0.24) lower mean SBP (Table 2). Multivariable

adjusted mean values of SBP across increasing SD intervals of NO_3TSS were 118 mm Hg (95% CI, 114–122), 115 mm Hg (95% CI, 112–118), 115 mm Hg (95% CI, 112–118), and 112 mm Hg (95% CI, 108–116) (linear trend $P=0.02$) (Figure 2). The association between NO_3TSS and mean DBP was smaller but likewise inverse (-0.60 ; 95% CI, -1.54 to 0.33) (Table 2). Higher NO_3TSS was not associated with a higher prevalence ratio of hypertension (1.03; 95% CI, 0.89–1.20).

Sensitivity analyses defining hypertension using old thresholds of SBP ≥ 140 mm Hg or DBP ≥ 90 mm Hg³⁶ found similar results: the prevalence ratio for hypertension was 1.21 (95% CI, 0.91–1.61). Results for SBP and DBP among patients with normotension using these thresholds are presented in Table S5.

Exploratory Analyses for Individual Nitrate-Reducing Bacterial Species

Upon examination of the associations between the individual nitrate-reducing bacterial taxa and cardiometabolic outcomes, a few significant associations were found (Tables S6 and S7). Higher relative abundance of *Neisseria flavescens* and

Table 2. Mean Difference in SBP and DBP for Every 1-SD Increase in NO₃TSS in the Full Sample and in Patients Without Hypertension

Model	All Patients (N=281)		Patients With Normotension (n=187)	
	SBP, mm Hg	DBP, mm Hg	SBP, mm Hg	DBP, mm Hg
	NO ₃ TSS (1 SD)	NO ₃ TSS (1 SD)	NO ₃ TSS (1 SD)	NO ₃ TSS (1 SD)
1	-1.25 (-2.68 to 0.18)	-0.59 (-1.71 to 0.52)	-1.70 (-3.00 to -0.40)	-0.69 (-1.58 to 0.19)
2	-0.87 (-2.22 to 0.49)	-0.31 (-1.37 to 0.76)	-1.60 (-2.84 to -0.34)	-0.62 (-1.51 to 0.27)
3	-0.76 (-2.07 to 0.55)	-0.23 (-1.28 to 0.81)	-1.51 (-2.74 to -0.27)	-0.60 (-1.49 to 0.29)
4	-0.81 (-2.12 to 0.50)	-0.26 (-1.31 to 0.79)	-1.51 (-2.74 to -0.27)	-0.60 (-1.49 to 0.29)
5	-0.66 (-2.04 to 0.72)	-0.12 (-1.22 to 0.98)	-1.53 (-2.82 to -0.24)	-0.60 (-1.54 to 0.33)
6	-0.74 (-2.17 to 0.69)	-0.35 (-1.49 to 0.79)	-1.85 (-3.10 to -0.61)	-0.69 (-1.63 to 0.25)

Model 1: crude (n=281; n=187 for without hypertension). Model 2: adjusted for age, sex, race, and education (n=281; n=187). Model 3: adjusted for age, sex, race, education, body mass index (BMI), and smoking (n=281; n=187). Model 4: adjusted for age, sex, race, education, BMI, smoking, and percentage of periodontal sites with attachment loss ≥ 3 mm (n=280; n=187). Model 5: adjusted for age, sex, race, education, BMI, smoking, percentage of periodontal sites with attachment loss ≥ 3 mm, and dietary pattern (n=253; n=170). Model 6: adjusted for age, sex, race, education, BMI, smoking, percentage of periodontal sites with attachment loss ≥ 3 mm, dietary pattern, alcohol use, and physical activity (n=226; n=155). DBP indicates diastolic blood pressure; NO₃TSS indicates nitrate-reducing taxa summary score; SBP, systolic blood pressure.

Haemophilus parainfluenzae were associated with lower insulin resistance and mean SBP, and lower plasma glucose, SBP, and DBP, respectively. While other taxa such as *Actinomyces naeslundii*, *Actinomyces viscosus*, *Capnocytophaga sputigena*, and *Neisseria sicca* also had some individual associations with lower cardiometabolic outcomes, no bacterial species was consistently associated across the different cardiometabolic outcomes.

Discussion

Among a sample of young diabetes mellitus-free individuals, we found higher relative abundance of nitrate-reducing oral bacteria to be associated with lower insulin resistance and plasma glucose in all participants, and with mean SBP in participants with normotension only, cross-sectionally. These results inform the potential influence of oral bacteria on cardiometabolic outcomes via the enterosalivary nitrate-nitrite-NO pathway of NO generation, and add important knowledge to the nascent literature in this area in a number of meaningful ways.

Unlike previous studies, this study directly examines a broad set of putative nitrate-reducing organisms in relation to cardiometabolic outcomes in a population-based observational setting. Few studies have directly measured the oral microbiota when examining the enterosalivary pathway of NO generation with health outcomes.^{20,21} Of these, none have examined the outcomes of insulin resistance and plasma glucose, and our study utilizes the largest sample size to date. Our findings demonstrate that a meaningful proportion ($\approx 20\%$) of oral taxa are potentially nitrate-reducing, while also showing substantial between-person variation in the relative abundance of nitrate-reducing bacteria. Moreover, the

results suggest that higher levels of nitrate-reducing organisms might confer health benefits across the population distribution of bacterial levels. Thus, if this relationship were causal, interventions to manipulate nitrate-reducing bacterial levels may be a useful treatment modality to improve cardiometabolic health, even in younger, generally healthy populations. The difference of ≈ 3 to 6 mm Hg in mean SBP observed between the highest and lowest SD intervals of NO₃TSS in our study is comparable to the estimated effects (5.7 mm Hg for SBP and 3.1 mm Hg for DBP) of first-line antihypertensive medications.³⁷ Importantly, a 4.4-mm Hg reduction in SBP has been estimated to reduce the risk of cardiovascular events by as much as 14%.³⁷ The smaller differences in insulin resistance and fasting plasma glucose currently observed between SD intervals of nitrate-reducing bacteria have not previously been associated with an increased conversion to overt diabetes mellitus or cardiovascular disease incidence. But future studies with longer follow-up times that allow for the development of greater impairment of glucose regulation may yield greater clinical relevance.

The lack of an association found between NO₃TSS and prediabetes mellitus or hypertension is inconsistent with the findings for insulin resistance, glucose, and BP in our study. It is possible that nitrate-reducing bacteria may be most relevant in the early preclinical stages of disease development, before more advanced pathophysiological alterations (eg, reduced β -cell function or increased arterial stiffness) occur and environmental risk factors lose importance. Furthermore, behavioral changes (eg, improved diet and activity levels) or medical therapies following the diagnosis of hypertension, prediabetes mellitus, or other comorbidities (eg, high cholesterol) could favorably influence both the nitrate-reducing bacteria and cardiometabolic health, masking these

associations. This notion is supported by the observation that NO₃TSS was most strongly associated with SBP among individuals with normotension only. Alternatively, these observations may simply be the result of chance, and replication in future studies will be important.

Study Limitations

Some important limitations should be noted. Because of the cross-sectional design of our study, reverse causation is possible. Insulin resistance, plasma glucose, and BP levels could all influence microbial community composition, even in the clinically normal range. High salivary glucose is associated with a shift in the composition of the oral microbiome, although the direct influence on nitrate-reducing bacteria is unknown and studies have mostly considered only glucose levels in the diabetic range.³⁸

Measurement error in the assessment of nitrate-reducing bacteria might have diluted the strength of association as the day-to-day stability of the oral microbiome is unclear.³⁹ ORIGINS only measured 20 of the 28 bacterial species previously identified as being associated with nitrate-reduction capacity. Since there are many more oral bacteria with nitrate-reducing capacity, it is also likely that not all relevant nitrate-reducing bacteria have been identified, as only 2 studies have sought to identify the key contributors to oral nitrate-reduction.^{25,26} Additionally, strain-level variation within the same species, horizontal gene transfer between bacteria,⁴⁰ and different rates of nitrate-reduction across taxa^{25,26} may all result in the misclassification of the individual's nitrate-reducing capacity. Metagenomics to directly assess the genes (eg, *narG*, *narL*, *napC*, *napB*) encoding for the nitrate-reductase produced by bacteria,²⁶ or metatranscriptomics measuring gene expression, may better capture the nitrate-reducing capacity. Likewise, including a measure of plasma or salivary nitrite together with the bacteria measures can further support the increase in nitrite production through these bacteria and may be useful for mediation analyses. Although the amount of salivary nitrate reduced to nitrite by oral bacteria appears to be substantial and dose-dependent,^{19,21} the percentage that reaches the systemic circulation as plasma nitrite is unclear but appears to be less, and a possible saturation threshold has been suggested that needs further examination.^{17,21} Nevertheless, misclassification of the nitrate-reducing bacteria exposure is likely nondifferential by the outcomes, biasing the estimates towards the null.

Nitrate-reducing bacteria are present in various sites of the oral cavity. Our study was only able to examine subgingival bacteria when the tongue is believed to be the main site of bacterial nitrate reduction in the mouth.²⁵ The oral cavity is thought to contain distinct niches of microbiota with varying microbial diversity and composition,⁴¹ and it is unknown

whether levels of nitrate-reducing bacteria in the subgingival plaque serve as a reasonable proxy for levels on the tongue. The assessment of subgingival microbiota from one periodontal site per participant, as in this study, is also highly likely to have increased measurement error of the full-mouth exposure to nitrate-reducing taxa, which would bias the results towards the null. Future studies that can directly assess nitrate-reducing bacteria on the tongue will be important.

Our results also do not account for pathways involving the gut. As contiguous parts of the digestive tract, the gut, and oral microbiota likely influence one another.⁴² The gut microbiome is also capable of nitrate reduction, although its contribution to circulating nitrite is likely minor, as the main site of nitrate to nitrite reduction occurs in the mouth, and dietary nitrate is mostly absorbed from the proximal intestine into the circulation, bypassing the nitrate-reducing bacteria residing more distally. However, gut bacteria (eg, *Lactobacilli* sp., *Bifidobacterium*) can directly produce NO, potentially influencing blood flow and mucus generation, and thus the uptake and bioavailability of nitrate and nitrite.¹⁰ Gastric pH is also of relevance with high levels of NO formed nonenzymatically in the acidic stomach from swallowed nitrite and acid-reducing medication shown to attenuate the BP-lowering effects of nitrate.⁴³ Furthermore, the gut microbiome produces hydrogen sulfide (H₂S), another physiologically important gaseous signaling molecule, involved in the formation of NO from nitrite in the intestine and systemic tissues.¹⁰ Evidence suggests that an interplay of H₂S and NO has cardiovascular effects.⁴⁴ Thus, the gut microbiome may modify nitrite and NO bioavailability, and more research is needed to fully understand the role of the gut microbiome in the nitrate-nitrite-NO enterosalivary pathway.

Future studies that consider the role of oral nitrite reduction will also be important. Oral bacteria can further reduce salivary nitrite to NO, influencing the amount of bioavailable salivary nitrite swallowed and absorbed into the systemic circulation.^{25,26} Thus, the optimal oral bacterial community for NO generation may be one that allows for nitrite accumulation, and the ratio of the nitrate versus nitrite-reducing capacity of the oral microbiome, the exposure of greatest interest.^{26,45} Furthermore, a correlation between higher bacterial nitrite-reductase gene abundance and lower resting SBP was recently found, suggesting that orally produced NO may have systemic effects on vasodilation as well.⁴⁵ However, few other studies have specifically explored the nitrite-reducing capacity of the oral cavity, and to the best of our knowledge the key bacterial species contributing to nitrite reduction in the mouth have yet to be identified.

Finally, although frequent regular mouthwash use has been associated with an increased risk of prediabetes mellitus,⁴⁶ the ORIGINS questionnaire did not contain information on mouthwash use and we were unable to control for mouthwash use.

Study Strengths

Despite these limitations, several strengths should be noted. ORIGINS collected a robust set of risk factor data allowing for comprehensive control for hypothesized confounders. The use of next-generation sequencing techniques allowed for more precise identification of a larger set of nitrate-reducing bacteria, and the relatively young cohort, free of diabetes mellitus and other clinical cardiovascular diseases, minimizes reverse causality.

Conclusions

This is one of the first studies to directly test the hypothesis of a priori-identified nitrate-reducing oral bacteria affecting cardiometabolic outcomes. Our results support the hypothesis that oral nitrate-reducing bacteria play a beneficial role in BP regulation and insulin resistance. Future longitudinal studies with enhanced assessment of nitrate-reducing bacterial exposure predicting progression of cardiometabolic risk biomarkers and incident clinical disease will improve temporal inference necessary to inform causality and inform the development of future intervention studies that could manipulate oral nitrate-reducing capacity.

Acknowledgments

We thank the following individuals for their invaluable contributions to this research: Consuelo McLaughlin, Bennett Batista, and Victor Rivera for their assistance with recruitment; Romanita Celenti for her efforts in performing phlebotomy and processing and analyzing plaque samples; Aleksandra Zuk and Garazi Zulaika for their leadership and excellent program coordination; and Drs Nidhi Arora, Ashwata Pokherel, Publio Silfa, and Thomas Spinell for their skilled examinations and essential participant engagement. We are also profoundly grateful to the ORIGINS participants, for their participation in this research.

Sources of Funding

This research was supported by National Institutes of Health (NIH) grants R00 DE018739, R21 DE022422, and R01 DK 102932 to Dr Demmer. This publication was also supported by the National Center for Advancing Translational Sciences, NIH, through grant number UL1TR001873. The content is solely the responsibility of the authors and does not necessarily represent the official views of the NIH.

Disclosures

None.

References

1. Le Chatelier E, Nielsen T, Qin J, Prifti E, Hildebrand F, Falony G, Almeida M, Arumugam M, Batto JM, Kennedy S, Leonard P, Li J, Burgdorf K, Grarup N,

- Jorgensen T, Brandslund I, Nielsen HB, Juncker AS, Bertalan M, Levenez F, Pons N, Rasmussen S, Sunagawa S, Tap J, Tims S, Zoetendal EG, Brunak S, Clement K, Dore J, Kleerebezem M, Kristiansen K, Renault P, Sicheritz-Ponten T, de Vos WM, Zucker JD, Raes J, Hansen T, Meta HITc, Bork P, Wang J, Ehrlich SD, Pedersen O. Richness of human gut microbiome correlates with metabolic markers. *Nature*. 2013;500:541–546.
2. Demmer RT, Breskin A, Rosenbaum M, Zuk A, LeDuc C, Leibel R, Paster B, Desvarieux M, Jacobs DR Jr, Papapanou PN. The subgingival microbiome, systemic inflammation and insulin resistance: the Oral Infections, Glucose Intolerance and Insulin Resistance Study (ORIGINS). *J Clin Periodontol*. 2017;44:255–265.
3. Tai N, Wong FS, Wen L. The role of gut microbiota in the development of type 1, type 2 diabetes mellitus and obesity. *Rev Endocr Metab Disord*. 2015;16:55–65.
4. Demmer RT, Jacobs DR Jr, Singh R, Zuk A, Rosenbaum M, Papapanou PN, Desvarieux M. Periodontal bacteria and prediabetes prevalence in ORIGINS: the Oral Infections, Glucose Intolerance, and Insulin Resistance Study. *J Dent Res*. 2015;94:201s–211s.
5. Pevsner-Fischer M, Blacher E, Tatrovsky E, Ben-Dov IZ, Elinav E. The gut microbiome and hypertension. *Curr Opin Nephrol Hypertens*. 2017;26:1–8.
6. Desvarieux M, Demmer RT, Jacobs DR Jr, Rundek T, Boden-Albala B, Sacco RL, Papapanou PN. Periodontal bacteria and hypertension: the oral infections and vascular disease epidemiology study (INVEST). *J Hypertens*. 2010;28:1413–1421.
7. Hajishengallis G. Periodontitis: from microbial immune subversion to systemic inflammation. *Nat Rev Immunol*. 2015;15:30–44.
8. Lundberg JO, Weitzberg E, Gladwin MT. The nitrate-nitrite-nitric oxide pathway in physiology and therapeutics. *Nat Rev Drug Discov*. 2008;7:156–167.
9. Sansbury BE, Hill BG. Regulation of obesity and insulin resistance by nitric oxide. *Free Radic Biol Med*. 2014;73:383–399.
10. Koch CD, Gladwin MT, Freeman BA, Lundberg JO, Weitzberg E, Morris A. Enterosalivary nitrate metabolism and the microbiome: intersection of microbial metabolism, nitric oxide and diet in cardiac and pulmonary vascular health. *Free Radic Biol Med*. 2017;105:48–67.
11. Ashor AW, Lara J, Siervo M. Medium-term effects of dietary nitrate supplementation on systolic and diastolic blood pressure in adults: a systematic review and meta-analysis. *J Hypertens*. 2017;35:1353–1359.
12. Jiang H, Torregrossa AC, Potts A, Pierini D, Aranke M, Garg HK, Bryan NS. Dietary nitrite improves insulin signaling through GLUT4 translocation. *Free Radic Biol Med*. 2014;67:51–57.
13. Carlström M, Larsen FJ, Nyström T, Hezel M, Borniquel S, Weitzberg E, Lundberg JO. Dietary inorganic nitrate reverses features of metabolic syndrome in endothelial nitric oxide synthase-deficient mice. *Proc Natl Acad Sci USA*. 2010;107:17716–17720.
14. Beals JW, Binns SE, Davis JL, Giordano GR, Klochak AL, Paris HL, Schweder MM, Peltonen GL, Scalzo RL, Bell C. Concurrent beet juice and carbohydrate ingestion: influence on glucose tolerance in obese and nonobese adults. *J Nutr Metab*. 2017;2017:6436783.
15. Lidder S, Hunt J, Omar S, Webb A. Acute effects of dietary nitrate on glucose handling and insulin levels during an oral glucose tolerance test in healthy subjects [Abstract]. *Nitric Oxide*. 2011;24:S32.
16. Ashor AW, Chowdhury S, Oggioni C, Qadir O, Brandt K, Ishaq A, Mathers JC, Saretzki G, Siervo M. Inorganic nitrate supplementation in young and old obese adults does not affect acute glucose and insulin responses but lowers oxidative stress. *J Nutr*. 2016;146:2224–2232.
17. Govoni M, Jansson EA, Weitzberg E, Lundberg JO. The increase in plasma nitrite after a dietary nitrate load is markedly attenuated by an antibacterial mouthwash. *Nitric Oxide*. 2008;19:333–337.
18. Woessner M, Smoliga JM, Tarzia B, Stabler T, Van Bruggen M, Allen JD. A stepwise reduction in plasma and salivary nitrite with increasing strengths of mouthwash following a dietary nitrate load. *Nitric Oxide*. 2016;54:1–7.
19. Kapil V, Haydar SM, Pearl V, Lundberg JO, Weitzberg E, Ahluwalia A. Physiological role for nitrate-reducing oral bacteria in blood pressure control. *Free Radic Biol Med*. 2013;55:93–100.
20. Vanhatalo A, Blackwell JR, L'Heureux JE, Williams DW, Smith A, van der Giezen M, Winyard PG, Kelly J, Jones AM. Nitrate-responsive oral microbiome modulates nitric oxide homeostasis and blood pressure in humans. *Free Radic Biol Med*. 2018;124:21–30.
21. Burleigh MC, Liddle L, Monaghan C, Mugeridge DJ, Sculthorpe N, Butcher JP, Henriquez FL, Allen JD, Easton C. Salivary nitrite production is elevated in individuals with a higher abundance of oral nitrate-reducing bacteria. *Free Radic Biol Med*. 2018;120:80–88.
22. The Forsyth Institute. Human oral microbe identification using next generation sequencing (HOMINGS). Published February 2008. Available at: <http://homings.forsyth.org/sampleprep.html>. Accessed December 29, 2017.

23. Gomes BP, Berber VB, Kokaras AS, Chen T, Paster BJ. Microbiomes of endodontic-periodontal lesions before and after chemomechanical preparation. *J Endod*. 2015;41:1975–1984.
24. Mougeot JL, Stevens CB, Cotton SL, Morton DS, Krishnan K, Brennan MT, Lockhart PB, Paster BJ, Bahrani Mougeot FK. Concordance of HOMIM and HOMINGS technologies in the microbiome analysis of clinical samples. *J Oral Microbiol*. 2016;8:30379.
25. Doel JJ, Benjamin N, Hector MP, Rogers M, Allaker RP. Evaluation of bacterial nitrate reduction in the human oral cavity. *Eur J Oral Sci*. 2005;113:14–19.
26. Hyde ER, Andrade F, Vaksman Z, Parthasarathy K, Jiang H, Parthasarathy DK, Torregrossa AC, Tribble G, Kaplan HB, Petrosino JF, Bryan NS. Metagenomic analysis of nitrate-reducing bacteria in the oral cavity: implications for nitric oxide homeostasis. *PLoS One*. 2014;9:e88645.
27. Palmer RJ, Cotton SL, Kokaras AS, Gardner P, Grisius M, Pelayo E, Warner B, Paster BJ, Alevizos I. Analysis of oral bacterial communities: comparison of HOMINGS with a tree-based approach implemented in QIIME. *J Oral Microbiol*. 2019;11:1586413.
28. Morgan XC, Kabakchiev B, Waldron L, Tyler AD, Tickle TL, Milgrom R, Stempak JM, Gevers D, Xavier RJ, Silverberg MS, Huttenhower C. Associations between host gene expression, the mucosal microbiome, and clinical outcome in the pelvic pouch of patients with inflammatory bowel disease. *Genome Biol*. 2015;16:67.
29. Matthews DR, Hosker JP, Rudenski AS, Naylor BA, Treacher DF, Turner RC. Homeostasis model assessment: insulin resistance and beta-cell function from fasting plasma glucose and insulin concentrations in man. *Diabetologia*. 1985;28:412–419.
30. American Diabetes Association. 2. Classification and diagnosis of diabetes. *Diabetes Care*. 2017;40:S11–S24.
31. Whelton PK, Carey RM, Aronow WS, Casey DE, Collins KJ, Himmelfarb CD, DePalma SM, Gidding S, Jamerson KA, Jones DW, MacLaughlin EJ, Muntner P, Ovbigele B, Smith SC, Spencer CC, Stafford RS, Taler SJ, Thomas RJ, Williams KA, Williamson JD, Wright JT. 2017 ACC/AHA/AAPA/ABC/ACPM/AGS/APhA/ASH/ASPC/NMA/PCNA guideline for the prevention, detection, evaluation, and management of high blood pressure in adults: a report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines. *Hypertension*. 2018;71:e13–e115.
32. Thai A, Papapanou PN, Jacobs DR Jr, Desvarieux M, Demmer RT. Periodontal infection and cardiorespiratory fitness in younger adults: results from continuous National Health and Nutrition Examination Survey 1999–2004. *PLoS One*. 2014;9:e92441.
33. Chiuve SE, Fung TT, Rimm EB, Hu FB, McCullough ML, Wang M, Stampfer MJ, Willett WC. Alternative dietary indices both strongly predict risk of chronic disease. *J Nutr*. 2012;142:1009–1018.
34. Zou G. A modified Poisson regression approach to prospective studies with binary data. *Am J Epidemiol*. 2004;159:702–706.
35. Eke PI, Page RC, Wei L, Thornton-Evans G, Genco RJ. Update of the case definitions for population-based surveillance of periodontitis. *J Periodontol*. 2012;83:1449–1454.
36. Chobanian AV, Bakris GL, Black HR, Cushman WC, Green LA, Izzo J, Joseph L, Jones DW, Materson BJ, Oparil S, Wright J, Jackson T, Roccella EJ; National Heart, Lung, and Blood Institute Joint National Committee on Prevention, Detection, Evaluation, and Treatment of High Blood Pressure; National High Blood Pressure Education Program Coordinating Committee. The seventh report of the Joint National Committee on Prevention, Detection, Evaluation, and Treatment of High Blood Pressure: the JNC 7 report. *JAMA*. 2003;289:2560–2571.
37. Law MR, Morris JK, Wald NJ. Use of blood pressure lowering drugs in the prevention of cardiovascular disease: meta-analysis of 147 randomised trials in the context of expectations from prospective epidemiological studies. *BMJ*. 2009;338:b1665.
38. Goodson JM, Hartman ML, Shi P, Hasturk H, Yaskell T, Vargas J, Song X, Cugini M, Barake R, Alsmadi O, Al-Mutawa S, Ariga J, Soparkar P, Behbehani J, Behbehani K. The salivary microbiome is altered in the presence of a high salivary glucose concentration. *PLoS One*. 2017;12:e0170437.
39. Vogtmann E, Hua X, Zhou L, Wan Y, Suman S, Zhu B, Dagnall CL, Hutchinson A, Jones K, Hicks BD, Sinha R, Shi J, Abnet CC. Temporal variability of oral microbiota over 10 months and the implications for future epidemiologic studies. *Cancer Epidemiol Biomarkers Prev*. 2018;27:594–600.
40. Roberts AP, Kreth J. The impact of horizontal gene transfer on the adaptive ability of the human oral microbiome. *Front Cell Infect Microbiol*. 2014;4:124.
41. Xu X, He J, Xue J, Wang Y, Li K, Zhang K, Guo Q, Liu X, Zhou Y, Cheng L, Li M, Li Y, Li Y, Shi W, Zhou X. Oral cavity contains distinct niches with dynamic microbial communities. *Environ Microbiol*. 2015;17:699–710.
42. Ding T, Schloss PD. Dynamics and associations of microbial community types across the human body. *Nature*. 2014;509:357–360.
43. Montenegro MF, Sundqvist ML, Larsen FJ, Zhuge Z, Carlstrom M, Weitzberg E, Lundberg JO. Blood pressure-lowering effect of orally ingested nitrite is abolished by a proton pump inhibitor. *Hypertension*. 2017;69:23–31.
44. Nagpure BV, Bian JS. Interaction of hydrogen sulfide with nitric oxide in the cardiovascular system. *Oxid Med Cell Longev*. 2016;2016:6904327.
45. Tribble GD, Angelov N, Weltman R, Wang BY, Eswaran SV, Gay IC, Parthasarathy K, Dao DH, Richardson KN, Ismail NM, Sharina IG, Hyde ER, Ajami NJ, Petrosino JF, Bryan NS. Frequency of tongue cleaning impacts the human tongue microbiome composition and enterosalivary circulation of nitrate. *Front Cell Infect Microbiol*. 2019;9:39.
46. Joshipura KJ, Munoz-Torres FJ, Morou-Bermudez E, Patel RP. Over-the-counter mouthwash use and risk of pre-diabetes/diabetes. *Nitric Oxide*. 2017;71:14–20.

SUPPLEMENTAL MATERIALS

Supplemental Methods

Information on the potential confounders age, sex, race (non-Hispanic Black, non-Hispanic White, Hispanic, Other), education (high school completion, college or vocational training, advanced degrees), cigarette smoking status (current, former or never smoking), leisure-time physical activity, dietary patterns, and alcohol use (non-drinker, ≤ 1 drink/day, > 1 drink/day) was obtained through detailed risk factor survey questionnaires.

Physical activity: Frequency and intensity of physical activity assessed in the questionnaire were operationalized by calculating metabolic equivalents (METs)¹. Total METs per week were calculated $[(\text{Number of times engaged in the activity in past 30 days} * \text{Average duration of activity} * \text{MET score for activity})/30 \text{ days}] * 7 \text{ days}$, and categorized into 4 categories according to the 2008 Physical Activity Guidelines for Americans: no physical activity reported, low (0 to <500 MET min/week), moderate (500 to $<1,000$ MET min/week), or high ($\geq 1,000$ MET min/week)².

Alcohol consumption: Alcohol use was operationalized from the questionnaires as average number of drinks/day and categorized into 3 categories (non-drinker, ≤ 1 drink/day, > 1 drink/day).

Body mass index (BMI): BMI was calculated as weight (in kilograms)/height (meters²) obtained from in-person physical assessments and operationalized as a continuous measure, as categories of BMI (underweight, normal, overweight, obese) showed a linear relationship with the outcomes.

Periodontal status: Previous methodological studies suggest that clinical periodontal metrics such as percent of probing sites with attachment loss ≥ 3 mm relate better to extra-oral disease than the periodontitis diagnosis classifications^{3,4}. This is especially so in healthy populations where low threshold periodontal parameters correlate more closely with bacteria exposure than more severe periodontal measures⁵. In addition, the percent of probing sites with attachment loss ≥ 3 mm is a more nuanced continuous measure of disease, with variation even in mild periodontal disease. Sensitivity analyses using Centers for Disease Control and Prevention/American Academy of Periodontology (CDC/AAP) diagnosis classification (none/mild vs. moderate/severe)⁶ showed no appreciable difference from our main results.

REFERENCES

1. Ainsworth BE, Haskell WL, Leon AS, Jacobs DR, Jr., Montoye HJ, Sallis JF, Paffenbarger RS, Jr. Compendium of physical activities: Classification of energy costs of human physical activities. *Med Sci Sports Exerc.* 1993;25:71-80
2. U.S. Department of Health and Human Services. 2008 physical activity guidelines for americans. 2008;2017
3. Beck JD, Offenbacher S. Relationships among clinical measures of periodontal disease and their associations with systemic markers. *Ann Periodontol.* 2002;7:79-89
4. Demmer RT, Kocher T, Schwahn C, Volzke H, Jacobs DR, Jr., Desvarieux M. Refining exposure definitions for studies of periodontal disease and systemic disease associations. *Community Dent Oral Epidemiol.* 2008;36:493-502
5. Demmer RT, Papapanou PN, Jacobs DR, Jr., Desvarieux M. Evaluating clinical periodontal measures as surrogates for bacterial exposure: The oral infections and vascular disease epidemiology study (invest). *BMC Med Res Methodol.* 2010;10:2
6. Eke PI, Page RC, Wei L, Thornton-Evans G, Genco RJ. Update of the case definitions for population-based surveillance of periodontitis. *J Periodontol.* 2012;83:1449-1454

Figure S1. Spearman correlations of the relative abundance levels for 20 nitrate-reducing subgingival bacterial species (n=281)

	<i>A.n</i>	<i>A.o</i>	<i>A.v</i>	<i>C.s</i>	<i>C.d</i>	<i>C.m</i>	<i>E.c</i>	<i>H.p</i>	<i>N.f</i>	<i>N.s</i>	<i>N.subflava</i>	<i>P.m</i>	<i>P.s</i>	<i>P.a</i>	<i>R.d</i>	<i>R.m</i>	<i>S.n</i>	<i>V.d</i>	<i>V.p</i>	<i>V.a</i>
<i>Actinomyces naeslundii</i>	1.00	0.02	0.20	0.19	0.31	0.44	0.16	0.07	0.11	0.04	0.04	-0.01	-0.01	-0.02	0.08	-0.06	0.24	0.04	0.02	-0.06
<i>Actinomyces odontolyticus</i>	0.02	1.00	0.05	0.06	0.08	0.01	0.02	0.25	0.15	0.06	0.03	0.28	0.14	-0.06	0.02	0.37	-0.15	0.09	-0.05	0.29
<i>Actinomyces viscosus</i>	0.20	0.05	1.00	-0.08	0.16	-0.06	-0.15	0.04	-0.04	-0.08	0.02	0.02	-0.05	0.06	0.39	0.17	-0.13	0.24	0.21	-0.03
<i>Capnocytophaga sputigena</i>	0.19	0.06	-0.08	1.00	0.27	0.24	0.45	0.22	0.21	0.27	0.04	0.09	-0.02	0.02	-0.07	-0.07	0.21	0.03	-0.02	-0.09
<i>Corynebacterium durum</i>	0.31	0.08	0.16	0.27	1.00	0.22	0.05	0.26	0.25	0.08	0.01	-0.01	-0.05	0.06	0.36	0.05	-0.01	-0.03	-0.08	-0.06
<i>Corynebacterium matruchotii</i>	0.44	0.01	-0.06	0.24	0.22	1.00	0.34	-0.06	0.10	-0.04	0.01	0.07	0.00	-0.11	-0.09	-0.22	0.43	0.10	0.07	-0.09
<i>Eikenella corrodens</i>	0.16	0.02	-0.15	0.45	0.05	0.34	1.00	0.07	0.18	0.18	0.06	-0.02	0.04	-0.04	-0.17	-0.20	0.32	-0.01	-0.05	-0.14
<i>Haemophilus parainfluenzae</i>	0.07	0.25	0.04	0.22	0.26	-0.06	0.07	1.00	0.36	0.23	0.15	0.25	-0.02	0.13	0.11	0.40	-0.17	0.13	0.05	0.18
<i>Neisseria flavescens</i>	0.11	0.15	-0.04	0.21	0.25	0.10	0.18	0.36	1.00	0.14	0.03	0.09	0.05	0.09	0.04	0.23	0.01	-0.07	-0.07	0.09
<i>Neisseria sicca</i>	0.04	0.06	-0.08	0.27	0.08	-0.04	0.18	0.23	0.14	1.00	0.10	0.03	-0.09	0.00	-0.08	0.03	-0.08	0.02	-0.01	-0.05
<i>Neisseria subflava</i>	0.04	0.03	0.02	0.04	0.01	0.01	0.06	0.15	0.03	0.10	1.00	-0.07	-0.09	-0.01	0.03	-0.02	-0.03	0.05	-0.02	-0.04
<i>Prevotella melaninogenica</i>	-0.01	0.28	0.02	0.09	-0.01	0.07	-0.02	0.25	0.09	0.03	-0.07	1.00	0.36	0.01	-0.06	0.27	0.12	0.33	0.25	0.45
<i>Prevotella salivae</i>	-0.01	0.14	-0.05	-0.02	-0.05	0.00	0.04	-0.02	0.05	-0.09	-0.09	0.36	1.00	-0.03	-0.04	0.17	0.19	0.09	0.08	0.38
<i>Propionibacterium acnes</i>	-0.02	-0.06	0.06	0.02	0.06	-0.11	-0.04	0.13	0.09	0.00	-0.01	0.01	-0.03	1.00	0.06	0.16	-0.08	-0.04	0.00	0.04
<i>Rothia dentocariosa</i>	0.08	0.02	0.39	-0.07	0.36	-0.09	-0.17	0.11	0.04	-0.08	0.03	-0.06	-0.04	0.06	1.00	0.20	-0.23	0.10	0.08	0.04
<i>Rothia mucilaginosa</i>	-0.06	0.37	0.17	-0.07	0.05	-0.22	-0.20	0.40	0.23	0.03	-0.02	0.27	0.17	0.16	0.20	1.00	-0.24	0.03	0.03	0.38
<i>Selenomonas noxia</i>	0.24	-0.15	-0.13	0.21	-0.01	0.43	0.32	-0.17	0.01	-0.08	-0.03	0.12	0.19	-0.08	-0.23	-0.24	1.00	0.12	0.14	-0.03
<i>Veillonella dispar</i>	0.04	0.09	0.24	0.03	-0.03	0.10	-0.01	0.13	-0.07	0.02	0.05	0.33	0.09	-0.04	0.10	0.03	0.12	1.00	0.61	0.24
<i>Veillonella parvula</i>	0.02	-0.05	0.21	-0.02	-0.08	0.07	-0.05	0.05	-0.07	-0.01	-0.02	0.25	0.08	0.00	0.08	0.03	0.14	0.61	1.00	0.22
<i>Veillonella atypica</i>	-0.06	0.29	-0.03	-0.09	-0.06	-0.09	-0.14	0.18	0.09	-0.05	-0.04	0.45	0.38	0.04	0.04	0.38	-0.03	0.24	0.22	1.00

Correlations with a $p < 0.05$ in italics.

Actinomyces naeslundii (*A.n*); *Actinomyces odontolyticus* (*A.o*); *Actinomyces viscosus* (*A.v*); *Capnocytophaga sputigena* (*C.s*); *Corynebacterium durum* (*C.d*); *Corynebacterium matruchotii* (*C.m*); *Eikenella corrodens* (*E.c*); *Haemophilus parainfluenzae* (*H.p*); *Neisseria flavescens* (*N.f*); *Neisseria sicca* (*N.s*); *Neisseria subflava* (*N.subflava*); *Prevotella melaninogenica* (*P.m*); *Prevotella salivae* (*P.s*); *Propionibacterium acnes* (*P.a*); *Rothia dentocariosa* (*R.d*); *Rothia mucilaginosa* (*R.m*); *Selenomonas noxia* (*S.n*); *Veillonella dispar* (*V.d*); *Veillonella parvula* (*V.p*); *Veillonella atypica* (*V.a*).

Table S1. The list of previously identified oral bacterial species or genera with potential nitrate-reducing capacity.

Genera ²	Species
<i>Actinomyces</i>	<i>Actinomyces naeslundii</i> ¹
<i>Brevibacillus</i>	<i>Actinomyces odontolyticus</i> ^{1,2}
<i>Fusobacterium</i>	<i>Actinomyces oris/Actinomyces naeslundii</i> <i>genospecies-2</i> ²
<i>Granulicatella</i>	<i>Actinomyces viscius</i> ^{1,2}
<i>Haemophilus</i>	<i>Brevibacillus brevis/ Bacillus brevis</i> ²
<i>Leptotrichia</i>	<i>Capnocytophaga sputigena</i> ¹
<i>Neisseria</i>	<i>Corynebacterium durum</i> ¹
<i>Porphyromonas</i>	<i>Corynebacterium matruchotii</i> ¹
<i>Prevotella</i>	<i>Eikenella corrodens</i> ¹
<i>Veillonella</i>	<i>Granulicatella adiacens</i> ^{1,2}
Unclassified genus of <i>Gemellaceae</i> family	<i>Haemophilus parainfluenzae</i> ^{1,2}
	<i>Haemophilus segnis</i> ¹
	<i>Microbacterium oxydans</i> ¹
	<i>Neisseria flavescens</i> ²
	<i>Neisseria mucosa</i> ²
	<i>Neisseria sicca</i> ²
	<i>Neisseria subflava</i> ²
	<i>Prevotella melaninogenica</i> ²
	<i>Prevotella salivae</i> ²
	<i>Propionibacterium acnes</i> ¹
	<i>Rothia dentocariosa</i> ¹
	<i>Rothia mucilaginosa</i> ¹
	<i>Staphylococcus epidermidis</i> ¹
	<i>Staphylococcus hemolyticus</i> ¹
	<i>Selenomonas noxia</i> ¹
	<i>Veillonella dispar</i> ^{1,2}
	<i>Veillonella parvula</i> ²
	<i>Veillonella atypica</i> ^{1,2}

¹ Doel JJ, Benjamin N, Hector MP, Rogers M, Allaker RP. Evaluation of bacterial nitrate reduction in the human oral cavity. *European journal of oral sciences*. 2005;113:14-19.

² Hyde ER, Andrade F, Vaksman Z, Parthasarathy K, Jiang H, Parthasarathy DK, Torregrossa AC, Tribble G, Kaplan HB, Petrosino JF, others. Metagenomic analysis of nitrate-reducing bacteria in the oral cavity: Implications for nitric oxide homeostasis. *PLoS One*. 2014;9:e88645.

Table S2. Spearman correlations between the mean relative abundance for each of the 20 nitrate-reducing taxa and summary scores with (i.e. NO₃TSS) or without standardization.

Taxa	Mean Relative abundance	Correlation coefficient (p-value) Summary score with standardization	Correlation coefficient (p-value) Summary score without standardization
<i>Rothia dentocariosa</i>	7.91%	0.16 (<0.001)	0.63 (<0.001)
<i>Corynebacterium matruchotii</i>	4.45%	0.24 (<0.001)	0.18 (0.003)
<i>Veillonella parvula</i>	2.46%	0.32 (<0.001)	0.22 (<0.001)
<i>Haemophilus parainfluenzae</i>	1.35%	0.46(<0.001)	0.25 (<0.001)
<i>Corynebacterium durum</i>	0.96%	0.32 (<0.001)	0.33 (<0.001)
<i>Actinomyces naeslundii</i>	0.79%	0.32 (<0.001)	0.17 (0.004)
<i>Rothia mucilaginosa</i>	0.60%	0.28 (<0.001)	0.15 (0.01)
<i>Veillonella atypica</i>	0.56%	0.35 (<0.001)	0.14 (0.02)
<i>Prevotella melaninogenica</i>	0.49%	0.48 (<0.001)	0.16 (0.006)
<i>Selenomonas noxia</i>	0.32%	0.20 (<0.001)	-0.07 (0.25)
<i>Neisseria flavescens</i>	0.26%	0.30 (<0.001)	0.09 (0.12)
<i>Capnocytophaga sputigena</i>	0.11%	0.34 (<0.001)	-0.02 (0.77)
<i>Eikenella corrodens</i>	0.09%	0.22 (<0.001)	-0.13 (0.02)
<i>Veillonella dispar</i>	0.03%	0.42 (<0.001)	0.31 (<0.001)
<i>Prevotella salivae</i>	0.02%	0.27 (<0.001)	-0.02 (0.72)
<i>Actinomyces odontolyticus</i>	0.01%	0.36 (<0.001)	0.15 (0.01)
<i>Actinomyces viscosus</i>	0.01%	0.29 (<0.001)	0.33 (<0.001)
<i>Neisseria subflava</i>	0.00%	0.14 (0.02)	0.04 (0.52)
<i>Neisseria sicca</i>	0.00%	0.21 (<0.001)	-0.06 (0.32)
<i>Propionibacterium acnes</i>	0.00%	0.12 (0.04)	-0.01 (0.90)

Taxa are listed in descending order of mean relative abundance.

The summary score without standardization was created by summing the relative abundances across the 20 individual nitrate-reducing bacteria for a total relative abundance of nitrate-reducing bacteria.

A summary score created without standardization of taxa is generally correlated only with the most abundant taxa. In contrast, the standardized summary score of nitrate-reducing bacteria (NO₃TSS) is more consistently correlated with the relative abundances of individual nitrate-reducing taxa.

Table S3. Associations between potential confounders and A) nitrate-reducing bacterial summary score (NO₃TSS), insulin resistance, B) fasting glucose, blood pressure, C) prediabetes and hypertension.

A)

Variable	Regression coefficient with NO ₃ TSS as dependent variable		Regression coefficient with insulin resistance as dependent variable	
	Crude Mean difference (95% CI)	<i>P</i> value	Crude Mean difference (95% CI)	<i>P</i> value
Age (5 years)	-0.11 (-0.43, 0.22)	0.52	0.06 (0.03, 0.09)	<0.001
Female	0.42 (-1.11, 1.96)	0.59	0.09 (-0.07, 0.25)	0.26
Race/ethnicity		0.26		<0.001
Hispanic	Ref	-	Ref	-
Non-Hispanic White	1.56 (-0.06, 3.18)	0.06	-0.44 (-0.61, -0.28)	<0.001
Black	0.001 (-1.76, 1.76)	1.00	-0.12 (-0.30, 0.05)	0.17
Other	-0.02 (-1.98, 1.94)	0.98	-0.24 (-0.44, -0.04)	0.02
Education		0.07		0.005
< Bachelor's Degree	Ref	-	Ref	-
Bachelor's Degree	1.62 (0.18, 3.06)	0.03	-0.19 (-0.33, -0.04)	0.02
> Bachelor's Degree	1.56 (-0.16, 3.29)	0.08	-0.28 (-0.46, -0.10)	0.002
Smoking Status		0.70		0.85
Never	Ref	-	Ref	-
Former	-0.09 (-2.07, 1.89)	0.93	-0.01 (-0.22, 0.19)	0.89
Current	-0.97 (-3.21, 1.26)	0.39	-0.07 (-0.30, 0.17)	0.57
BMI (kg/m²)	-0.07 (-0.17, 0.04)	0.21	0.04 (0.03, 0.05)	<0.001
BMI categories		0.51		<0.001
Underweight/Normal	Ref	-	Ref	-
Overweight	-0.43 (-1.89, 1.02)	0.56	0.35 (0.21, 0.48)	<0.001
Obese	-0.96 (-2.59, 0.67)	0.25	0.63 (0.47, 0.78)	<0.001
Percent of sites with attachment loss ≥3mm (10% increase)	-0.53 (-0.97, -0.09)	0.02	0.03 (-0.02, 0.08)	0.20
Average alcohol consumption		0.06		0.26
None	Ref	-	Ref	-
≤1 drink/day	1.06 (-1.59, 3.71)	0.43	-0.13 (-0.41, 0.15)	0.38
> 1 drink/day	3.09 (0.02, 6.17)	0.05	-0.26 (-0.58, 0.07)	0.12
Physical activity METS min/week		0.61		0.21
None	Ref	-	Ref	-
Low	0.59 (-1.55, 2.73)	0.59	0.05 (-0.17, 0.27)	0.67
Moderate	1.32 (-0.69, 3.32)	0.20	0.02 (-0.19, 0.23)	0.84
High	0.19 (-1.32, 1.70)	0.80	-0.13 (-0.29, 0.03)	0.11
Alternate Healthy Eating Index Score (AHEI)	0.04 (-0.02, 0.10)	0.19	-0.01 (-0.01, -0.003)	0.004

n=281; n=1 missing for percent of probing sites with attachment loss ≥ 3 mm; n=25 missing for alcohol use; n=6 missing for physical activity, n=27 missing for dietary pattern score. **Bold face** used for p <0.20. Insulin resistance as measured by the natural log-transformed Homeostasis Model Assessment for Insulin Resistance (HOMA-IR).

*BMI indicates body mass index; METS, metabolic equivalents.

B)

Variable	Regression coefficient with plasma glucose as dependent variable		Regression coefficient with systolic blood pressure as dependent variable		Regression coefficient with diastolic blood pressure as dependent variable	
	Crude Mean difference (95% CI)	P value	Crude Mean difference (95% CI)	P value	Crude Mean difference (95% CI)	P value
Age (5 years)	1.25 (0.82, 1.68)	<0.001	1.10 (0.38, 1.82)	0.003	1.25 (0.70, 1.80)	<0.001
Female	-1.78 (-3.93, 0.37)	0.10	-6.81 (-10.20, -3.42)	<0.001	-3.37 (-6.05, -0.69)	0.01
Race/ethnicity		0.002		0.006		0.005
Hispanic	Ref	-	Ref	-	Ref	-
Non-Hispanic White	3.50 (-5.74, -1.26)	0.002	-3.82 (-7.44, -0.19)	0.04	-3.17 (-5.99, -0.36)	0.03
Black	-2.77 (-5.19, -0.34)	0.03	3.04 (-0.90, 6.98)	0.13	3.09 (0.03, 6.14)	0.05
Other	1.08 (-1.63, 3.78)	0.43	-4.29 (-8.67, 0.08)	0.05	-1.42 (-4.82, 1.98)	0.41
Education		0.05		0.002		0.01
< Bachelor's Degree	Ref	-	Ref	-	Ref	-
Bachelor's Degree	-2.52 (-4.55, -0.49)	0.01	-3.99 (-0.72, -7.26)	0.02	-3.43 (-5.96, -0.90)	0.008
> Bachelor's Degree	-1.24 (-3.66, 1.18)	0.31	-6.87 (-10.72, -3.01)	<0.001	-3.78 (-6.80, -0.76)	0.01
Smoking Status		0.52		0.22		0.29
Never	Ref	-	Ref	-	Ref	-
Former	1.08 (-1.70, 3.85)	0.45	3.81 (-0.65, 8.28)	0.09	2.18 (-1.29, 5.66)	0.22
Current	1.50 (-1.64, 4.64)	0.35	1.67 (-3.39, 6.72)	0.52	2.26 (-1.67, 6.19)	0.26
BMI (kg/m²)	0.28 (0.14, 0.43)	<0.001	0.64 (0.41, 0.86)	<0.001	0.42 (0.25, 0.60)	<0.001
BMI categories		<0.001		<0.001		
Underweight/Normal	Ref	-	Ref	-	Ref	-
Overweight	2.20 (0.21, 4.20)	0.03	4.92 (1.78, 8.05)	0.002	3.14 (0.68, 5.60)	0.01
Obese	4.20 (2.07, 6.54)	<0.001	9.92 (6.40, 13.43)	<0.001	7.00 (4.25, 9.76)	<0.001
Percent of sites with attachment loss ≥3mm (10% increase)	1.11 (0.50, 1.72)	<0.001	1.35 (0.36, 2.33)	0.007	1.22 (0.46, 1.99)	0.002
Average alcohol consumption		0.06		0.92		0.48
None	Ref	-	Ref	-	Ref	-
≤ 1 drink/day	-3.57 (-7.24, 0.09)	0.06	-1.25 (-7.45, 4.95)	0.69	-2.56 (-7.31, 2.18)	0.29

> 1 drink/day	-5.14 (-9.39, -0.89)	0.02	-0.97 (-8.16, 6.23)	0.79	-3.38 (-8.89, 2.13)	0.23
Physical activity		0.20		0.01		0.03
METS min/week						
None	Ref	-	Ref	-	Ref	-
Low	-1.17 (-4.13, 1.80)	0.44	-5.31 (-10.08, -0.56)	0.03	-4.72 (-8.35, -1.08)	0.01
Moderate	-0.65 (-3.42, 2.13)	0.65	3.76 (-0.70, 8.22)	0.10	1.22 (-2.19, 4.62)	0.48
High	-2.26 (-4.35, -0.17)	0.03	-0.26 (-3.61, 3.10)	0.88	-1.16 (-3.72, 1.40)	0.37
Alternate Healthy Eating	-0.06 (-0.15, 0.02)	0.13	-0.17 (-0.30, -0.04)	0.01	-0.16 (-0.26, -0.06)	0.002
Index Score (AHEI)						

n=281; n=1 missing for percent of probing sites with attachment loss ≥ 3 mm; n=25 missing for alcohol use; n=6 missing for physical activity, n=27 missing for dietary pattern score. **Bold face** used for $p < 0.20$.

*BMI indicates body mass index; METS, metabolic equivalents.

C)

Variable	Regression coefficient (β)* with prediabetes as dependent variable	P value	Regression coefficient (β)* with hypertension as dependent variable	P value
Age (5 years)	0.38 (0.28, 0.49)	<0.001	0.17 (0.10, 0.25)	<0.001
Female	0.53 (-0.21, 1.28)	0.16	-0.35 (-0.70, -0.001)	0.05
Race/ethnicity		<0.001		0.001
Hispanic	Ref	-	Ref	-
Non-Hispanic White	-2.64 (-4.60, -0.67)	0.01	-0.78 (-1.36, -0.20)	0.008
Black	0.16 (-0.40, 0.72)	0.57	0.19 (-0.17, 0.56)	0.30
Other	-0.33 (-1.13, 0.47)	0.42	-0.58 (-1.23, 0.07)	0.08
Education		<0.001		0.04
< Bachelor's Degree	Ref	-	Ref	-
Bachelor's Degree	-0.99 (-1.55, -0.43)	<0.001	-0.33 (-0.68, 0.03)	0.07
> Bachelor's Degree	-1.21 (-2.03, -0.39)	0.004	-0.59 (-1.09, -0.09)	0.02
Smoking Status		0.33		0.53
Never	Ref	-	Ref	-
Former	0.47 (-0.16, 1.10)	0.14	0.28 (-0.16, 0.72)	0.21

Current	-0.35 (-1.45, 0.75)	0.53	0.001 (-0.60, 0.60)	0.99
BMI (kg/m²)	0.06 (0.03, 0.09)	<0.001	0.05 (0.04, 0.07)	<0.0001
BMI categories		0.002		<0.0001
Underweight/Normal	Ref	-	Ref	-
Overweight	0.69 (0.01, 1.37)	0.05	0.51 (0.06, 0.95)	0.02
Obese	1.17 (0.52, 1.82)	<0.001	0.96 (0.55, 1.37)	<0.0001
Percent of sites with attachment loss \geq3mm (10% increase)	0.27 (0.15, 0.39)	<0.001	0.16 (0.07, 0.24)	0.0003
Average alcohol consumption		0.69		0.56
None	Ref	-	Ref	-
\leq 1 drink/day	-0.34 (-1.25, 0.58)	0.47	0.11 (-0.66, 0.87)	0.79
> 1 drink/day	-0.55 (-1.74, 0.63)	0.36	-0.19 (-1.12, 0.74)	0.69
Physical activity METS min/week		0.02		0.22
None	Ref	-	Ref	-
Low	-0.04 (-0.75, 0.67)	0.92	-0.49 (-1.22, 0.23)	0.18
Moderate	-0.22 (-0.95, 0.50)	0.54	0.23 (-0.24, 0.70)	0.33
High	-0.93 (-1.60, -0.25)	0.007	0.20 (-0.38, 0.41)	0.94
Alternate Healthy Eating Index Score (AHEI)	-0.004 (-0.03, 0.02)	0.77	-0.02 (-0.03, 0.0001)	0.05

n=281; n=280 for hypertension. n=1 missing for percent of probing sites with attachment loss \geq 3mm; n=25 missing for alcohol use; n=6 missing for physical activity, n=27 missing for dietary pattern score. **Bold face** used for p <0.20

* Beta coefficient of the multivariable regression, before exponentiation.

*BMI indicates body mass index; METS, metabolic equivalents.

Table S4. Characteristics of the 281 participants from the Oral Infections, Glucose Intolerance, and Insulin Resistance Study (ORIGINS) used in this study

Variable	Total sample N=281	Normotensive participants N=187	Hypertensive participants N=93
Age (years)	34 (10)	32 (9)	37 (11)
Female	78%	82%	72%
Race/ethnicity			
Hispanic	47%	43%	55%
Non-Hispanic White	22%	27%	12%
Black	17%	14%	25%
Other	13%	16%	9%
Education			
<Bachelor's Degree	32%	28%	42%
Bachelor's Degree	45%	47%	42%
>Bachelor's Degree	22%	26%	16%
Smoking Status			
Never	79%	81%	76%
Former	12%	10%	15%
Current	9%	9%	9%
BMI (kg/m²)	27.0 (6.1)	25.7 (5.3)	29.6 (6.8)
Periodontitis			
None/mild	42%	45%	36%
Moderate	52%	50%	56%
Severe	6%	5%	9%
% of sites with attachment loss ≥3mm	17 (14)	15 (13)	21 (16)
Average alcohol consumption			
None	7%	7%	6%
≤ 1 drink/day	79%	77%	83%
> 1 drink/day	14%	16%	11%
Physical activity (METS min/week)			
None	31%	31%	32%
Low	12%	15%	8%
Moderate	15%	13%	18%
High	41%	41%	42%
Alternate Healthy Eating Index Score (AHEI)	49 (12)	50 (12)	47 (12)
NO₃TSS	0 (5.42)	0.13 (5.25)	-0.29 (5.79)
Systolic blood pressure	117.4 (12.3)	112.1 (8.94)	128.2 (11.1)
Diastolic blood pressure	74.9 (9.6)	70.4 (6.02)	84.0 (8.91)
Plasma glucose	85.0 (7.63)	84.4 (7.26)	86.2 (8.25)
Plasma insulin	10.52 (8.49)	9.63 (9.02)	12.34 (7.06)

Insulin resistance (HOMA-IR)	2.26 (2.11)	2.05 (2.24)	2.68 (1.75)
Median Insulin Resistance (HOMA-IR)	1.75 (1.45)	1.61 (1.08)	2.33 (1.88)

Values presented in mean (standard deviation) or percentages unless otherwise stated, or median (interquartile range),

*BMI indicates body mass index; METS, metabolic equivalents; NO₃TSS, nitrate-reducing bacterial summary score; HOMA-IR, Homeostasis Model Assessment for Insulin Resistance.

For total sample: n=1 missing for percent of probing sites with attachment loss ≥ 3 mm; n=25 missing for alcohol use; n=6 missing for physical activity, n=27 missing for dietary pattern score.

For those without hypertension: none missing for periodontitis and percent of probing sites with attachment loss ≥ 3 mm; n=11, missing for alcohol use; n=17 missing for diet score; n=5 missing for physical activity.

For those with hypertension: n=1 missing for percent of probing sites with attachment loss ≥ 3 mm; n=13 missing for alcohol use, n=1 missing for physical activity; n=10 missing for diet score.

TABLE S5. Mean difference in systolic and diastolic blood pressure (mmHg) for every 1 standard deviation (STD) increase in nitrate-reducing bacterial summary score (NO₃TSS) in those without hypertension (n=242) according to the 2003 American Heart Association (AHA) Guideline's 140/90 mmHg cutoff for hypertension, and in those who were hypertensive and taking medication (n=13)

	NORMOTENSIVE (according to the 2003 AHA 140/90 mmHg criteria)		HYPERTENSIVE AND TAKING MEDICATIONS (n=13)	
	Systolic blood pressure (mmHg)	Diastolic blood pressure (mmHg)	Systolic blood pressure (mmHg)	Diastolic blood pressure (mmHg)
	NO₃TSS (1 STD)	NO₃TSS (1 STD)	NO₃TSS (1 STD)	NO₃TSS (1 STD)
M1	-2.02 (-3.32, -0.71)	-1.22 (-2.18, -0.27)	-1.43 (-7.97, 5.12)	-1.80 (-7.53, 3.93)
M2	-1.87 (-3.13, -0.61)	-1.10 (-2.05, -0.15)	2.54 (-4.02, 9.11)	5.00 (-0.77, 10.76)
M3	-1.71 (-2.94, -0.49)	-1.04 (-1.98, -0.10)	2.89 (-3.87, 9.66)	6.29 (-0.19, 12.77)
M4	-1.68 (-2.92, -0.45)	-1.00 (-1.94, -0.05)	3.12 (0.76, 5.48)	7.72 (3.69, 11.75)
M5	-1.55 (-2.85, -0.26)	-0.76 (-1.74, 0.22)	Not able to estimate due to small n	Not able to estimate due to small n
M6	-1.68 (-3.01, -0.35)	-0.97 (-1.97, 0.02)	Not able to estimate due to small n	Not able to estimate due to small n

Model 1: Crude (n=242; n=13)

Model 2: Adjusted for age, sex, race, education (n=242; n=13)

Model 3: Adjusted for age, sex, race, education, BMI, smoking (n=242; n=13)

Model 4: Adjusted for age, sex, race, education, BMI, smoking, percent of periodontal sites with attachment loss ≥ 3 mm (n=242; n=13)

Model 5: Adjusted for age, sex, race, education, BMI, smoking, percent of periodontal sites with attachment loss ≥ 3 mm, and dietary pattern (n=220)

Model 6: Adjusted for age, sex, race, education, BMI, smoking, percent of periodontal sites with attachment loss ≥ 3 mm, dietary pattern, alcohol use, and physical activity (n=198)

* **BMI indicates body mass index.**

TABLE S6. Mean difference in natural log-transformed Homeostasis Model Assessment for Insulin Resistance (lnHOMA-IR) and plasma glucose, and prevalence ratio for prediabetes, for every 1 standard deviation higher z-score arcsin-sqrt transformed relative abundance for the 20 individual nitrate-reducing bacteria taxa

Adjusted for age, sex, race, education, BMI, smoking, percent of sites with attachment loss ≥ 3 mm, and dietary pattern

Taxa	Insulin resistance	Lower 95% CI	Upper 95% CI	Glucose	Lower 95% CI	Upper 95% CI	Prediabetes	Lower 95% CI	Upper 95% CI
<i>Actinomyces naeslundii</i>	-0.07	-0.13	0.00	-0.44	-1.33	0.45	0.73	0.50	1.06
<i>Actinomyces odontolyticus</i>	-0.03	-0.09	0.03	-0.33	-1.17	0.52	0.85	0.67	1.09
<i>Actinomyces viscosus</i>	-0.05	-0.11	0.02	-0.72	-1.63	0.19	0.65	0.46	0.93
<i>Capnocytophaga sputigena</i>	-0.03	-0.09	0.03	-0.76	-1.59	0.07	0.65	0.45	0.95
<i>Corynebacterium durum</i>	-0.06	-0.12	0.01	-0.70	-1.61	0.22	0.72	0.49	1.06
<i>Corynebacterium matruchotii</i>	-0.05	-0.12	0.01	0.04	-0.84	0.92	0.94	0.66	1.34
<i>Eikenella corrodens</i>	-0.03	-0.09	0.03	-0.30	-1.16	0.55	0.97	0.73	1.29
<i>Haemophilus parainfluenzae</i>	-0.05	-0.11	0.02	-0.96	-1.88	-0.05	0.87	0.64	1.20
<i>Neisseria flavescens</i>	-0.08	-0.14	-0.02	-0.73	-1.57	0.12	1.04	0.86	1.27
<i>Neisseria sicca</i>	-0.05	-0.11	0.01	-0.03	-0.87	0.80	0.54	0.22	1.37
<i>Neisseria subflava</i>	-0.01	-0.09	0.06	0.09	-0.91	1.09	1.16	0.94	1.44
<i>Prevotella melaninogenica</i>	0.03	-0.03	0.09	-0.16	-1.00	0.68	1.09	0.79	1.51
<i>Prevotella salivae</i>	0.02	-0.04	0.08	-0.07	-0.93	0.78	1.24	0.98	1.57
<i>Propionibacterium acnes</i>	0.01	-0.05	0.07	0.47	-0.37	1.31	0.79	0.45	1.38
<i>Rothia dentocariosa</i>	0.00	-0.06	0.06	0.31	-0.59	1.21	1.10	0.87	1.40
<i>Rothia mucilaginosa</i>	-0.03	-0.10	0.03	-0.34	-1.22	0.54	0.80	0.49	1.28
<i>Selenomonas noxia</i>	-0.03	-0.09	0.04	-0.78	-1.66	0.11	1.04	0.80	1.35
<i>Veillonella atypica</i>	0.00	-0.06	0.06	0.01	-0.87	0.90	0.95	0.81	1.12
<i>Veillonella dispar</i>	-0.01	-0.07	0.06	-0.21	-1.07	0.65	0.83	0.60	1.17
<i>Veillonella parvula</i>	0.01	-0.05	0.07	-0.03	-0.89	0.83	1.02	0.77	1.37

p < 0.05 in bold, BMI indicates body mass index.

TABLE S7. Mean difference in mean systolic and diastolic blood pressure, and prevalence ratio for hypertension, for every 1 standard deviation higher z-score arcsin-sqrt transformed relative abundance for the 20 individual nitrate-reducing bacteria taxa.

Adjusted for age, sex, race, education, BMI, smoking, percent of sites with attachment loss ≥ 3 mm, and dietary pattern

Taxa	Systolic blood pressure	Lower 95% CI	Upper 95% CI	Diastolic blood pressure	Lower 95% CI	Upper 95% CI	Hypertension	Lower 95% CI	Upper 95% CI
<i>Actinomyces naeslundii</i>	1.21	-0.18	2.59	0.10	-0.90	1.09	1.17	1.03	1.32
<i>Actinomyces odontolyticus</i>	-1.16	-2.35	0.03	0.27	-0.59	1.13	0.99	0.84	1.17
<i>Actinomyces viscosus</i>	-0.35	-1.65	0.95	-0.28	-1.21	0.66	1.06	0.88	1.28
<i>Capnocytophaga sputigena</i>	-0.05	-1.35	1.26	0.56	-0.37	1.49	0.98	0.86	1.11
<i>Corynebacterium durum</i>	0.05	-1.31	1.41	-0.41	-1.38	0.56	1.19	0.99	1.42
<i>Corynebacterium matruchotii</i>	-0.72	-1.95	0.52	-0.37	-1.26	0.51	0.93	0.77	1.11
<i>Eikenella corrodens</i>	0.15	-1.26	1.57	0.51	-0.49	1.52	1.02	0.89	1.18
<i>Haemophilus parainfluenzae</i>	-2.59	-3.99	-1.18	-1.41	-2.43	-0.39	1.07	0.90	1.28
<i>Neisseria flavescens</i>	-1.35	-2.58	-0.12	-0.63	-1.52	0.26	0.97	0.77	1.22
<i>Neisseria sicca</i>	-1.37	-2.40	-0.35	-1.13	-1.86	-0.40	0.92	0.79	1.07
<i>Neisseria subflava</i>	-0.10	-1.33	1.12	-0.48	-1.35	0.39	1.06	0.87	1.27
<i>Prevotella melaninogenica</i>	0.29	-0.82	1.39	0.03	-0.76	0.82	0.87	0.72	1.06
<i>Prevotella salivae</i>	0.29	-0.86	1.45	0.06	-0.77	0.89	1.01	0.85	1.20
<i>Propionibacterium acnes</i>	0.29	-0.80	1.38	0.73	-0.04	1.50	0.91	0.74	1.12
<i>Rothia dentocariosa</i>	-0.13	-1.39	1.12	0.41	-0.48	1.31	1.04	0.87	1.24
<i>Rothia mucilaginosa</i>	-0.86	-2.07	0.35	-0.75	-1.61	0.11	0.94	0.74	1.18
<i>Selenomonas noxia</i>	-0.08	-1.27	1.11	0.10	-0.75	0.95	0.90	0.74	1.09
<i>Veillonella atypica</i>	0.16	-1.91	2.24	0.72	-0.76	2.20	1.02	0.96	1.08
<i>Veillonella dispar</i>	-0.26	-1.53	1.00	-0.16	-1.07	0.74	1.05	0.89	1.24
<i>Veillonella parvula</i>	-0.99	-2.44	0.45	-0.53	-1.57	0.51	1.11	0.96	1.28

p <0.05 in bold, BMI indicates body mass index. Results for systolic and diastolic blood pressure for the participants without hypertension, results for hypertension in the full sample.