

Poster presentation

Open Access

The immunological basis of tumor therapy by targeted delivery of TNF α to tumor vessels

Enrica Balza¹, Lorenzo Mortara², Francesca Sassi¹, Stefano Monteghirfo¹, Barbara Carnemolla¹, Patrizia Castellani¹, Dario Neri³, Roberto Accollo^{*3}, Luciano Zardi⁴ and Laura Borsi¹

Address: ¹Department of Translational Oncology, Istituto Nazionale per la Ricerca sul Cancro, Genoa, Italy, ²Department of Clinical and Biological Sciences, School of Medicine, University of Insubria, Varese, Italy, ³Department of Chemistry and Applied Biosciences, Institute of Pharmaceutical Sciences, Swiss Federal Institute of Technology, Zurich, Switzerland and ⁴Unit of Innovative Therapies, Istituto Giannina Gaslini, Centro Biotecnologie Avanzate, Genoa, Italy

* Corresponding author

from 2006 International Meeting of The Institute of Human Virology
Baltimore, USA. 17–21 November, 2006

Published: 21 December 2006

Retrovirology 2006, **3**(Suppl 1):P1 doi:10.1186/1742-4690-3-S1-P1

© 2006 Balza et al; licensee BioMed Central Ltd.

L19mTNF α is a fusion protein constituted by the scFv L19 specific for the oncofetal ED-B domain of fibronectin and TNF α . Treatment with L19mTNF α , in combination with melphalan, induced complete tumor regression in 83% of BALB/c mice with WEHI-164 fibrosarcoma and 33% of animals with C51 colon carcinoma. All cured mice rejected challenges with the same tumor cells and, in a very high percentage of animals, also challenges with syngeneic tumor cells of different histological origin. In adoptive immunity transfer experiments the splenocytes from C51-cured mice protected 100% of naive mice both from C51 colon carcinoma and from WEHI-164 fibrosarcoma. The splenocytes from WEHI-164-cured mice protected 100% of mice from the fibrosarcoma and 80% from the C51 colon carcinoma. Similar results were also obtained in adoptive immunity transfer experiments using severely immunodepressed SCID mice. Experiments using depleted splenocytes showed that T cells play a major role in tumor rejection. These data demonstrate that the selective targeting of mTNF α to the tumor enhances its immunostimulatory properties to the point of generating a therapeutic immune response against different histologically unrelated syngeneic tumors. These findings predicate treatment approaches for cancer patients based on the targeted delivery of TNF α to tumor vasculature.