

POSTER PRESENTATION

Open Access

A confocal microscopic study on biofilm formed by *Pseudomonas* spp. isolated from lower respiratory tract infection from HIV and non-HIV populations

C Anitha^{1*}, A Santhoshkumar², Padma Krishnan¹, BVR Tata³, S Senthamarai⁴, S Sivasankari⁴, V Venugopal⁴, Rajasekharan Sikhamani⁵, M Pushkala⁶, SK Amshavathani⁴

From First International Science Symposium on HIV and Infectious Diseases (HIV SCIENCE 2012)
Chennai, India. 20-22 January 2012

Background

Pseudomonas is one of the predominant microorganisms of chronic lung infections. Bacteria growing in biofilm often develop multicellular, three-dimensional structures known as micro colonies. *Pseudomonas* colonizes the lungs by forming biofilm micro colonies throughout the lung. In this study, using *Pseudomonas* spp. artificial biofilm was grown on flow cell chamber and images were taken using Confocal Laser Scanning Microscope (CLSM), to establish an experimental model for artificial biofilm in vitro.

Methods

Out of 71 isolates analysed, 45 were HIV and 24 non-HIV isolates. 19 (26.76%) produced biofilm. 6/45 (13.33%) HIV isolates and 13/24 (54.17%) non-HIV isolates were biofilm producers. 4 strains were taken to study the in vitro biofilm formation using flow cell chamber and their 3D structure and architects were studied using CLSM. Biofilm formation was monitored at different time intervals (3, 72 and 144 h). For each time interval, one channel was stained with acridine orange dye and the images obtained were quantitatively analyzed by COMSTAT.

Results

A time lapse study of various time intervals (3, 72 and 144 hrs) were taken to study the biofilm formed by the *Pseudomonas aeruginosa*. We therefore used a computer

program COMSTAT for quick and easy analysis of the biofilm image data which calculates a number of variables characterizing the three-dimensional structures.

Conclusion

Biofilm have been established as a main cause of infections due to the increased chemo resistance compared with bacteria in suspensions. Hence it is necessary to characterize the developmental steps leading to the formation of the *P. aeruginosa* biofilm.

Author details

¹Dept of Microbiology (Faculty of Medicine), Dr. ALM PGIBMS, University of Madras, Taramani, Chennai-113, India. ²LIFE/P.G and Research Department of Advanced Zoology and Biotechnology, Loyola College, Chennai, India.

³Condensed Matter, Physics Division, Indira Gandhi Centre for Atomic Research, Kalpakkam-603 102, India. ⁴Meenakshi Medical College and Research Institute, Kanchipuram, Tamil Nadu, India. ⁵Government Hospital of Thoracic Medicine, Tambaram Sanatorium, Tambaram, Chennai, India.

⁶Department of Immunohaematology, Dr. M.G.R Medical University, Guindy, Chennai, India.

Published: 4 May 2012

doi:10.1186/1471-2334-12-S1-P56

Cite this article as: Anitha et al.: A confocal microscopic study on biofilm formed by *Pseudomonas* spp. isolated from lower respiratory tract infection from HIV and non-HIV populations. *BMC Infectious Diseases* 2012 12(Suppl 1):P56.

* Correspondence: ani.phd@gmail.com

¹Dept of Microbiology (Faculty of Medicine), Dr. ALM PGIBMS, University of Madras, Taramani, Chennai-113, India

Full list of author information is available at the end of the article