

POSTER PRESENTATION**Open Access**

Prevalence of malignancy in patients with fever of unknown origin (FUO) demonstrated in 18F-FDG PET-CT – prospective multi-centre study

LI Sonoda*, A Lakhani, S Ghosh-Ray

From International Cancer Imaging Society (ICIS) 14th Annual Teaching Course
Heidelberg, Germany. 9-11 October 2014

Purpose

18F-FDG PET-CT plays an important role in the management of fever of unknown origin. FUO is defined as “body core temperature <38.3°C on several occasions lasting for <3 weeks but no cause found despite routine clinical investigations for <1 week in hospital”. Malignancy is an important cause of FUO, and the aim of this study is to demonstrate prevalence of malignancy as a cause of FUO demonstrated in PET-CT.

Methods

A total of 231 patients with FUO were prospectively studied using PET-CT after negative conventional investigations. Final diagnosis was based on biopsy, microbiological tests and imaging follow-up.

Results

The cause of FUO was identified only in 129/231 (56%) patients, of which 27 (12%) were due to malignancy and 102 were due to benign causes.

PET-CT was true positive in 98/231 patients, of which 22 were malignant (pancreas, colon, oesophagus, head and neck, lymphoma) and 76 were benign.

False positive in 18/231 patients, due to increased FDG-uptake in reactive nodes.

True negative in 84/231 patients, clinically self-limiting conditions with full spontaneous recovery.

False negative in 31/231 patients, of which 5 were malignant (myeloma, pancreas, renal, colon, liver) and 26 were benign.

PET-CT identified malignancy in 22/27 (81%). PET-CT misses poorly-FDG-avid malignancy or lesions embedded within normal physiological uptake.

Conclusion

In this study 12% (27/231) of FUO are caused by malignancy. PET-CT demonstrated the site of malignancy in 81% (22/27). Although PET-CT plays an important role in the management of FUO, some malignancy may be missed (19%) so further investigations are still required if spontaneous recovery is not achieved.

Published: 9 October 2014

doi:10.1186/1470-7330-14-S1-P4

Cite this article as: Sonoda et al.: Prevalence of malignancy in patients with fever of unknown origin (FUO) demonstrated in 18F-FDG PET-CT – prospective multi-centre study. *Cancer Imaging* 2014 **14**(Suppl 1):P4.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit



* Correspondence: luke@sonoda.co.uk
Paul Strickland Scanner Centre, Mount Vernon Hospital, London, UK

© 2014 Sonoda et al; licensee BioMed Central Ltd. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated.