

Ethyl 1-oxo-1,2,3,4-tetrahydro-9H-carbazole-3-carboxylate

Tuncer Hökelek,^{a*} Hakan Dal,^b Barış Tercan,^c Mustafa Göçmentürk^d and Yavuz Ergün^d

^aDepartment of Physics, Hacettepe University, 06800 Beytepe, Ankara, Turkey,

^bDepartment of Chemistry, Faculty of Science, Anadolu University, 26470

Yenibağlar, Eskişehir, Turkey, ^cDepartment of Physics, Karabük University, 78050

Karabük, Turkey, and ^dDepartment of Chemistry, Faculty of Arts and Sciences,

Dokuz Eylül University, Tinaztepe, 35160 Buca-İzmir, Turkey

Correspondence e-mail: merzifon@hacettepe.edu.tr


Received 19 June 2009; accepted 22 June 2009

Key indicators: single-crystal X-ray study; $T = 100\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.004\text{ \AA}$; R factor = 0.075; wR factor = 0.221; data-to-parameter ratio = 17.0.

The title compound, $C_{15}H_{15}NO_3$, contains a carbazole skeleton with an ethoxycarbonyl group at the 3 position. In the indole ring system, the benzene and pyrrole rings are nearly coplanar, forming a dihedral angle of $1.95(8)^\circ$. The cyclohexenone ring has an envelope conformation. In the crystal structure, pairs of strong $\text{N}-\text{H}\cdots\text{O}$ hydrogen bonds link the molecules into centrosymmetric dimers with $R_2^2(10)$ ring motifs. $\pi-\pi$ contacts between parallel pyrrole rings [centroid-centroid distance = $3.776(2)\text{ \AA}$] may further stabilize the structure. A weak $\text{C}-\text{H}\cdots\pi$ interaction is also observed.

Related literature

For tetrahydrocarbazole derivatives as synthetic precursors of cyclic indole-type alkaloids of biological interest, see: Abraham (1975); Phillipson & Zenk (1980); Saxton (1983). The title compound is used in the synthesis of a precursor for the synthesis of the anti-tumor drug ellipticine (Ergün *et al.*, 2004). *Murraya L.* (Rutaceae) is a genus of shrubs or small trees from Southern Asia (Chang, 1977) from which carbazole alkaloids have been isolated (Chakraborty & Roy, 1991). For the biological activity of carbazole alkaloids, see: Kondo *et al.* (1986); Te Paske *et al.* (1989*a,b*). For related structures, see: Çaylak *et al.* (2007); Uludağ *et al.* (2009). For bond-length data, see: Allen *et al.* (1987). For ring-motifs, see: Bernstein *et al.* (1995).



Experimental

Crystal data

$C_{15}H_{15}NO_3$	$V = 1230.69(12)\text{ \AA}^3$
$M_r = 257.28$	$Z = 4$
Monoclinic, $P2_1/c$	Mo $K\alpha$ radiation
$a = 5.6811(3)\text{ \AA}$	$\mu = 0.10\text{ mm}^{-1}$
$b = 8.7378(5)\text{ \AA}$	$T = 100\text{ K}$
$c = 24.8310(14)\text{ \AA}$	$0.45 \times 0.20 \times 0.15\text{ mm}$
$\beta = 93.208(4)^\circ$	

Data collection

Bruker Kappa APEXII CCD area-detector diffractometer	9148 measured reflections
Absorption correction: multi-scan (<i>SADABS</i> ; Bruker, 2005)	3004 independent reflections
$T_{\min} = 0.958$, $T_{\max} = 0.983$	2145 reflections with $I > 2\sigma(I)$
	$R_{\text{int}} = 0.059$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.075$	H atoms treated by a mixture of independent and constrained refinement
$wR(F^2) = 0.221$	$\Delta\rho_{\max} = 1.25\text{ e \AA}^{-3}$
$S = 1.05$	$\Delta\rho_{\min} = -0.42\text{ e \AA}^{-3}$
3004 reflections	
177 parameters	

Table 1

Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{N9}-\text{H9}\cdots\text{O1}^{\text{i}}$	0.82 (4)	2.08 (4)	2.834 (3)	154 (4)
$\text{C4}-\text{H4A}\cdots\text{Cg3}^{\text{ii}}$	0.99	2.75	3.727 (3)	171

Symmetry codes: (i) $-x + 1, -y, -z + 1$; (ii) $-x, -y + 2, -z$. Cg3 is the centroid of the CSA/C8/C8A ring.

Data collection: *APEX2* (Bruker, 2007); cell refinement: *SAINT* (Bruker, 2007); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3 for Windows* (Farrugia, 1997) and *Mercury* (Macrae *et al.*, 2006); software used to prepare material for publication: *WinGX* (Farrugia, 1999) and *PLATON* (Spek, 2009).

The authors are indebted to Anadolu University and the Medicinal Plants and Medicine Research Centre of Anadolu University, Eskişehir, Turkey, for the use of X-ray diffractometer.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: XU2543).

References

- Abraham, D. J. (1975). *The Catharanthus Alkaloids*, edited by W. I. Taylor & N. R. Fransworth, chs. 7 and 8. New York: Marcel Decker.
- Allen, F. H., Kennard, O., Watson, D. G., Brammer, L., Orpen, A. G. & Taylor, R. (1987). *J. Chem. Soc. Perkin Trans. 2*, pp. S1–19.
- Bernstein, J., Davis, R. E., Shimonini, L. & Chang, N.-L. (1995). *Angew. Chem. Int. Ed. Engl.* **34**, 1555–1573.
- Bruker (2005). *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Bruker (2007). *APEX2* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Çaylak, N., Hökelek, T., Uludağ, N. & Patir, S. (2007). *Acta Cryst. E***63**, o3913–o3914.
- Chakraborty, D. P. & Roy, S. (1991). *Progress in the Chemistry of Organic Natural Products*, edited by W. Herz, G. W. Kirby, W. Steglich & Ch. Tamm, Vol. 57, pp. 71–152. Wien, New York: Springer-Verlag.
- Chang, C. E. (1977). *Flora of Taiwan*, Vol. 3, pp. 520–523. Taipei, Taiwan: Poch Publishing Co. Ltd.
- Ergün, Y., Patir, S. & Okay, G. (2004). *Synth. Commun.* **34**, 435–442.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Farrugia, L. J. (1999). *J. Appl. Cryst.* **32**, 837–838.
- Kondo, S., Katayama, M. & Marumo, S. (1986). *J. Antibiot.* **39**, 727–730.
- Macrae, C. F., Edgington, P. R., McCabe, P., Pidcock, E., Shields, G. P., Taylor, R., Towler, M. & van de Streek, J. (2006). *J. Appl. Cryst.* **39**, 453–457.
- Phillipson, J. D. & Zenk, M. H. (1980). *Indole and Biogenetically Related Alkaloids*, ch 3. New York: Academic Press.
- Saxton, J. E. (1983). Editor. *Heterocyclic Compounds*, Vol. 25, *The Monoterpene Indole Alkaloids*, chs. 8 and 11. New York: Wiley.
- Sheldrick, G. M. (2008). *Acta Cryst. A***64**, 112–122.
- Spek, A. L. (2009). *Acta Cryst. D***65**, 148–155.
- Te Paske, M. R., Gloer, J. B., Wicklow, D. T. & Dowd, P. F. (1989a). *J. Org. Chem.* **54**, 4743–4746.
- Te Paske, M. R., Gloer, J. B., Wicklow, D. T. & Dowd, P. F. (1989b). *Tetrahedron Lett.* **30**, 5965–5968.
- Uludağ, N., Öztürk, A., Hökelek, T. & Erdoğan, Ü. I. (2009). *Acta Cryst. E***65**, o595–o596.

supplementary materials

Acta Cryst. (2009). E65, o1702-o1703 [doi:10.1107/S160053680902385X]

Ethyl 1-oxo-1,2,3,4-tetrahydro-9H-carbazole-3-carboxylate

T. Hökelek, H. Dal, B. Tercan, M. Göçmentürk and Y. Ergün

Comment

Tetrahydrocarbazole derivatives can be considered to be synthetic precursors of cyclic indole-type alkaloids of biological interest (Abraham, 1975; Phillipson & Zenk, 1980; Saxton, 1983). The title compound was used in the synthesis of the precursor compound for the synthesis of anti-tumor drug ellipticine (Ergün *et al.*, 2004). *Murraya L.* (Rutaceae) is a genus of shrubs or small trees from Southern Asia (Chang, 1977). The main constituent of this genus include carbazole alkaloids (Chakraborty & Roy, 1991). Several biological properties have been reported for carbazole alkaloids including antibiotic, cytotoxic and antiviral activities (Kondo *et al.*, 1986; Te Paske *et al.*, 1989*a,b*). The title compound may also be used as a precursor in the synthesis of *Murraya* alkaloids. The present study was undertaken to ascertain its crystal structure.

The molecule of the title compound (Fig. 1) contains a carbazole skeleton with a carboxyethyl group at position 3, where the bond lengths (Allen *et al.*, 1987) and angles are within normal ranges.

An examination of the deviations from the least-squares planes through individual rings shows that rings B (C4a/C5a/C8a/N9/C9a) and C (C5a/C5—C8/C8a) are nearly coplanar [with a maximum deviation of -0.028 (3) Å for atom C4a] with dihedral angle of A/B = 1.95 (8)°. Ring A (C1—C4/C4a/C9a) adopts envelope conformation with atom C3 displaced by 0.527 (3) Å from the plane of the other rings atoms, as in 3a,4,10,10b-tetrahydro-2H-furo[2,3-*a*]carbazol-5(3H)-one (Çaylak *et al.*, 2007) and 3,3-ethylenedithio-3,3a,4,5,10,10b-hexahydro-2H-furo[2,3-*a*]carbazole (Uludağ *et al.*, 2009).

In the crystal structure, pairs of strong intermolecular N—H···O hydrogen bonds (Table 1) link the molecules into centrosymmetric dimers with $R^2_2(10)$ ring motifs (Bernstein *et al.*, 1995) (Fig. 2), in which they may be effective in the stabilization of the structure. The π – π contact between the pyrrole rings, $Cg2—Cg2^i$, [symmetry code: (i) $-x, 1 - y, -z$, where $Cg2$ is centroid of the ring B (C4a/C5a/C8a/N9/C9a)] may further stabilize the structure, with centroid-centroid distance of 3.776 (2) Å. There also exists a weak C—H··· π interaction (Table 1).

Experimental

For the preparation of the title compound, a solution of ethyl 1,2,3,4-tetrahydro-9H-carbazole-3-carboxylate (5.00 g, 20.5 mmol) in methanol (25 ml) was added dropwise to a solution of periodic acid (9.35 g, 41.0 mmol) in methanol-water (1:1, 100 ml) at 273 K. The reaction mixture was stirred for 1 h at 273 K, then stirring was continued for a further 1 h at room temperature. The solvent was evaporated, then the residue was dissolved in chloroform and washed first with sodium carbonate (10%, 50 ml) and then with sodium bisulfite (10%, 50 ml). The organic layer was dried over anhydrous magnesium sulfate and the solvent was evaporated. The residue was chromatographed on silica gel using ethyl acetate and crystallized from methanol (yield: 3.17 g, 67%, m.p. 411 K).

supplementary materials

Refinement

The highest peak in the final difference electron-density map is apart 0.94 Å from atom C3. Atom H9 (for NH) was located in difference Fourier map and refined isotropically. The remaining H atoms were positioned geometrically, with C—H = 0.95, 1.00, 0.99 and 0.98 Å for aromatic, methine, methylene and methyl H, respectively, and constrained to ride on their parent atoms, with $U_{\text{iso}}(\text{H}) = xU_{\text{eq}}(\text{C})$, where $x = 1.5$ for methyl H and $x = 1.2$ for all other H atoms.

Figures




Fig. 1. The molecular structure of the title molecule with the atom-numbering scheme. The displacement ellipsoids are drawn at the 50% probability level.




Fig. 2. A partial packing diagram for the title compound. Hydrogen bonds are shown as dashed lines. H atoms not involved in hydrogen bonding have been omitted for clarity.

Ethyl 1-oxo-1,2,3,4-tetrahydro-9*H*-carbazole-3-carboxylate

Crystal data

C ₁₅ H ₁₅ NO ₃	$F_{000} = 544$
$M_r = 257.28$	$D_x = 1.389 \text{ Mg m}^{-3}$
Monoclinic, $P2_1/c$	Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
Hall symbol: -P 2ybc	Cell parameters from 3480 reflections
$a = 5.6811 (3) \text{ \AA}$	$\theta = 1.6\text{--}28.3^\circ$
$b = 8.7378 (5) \text{ \AA}$	$\mu = 0.10 \text{ mm}^{-1}$
$c = 24.8310 (14) \text{ \AA}$	$T = 100 \text{ K}$
$\beta = 93.208 (4)^\circ$	Rod-shaped, colorless
$V = 1230.69 (12) \text{ \AA}^3$	$0.45 \times 0.20 \times 0.15 \text{ mm}$
$Z = 4$	

Data collection

Bruker Kappa APEXII CCD area-detector diffractometer	3004 independent reflections
Radiation source: fine-focus sealed tube	2145 reflections with $I > 2\sigma(I)$
Monochromator: graphite	$R_{\text{int}} = 0.059$
$T = 100 \text{ K}$	$\theta_{\text{max}} = 28.3^\circ$
φ and ω scans	$\theta_{\text{min}} = 1.6^\circ$
Absorption correction: multi-scan (SADABS; Bruker, 2005)	$h = -7 \rightarrow 7$

$T_{\min} = 0.958$, $T_{\max} = 0.983$
9148 measured reflections

$k = -11 \rightarrow 9$
 $l = -32 \rightarrow 28$

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.075$	H atoms treated by a mixture of independent and constrained refinement
$wR(F^2) = 0.221$	$w = 1/[\sigma^2(F_o^2) + (0.1083P)^2 + 1.2571P]$ where $P = (F_o^2 + 2F_c^2)/3$
$S = 1.05$	$(\Delta/\sigma)_{\max} < 0.001$
3004 reflections	$\Delta\rho_{\max} = 1.25 \text{ e \AA}^{-3}$
177 parameters	$\Delta\rho_{\min} = -0.42 \text{ e \AA}^{-3}$
Primary atom site location: structure-invariant direct methods	Extinction correction: none

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
O1	0.4546 (3)	0.0609 (2)	0.42291 (7)	0.0294 (5)
O2	-0.0619 (5)	0.3051 (4)	0.27554 (9)	0.0775 (11)
O3	-0.2959 (5)	0.4520 (3)	0.32045 (8)	0.0603 (8)
C1	0.2768 (4)	0.1407 (3)	0.42209 (10)	0.0245 (5)
C2	0.1783 (5)	0.2133 (4)	0.37051 (10)	0.0316 (6)
H2A	0.1988	0.1406	0.3405	0.038*
H2B	0.2727	0.3056	0.3634	0.038*
C3	-0.0831 (5)	0.2597 (4)	0.36991 (11)	0.0355 (7)
H3	-0.1758	0.1623	0.3689	0.043*
C4	-0.1525 (4)	0.3454 (3)	0.41897 (10)	0.0263 (6)
H4A	-0.0978	0.4527	0.4171	0.032*
H4B	-0.3264	0.3461	0.4202	0.032*
C4A	-0.0453 (4)	0.2708 (3)	0.46893 (10)	0.0229 (5)
C5	-0.2963 (4)	0.3489 (3)	0.54916 (10)	0.0270 (6)
H5	-0.4152	0.4041	0.5289	0.032*

supplementary materials

C5A	-0.1107 (4)	0.2805 (3)	0.52332 (9)	0.0232 (5)
C6	-0.3046 (5)	0.3353 (3)	0.60418 (11)	0.0313 (6)
H6	-0.4315	0.3802	0.6219	0.038*
C7	-0.1266 (5)	0.2553 (3)	0.63473 (10)	0.0314 (6)
H7	-0.1334	0.2502	0.6728	0.038*
C8	0.0550 (5)	0.1850 (3)	0.61069 (10)	0.0283 (6)
H8	0.1737	0.1313	0.6315	0.034*
C8A	0.0608 (4)	0.1948 (3)	0.55452 (10)	0.0243 (5)
N9	0.2174 (4)	0.1343 (3)	0.52100 (8)	0.0240 (5)
H9	0.335 (7)	0.082 (4)	0.5280 (15)	0.050 (11)*
C9A	0.1539 (4)	0.1807 (3)	0.46917 (9)	0.0233 (5)
C10	-0.1422 (5)	0.3404 (4)	0.31657 (11)	0.0343 (7)
C11	-0.3501 (8)	0.5424 (4)	0.27171 (14)	0.0588 (11)
H11A	-0.3926	0.6478	0.2821	0.071*
H11B	-0.2082	0.5483	0.2504	0.071*
C12	-0.5423 (7)	0.4765 (6)	0.23878 (18)	0.0713 (13)
H12A	-0.5770	0.5416	0.2072	0.107*
H12B	-0.6825	0.4693	0.2599	0.107*
H12C	-0.4976	0.3741	0.2269	0.107*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
O1	0.0278 (9)	0.0346 (11)	0.0253 (9)	0.0038 (8)	-0.0018 (7)	0.0018 (8)
O2	0.0752 (18)	0.139 (3)	0.0185 (10)	0.0649 (19)	0.0037 (11)	0.0088 (13)
O3	0.103 (2)	0.0541 (16)	0.0224 (10)	0.0415 (15)	-0.0067 (11)	0.0049 (10)
C1	0.0240 (11)	0.0279 (14)	0.0213 (11)	-0.0035 (10)	-0.0028 (9)	0.0015 (10)
C2	0.0282 (13)	0.0451 (18)	0.0215 (12)	0.0037 (12)	0.0013 (10)	0.0077 (11)
C3	0.0433 (16)	0.0393 (18)	0.0234 (13)	0.0103 (13)	-0.0034 (11)	-0.0001 (11)
C4	0.0259 (12)	0.0319 (15)	0.0211 (11)	0.0004 (11)	0.0000 (9)	0.0074 (10)
C4A	0.0244 (11)	0.0230 (13)	0.0209 (11)	-0.0034 (10)	-0.0032 (9)	0.0035 (9)
C5	0.0274 (12)	0.0279 (14)	0.0254 (12)	-0.0009 (10)	-0.0016 (9)	0.0029 (10)
C5A	0.0267 (12)	0.0223 (13)	0.0200 (11)	-0.0048 (10)	-0.0037 (9)	0.0025 (9)
C6	0.0334 (13)	0.0349 (16)	0.0259 (13)	0.0017 (12)	0.0033 (10)	-0.0001 (11)
C7	0.0415 (15)	0.0333 (16)	0.0193 (11)	-0.0027 (12)	-0.0009 (10)	0.0013 (11)
C8	0.0353 (13)	0.0285 (15)	0.0203 (12)	-0.0019 (11)	-0.0062 (10)	0.0020 (10)
C8A	0.0268 (12)	0.0240 (13)	0.0213 (11)	-0.0033 (10)	-0.0040 (9)	0.0007 (10)
N9	0.0278 (11)	0.0250 (12)	0.0186 (10)	0.0007 (9)	-0.0043 (8)	0.0003 (8)
C9A	0.0274 (12)	0.0237 (13)	0.0183 (11)	-0.0049 (10)	-0.0035 (9)	0.0024 (9)
C10	0.0376 (14)	0.0441 (18)	0.0207 (12)	0.0060 (13)	-0.0035 (10)	0.0014 (11)
C11	0.101 (3)	0.040 (2)	0.0337 (17)	0.013 (2)	-0.0184 (18)	0.0118 (14)
C12	0.053 (2)	0.083 (3)	0.076 (3)	-0.005 (2)	-0.022 (2)	0.041 (2)

Geometric parameters (\AA , $^\circ$)

O1—C1	1.227 (3)	C6—C5	1.375 (4)
O2—C10	1.180 (3)	C6—H6	0.9500
O3—C10	1.315 (4)	C7—C6	1.415 (4)
O3—C11	1.464 (4)	C7—C8	1.366 (4)

C1—C9A	1.438 (3)	C7—H7	0.9500
C2—C1	1.508 (3)	C8—C8A	1.400 (3)
C2—H2A	0.9900	C8—H8	0.9500
C2—H2B	0.9900	C8A—N9	1.359 (3)
C3—C2	1.539 (4)	C8A—C5A	1.423 (3)
C3—C4	1.501 (4)	N9—H9	0.82 (4)
C3—C10	1.521 (4)	C9A—C4A	1.378 (4)
C3—H3	1.0000	C9A—N9	1.378 (3)
C4—C4A	1.500 (3)	C11—C12	1.446 (6)
C4—H4A	0.9900	C11—H11A	0.9900
C4—H4B	0.9900	C11—H11B	0.9900
C5—C5A	1.399 (4)	C12—H12A	0.9800
C5—H5	0.9500	C12—H12B	0.9800
C5A—C4A	1.423 (3)	C12—H12C	0.9800
C10—O3—C11	116.7 (3)	C5—C6—H6	119.6
O1—C1—C2	121.3 (2)	C7—C6—H6	119.6
O1—C1—C9A	124.2 (2)	C6—C7—H7	119.3
C9A—C1—C2	114.5 (2)	C8—C7—C6	121.5 (2)
C1—C2—C3	115.5 (2)	C8—C7—H7	119.3
C1—C2—H2A	108.4	C7—C8—C8A	118.0 (2)
C1—C2—H2B	108.4	C7—C8—H8	121.0
C3—C2—H2A	108.4	C8A—C8—H8	121.0
C3—C2—H2B	108.4	N9—C8A—C5A	108.9 (2)
H2A—C2—H2B	107.5	N9—C8A—C8	129.8 (2)
C2—C3—H3	106.4	C8—C8A—C5A	121.4 (2)
C4—C3—C2	114.9 (2)	C8A—N9—C9A	108.1 (2)
C4—C3—C10	114.8 (2)	C8A—N9—H9	130 (3)
C4—C3—H3	106.4	C9A—N9—H9	122 (3)
C10—C3—C2	107.3 (2)	N9—C9A—C1	125.0 (2)
C10—C3—H3	106.4	C4A—C9A—N9	110.1 (2)
C3—C4—H4A	109.7	C4A—C9A—C1	124.8 (2)
C3—C4—H4B	109.7	O2—C10—O3	123.2 (3)
C4A—C4—C3	110.0 (2)	O2—C10—C3	123.6 (3)
C4A—C4—H4A	109.7	O3—C10—C3	113.3 (2)
C4A—C4—H4B	109.7	O3—C11—H11A	109.3
H4A—C4—H4B	108.2	O3—C11—H11B	109.3
C5A—C4A—C4	130.1 (2)	C12—C11—O3	111.7 (3)
C9A—C4A—C4	123.2 (2)	C12—C11—H11A	109.3
C9A—C4A—C5A	106.7 (2)	C12—C11—H11B	109.3
C5A—C5—H5	120.4	H11A—C11—H11B	107.9
C6—C5—C5A	119.2 (2)	C11—C12—H12A	109.5
C6—C5—H5	120.4	C11—C12—H12B	109.5
C4A—C5A—C8A	106.1 (2)	C11—C12—H12C	109.5
C5—C5A—C4A	134.8 (2)	H12A—C12—H12B	109.5
C5—C5A—C8A	119.1 (2)	H12A—C12—H12C	109.5
C5—C6—C7	120.8 (3)	H12B—C12—H12C	109.5
C11—O3—C10—O2	-5.3 (5)	C5—C5A—C4A—C4	5.2 (5)
C11—O3—C10—C3	175.9 (3)	C5—C5A—C4A—C9A	-177.5 (3)

supplementary materials


C10—O3—C11—C12	88.6 (5)	C8A—C5A—C4A—C4	-176.3 (3)
O1—C1—C9A—N9	-0.6 (4)	C8A—C5A—C4A—C9A	1.0 (3)
O1—C1—C9A—C4A	179.6 (2)	C7—C6—C5—C5A	0.9 (4)
C2—C1—C9A—N9	-176.9 (2)	C8—C7—C6—C5	-2.0 (4)
C2—C1—C9A—C4A	3.2 (4)	C6—C7—C8—C8A	0.2 (4)
C3—C2—C1—O1	158.9 (3)	C7—C8—C8A—N9	-178.9 (3)
C3—C2—C1—C9A	-24.5 (4)	C7—C8—C8A—C5A	2.7 (4)
C4—C3—C2—C1	46.9 (4)	N9—C8A—C5A—C4A	-1.2 (3)
C10—C3—C2—C1	175.9 (2)	N9—C8A—C5A—C5	177.5 (2)
C2—C3—C4—C4A	-44.0 (3)	C8—C8A—C5A—C4A	177.5 (2)
C10—C3—C4—C4A	-169.1 (2)	C8—C8A—C5A—C5	-3.7 (4)
C2—C3—C10—O2	36.0 (5)	C8—C8A—N9—C9A	-177.6 (3)
C2—C3—C10—O3	-145.3 (3)	C5A—C8A—N9—C9A	1.0 (3)
C4—C3—C10—O2	165.0 (3)	C1—C9A—N9—C8A	179.8 (2)
C4—C3—C10—O3	-16.3 (4)	C4A—C9A—N9—C8A	-0.4 (3)
C3—C4—C4A—C5A	-159.6 (3)	N9—C9A—C4A—C4	177.1 (2)
C3—C4—C4A—C9A	23.5 (4)	N9—C9A—C4A—C5A	-0.4 (3)
C6—C5—C5A—C4A	-179.8 (3)	C1—C9A—C4A—C4	-3.1 (4)
C6—C5—C5A—C8A	1.8 (4)	C1—C9A—C4A—C5A	179.4 (2)

Hydrogen-bond geometry (\AA , $^\circ$)

$D\text{—H}\cdots A$	$D\text{—H}$	$H\cdots A$	$D\cdots A$	$D\text{—H}\cdots A$
N9—H9 \cdots O1 ⁱ	0.82 (4)	2.08 (4)	2.834 (3)	154 (4)
C4—H4A \cdots Cg3 ⁱⁱ	0.99	2.75	3.727 (3)	171

Symmetry codes: (i) $-x+1, -y, -z+1$; (ii) $-x, -y+2, -z$.

Fig. 1



supplementary materials

Fig. 2

