

Impact of White Matter Lesions on Depression in the Patients with Alzheimer's Disease

Jung Jae Lee¹, Eun Young Lee¹, Seok Bum Lee¹, Joon Hyuk Park², Tae Hui Kim³, Hyun-Ghang Jeong⁴, Jae Hyoung Kim⁵, Ji Won Han⁶, and Ki Woong Kim^{6,7,8} ✉

¹Department of Psychiatry, Dankook University College of Medicine, Cheonan, Republic of Korea

²Department of Psychiatry, Jeju National University School of Medicine, Jeju National University Hospital, Jeju, Republic of Korea

³Department of Psychiatry, Yonsei University Wonju Severance Christian Hospital, Wonju, Republic of Korea

⁴Department of Psychiatry, Korea University Guro Hospital, Korea University College of Medicine, Seoul, Republic of Korea

⁵Department of Radiology, Seoul National University Bundang Hospital, Seongnam, Republic of Korea

⁶Department of Neuropsychiatry, Seoul National University Bundang Hospital, Seongnam, Republic of Korea

⁷Department of Psychiatry, Seoul National University College of Medicine, Seoul, Republic of Korea

⁸Department of Brain and Cognitive Science, Seoul National University College of Natural Sciences, Seoul, Republic of Korea

Objective Comorbid depression is common in patients with Alzheimer's disease (AD). An increase in white matter lesions (WMLs) has been associated with depression in both elderly individuals with normal cognition and patients with Alzheimer's disease. We investigated whether the severity and location of WMLs influence the association between WMLs and comorbid depression in AD.

Methods We enrolled 93 AD patients from Seoul National University Bundang Hospital. We administered both the Mini International Neuropsychiatric Inventory (MINI) and the Korean version of the Consortium to Establish a Registry for Alzheimer's Disease Assessment Packet (CERAD-K) clinical and neuropsychological battery. Subjects also underwent brain magnetic resonance imaging (MRI). We diagnosed AD according to the criteria of the National Institute of Neurological and Communicative Disorders and Stroke and the Alzheimer's Disease and Related Disorders Association. We diagnosed depressive disorders according to the DSM-IV diagnostic criteria, and evaluated the severity of depressive symptoms using the Korean version of the Geriatric Depression Scale (GDS-K). We quantified the WML volumes from the brain MRI using a fully automated segmentation algorithm.

Results The log of the WML volume in the frontal lobe was significantly associated with depressive disorders (odds ratio=1.905, 95% CI=1.027-3.533, $p=0.041$), but not with the severity of depressive symptoms as measured by the GDS-K.

Conclusion The WML volume in the frontal lobe conferred a risk of comorbid depressive disorders in AD, which implies that comorbid depression in AD may be attributed to vascular causes.

Psychiatry Investig 2015;12(4):516-522

Key Words Alzheimer's disease, White matter lesions, Depression.

INTRODUCTION

White matter lesions (WMLs) are commonly found in the T2-weighted magnetic resonance imaging (MRI) or fluid-attenuated inversion recovery (FLAIR) scans of normal elderly people.¹ WMLs also have been found in Alzheimer's disease (AD),² vascular dementia,^{3,4} and depression,⁵⁻⁷ and have been

associated with vascular risk factors such as hypertension and diabetes.^{8,9} Neuropathologically, WMLs are characterized by the demyelination of white matter axons, mild fibrillary gliosis, and fibrohyaline thickening of small vessels that supply the white matter.^{10,11} In particular, irregular periventricular WMLs and deep WMLs have been associated with microcystic infarcts and patchy rarefaction of the myelin, which are ischemic in nature.^{12,13} These findings support a prominent role of vascular factors in the pathogenesis of WMLs.

Pathological evidence suggests that WMLs may play a role in the clinical symptoms of AD. Individuals with more WMLs have a higher risk of developing AD.^{14,15} In addition, WMLs are associated with cognitive dysfunction and the severity of the dementia.^{16,17} However, the relationship between depression and WMLs in AD remains unclear. Several studies have

Received: April 14, 2015 Revised: May 16, 2015

Accepted: May 16, 2015 Available online: September 30, 2015

✉ Correspondence: Ki Woong Kim, MD, PhD

Department of Neuropsychiatry, Seoul National University Bundang Hospital, 166 Gumi-ro, Bundang-gu, Seongnam 463-707, Republic of Korea
Tel: +82-31-787-7432, Fax: +82-31-787-4058, E-mail: kwkimmd@snu.ac.kr

© This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

explored the relationship between WMLs and depression. WMLs, particularly in the frontal area, have been found to be associated with higher depression scores.^{18,20} Other studies, however, did not find such a relationship.²¹⁻²³ The heterogeneity of the study populations, the methods by which depression was assessed, and the measurements of the WML volumes might contribute to these inconsistent results. In most studies, different semi-quantitative visual rating scales, such as Faze-ka's scale and Schelton's scale, were used.^{18,19,21-23} In one study, an automated volumetric method was used to assess WML volumes, but depression was assessed by rating scales rather than by clinical diagnosis.²⁰ In other studies, depression was assessed by clinical diagnosis, but WMLs were assessed by visual rating scales.^{21,22}

The present study avoided some of the limitations of these previous studies by 1) using automated quantitative volumetric methods to assess the severity and location of WMLs; 2) using both a structured clinical interview and a self-report questionnaire to diagnose the presence and severity of depression; and 3) recruiting only AD patients to avoid a confounding effect from a heterogeneous study population.

The purpose of this study was to explore the relationship between depression and the severity and location of WMLs in patients with AD. We tested the hypothesis that larger WML volumes and frontal WMLs in particular are associated with depressive mood and clinician-diagnosed depression in patients with AD.

METHODS

Subjects

We recruited AD patients from a dementia clinic (Seoul National University Bundang Hospital, SNUBH) and communities [the Korean Longitudinal Study on Health and Aging (KLoSHA)²⁴ and the nationwide dementia screening and registry program (DSRP)]. All subjects underwent standardized clinical interviews and neurological and physical examinations, which were administered by four geriatric psychiatrists with advanced training in neuropsychiatry and dementia research according to the protocol of the Korean version of the Consortium to Establish a Registry for Alzheimer's Disease clinical assessment battery (CERAD-K),²⁵ the standardized TIA/stroke form from the Atherosclerosis Risk in Communities Study (ARIC),²⁶ and the Mini International Neuropsychiatric Interview (MINI),²⁷ to diagnose dementia, stroke, and major psychiatric disorders, respectively.

Dementia was diagnosed in these participants in accordance with the Diagnostic and Statistical Manual of Mental Disorders, Fourth Edition (DSM-IV) criteria,²⁸ and probable or possible AD was diagnosed according to the National Insti-

tute of Neurological and Communicative Disorders and Stroke and the Alzheimer's Disease and Related Disorders Association (NINCDS-ADRDA) criteria.²⁹ A total of 93 subjects met the inclusion criteria. Sixty-six (71%) subjects had probable AD and 27 (29%) had possible AD.

All the participants were fully informed regarding their study participation and either the participant or a legal guardian provided written informed consent. The study protocol was approved by the Institutional Review Board of SNUBH.

Assessment of depression

Depression was evaluated based on information provided from the patients as well as information provided by their primary caregiver. First, we administered a self-report questionnaire, the Korean version of the Geriatric Depression Scale (GDS-K), to assess subjective depressive mood.³⁰ Second, research-trained psychiatrists interviewed the patients and(or) primary caregivers using the MINI to clinically diagnose depression.²⁷ The severity of depression was rated from very mild (subsyndromal depression) to moderate-to-severe depression (e.g., minor and major depressive disorder, dysthymic disorder).

Major depressive disorder (MDD) and dysthymic disorder were diagnosed according to the DSM-IV criteria, and minor depressive disorder (MnDD) was diagnosed according to the research criteria proposed in the DSM-IV. Subsyndromal depression (SSD) was diagnosed according to the following operational criteria: 1) two or more symptoms listed in the DSM-IV criterion A for a major depressive episode occurring during the same two-week period; 2) at least one symptom should be depressed mood or anhedonia; 3) each depressive symptom should be present for more than half of one day or more than seven days during the 2 weeks; and 4) symptoms do not meet the criteria for a diagnosis of MDD or MnDD. Subjects were divided into two diagnostic groups: 1) Depression, which was defined as evidence of MDD, MnDD, dysthymic disorder, or SSD or 2) No depression, which was defined as no current evidence of significant depressed mood.

Assessment of neuropsychological functions

Global cognitive function was evaluated using the Mini-Mental Status Examination (MMSE).^{25,31} To assess memory function, the word list memory test (WLMT), the word list recall test (WLRT), and the word list recognition test (WLRcT) from the Korean version of the CERAD Neuropsychological Assessment Battery (CERAD-K-N) were administered. Language function was evaluated using the verbal fluency test (VFT) and the 15-item modified Boston Naming Test (mBNT) from the CERAD-K-N.^{25,31} Visuospatial function was evaluated using the constructional praxis test (CPT) from the CERAD-K-N.^{25,31} Frontal function was evaluated using the trail making test-A/

B (TMT-A/B) from the CERAD-K-N.^{25,31} All the assessments were performed at SNUBH

Assessment of WMLs

T1-weighted MR and FLAIR images were acquired on a 3.0 Tesla MRI scanner system (Achieva, Philips Medical System, The Netherlands) at SNUBH. The T1 images were acquired using the following protocol: magnetic field=3.0 Tesla; sequence=SPGR; repetition time/echo time=22/4 msec; number of excitations=1; flip angle=40°; field of view=240 mm; plane=sagittal; matrix=256×256 mm; thickness=1 mm; inter-slice gap=none; slice number=120. The FLAIR images were acquired using the following protocol: magnetic field=3.0 Tesla; repetition time/echo time=9900/160 msec; number of excitations=1; flip angle=90°; field of view=240 mm; axial plane matrix=256×256 mm; thickness=3 mm; and inter-slice gap=none.

All the processes were implemented in Matlab 2010a (The MathWorks, Inc., Natick, MA, USA). WMLs were automatically segmented from the FLAIR images according to the method adopted by Yoo et al.³² A study-specific T1 template (SST) was constructed for accurate spatial normalization using the DARTEL toolbox in SPM8 (Wellcome Trust Centre for Neuroimaging, London, UK; <http://www.fil.ion.ucl.ac.uk/spm>). All the T1 images, as well as the FLAIR and segmented WML images, were spatially transformed to the SST.

To define the regions to be classified as WMLs, a mask-labeled region was manually delineated from the MNI standard space by one neurological expert. Frontal, temporal, parietal, and occipital WML regions were defined. The MNI152 standard T1 image as well as the mask-labeled region was spatially transformed to the SST. Sub-classified WML images were obtained by applying the transformed mask to the segmented WML images through voxel-wise image multiplication for each individual WMH image. The sub-classified WMH volumes then were computed by counting the voxels from the labeled WML images.

Statistical analysis

We computed descriptive statistics on the demographic variables and disease-related measures using independent t-tests for each continuous variable and chi-square tests for each categorical variable to examine the differences between the groups. The neuropsychological measures between the groups were compared by an analysis of variance with adjustments for age, education, and GDS-K scores, which are well-established risk factors for cognitive dysfunction.^{31,33,34} We used a logarithmic transformation to produce normally distributed WML volumes. The associations between the log-transformed WML volumes and clinical variables were examined using

Pearson's correlation coefficient or Spearman's rank-order correlation coefficient (r) as appropriate. We also performed multivariate logistic regression analyses. First, we conducted unadjusted analyses examining the relationship between log WML volumes and the dependent variables. Second, we included two sets of adjustment variables: 1) those characteristics well-known to influence depression, i.e., age, gender, and level of education^{35,36} and 2) vascular risk factors (e.g., cardiovascular disease, hypertension, diabetes)³⁷ and prior depression history.^{38,39} Due to the high correlation between the total WML volume and regional WML volumes, separate models were generated to analyze the relationship between depression and WML volume by location. For all the analyses, a two-tailed p-value of less than 0.05 was considered statistically significant. All the analyses were conducted using SPSS version 12.0 (SPSS Inc., Chicago, IL, USA).

RESULTS

The clinical and demographic characteristics of the participants are summarized in Table 1. Of the 93 subjects, 25 were diagnosed with depression and 68 were assigned to the no depression group. No differences between the groups were observed on any clinical or demographic variables except GDS-K scores. The neuropsychological performance of the participants is summarized in Table 2. No differences were observed between the depression and no depression groups.

We performed multiple logistic regressions to assess the relationship between clinician-diagnosed depression and the total and regional WML volumes with adjustments for potential confounding effects of vascular risk factors and residual clinical variables. The total log WML volume was not significantly associated with the presence of clinician-diagnosed depression (odds ratio 1.820, 95% CI 0.962–3.443, $p=0.066$). When the WML volumes were analyzed separately with respect to location, the log frontal WML volume was significantly associated with the presence of clinician-diagnosed depression (odds ratio 1.905, 95% CI=1.027–3.533, $p=0.041$), but no significant relationship was observed in any other WML region (Table 3).

Table 4 shows the relationship between the GDS-K scores as well as other clinical variables and the log total or regional WML volumes. The GDS-K scores were not associated with either the log total WML volume ($p=0.369$, Pearson's chi-squared test) or the log frontal WML volume ($p=0.410$, Pearson's chi-squared test).

DISCUSSION

Depression has been recognized as one of the most fre-

Table 1. Sociodemographic characteristics of the study subjects

	Depression (N=25)	No depression (N=68)	p-value*
Age (years), mean±SD	76.12±7.61	77.63±7.09	0.373
Female, N (%)	18 (72.0)	49 (72.1)	0.996
Education (years), mean±SD	8.16±6.14	7.93±5.60	0.863
CDR-SOB	3.94±2.79	4.45±3.30	0.464
GDS-K, mean±SD	17.56±6.25	11.07±5.26	<0.001
Cardiovascular disease, N (%)	2 (8.0)	7 (10.3)	0.740
Hypertension, N (%)	13 (52.0)	36 (52.9)	0.936
Diabetes, N (%)	6 (24.0)	14 (20.6)	0.723
Head trauma, N (%)	3 (12.0)	6 (8.8)	0.646
Kidney disease, N (%)	0 (0.0)	4 (5.9)	0.215
Prior depression, N (%)	7 (28.0)	17 (25.0)	0.769

*t-tests for continuous variables and chi-square tests for categorical variables. GDS-K: Korean version of the Geriatric Depression Scale, CDR-SOB: Clinical Dementia Rating Scale-Sum of Boxes

Table 2. Neurocognitive function of the participants according to their depression diagnosis

	Depression N=25	No depression N=68	p-value*
Word Fluency	8.3±3.2	8.4±3.7	0.787
Boston Naming	8.0±3.0	7.7±3.2	0.892
MMSE-KC	17.9±5.1	16.6±5.5	0.348
Word List Memory	9.3±3.0	8.5±4.0	0.491
Constructional Praxis	7.7±2.5	8.1±2.3	0.304
Word Delayed Recall	1.6±1.5	1.2±1.6	0.429
Word Recognition	5.0±3.2	4.5±2.8	0.502
Constructional Recall	1.3±1.7	1.3±2.1	0.747
Trail Making A	173.7±106.4	171.0±111.6	0.455
Trail Making B	280.2±34.2	277.2±53.4	0.296

*ANOVA adjusted for age, sex, and education. MMSE-KC: Korean version of the Mini-Mental Status Examination

Table 3. Multiple logistic regression analysis of the association between clinician-diagnosed depression and the log of the white matter lesion volume

	Log WML	Odds ratio (95% CI)	p-value
Model 1	Total	1.820 (0.962–3.443)	0.066
Model 2	Frontal	1.905 (1.027–3.533)	0.041
Model 3	Temporal	1.611 (0.962–2.699)	0.070
Model 4	Parietal	1.146 (0.850–1.544)	0.797
Model 5	Occipital	1.092 (0.754–1.581)	0.643

Each model adjusted for age, sex, education, cardiovascular disease, hypertension, diabetes, and prior depression. WML: white matter lesion volume (mL)

quent psychiatric comorbidities of AD, and previous studies have suggested that 20% to 50% of patients with AD suffer from depression.^{40–42} The importance of depression in AD is underscored by its negative impact on the patient's quality of life,⁴³ cognitive ability,⁴⁴ and activities of daily living⁴⁵ as well as the caregiver's stress and psychological comorbidity.⁴⁶ Short-term mortality in elderly subjects with less severe dementia

Table 4. Correlations between the Geriatric Depression Scale and the log white matter lesion volume and other sociodemographic factors

Factors	Correlation coefficient		p-value
	Pearson (r)	Spearman (ρ)	
Age	-0.099		0.346
Gender	-0.124		0.235
Education	-0.174		0.096
MMSE-KC	-0.170		0.112
Cardiovascular disease		0.018	0.862
Hypertension		-0.161	0.124
Diabetes		0.068	0.515
Prior depression		0.259	0.012
Log WML			
Total	0.094		0.369
Frontal	0.086		0.410
Temporal	0.072		0.495
Parietal	0.034		0.749
Occipital	0.091		0.387

WML: white matter lesion volume (mL), MMSE-KC: Korean version of the Mini-Mental Status Examination

is predicted by the presence of subsyndromal depression and mood symptoms.⁴⁷ Identifying and treating patients with depression in AD may reverse this associated morbidity.

The etiologies and pathophysiology of depression in AD remains unclear. Depression can be organic in nature due to cerebrovascular changes.^{5,48} The neurodegenerative process may directly contribute to the development of depression. Several studies have found that depression has been associated with the loss of noradrenergic or serotonergic nuclei^{49,50} or as a consequence of low-grade inflammation in the brain.⁵¹

In our study, the WML volume, especially the frontal WML volume, was positively and independently correlated with depression after adjusting for vascular risk factors. This result is consistent with some previous studies.^{18–20} The WMLs are a

surrogate marker of small-vessel vascular disease resulting from ischemic damage due to chronic hypoperfusion, whereas mild periventricular WMLs are non-ischemic in origin. Recently, Brickman et al.⁵² reported that reduced cerebral autoregulation was associated with an increased WML volume. There is growing evidence that frontal and subcortical WMLs are implicated in late-life depression and influence the outcome, possibly via the disruption of frontal-subcortical circuits.^{2,53,54} These findings, together with those in the present study, suggest that vascular depression can be considered an etiology of depression in AD. WMLs may affect mood symptoms via the disruption of intracerebral connectivity, compromising efficient neuronal communication. The prefrontal cortex and its extensive cortical-cortical and cortical-subcortical connectivity is thought to play a central role in motivation-based depressive symptoms in dementia.⁵⁵

The results from previous studies have been inconsistent with respect to the relationship between WMLs and depression in AD. Several factors may contribute to these inconsistencies.

First, different visual rating scales were used across studies,^{18,19,21-23} and therefore scores from different rating scales are not directly comparable. Different visual rating scales can produce inconsistent results among neuroimaging studies.⁵⁶ Furthermore, visual rating scales are less sensitive than volumetric methods.⁵⁷ Soennesyn et al.²⁰ used both automated volumetric WML measurement and a visual rating scale, and reported that the automated volumetric measure of the total and frontal WMLs but not the visual rating scale scores were associated with the severity of depressive symptoms. This finding suggests that visual rating scales inevitably have some limitations such as data nonlinearity, lack of sensitivity to small changes, and susceptibility to ceiling effects.^{8,58,59}

Second, the methods to assess depression were different across studies. Diagnosing depression in AD remains problematic for several reasons. First, the symptoms of depression in AD overlap with other neuropsychiatric manifestations of dementia. The use of rating scales in a patient with dementia without consideration of this overlap in nonspecific depressive symptoms may result in an erroneous assignment of a depression diagnosis in AD.⁶⁰ Second, another complicating factor in the diagnosis of depression in AD is the patient's limited ability to communicate distress coherently. Patients with dementia cannot always accurately report their own symptoms, and therefore may underreport their symptoms.⁶¹ AD patients may also perceive themselves as less depressed than do their caregivers or a clinician.⁶² Third, depressive symptoms in patients with AD differ from those in depressed but otherwise cognitively normal elderly patients. Patients with AD appear to have more motivational symptoms and delusions and

less commonly experience sadness, sleep disturbance, and appetite loss.^{41,63,64} Fourth, there are currently no standardized methods to diagnose depression in AD. The recommendation for diagnosing depression in AD is to use a structured psychiatric interview; depression scales should be used to rate the severity of the depression but not to diagnose. In our study, we used both a self-rating depression questionnaire and a structured psychiatric interview to assess depression. Interestingly, we found a significant association between clinician-diagnosed depression and WML volumes but failed to show such a relationship between self-reported depression scores and WML volumes.

The heterogeneity of the study populations may also contribute to the inconsistent results across studies. Some studies included only patients with AD,^{21,22} but other studies included patients with other types of dementia such as vascular dementia, dementia with Lewy bodies, and frontotemporal dementia as well as AD.^{18-20,23}

The prevention or reduction of WMHs could lessen the burden of depression and depressive symptoms in patients with dementia. The presence of WMLs, in particular in the frontal area, is associated with depression in AD, which implies a common pathophysiology of depression caused by disruption to the frontal-subcortical circuits. Understanding the pathogenesis is important and may offer important insights into therapeutic options for the treatment and prevention of depression in AD.

To our knowledge, this is the first study to investigate the association between depression and WMLs in AD using a standardized structured psychiatric interview to assess depression and an automated quantitative measurement to assess the severity and location of WMLs. However, this study was cross-sectional in design and the depression group was relatively small, which may warrant a prospective study on a larger sample to confirm the positive correlation between WMLs in AD and depression and investigate its clinical significance.

Acknowledgments

This study was supported by grant no. 04-2008-001 from the SNUBH Research Fund and by a grant from the Korean Health Technology R&D Project, Ministry of Health and Welfare, Republic of Korea [grant number HI09C1379 (A092077)].

REFERENCES

- Schmidt R, Fazekas F, Offenbacher H, Dusek T, Zach E, Reinhart B, et al. Neuropsychologic correlates of MRI white matter hyperintensities: a study of 150 normal volunteers. *Neurology* 1993;43:2490-2494.
- O'Brien J, Desmond P, Ames D, Schweitzer I, Harrigan S, Tress B. A magnetic resonance imaging study of white matter lesions in depression and Alzheimer's disease. *Br J Psychiatry* 1996;168:477-485.
- Almkvist O, Wahlund LO, Andersson-Lundman G, Basun H, Backman L. White-matter hyperintensity and neuropsychological functions in dementia and healthy aging. *Arch Neurol* 1992;49:626-632.

4. Schmidt R. Comparison of magnetic resonance imaging in Alzheimer's disease, vascular dementia and normal aging. *Eur Neurol* 1992;32:164-169.
5. Alexopoulos GS, Meyers BS, Young RC, Campbell S, Silbersweig D, Charlson M. 'Vascular depression' hypothesis. *Arch Gen Psychiatry* 1997;54:915-922.
6. Krishnan KR, Hays JC, Blazer DG. MRI-defined vascular depression. *Am J Psychiatry* 1997;154:497-501.
7. Lesser IM, Hill-Gutierrez E, Miller BL, Boone KB. Late-onset depression with white matter lesions. *Psychosomatics* 1993;34:364-367.
8. Longstreth WT Jr, Manolio TA, Arnold A, Burke GL, Bryan N, Jungreis CA, et al. Clinical correlates of white matter findings on cranial magnetic resonance imaging of 3301 elderly people. The Cardiovascular Health Study. *Stroke* 1996;27:1274-1282.
9. de Leeuw FE, de Groot JC, Oudkerk M, Witteman JC, Hofman A, van Gijn J, et al. Hypertension and cerebral white matter lesions in a prospective cohort study. *Brain* 2002;125:765-772.
10. Englund E. Neuropathology of white matter changes in Alzheimer's disease and vascular dementia. *Dement Geriatr Cogn Disord* 1998;9(Suppl 1):6-12.
11. Ogata J. Vascular dementia: the role of changes in the vessels. *Alzheimer Dis Assoc Disord* 1999;13(Suppl 3):S55-S58.
12. Thomas AJ, O'Brien JT, Barber R, McMeekin W, Perry R. A neuropathological study of periventricular white matter hyperintensities in major depression. *J Affect Disord* 2003;76:49-54.
13. van Swieten JC, van den Hout JH, van Ketel BA, Hijdra A, Wokke JH van Gijn J. Periventricular lesions in the white matter on magnetic resonance imaging in the elderly. A morphometric correlation with arteriosclerosis and dilated perivascular spaces. *Brain* 1991;114(Pt 2):761-774.
14. Prins ND, van Dijk EJ, den Heijer T, Vermeer SE, Koudstaal PJ, Oudkerk M, et al. Cerebral white matter lesions and the risk of dementia. *Arch Neurol* 2004;61:1531-1534.
15. Vermeer SE, Prins ND, den Heijer T, Hofman A, Koudstaal PJ, Breteler MM. Silent brain infarcts and the risk of dementia and cognitive decline. *N Engl J Med* 2003;348:1215-1222.
16. Stout JC, Jernigan TL, Archibald SL, Salmon DP. Association of dementia severity with cortical gray matter and abnormal white matter volumes in dementia of the Alzheimer type. *Arch Neurol* 1996;53:742-749.
17. Diaz JF, Merskey H, Hachinski VC, Lee DH, Boniferno M, Wong CJ, et al. Improved recognition of leukoaraiosis and cognitive impairment in Alzheimer's disease. *Arch Neurol* 1991;48:1022-1025.
18. Barber R, Scheltens P, Gholkar A, Ballard C, McKeith I, Ince P, et al. White matter lesions on magnetic resonance imaging in dementia with Lewy bodies, Alzheimer's disease, vascular dementia, and normal aging. *J Neurol Neurosurg Psychiatry* 1999;67:66-72.
19. O'Brien J, Perry R, Barber R, Gholkar A, Thomas A. The association between white matter lesions on magnetic resonance imaging and non-cognitive symptoms. *Ann N Y Acad Sci* 2000;903:482-489.
20. Soennesyn H, Oppedal K, Greve OJ, Fritze F, Auestad BH, Nore SP, et al. White matter hyperintensities and the course of depressive symptoms in elderly people with mild dementia. *Dement Geriatr Cogn Dis Extra* 2012;2:97-111.
21. Clark LM, McDonald WM, Welsh-Bohmer KA, Siegler IC, Dawson DV, Tupler LA, et al. Magnetic resonance imaging correlates of depression in early- and late-onset Alzheimer's disease. *Biol Psychiatry* 1998;44:592-599.
22. Harrell LE, Duvall E, Folks DG, Duke L, Bartolucci A, Conboy T, et al. The relationship of high-intensity signals on magnetic resonance images to cognitive and psychiatric state in Alzheimer's disease. *Arch Neurol* 1991;48:1136-1140.
23. Lind K, Jonsson M, Karlsson I, Sjogren M, Wallin A, Edman A. Depressive symptoms and white matter changes in patients with dementia. *Int J Geriatr Psychiatry* 2006;21:119-125.
24. Park JH, Lim S, Lim JY, Han MK, Yoon IY, Kim JM, et al. An overview of the Korean longitudinal study on health and aging. *Psychiatry Investig* 2007;4:84-95.
25. Lee JH, Lee KU, Lee DY, Kim KW, Jhoo JH, Kim JH, et al. Development of the Korean version of the Consortium to Establish a Registry for Alzheimer's Disease Assessment Packet (CERAD-K): clinical and neuropsychological assessment batteries. *J Gerontol B Psychol Sci Soc Sci* 2002;57:47-53.
26. The Atherosclerosis Risk in Communities (ARIC) Study: design and objectives. The ARIC investigators. *Am J Epidemiol* 1989;129:687-702.
27. Yoo SW, Kim YS, Noh JS, Oh KS, Kim CH, Namkoong K. Validity of Korean version of the Mini-International Neuropsychiatric Interview. *Anxiety Mood* 2006;2:50-55.
28. American Psychiatric Association. *Diagnostic and Statistical Manual of Mental Disorders, Fourth Edition Revised (DSM-IV)*. Washington, DC: American Psychiatric Association; 1994.
29. McKhann G, Drachman D, Folstein M, Katzman R, Price D, Stadlan EM. Clinical diagnosis of Alzheimer's disease: report of the NINCDS-ADRDA Work Group under the auspices of Department of Health and Human Services Task Force on Alzheimer's Disease. *Neurology* 1984;34:939-944.
30. Cho MJ, Bae JN, Suh GH, Hahm BJ, Kim JK, Lee DW, et al. Validation of Geriatric Depression Scale, Korean Version (GDS) in the Assessment of DSM-III-R Major depression. *J Korean Neuropsychiatr Assoc* 1999;31:48-63.
31. Lee DY, Lee KU, Lee JH, Kim KW, Jhoo JH, Kim SY, et al. A normative study of the CERAD neuropsychological assessment battery in the Korean elderly. *J Int Neuropsychol Soc* 2004;10:72-81.
32. Yoo BI, Lee JJ, Han JW, Oh SY, Lee EY, MacFall JR, et al. Application of variable threshold intensity to segmentation for white matter hyperintensities in fluid attenuated inversion recovery magnetic resonance images. *Neuroradiology* 2014;56:265-281.
33. Lichtenberg PA, Ross T, Millis SR, Manning CA. The relationship between depression and cognition in older adults: a cross-validation study. *J Gerontol B Psychol Sci Soc Sci* 1995;50:25-32.
34. Wilson RS, Barnes LL, Mendes de Leon CF, Aggarwal NT, Schneider JS, Bach J, et al. Depressive symptoms, cognitive decline, and risk of AD in older persons. *Neurology* 2002;59:364-370.
35. Park JH, Kim KW, Kim MH, Kim MD, Kim BJ, Kim SK, et al. A nationwide survey on the prevalence and risk factors of late life depression in South Korea. *J Affect Disord* 2012;138:34-40.
36. Park JH, Lee JJ, Lee SB, Huh Y, Choi EA, Youn JC, et al. Prevalence of major depressive disorder and minor depressive disorder in an elderly Korean population: results from the Korean Longitudinal Study on Health and Aging (KLoSHA). *J Affect Disord* 2010;125:234-240.
37. Djernes JK. Prevalence and predictors of depression in populations of elderly: a review. *Acta Psychiatr Scand* 2006;113:372-387.
38. Alexopoulos GS. Depression in the elderly. *Lancet* 2005;365:1961-1970.
39. Teodorczuk A, O'Brien JT, Firbank MJ, Pantoni L, Poggesi A, Erkinjuntti T, et al. White matter changes and late-life depressive symptoms: longitudinal study. *Br J Psychiatry* 2007;191:212-217.
40. Lyketsos CG, Steinberg M, Tschanz JT, Norton MC, Steffens DC, Breitner JC. Mental and behavioral disturbances in dementia: findings from the Cache County Study on Memory in Aging. *Am J Psychiatry* 2000;157:708-714.
41. Olin JT, Katz IR, Meyers BS, Schneider LS, Lebowitz BD. Provisional diagnostic criteria for depression of Alzheimer disease: rationale and background. *Am J Geriatr Psychiatry* 2002;10:129-141.
42. Olin JT, Schneider LS, Katz IR, Meyers BS, Alexopoulos GS, Breitner JC, et al. Provisional diagnostic criteria for depression of Alzheimer disease. *Am J Geriatr Psychiatry* 2002;10:125-128.
43. Gonzalez-Salvador T, Lyketsos CG, Baker A, Hovanec L, Roques C, Brandt J, et al. Quality of life in dementia patients in long-term care. *Int J Geriatr Psychiatry* 2000;15:181-189.
44. Bassuk SS, Berkman LF, Wypij D. Depressive symptomatology and in-

- cident cognitive decline in an elderly community sample. *Arch Gen Psychiatry* 1998;55:1073-1081.
45. Lyketsos CG, Steele C, Baker L, Galik E, Kopunek S, Steinberg M, et al. Major and minor depression in Alzheimer's disease: prevalence and impact. *J Neuropsychiatry Clin Neurosci* 1997;9:556-561.
 46. Gonzalez-Salvador MT, Arango C, Lyketsos CG, Barba AC. The stress and psychological morbidity of the Alzheimer patient caregiver. *Int J Geriatr Psychiatry* 1999;14:701-710.
 47. Janzing JG, Bouwens JM, Teunisse RJ, Van't Hof MA, Zitman FG. The relationship between depression and mortality in elderly subjects with less severe dementia. *Psychol Med* 1999;29:979-983.
 48. O'Brien JT, Ames D. White matter lesions in depression and Alzheimer's disease. *Br J Psychiatry* 1996;169:671.
 49. Forstl H, Burns A, Luthert P, Cairns N, Lantos P, Levy R. Clinical and neuropathological correlates of depression in Alzheimer's disease. *Psychol Med* 1992;22:877-884.
 50. Zweig RM, Ross CA, Hedreen JC, Steele C, Cardillo JE, Whitehouse PJ, et al. The neuropathology of aminergic nuclei in Alzheimer's disease. *Ann Neurol* 1988;24:233-242.
 51. Leonard BE. Inflammation, depression and dementia: are they connected? *Neurochem Res* 2007;32:1749-1756.
 52. Brickman AM, Guzman VA, Gonzalez-Castellon M, Razlighi Q, Gu Y, Narkhede A, et al. Cerebral autoregulation, beta amyloid, and white matter hyperintensities are interrelated. *Neurosci Lett* 2015;592:54-58.
 53. Hickie I, Scott E, Wilhelm K, Brodaty H. Subcortical hyperintensities on magnetic resonance imaging in patients with severe depression--a longitudinal evaluation. *Biol Psychiatry* 1997;42:367-374.
 54. Greenwald BS, Kramer-Ginsberg E, Krishnan KR, Ashtari M, Auerbach C, Patel M. Neuroanatomic localization of magnetic resonance imaging signal hyperintensities in geriatric depression. *Stroke* 1998;29:613-617.
 55. Naarding P, de Koning I, van Kooten F, Janzing JG, Beekman AT, Koudstaal PJ. Post-stroke dementia and depression: frontosubcortical dysfunction as missing link? *Int J Geriatr Psychiatry* 2007;22:1-8.
 56. Mantyla R, Erkinjuntti T, Salonen O, Aronen HJ, Peltonen T, Pohjasvaara T, et al. Variable agreement between visual rating scales for white matter hyperintensities on MRI. Comparison of 13 rating scales in a poststroke cohort. *Stroke* 1997;28:1614-1623.
 57. van Straaten EC, Fazekas F, Rostrup E, Scheltens P, Schmidt R, Pantoni L, et al. Impact of white matter hyperintensities scoring method on correlations with clinical data: the LADIS study. *Stroke* 2006;37:836-840.
 58. Fazekas F, Chawluk JB, Alavi A, Hurtig HI, Zimmerman RA. MR signal abnormalities at 1.5 T in Alzheimer's dementia and normal aging. *AJR Am J Roentgenol* 1987;149:351-356.
 59. Melhem ER, Herskovits EH, Karli-Oguz K, Golay X, Hammoud DA, Fortman BJ, et al. Defining thresholds for changes in size of simulated T2-hyperintense brain lesions on the basis of qualitative comparisons. *AJR Am J Roentgenol* 2003;180:65-69.
 60. Burke WJ, Rubin EH, Morris JC, Berg L. Symptoms of "depression" in dementia of the Alzheimer type. *Alzheimer Dis Assoc Disord* 1988;2:356-362.
 61. Burke WJ, Roccaforte WH, Wengel SP, McArthur-Miller D, Folks DG, Potter JF. Disagreement in the reporting of depressive symptoms between patients with dementia of the Alzheimer type and their collateral sources. *Am J Geriatr Psychiatry* 1998;6:308-319.
 62. Teri L, Wagner AW. Assessment of depression in patients with Alzheimer's disease: concordance among informants. *Psychol Aging* 1991;6:280-285.
 63. Janzing JG, Hooijer C, van 't Hof MA, Zitman FG. Depression in subjects with and without dementia: a comparison using GMS-AGECAT. *Int J Geriatr Psychiatry* 2002;17:1-5.
 64. Zubenko GS, Zubenko WN, McPherson S, Spoor E, Marin DB, Farlow MR, et al. A collaborative study of the emergence and clinical features of the major depressive syndrome of Alzheimer's disease. *Am J Psychiatry* 2003;160:857-866.