



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.



## Letter to the Editor

**Delirium in a pregnant woman with SARS-CoV-2 infection in India***To the Editor*

Severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) infection is rapidly spreading all over the world affecting all aspects of health, most importantly the mental health. In the current healthcare crisis due to COVID-19 pandemic, immediate dissemination of evidence is a priority for empowering the healthcare providers and policy makers (Tandon, 2020). Currently, there is limited data on impact of COVID-19 on mental health of individuals residing in low income and middle-income countries (LMICs), especially pregnant women. Pregnant women are at increased risk of contracting COVID-19 and thus require special attention, especially while dealing with mental health issues. It is extremely challenging to manage the mental health problems of pregnant women with COVID-19 in India and other LMICs due to an inadequate health system infrastructure, lack of trained manpower and mental health services (Jungari, 2020). Several challenges are being faced in the COVID-19 hospitals in LMICs, especially for the management of pregnant and post-partum women with COVID-19. In a dedicated COVID-19 facility, we managed 885 pregnant women with COVID-19 wherein more than 600 women delivered. In these women, increased anxiety and psychological distress related to COVID-19 was observed (unpublished data). Three cases of post-partum psychosis associated with COVID-19 were successfully managed at our dedicated COVID-19 hospital (Mahajan et al., 2020; Subramanyam et al., 2020).

We report an unusual case of altered mental status with delirium in a 30-weeks pregnant woman positive for SARS-CoV-2 infection. She had preeclampsia, normal oxygen saturation and no other symptoms of COVID-19. Her agitation started 14 hs after admission and delivered vaginally 6 hs after first episode of agitation. In this case of delirium, restraining her extreme agitation was very challenging, especially while wearing personal protective equipments. Neurological and psychiatric evaluation confirmed a diagnosis of delirium, secondary to co-morbidities related to pregnancy [anaemia, severe preeclampsia, bad obstetric history (BOH) and preterm delivery] and SARS-CoV-2 infection. She had similar episode on Day-4 post-partum. There was no psychotic phenomenon like delusion or hallucinations ruling out post-partum psychosis. Her neurological examination including a CT scan was normal, ruling out posterior reversible encephalopathy syndrome (PRES) which is also known to be present as delirium in cases of severe preeclampsia (Schusse et al., 2013). Her symptoms improved gradually and got discharged from the hospital on Day-31 (Table 1).

In this case, pregnancy coupled with COVID-19 and co-morbidities such as severe preeclampsia, preterm labour induction, anaemia requiring blood transfusions could have caused various biochemical changes. This might have led to aberrant stress responses manifesting as delirium. It is worth to note that certain patients with COVID-19 may have non-specific neurological symptoms such as delirium, preceding symptoms of even fever and cough (Cipriani et al., 2020). Eclampsia and severe preeclampsia are considered as differential diagnosis in patients with presentation of agitation and delirium (Aftab and Shah, 2017). Delirium is an acute, complex, organic brain syndrome occurring secondary to multifactorial aetiology which can be either due to direct brain insult or aberrant stress responses to stressors like infections, surgical trauma, anxiety, etc. (Brockington, 2004). Pregnancy and puerperium also cause considerable physical and psychiatric modifications that impose significant amount of stress placing a woman in a situation of restructuring her role as a mother. This may lead to psychological symptoms or provoke various mental illnesses. The somatic alterations of pregnancy can lead to psychiatric problems and vice versa in different ways (Di Renzo, 2002). Acute agitation should be treated as an obstetric emergency as it can compromise the safety of the mother, the child or others present in the vicinity (Niforatos et al., 2019). Delirium as such is a rare disorder in pregnancy but was common in early 19th century with an incidence of 1/5000 deliveries. However, now it has become rare due to good antenatal care and labour analgesia (Brockington, 2004). Viral infection like SARS-CoV-2 with fever and hypoxemia may trigger delirium (Cipriani et al., 2020). Apart from lungs, central nervous system (CNS) has also been observed to be affected by SARS-CoV-2 infection (Mao et al., 2020). Neuropsychiatric manifestations including delirium are now recognized as presenting features of COVID-19 and delirium may be the only presenting symptom (Hosseini et al., 2020). However, it is not clear whether these neurological manifestations are due to direct injury to CNS or an indirect response to systemic inflammatory storm (Cipriani et al., 2020; Mao et al., 2020). Based on the emerging evidence of neuropsychiatric manifestations of COVID-19, further studies on its pathogenesis are required.

There are several challenges for managing the psychiatric emergencies in pregnant women with COVID-19 in LMICs. There is lack of trained counselors for addressing the multiple psychological problems of pregnant women with COVID-19, as they worry about transmission of infection to their child, anxiety about the safe delivery, fear of separation from newborn, etc. Unavailability of trained healthcare workers in maternity units on diagnosis and management of psychiatric emergencies, limited availability of mental health providers and services are some of the other challenges in LMICs.

There is an opportunity for healthcare providers engaged in maternity units for acquiring the skills on diagnosis and effective management of psychiatric disorders associated with viral infections including the current pandemic of COVID-19. Based on our experience, we recommend the training of healthcare workers on early recognition and appropriate treatment of psychiatric emergencies in pregnancy and post-partum period. There is a need for creating specialist mental health services and provision of a psychologist counselor for specifically dealing with pregnancy care in public healthcare system. Accurate information about COVID-19 as well as counseling should be provided to the pregnant and post-partum women for

<https://doi.org/10.1016/j.ajp.2020.102513>

Received 4 September 2020

Available online 11 December 2020

1876-2018/© 2020 Elsevier B.V. All rights reserved.

**Table 1**  
Socio-demographic and clinical characteristics of a woman with delirium and COVID-19 in India.

Parameters	Delirium with COVID-19
Age	34 years
Socio-economic Status	Low
Gravida (G) /Parity (P) /Living children (L)	G5 P1L0
Bad Obstetric History	Previous 3 spontaneous abortions and one still birth
H/O Contact with Positive Person/foreign travel	No
Indication for COVID-19 RT-PCR testing	Universal Screening as per existing local guidelines
Spontaneous Conception	Yes
Antenatal Comorbidities	Severe Preeclampsia, Anemia, BOH
Family and previous history of Psychiatric illness and substance abuse	No
Symptoms	Agitation, aggression, mental confusion, alteration of sleep, headache, loss of appetite, sleep disturbance, violent behavior, disorientation, Not cooperating. Did not allow per-vaginal examination, monitoring foetal heart sounds and blood pressure. During labour she became agitated, violent and pulled off intravenous catheter, abused and attempted biting the healthcare workers in maternity unit.
Significant clinical findings	Severe preeclampsia, Minimal pleural effusion
Hemoglobin% (g/dl)	8.6
Total Leucocyte Count	10,200/ $\mu$ L
Blood Group and Rh type	O Positive
Duration of Delirium Symptoms	5 days
Obstetric Management	Induction of Labour with dinoprostone gel, magnesium sulphate therapy for severe preeclampsia Inj. mannitol, anti-hypertensives and three PCV transfusion. Restrained by six attendants. Inj. Midazolam followed by Inj. haloperidol to control agitation. Tab. haloperidol in tapering doses over 11 days
Management	31 days, as baby was in NICU till then,
Duration of Hospital stay	Neonatal death (female) on Day 31, owing to extreme prematurity and low birth weight (1.270 kg).
Neonatal outcome	

BOH, Bad Obstetric History; NICU, Neonatal Intensive Care Unit, Inj, Injection; PCV, Packed Cell Volume.

reducing stress. The guidelines covering maternal health should have a provision for training the obstetricians in management of the psychiatric emergencies and use of short term psychotropic.

### Funding

The study is supported by intramural grant of ICMR-NIRRH (ICMR-NIRRH/RA/966/09-2020).

### Trial registration

PregCovid study is registered with Clinical Trial Registry of India (Registration no: CTRI/2020/05/025423)

### Ethics approval

The study was approved by the Ethics Committees of TNMC (No. ECARP/2020/63 dated 27.05.2020) and ICMR-NIRRH (IEC no. D/ICEC/Sci-53/55/2020 dated 04.06.2020).

### Contribution to authorship

NM and RG had full access to all of the data in the study and take responsibility for the integrity of the data and the accuracy of the data analysis.  
Concept and design: RG, NM  
Acquisition of data: MP SK, NM, AB  
Analysis, or interpretation of data: All authors  
Drafting of the manuscript: RP, NM, RG  
Critical revision of the manuscript for important intellectual content: NM, RG  
Statistical analysis: NM, RG  
Administrative and technical or material support: NM, SM, RG

### Declaration of Competing Interest

The authors declare that they have no conflict of interest.

### Acknowledgements

The authors acknowledge the Director General, ICMR and Network of National Registry of Pregnant women with COVID-19 in India (PregCovid Registry, CTRI/2020/05/025423). The Dean, TNMC, Faculties, Resident doctors in the Department of Psychiatry and Department of Obstetrics and Gynecology at TNMC, Mumbai are sincerely acknowledged. RG is an awardee of the DBT Wellcome India Alliance Clinical and Public Health Intermediate Fellowship (Grant no. IA/CPHI/18/1/503933).

## References

- Aftab, A., Shah, A.A., 2017. Behavioral emergencies: special considerations in the pregnant patient. *Psychiatr. Clin. North Am.* 40, 435–448. <https://doi.org/10.1016/j.psc.2017.05.017>.
- Brockington, I., 2004. *Diagnosis and management of post-partum disorders: a review.* *World Psychiatry* 3, 89–95.
- Cipriani, G., Danti, S., Nuti, A., Carlesi, C., Lucetti, C., Di Fiorino, M., 2020. A complication of coronavirus disease 2019: delirium. *Acta Neurol. Belg.* 120, 927–932. <https://doi.org/10.1007/s13760-020-01401-7>.
- Di Renzo, G.C., 2002. Pregnancy as delirium. *J. Matern. Fetal. Neonatal. Med.* 12, 145–148. <https://doi.org/10.1080/jmf.12.3.145.148>.
- Hosseini, A.A., Shetty, A.K., Sprigg, N., Auer, D.P., Constantinescu, C.S., 2020. Delirium as a presenting feature in COVID-19: neuroinvasive infection or autoimmune encephalopathy? *Brain Behav. Immun.* 88, 68–70. <https://doi.org/10.1016/j.bbi.2020.06.012>.
- Jungari, S., 2020. Maternal mental health in India during COVID-19. *Public Health* 185, 97–98. <https://doi.org/10.1016/j.puhe.2020.05.062>.
- Mahajan, N.N., Pednekar, R., Patil, S.R., Subramanyam, A.A., Rathi, S., Malik, S., Mohite, S.C., Shinde, G., Joshi, M., Kumbhar, P., Tilve, A., Lokhande, P.D., Srivastava, S.A., 2020. Preparedness, administrative challenges for establishing obstetric services, and experience of delivering over 400 women at a tertiary care COVID-19 hospital in India [WWW Document]. *Int. J. Gynecol. Obstet.* URL <https://obgyn.onlinelibrary.wiley.com/doi/abs/10.1002/ijgo.13338> (Accessed 8.25.20).
- Mao, L., Jin, H., Wang, M., Hu, Y., Chen, S., He, Q., Chang, J., Hong, C., Zhou, Y., Wang, D., Miao, X., Li, Y., Hu, B., 2020. Neurologic manifestations of hospitalized patients with coronavirus disease 2019 in Wuhan, China. *JAMA Neurol.* 77, 683–690. <https://doi.org/10.1001/jamaneurol.2020.1127>.
- Niforatos, J.D., Wanta, J.W., Shapiro, A.P., Yax, J.A., Viguera, A.C., 2019. How should I treat acute agitation in pregnancy? *Cleve Clin. J. Med.* 86, 243–247. <https://doi.org/10.3949/cjcm.86a.18041>.
- Schusse, C.M., Peterson, A.L., Caplan, J.P., 2013. Posterior reversible encephalopathy syndrome. *Psychosomatics* 54, 205–211. <https://doi.org/10.1016/j.psym.2013.01.014>.
- Subramanyam, A.A., Nachane, H.B., Mahajan, N.N., Shinde, S., D Mahale, S., Gajbhiye, R.K., 2020. Postpartum psychosis in mothers with SARS-CoV-2 infection: a case series from India. *Asian J. Psychiatr.* 102406. <https://doi.org/10.1016/j.ajp.2020.102406>.
- Tandon, R., 2020. COVID-19 and mental health: preserving humanity, maintaining sanity, and promoting health. *Asian J. Psychiatr.* 51, 102256. <https://doi.org/10.1016/j.ajp.2020.102256>.

Niraj N. Mahajan<sup>\*1</sup>

*Department of Obstetrics and Gynecology, Topiwala National Medical College & BYL Nair Charitable Hospital, Mumbai, 400 008, India*

Rahul K. Gajbhiye<sup>\*\*1</sup>

*ICMR-National Institute for Research in Reproductive Health, Mumbai, 400 012, India*

Rahi R. Pednekar

*Department of Obstetrics and Gynecology, Topiwala National Medical College & BYL Nair Charitable Hospital, Mumbai, 400 008, India*

Madhura P. Pophalkar

*Department of Obstetrics and Gynecology, Topiwala National Medical College & BYL Nair Charitable Hospital, Mumbai, 400 008, India*

Shweta N. Kesarwani

*Department of Obstetrics and Gynecology, Topiwala National Medical College & BYL Nair Charitable Hospital, Mumbai, 400 008, India*

Aishwarya V. Bhurke

*ICMR-National Institute for Research in Reproductive Health, Mumbai, 400 012, India*

Smita D. Mahale

*ICMR-National Institute for Research in Reproductive Health, Mumbai, 400 012, India*

<sup>\*</sup> Corresponding author at: Department of Obstetrics and Gynecology, B. Y. L. Nair Charitable Hospital and T. N. Medical College, Mumbai, 400008, India.

<sup>\*\*</sup> Corresponding author at: Scientist D & DBT Wellcome India Alliance, Clinical & Public Health Intermediate Fellow, Department of Clinical Research, ICMR-National Institute for Research in Reproductive Health, J M Street, Parel, Mumbai, 400012, India.

*E-mail address:* [nirajdr@hotmail.com](mailto:nirajdr@hotmail.com) (N.N. Mahajan).

*E-mail address:* [gajbhiyer@nirrh.res.in](mailto:gajbhiyer@nirrh.res.in) (R.K. Gajbhiye).

<sup>1</sup> This author contributed equally.