

## How to reduce the ecological impact of gastric peroral endoscopic myotomy: a small effort for the environment




It is already common knowledge that endoscopic procedures are responsible for the production of large amounts of waste, mostly owing to the usage of a high number of disposable instruments [1]. Gastric peroral endoscopic myotomy (G-POEM) is a well-known procedure used for the treatment of refractory gastroparesis [2]. Like most submucosal endoscopic procedures, this technique requires the injection of a premixed methylene blue/glycerol solution into the submucosa, in order to raise a mucosal bleb and create the space to make a safe mucosal cut. This step is usually performed with a routine single-use sclerotherapy needle [2]; however, a DualKnife could also be used for the same purpose, allowing the use of one disposable injection needle to be spared.

We report the case of a 31-year-old patient who was suffering from refractory gastroparesis secondary to diabetes mellitus type 1, who was referred to our center for G-POEM treatment. After the ideal mucosal entry site had been identified, we opted to use the DualKnife for the initial injection, as well as during the rest of the procedure (► **Video 1**). This was easily performed without any extra inconvenience or difficulty during the whole endoscopic intervention.

This case illustrates that, even if the DualKnife is shorter than the sclerotherapy needle, it can have the same efficacy and performance for submucosal injection as the latter. This approach has multiple advantages as it offers a time-saving during submucosal endoscopy by eliminating an instrument change and, most importantly, it leads to a significant reduction in waste production and the associated ecological footprint of the procedure.

Endoscopy\_UCTN\_Code\_TTT\_1AU\_2AF



► **Video 1** A method to reduce the ecological impact of gastric peroral endoscopic myotomy by performing the injection with the DualKnife, rather than with a sclerotherapy needle.

### Competing interests

The authors declare that they have no conflict of interest.

### The authors

**Mariana Figueiredo<sup>1</sup>, Clara Yzet<sup>1</sup>, Daniel Grinberg<sup>2,3</sup>, Thomas Lambin<sup>1</sup>, Pierre-Jean Cottinet<sup>4</sup>, Jérôme Rivory<sup>1</sup>, Mathieu Pioche<sup>1,4</sup>**

- 1 Gastroenterology and Endoscopy Unit, Pavillon L, Edouard Herriot Hospital, Lyon, France
- 2 Chirurgie Cardiaque, Edouard Herriot Hospital, Lyon, France
- 3 Équipe de matériaux et procédés d'élaboration, Institut National des Sciences Appliquées, Lyon, France
- 4 Inserm U1032, Labtau, Lyon, France

### Corresponding author

**Mariana Figueiredo, MD**

Service hépato-gastroentérologie, Hôpital Edouard Herriot, 5 place d'Arsonval, 69003 Lyon, France  
mariana.figueiredo.pro@gmail.com

### References

- [1] Nambur S, von Renteln D, Damianos J et al. Estimating the environmental impact of disposable endoscopic equipment and endoscopes. *Gut* 2022. doi:10.1136/gutjnl-2021-324729
- [2] Li L, Spandorfer R, Qu C et al. Gastric peroral endoscopic myotomy for refractory gastroparesis: a detailed description of the procedure, our experience, and review of the literature. *Surg Endosc* 2018; 32: 3421–3431

### Bibliography

*Endoscopy* 2022; 54: E899  
DOI 10.1055/a-1847-7490  
ISSN 0013-726X  
published online 1.7.2022  
© 2022. The Author(s).

This is an open access article published by Thieme under the terms of the Creative Commons Attribution-NonDerivative-NonCommercial License, permitting copying and reproduction so long as the original work is given appropriate credit. Contents may not be used for commercial purposes, or adapted, remixed, transformed or built upon. (<https://creativecommons.org/licenses/by-nc-nd/4.0/>)

Georg Thieme Verlag KG, Rüdigerstraße 14, 70469 Stuttgart, Germany

