

Predictors of sedentary status in overweight and obese patients with multiple chronic conditions: a cohort study

Mark H Joven
Ivana T Croghan
Stephanie M Quigg
Jon O Ebbert
Paul Y Takahashi

Division of Primary Care Internal
Medicine, Mayo Clinic, Rochester,
Minnesota, MN, USA

Purpose: Walking may improve health in obese patients with multimorbidity. We aimed to identify predictors associated with achieving <5000 steps per day after 4 months.

Patients and methods: We conducted a cohort study of 125 adult patients with a body mass index >25 kg/m² and ≥7 comorbidities. We evaluated potential predictors for <5000 steps per day using logistic regression and adjusting for age >65 years and sex.

Results: The mean (range) age was 63.6 (20.3–89.8) years. Daily step counts <5000 at baseline showed the highest risk of <5000 daily steps at 4 months (odds ratio [OR] 31.82, 95% confidence interval [CI]: 12.14–95.50). Other significant characteristics were physical quality of life (OR 6.21, 95% CI: 2.32–18.54), gait speed <1 m/s (OR 2.57, 95% CI: 1.18–5.71), age ≥65 years (OR 2.21, 95% CI: 1.05–4.77), waist circumference ≥102 cm (OR 2.48, 95% CI: 1.05–6.06), and body mass index ≥30 kg/m² (OR 2.69, 95% CI: 1.20–6.26).

Conclusion: New models to increase walking may be required for higher-risk patients.

Keywords: multimorbidity, obesity, pedometer, walking

Introduction

The prevalence of obesity, defined as a body mass index (BMI) ≥30 kg/m², has a mean (95% confidence interval [CI]) of 34.9% (32.0%–37.9%) among US adults aged ≥20 years. The National Health and Nutrition Examination Survey (NHANES) from 2011 to 2012 reported that, on average, 68.5% (95% CI, 65.2%–71.6%) of persons were either overweight (BMI ≥25 but <30 kg/m²) or obese.¹ Obesity is associated with metabolic syndrome, characterized by dyslipidemia (e.g., high level of triglycerides, low level of high-density lipoprotein cholesterol [HDL-C]), impaired fasting glucose, and hypertension. Metabolic syndrome increases the risks of cardiovascular disease and type 2 diabetes mellitus,² and sedentary behaviors increase the risks of metabolic syndrome,³ type 2 diabetes mellitus, cardiovascular disease, and death.⁴ All-cause mortality risk increases progressively as BMI exceeds 25.0 kg/m².⁵

Lower levels of physical activity have been associated in 9% of premature death. This is projected to increase in the coming years.⁶ It is estimated to be the cause of 6% of coronary artery disease and about 7% of type 2 diabetes mellitus worldwide.⁶ In contrast, higher levels of physical activity have been associated with lower rates of chronic disease, cardiovascular disease, and type 2 diabetes mellitus.^{6,7} Reductions in blood pressure (decrease in systolic blood pressure by 2–5 mmHg and in diastolic blood pressure by 1–4 mmHg), as well as decreases in levels of low-density lipoprotein cholesterol and non-HDL-C, have been observed with aerobic physical activity.⁷ In a

Correspondence: Paul Y Takahashi
Division of Primary Care Internal
Medicine, Mayo Clinic, 200 First St SW,
Rochester, MN 55905, USA
Email takahashi.paul@mayo.edu

population-based sample of 60-year-old persons, increased physical activity was associated with improved cardiovascular health.⁸ Considerable interest exists about increasing the aerobic activity of patients with medical comorbidities. Increasing step counts can have positive effects on health. When patients used a pedometer, they had decreased BMI by a mean (95% CI) of 0.38 (0.05–0.72) kg/m² and systolic blood pressure by a mean (95% CI) of 3.8 (1.7–5.9) mmHg.⁹ Among patients with musculoskeletal concerns, an increase of 1950 steps per day showed improvements in disability, strength, and pain scores.¹⁰ However, the positive effects of an increased step count on health outcomes have not been seen in all studies. The ASUKI Step study, a 6-month pedometer-based intervention performed at the Arizona State University in the Phoenix area of Arizona and the Karolinska Institutet in Stockholm, Sweden, showed no significant change in BMI, but a decrease of 3.0 cm in waist circumference was observed at the end of the trial.¹¹

The available literature is limited about barriers to physical activity. A survey of 5167 persons in Canada aged 15–79 years found that educational attainment, family income, self-rated health, intentions, self-efficacy scores, perceived barriers, perceived health benefits, and facility availability were factors associated with physical activity behaviors.¹² A cross-sectional study of overweight and obese persons in Iran with type 2 diabetes mellitus identified barriers to physical activity that included negative attitude toward physical activity, discouragement, physical problems, cost, and environmental factors.¹³ Barriers more specific to older obese and overweight persons with multiple chronic conditions, have not been identified.

We previously performed a randomized controlled trial of goal setting and pedometer use among obese patients with multiple chronic conditions.¹⁴ Step counts (the primary outcome) did not increase significantly with this intervention ($p=0.08$). Patient factors likely had a role in preventing engagement in increased activity, but these factors have not been elucidated. Our primary aim was to identify risk factors for walking <5000 steps per day for overweight and obese patients with multiple chronic health conditions. Identifying these risk factors may guide future interventions to increase step count and activity.

Research design

Study design

We conducted a secondary analysis of a published clinical trial¹⁴ using a cohort design. In this trial, patients were

randomly assigned to either immediate pedometer use plus behavioral goal setting in the form of SMART (specific, meaningful, action-based, realistic, and timely) goals or delayed pedometer use plus delayed goal setting. Patients in the delayed group received the intervention with pedometer and SMART goals at 2 months. The SMART goals were used from previously published principles of formulating personalized and specific goals.¹⁵ This model is based on the transtheoretical model of change where the SMART goals help move the participant to action. The study was reviewed and approved by the Mayo Clinic Institutional Review Board.

Setting

The initial clinical study was completed at a clinical center in Rochester, Minnesota (Clinical Trial Number: NCT01833507). Study enrollment occurred from May 1, 2013, through September 10, 2014. Patients had a face-to-face visit with the study coordinator, and written informed consent was obtained.¹⁶

Participants

Study participants were 130 community-dwelling persons who were overweight or obese (BMI, 25.0–29.9 kg/m² and ≥ 30 kg/m², respectively) and had ≥ 7 comorbid chronic conditions based on outpatient and inpatient billing codes. All enrolled participants received primary care through Mayo Clinic to reduce referral bias. This disease burden of 7 or more conditions correlates with extended or complex tiering (tier 3 or 4) based on the Minnesota Tiering Model, a clinical risk stratification method.¹⁷ This was calculated by a trained medical staff not involved in the study. Participants who were excluded were either enrolled in hospice or wheelchair-bound, had uncontrolled depression (Patient Health Questionnaire 9 [PHQ-9] scores ≥ 10),¹⁸ or had documented dementia in accordance with the Short Test of Mental Status^{19,20} (scores $\leq 29/38$). Patients were excluded if they lived in a skilled nursing or correctional facility. Patients were also excluded if they refused medical record review. All patients who had available step data at the end of the original study were included in the present study.

Outcome

The primary outcome was attainment of <5000 steps per day on average, as recorded by a pedometer at 4 months. Achieving <5000 steps per day has been considered to be sedentary status by some investigators²¹ and thus, the determination of a cut-off of 5000 steps. All participants were seen at baseline and had baseline information collected.

Predictors

The demographic, medical, and anthropomorphic data were collected. These included age, race, sex, elder risk assessment (ERA) score,²² BMI, baseline systolic blood pressure, and waist circumference. Age was dichotomized as <65 years and ≥65 years on the basis of potential enrollment in Medicare. We dichotomized ERA score into ≥16, which represented the highest risk of hospital admission.²² We used a BMI cutoff score of ≥30 kg/m² to represent persons with obesity.²³ We used cutoffs of systolic blood pressure >150 mmHg²⁴ and waist circumference of >102 cm, as well as a cutoff of <50 for both the physical component summary (PCS) and the mental component summary (MCS) of the 12-Item Short Form Survey (SF-12).²⁵ For grip strength, our cutoff was <25 kg. For 4 m walk time, we dichotomized at <4 s for a calculated gait speed of 1.0 m per s.²⁶

The team obtained baseline average steps per day, grip strength, and gait speed to identify potential factors at baseline that may affect sedentary status after 2 or 4 months of pedometer use. Grip strength was measured in pound per 1 inch squared using a hydraulic hand dynamometer (Baseline, Fabrication Enterprise, White Plains, NY, USA). Handheld dynamometers are considered a practical modality to assess muscle strength.²⁷ The immediate pedometer group used a pedometer for 4 months (Omron HJ-112; Omron Healthcare, Inc., Bannockburn, IL, USA); the delayed pedometer group used the pedometer for 2 months.^{28,29} Length of actual pedometer use was also assessed. Research Electronic Data Capture (Mayo Clinic, Rochester, MN, USA) was used to computerize the collected data that were initially logged in paper.³⁰

Patients completed the PHQ-9¹⁸ and the SF-12.²⁵ The PHQ-9 assesses mood; scores ≥9 indicate a possible depressive disorder.¹⁸ Physical and mental quality of life (QoL) was measured by SF-12 Health Survey at baseline, a valid measure of physical and mental functioning, as well as overall health-related QoL. The SF-12 is a questionnaire that contains 2 components of health summary – PCS and MCS – with higher scores indicating better perception of health.^{25,31,32} The SF-12 uses a normalized mean of 50 for each subscore and 10 points for SD.²⁵

Statistical methods

The present analysis was conducted on patients who had available step data. For power calculations, we used all available information. We compared the demographic characteristics, biometric information, pedometer use, and self-reported data between the 2 groups at baseline. For continuous variables, we reported mean and SD for the over-

all cohort and for persons who had ≥5000 steps and those with <5000 steps. We also categorized each predictor into a binary 2-group method. We split age into >65 years and ≤65 years. We presented descriptive data for both continuous and dichotomized information. We used available information for determination of significance of predictors. We compared persons with <5000 steps per day to those with ≥5000 steps using logistic regression unadjusted and adjusted for age >65 years and sex. We calculated odds ratios (ORs) with respective 95% CIs. *p*-values were obtained for each measurement to determine statistical significance, with *p*<0.05 considered significant. All analyses were conducted using JMP statistical software (SAS Institute Inc., Cary, NC, USA).

Results

Participants

Of 244 potential participants, 130 patients consented in the initial study, with 125 patients with step data for analysis (Table 1). Patients who had <5000 steps per day had a mean (SD) age of 66.4 (12.3) years versus 59.6 (16.8) years for those with ≥5000 steps. Of the patient population, 98% were white and 73% were women, and 71 of the 125 patients (57%) had <5000 steps daily at the end of 4 months.

Predictors of <5000 steps per day

After adjusting for age >65 years and sex, we found 6 predictors of <5000 steps per day (Table 2). We observed that patients aged >65 years had higher odds of <5000 steps (adjusted OR 2.21, 95% CI: 1.05–4.77) compared with patients aged ≤65 years. BMI ≥30 kg/m² also had higher odds of <5000 steps per day (adjusted OR 2.69, 95% CI: 1.20–6.26). Waist circumference >102 cm showed higher odds of <5000 steps (adjusted OR 2.48, 95% CI: 1.05–6.06) compared with waist circumference ≤102 cm. No association was found with white race, female sex, or systolic blood pressure >140 mmHg and <5000 steps per day.

For functional and emotional predictors, the strongest predictor of <5000 steps involved having <5000 steps at 2 months of measurement (adjusted OR 31.83, 95% CI: 12.14–95.50) (Table 2). Thus, patients who had <5000 steps at the midway measurement point were likely to continue to have <5000 steps. We did not observe an association between the length of pedometer wearing time and the status of <5000 steps at 4 months, an observation consistent with the original trial findings.¹⁴ A self-rated physical QoL score <50 had higher odds of <5000 steps at 4 months (adjusted OR 6.21, 95% CI: 2.32–18.54). Patients who required >4 s to walk 4 m (gait speed, <1 m per s) also had higher odds

Table 1 Demographic and biometric characteristics at baseline

Characteristic ^a	Overall (N=125) ^b	Patients who had ≥5000 steps per day at 4 months (n=54) ^b	Patients who had <5000 steps per day at 4 months (n=71) ^b
Demographic			
Age, mean (SD), year	63.5 (15)	59.6 (16.8)	66.4 (12.8)
Age >65 years	63 (50)	22 (41)	41 (58)
White race	122 (98)	53 (98)	69 (97)
Female sex	91 (73)	38 (70)	53 (75)
ERA score, mean (SD)	6.3 (5.4)	4.9 (4.9)	7.3 (5.6)
ERA score ≥16	8 (6)	1 (2)	7 (10)
Biometric			
BMI, mean (SD), kg/m ²	33.8 (5.9)	32.4 (4.6)	34.8 (6.6)
BMI >30 kg/m ²	86 (69)	32 (59)	54 (76)
Systolic blood pressure, mean (SD), mmHg	130 (14.1)	127.6 (13.1)	131.9 (14.7)
Systolic blood pressure >140 mmHg	31/124 (25)	11 (20)	20/70 (29)
Waist circumference, mean (SD), cm	113 (15)	109 (13)	117 (16)
Waist circumference >102 cm	93 (74)	36 (67)	57 (80)
Functional and emotional status			
Pedometer intervention at 4 months	63 (50)	29 (54)	34 (48)
PHQ-9, mean (SD)	2.9 (2.6)	2.9 (2.8)	2.9 (2.4)
Physical QoL score, mean (SD), PCS	40.8 (9.4)	44.2 (8.8)	38.3 (9.1)
Physical QoL score <50 PCS	97/123 (79)	34/53 (64)	63/70 (90)
Mental QoL score, mean (SD) MCS	54.1 (8.1)	53.8 (8.3)	54.3 (8.0)
Mental QoL score <50 MCS	31/123 (25)	13/53 (25)	18/70 (26)
Steps at 2 months, mean (SD)	4895 (2806)	6867 (2228)	3395 (2169)
Steps <5000 at 2 months	74 (59)	11 (20)	63 (89)
Grip strength, mean (SD), kg	24.7 (9.0)	26.2 (9.0)	23.5 (9.0)
Grip strength <25 kg	70 (56)	28 (52)	42 (59)
Gait speed over 4 m, mean (SD), s	4.5 (1.3)	4.0 (0.9)	4.9 (1.4)
Gait time >4 s	73 (58)	24 (44)	49 (69)

Notes: ^aValues are presented as number and percentage of patients unless specified otherwise. ^bWhere the number of patients is less than the number cited in the column heading, the patient number is listed as a denominator.

Abbreviations: BMI, body mass index; ERA, elder risk assessment; lb/in², pound per inch squared; MCS, mental component summary; PCS, physical component summary; PHQ-9, Patient Health Questionnaire 9; QoL, quality of life.

Table 2 Clinical characteristics and odds of achieving <5000 steps per day at 4 months

Characteristic ^a	OR (95% CI)	Unadjusted p-value ^b	Age- (>65 years) and sex-adjusted OR (95% CI)	Adjusted p-value ^b
Age >65 years	1.99 (0.97–4.12)	0.06	2.21 (1.05–4.77)	0.04 ^c
White race	0.65 (0.06–7.37)	0.72	0.49 (0.02–5.35)	0.55
Female sex	1.24 (0.56–2.74)	0.59	1.57 (0.68–3.67)	0.29
ERA score ≥16	1.98 (0.97–4.12)	0.06	5.17 (0.86–99.1)	0.07
BMI >30 kg/m ²	2.18 (1.02–4.77)	0.045 ^c	2.69 (1.20–6.26)	0.02 ^c
Systolic blood pressure >140 mmHg	1.56 (0.68–3.72)	0.30	1.40 (0.58–3.55)	0.46
Waist circumference >102 cm	2.19 (0.97–5.10)	0.06	2.48 (1.05–6.06)	0.04 ^c
Pedometer intervention at 4 months	0.79 (0.39–1.61)	0.52	0.80 (0.37–1.62)	0.50
Physical QoL score <50 PCS	5.02 (2.00–13.99)	<0.001 ^c	6.21 (2.32–18.54)	<0.001 ^c
Mental QoL score <50 MCS	1.07 (0.47–2.46)	0.88	1.19 (0.51–2.84)	0.69
Steps <5000 at 2 months	30.78 (12.05–88.39)	<0.001 ^c	31.83 (12.14–95.50)	<0.001 ^c
Grip strength <25 kg	1.34 (0.66–2.76)	0.41	0.93 (0.38–2.29)	0.88
Time >4 s to walk 4 m	2.78 (1.34–5.88)	0.006 ^c	2.57 (1.18–5.71)	0.02 ^c

Notes: ^aCharacteristics are at baseline with steps at 2 months of intervention. ^bStatistical method was logistic regression and multivariable logistic regression. ^cSignificant value after adjustment for age and sex.

Abbreviations: BMI, body mass index; CI, confidence interval; ERA, elder risk assessment; MCS, mental component summary; OR, odds ratio; PCS, physical component summary; QoL, quality of life.

of <5000 steps (adjusted OR 2.57, 95% CI: 1.18–5.71) than those who required ≤ 4 s. Grip strength and self-rated mental QoL score were not associated with <5000 steps.

Discussion

In our cohort of overweight and obese patients with multiple chronic conditions, we observed that patients who were walking <5000 steps at 2 months after study initiation had a more than 30-fold increased likelihood of continuing to walk <5000 steps at the end of 4 months. We found that patients with slower walking speed and lower self-rated physical QoL score were also more likely to have <5000 steps, with adjusted ORs more than 2.5- and 6-fold greater, respectively. In addition, patients aged ≥ 65 years with a BMI ≥ 30 kg/m² and waist circumference ≥ 102 cm had more than a 2-fold increased risk of taking <5000 steps daily. These novel findings indicate that patients who are not walking, feeling physically poor, older, heavier, and walking slowly are more likely to not walk 5000 steps per day after 4 months. We did not observe a difference among patients who had 4 months of pedometer use compared with 2 months of pedometer use. Thus, patients most in need of increasing their step counts are those who are not achieving 5000 steps per day currently.

Using step count as an outcome seems reasonable as patients and health care groups attempt to quantify physical activity. Some investigators have suggested walking ~5000 steps a day as a sedentary lifestyle index.³³ Among persons with age in the 40s, walking <5000 steps per day is associated with obesity.³⁴ The Tasped Prospective Cohort Study had a total of 2576 mostly overweight or normal weight patients (44.1% and 32.1%, respectively) from 3 cohort studies (mean age, 58.8 years) who were observed for over 10 years in 90% of participants.³⁵ Higher step counts linearly correlated with lower all-cause mortality rates (adjusted hazard ratio 0.94, 95% CI: 0.90–0.98 per 1000 steps; $p=0.004$). Increasing baseline steps per day from <1000 to a 10,000-step guideline produced a mean (95% CI) decline in mortality risk of 46% (18%–65%).

Our observation about poor physical QoL score, slower gait speed, and lower step count is consistent with studies in different patient populations. In an Australian sample of 1232 inactive patients and among the patient group aged 60–78 years, injury and poor health were the most frequently cited barriers to physical activity.³⁶ This finding differed from the groups aged 18–39 years, who stated that time was the biggest barrier to physical activity. A descriptive report of 75 American Indians (mean [SD] age, 58.5 [5.7] years), 81.3%

of whom had at least 1 self-reported chronic disease, showed that poor physical health, measured by SF-12 PCS, strongly correlated with various physical outcomes. Thus, perceived physical health and observed activity such as walking steps and gait speed may be risk factors for walking <5000 steps for obese patients with multiple chronic conditions. In our study, the mental QoL scores did not correlate with patients who achieved <5000 steps per day (adjusted p -value, 0.69). This result is consistent with some reported studies (Mansi et al,³⁷ and Humphreys et al³⁸). In other studies, depressed mood was associated with lower physical activity level.^{39,40} This lack of effect of mental QoL was not completely expected or explained.

In the present study, we did not observe that either short or extended pedometer use affected activity status. We hypothesize that directly addressing sedentary behaviors needs to be a component of any activity intervention. A meta-analysis of 33 studies by Prince and colleagues⁴¹ showed that interventions focusing on sedentary behaviors resulted in clinically meaningful decreases in sedentary time. Focus on physical activity and sedentary behaviors or physical activity components alone showed only modest reductions in sedentary time. Pedometer use with SMART goals did not affect step status in our clinical trial.¹⁴ Age >65 years, BMI >35 kg/m², and waist circumference >102 cm were also associated with <5000 steps at 4 months. The NHANES from 2005 to 2006 found an inverse relationship between step count and BMI and waist circumference in a cross-section of US patients.⁴² Thus, findings from an obese participant population with multiple chronic conditions mirror findings from the general US population.

These results have practical clinical applications. Providers routinely counsel overweight and obese patients on lifestyle modifications to improve and increase exercise, which is consistent with US Preventive Services Task Force guidelines.⁴³ Providers should evaluate patient clinical characteristics to tailor the advice and therapeutic options. Obese or self-rated physically ill patients may have more problems implementing lifestyle interventions. Chronic disease and advanced age can negatively affect physical function.³² Further clinically practical research is needed to improve physical activity in this population.

The present study has practical limitations. Its sample size was small, and our findings have not been replicated in a different cohort. Data were collected prospectively, and data collection was complete. However, a risk of measurement error exists in the outcomes and predictors. In particular, the continuous application of the pedometer was not verified,

and we cannot be certain it was used or worn correctly. An element of volunteer bias is possible because this cohort was taken from patients who volunteered for a prospective clinical trial. The study population was mostly white, which is reflective of the local population in the Upper Midwest of the USA.⁴⁴ Despite these limitations, the study provides some initial utility in predicting who is at risk of <5000 steps after initiation of pedometer use and SMART goal setting.

Conclusion

Patients with step counts <5000 at 2 months, a physical QoL score <50, and a time to walk 4 m >4 s had increased odds of walking <5000 steps daily at 4 months. Age >65 years, BMI >30 kg/m², and waist circumference >102 cm also resulted in higher odds of <5000 steps. Patients who need help the most with their health (e.g., older, <5000 steps daily, obese, poor physical QoL) are those who are most likely to continue at <5000 steps per day. The use of pedometers and goal setting strategies might be one of prevention or activity maintenance for those patients who are already active. Different strategies may be required for patients with lower step counts or higher BMI.

Acknowledgments

We acknowledge the Mayo Clinic Division of Primary Care Internal Medicine and the Department of Medicine for their support in this research. We acknowledge the Center for Clinical and Translational Science grant support (UL1 TR000135) for the use of REDCap.

Disclosure

The authors report no conflicts of interest in this work.

References

- Ogden CL, Carroll MD, Kit BK, Flegal KM. Prevalence of childhood and adult obesity in the United States, 2011–2012. *J Am Med Assoc.* 2014;311(8):806–814.
- Alberti KG, Eckel RH, Grundy SM, et al; International Diabetes Federation Task Force on Epidemiology and Prevention; National Heart, Lung, and Blood Institute; American Heart Association; World Heart Federation; International Atherosclerosis Society; International Association for the Study of Obesity. Harmonizing the metabolic syndrome: a joint interim statement of the International Diabetes Federation Task Force on Epidemiology and Prevention; National Heart, Lung, and Blood Institute; American Heart Association; World Heart Federation; International Atherosclerosis Society; and International Association for the Study of Obesity. *Circulation.* 2009;120(16):1640–1645.
- Edwardson CL, Gorely T, Davies MJ, et al. Association of sedentary behaviour with metabolic syndrome: a meta-analysis. *PLoS One.* 2012;7(4):e34916.
- Wilmot EG, Edwardson CL, Achana FA, et al. Sedentary time in adults and the association with diabetes, cardiovascular disease and death: systematic review and meta-analysis. *Diabetologia.* 2012;55(11):2895–2905. Erratum in: *Diabetologia.* 2013;56(4):942–943.
- Global BMI Mortality Collaboration. Body-mass index and all-cause mortality: individual-participant-data meta-analysis of 239 prospective studies in four continents. *Lancet.* 2016;388(10046):776–786.
- Lee IM, Shiroma EJ, Lobelo F, Puska P, Blair SN, Katzmarzyk PT; Lancet Physical Activity Series Working Group. Effect of physical inactivity on major non-communicable diseases worldwide: an analysis of burden of disease and life expectancy. *Lancet.* 2012;380(9838):219–229.
- Eckel RH, Jakicic JM, Ard JD, et al; American College of Cardiology/American Heart Association Task Force on Practice Guidelines. 2013 AHA/ACC guideline on lifestyle management to reduce cardiovascular risk: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines. *J Am Coll Cardiol.* 2014;63(25 Pt B):2960–2984. Erratum in: *J Am Coll Cardiol.* 2014;63(25 Pt B):3027–3028.
- Eklblom-Bak E, Eklblom B, Vikstrom M, de Faire U, Hellenius ML. The importance of non-exercise physical activity for cardiovascular health and longevity. *Br J Sports Med.* 2014;48(3):233–238.
- Bravata DM, Smith-Spangler C, Sundaram V, et al. Using pedometers to increase physical activity and improve health: a systematic review. *J Am Med Assoc.* 2007;298(19):2296–2304.
- Mansi S, Milosavljevic S, Baxter GD, Tumilty S, Hendrick P. A systematic review of studies using pedometers as an intervention for musculoskeletal diseases. *BMC Musculoskelet Disord.* 2014;15(1):231.
- Walker JR, Soroush A, Ainsworth BE, Belyea M, Swan PD, Yngve A. US cohort differences in body composition outcomes of a 6-month pedometer-based physical activity intervention: the ASUKI Step Study. *Asian J Sports Med.* 2014;5(4):e25748.
- Pan SY, Cameron C, Desmeules M, Morrison H, Craig CL, Jiang X. Individual, social, environmental, and physical environmental correlates with physical activity among Canadians: a cross-sectional study. *BMC Public Health.* 2009;9(1):21.
- Halali F, Mahdavi R, Asghari Jafarabadi M, Mobasseri M, Namazi N. A cross-sectional study of barriers to physical activity among overweight and obese patients with type 2 diabetes in Iran. *Health Soc Care Community.* 2016;24(5):e92–e100.
- Takahashi PY, Quigg SM, Croghan IT, Schroeder DR, Ebbert JO. Effect of pedometer use and goal setting on walking and functional status in overweight adults with multimorbidity: a crossover clinical trial. *Clin Interv Aging.* 2016;11:1099–1106.
- Bovend'Eerd T, Botell RE, Wade DT. Writing SMART rehabilitation goals and achieving goal attainment scaling: a practical guide. *Clin Rehabil.* 2009;23(4):352–361. Erratum in: *Clin Rehabil.* 2010;24(4):382.
- Takahashi PY, Naessens JM, Peterson SM, et al. Short-term and long-term effectiveness of a post-hospital care transitions program in an older, medically complex population. *Healthcare (Amsterdam).* 2016;4(1):30–35.
- Haas LR, Takahashi PY, Shah ND, et al. Risk-stratification methods for identifying patients for care coordination. *Am J Manag Care.* 2013;19(9):725–732.
- Kroenke K, Spitzer RL, Williams JB. The PHQ-9: validity of a brief depression severity measure. *J Gen Intern Med.* 2001;16(9):606–613.
- Kokmen E, Smith GE, Petersen RC, Tangalos E, Ivnik RC. The short test of mental status: correlations with standardized psychometric testing. *Arch Neurol.* 1991;48(7):725–728.
- Kokmen E, Naessens JM, Offord KP. A short test of mental status: description and preliminary results. *Mayo Clin Proc.* 1987;62(4):281–288.
- Tudor-Locke C, Hatano Y, Pangrazi RP, Kang M. Revisiting “how many steps are enough?” *Med Sci Sports Exerc.* 2008;40(7 Suppl):S537–S543.
- Crane SJ, Tung EE, Hanson GJ, Cha S, Chaudhry R, Takahashi PY. Use of an electronic administrative database to identify older community dwelling adults at high-risk for hospitalization or emergency department visits: the elders risk assessment index. *BMC Health Serv Res.* 2010;10(1):338.
- World Health Organization. BMI classification. [Internet]. 2014. Available from: http://apps.who.int/bmi/index.jsp?introPage=intro_3.html. Accessed January 1, 2017.

24. James PA, Oparil S, Carter BL, et al. 2014 evidence-based guideline for the management of high blood pressure in adults: report from the panel members appointed to the Eighth Joint National Committee (JNC 8). *J Am Med Assoc.* 2014;311(5):507–520. Erratum in: *J Am Med Assoc.* 2014;311(17):1809.
25. Ware J Jr, Kosinski M, Keller SD. A 12-item short-form health survey: construction of scales and preliminary tests of reliability and validity. *Med Care.* 1996;34(3):220–233.
26. Fielding RA, Vellas B, Evans WJ, et al; International Working Group on Sarcopenia. Sarcopenia: an undiagnosed condition in older adults: current consensus definition: prevalence, etiology, and consequences. *J Am Med Dir Assoc.* 2011;12(4):249–256.
27. Stark T, Walker B, Phillips JK, Fejer R, Beck R. Hand-held dynamometry correlation with the gold standard isokinetic dynamometry: a systematic review. *PM R.* 2011;3(5):472–479.
28. Schneider PL, Crouter SE, Lukajic O, Bassett DR Jr. Accuracy and reliability of 10 pedometers for measuring steps over a 400-m walk. *Med Sci Sports Exerc.* 2003;35(10):1779–1784.
29. Holbrook EA, Barreira TV, Kang M. Validity and reliability of Omron pedometers for prescribed and self-paced walking. *Med Sci Sports Exerc.* 2009;41(3):670–674.
30. Harris PA, Taylor R, Thielke R, Payne J, Gonzalez N, Conde JG. Research electronic data capture (REDCap): a metadata-driven methodology and workflow process for providing translational research informatics support. *J Biomed Inform.* 2009;42(2):377–3781.
31. Ware JE, Kosinski M, Keller SD; New England Medical Center Hospital Health Institute. *SF-12: How to Score the SF-12 Physical and Mental Health Summary Scales.* 2nd ed. Boston, MA: Health Institute, New England Medical Center; 1995.
32. Hopman WM, Harrison MB, Coe H, Friedberg E, Buchanan M, VanDenKerkhof EG. Associations between chronic disease, age and physical and mental health status. *Chronic Dis Can.* 2009;29(3):108–116.
33. Tudor-Locke C, Bassett DR Jr. How many steps/day are enough? Preliminary pedometer indices for public health. *Sports Med.* 2004;34(1):1–8.
34. Tudor-Locke C, Ainsworth BE, Whitt MC, Thompson RW, Addy CL, Jones DA. The relationship between pedometer-determined ambulatory activity and body composition variables. *Int J Obes Relat Metab Disord.* 2001;25(11):1571–1578.
35. Dwyer T, Pezic A, Sun C, et al. Objectively measured daily steps and subsequent long term all-cause mortality: the tasped Prospective Cohort Study. *PLoS One.* 2015;10(11):e0141274. Erratum in: *PLoS One.* 2015;10(12):e0146202.
36. Booth ML, Bauman A, Owen N, Gore CJ. Physical activity preferences, preferred sources of assistance, and perceived barriers to increased activity among physically inactive Australians. *Prev Med.* 1997;26(1):131–137.
37. Mansi S, Milosavljevic S, Tumilty S, Hendrick P, Higgs C, Baxter DG. Investigating the effect of a 3-month workplace-based pedometer-driven walking programme on health-related quality of life in meat processing workers: a feasibility study within a randomized controlled trial. *BMC Public Health.* 2015;15(1):410.
38. Humphreys DK, Goodman A, Ogilvie D. Associations between active commuting and physical and mental wellbeing. *Prev Med.* 2013;57(2):135–139.
39. Alosco ML, Spitznagel MB, Miller L, et al. Depression is associated with reduced physical activity in persons with heart failure. *Health Psychol.* 2012;31(6):754–762.
40. Yoshiuchi K, Nakahara R, Kumano H, et al. Yearlong physical activity and depressive symptoms in older Japanese adults: cross-sectional data from the Nakanajo study. *Am J Geriatr Psychiatry.* 2006;14(7):621–624.
41. Prince SA, Saunders TJ, Gresty K, Reid RD. A comparison of the effectiveness of physical activity and sedentary behaviour interventions in reducing sedentary time in adults: a systematic review and meta-analysis of controlled trials. *Obes Rev.* 2014;15(11):905–919.
42. Tudor-Locke C, Schuna JM Jr, Han HO, et al. Step-based physical activity metrics and cardiometabolic risk: NHANES 2005-2006. *Med Sci Sports Exerc.* 2017;49(2):283–291.
43. LeFevre ML; US Preventive Services Task Force. Behavioral counseling to promote a healthful diet and physical activity for cardiovascular disease prevention in adults with cardiovascular risk factors: US Preventive Services Task Force Recommendation Statement. *Ann Intern Med.* 2014;161(8):587–593.
44. St Sauver JL, Grossardt BR, Leibson CL, Yawn BP, Melton LJ 3rd, Rocca WA. Generalizability of epidemiological findings and public health decisions: an illustration from the Rochester Epidemiology Project. *Mayo Clin Proc.* 2012;87(2):151–160.

Pragmatic and Observational Research

Publish your work in this journal

Pragmatic and Observational Research is an international, peer-reviewed, open access journal that publishes data from studies designed to reflect more closely medical interventions in real-world clinical practice compared with classical randomized controlled trials (RCTs). The manuscript management system is completely online and includes a very quick and fair peer-review

Submit your manuscript here: <https://www.dovepress.com/pragmatic-and-observational-research-journal>

system. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Dovepress