



Review article

Discovery of bioactive nitrated lipids and nitro-lipid-protein adducts using mass spectrometry-based approaches

Tânia Melo^{a,b}, Javier-Fernando Montero-Bullón^a, Pedro Domingues^a, M. Rosário Domingues^{a,b,*}^a Centro de Espectrometria de Massa, Departamento de Química e QOPNA, Universidade de Aveiro, Campus Universitário de Santiago, 3810-193 Aveiro, Portugal^b Departamento de Química & CESAM & ECOMARE, Universidade de Aveiro, 3810-193 Aveiro, Portugal

A B S T R A C T

Nitro-fatty acids (NO₂-FA) undergo reversible Michael addition reactions with cysteine and histidine residues leading to the post-translational modification (PTM) of proteins. This electrophilic character of NO₂-FA is strictly related to their biological roles. The NO₂-FA-induced PTM of signaling proteins can lead to modifications in protein structure, function, and subcellular localization. The nitro lipid-protein adducts trigger a series of downstream signaling events that culminates with anti-inflammatory, anti-hypertensive, and cytoprotective effects mediated by NO₂-FA. These lipoxidation adducts have been detected and characterized both in model systems and in biological samples by using mass spectrometry (MS)-based approaches. These MS approaches allow to unequivocally identify the adduct together with the targeted residue of modification. The identification of the modified proteins allows inferring on the possible impact of the NO₂-FA-induced modification. This review will focus on MS-based approaches as valuable tools to identify NO₂-FA-protein adducts and to unveil the biological effect of this lipoxidation adducts.

1. Introduction

During the last decade, nitrated lipid gained the interest of the scientific community, as new endogenous signaling molecules with important regulatory role in health and disease. Research on this is aimed at understanding the reactivity of reactive nitrogen species (RNS) with lipids, to unravel their occurrence in vivo and their biological roles. Among nitrated and nitroxidized lipids identified so far, the nitro-fatty acids (NO₂-FA) are best-known products of RNS. These products have been widely detected in several tissues [1–5] and biofluids [6–12], and are nowadays a hot topic in nitro lipidomics. NO₂-FA are considered important bioactive molecules and have been associated with anti-inflammatory [6,13–24], anti-hypertensive [25–32], and anti-thrombotic properties [31,33] and cytoprotective effects [2,34–37]. More recently, other nitrated and nitroxidized lipids [1,6–8,13,38] and also nitro derivatives of phospholipids (PL) [39,40] and triglycerides (TAG) [41] have been detected in biological samples and were associated with protective and beneficial effects, but they are scarcely studied. Also, esterified forms of NO₂-FA have been found as they can be generated either by direct nitration of the esterified fatty acyl moiety or by the incorporation of NO₂-FA [41].

NO₂-FA are also known as nitroalkenes derivatives of fatty acids since it includes a nitro group linked to the double bond (alkene group) of the unsaturated fatty acyl chain, and the nitro-alkene moiety makes

these derivatives highly reactive with electrophilic properties. These endogenous electrophilic lipids are capable to covalently link to proteins, via Michael addition [42], leading to the formation of lipoxidation adducts. This type of post-translational modification (PTM) of proteins can modulate protein function, which underlies some of the biological roles attributed to the NO₂-FA (Fig. 1). Some of these PTMs are shown to elicit a protective effect, which may provide clues for new therapeutic strategies and new drugs.

Detection of NO₂-FA and especially their lipoxidation adducts are still a challenge that is mostly addressed by MS approaches. MS-based approaches have been extensively applied in the study of NO₂-FA-protein adducts [4,7,34,42–44], providing detailed structural information of these adducts both in vitro and in vivo. LC-MS and MS/MS-based proteomics approaches have been performed to characterize the NO₂-FA protein adducts and the sites of adduction [34,42,45–48]. Very recently peptide adducts were also reported for NO₂-FA esterified in phospholipids using biomimetic in vitro studies and MS approaches [49].

In this review we will discuss the formation and type of nitrated FA found in biological systems, their structure and reactivity with proteins and characterization by a MS-based proteomic and lipidomic approach that allowed to disclose possible biological roles associate with nitrated lipids-protein adducts.

* Corresponding author at: University of Aveiro Campus Universitário de Santiago 3810-193 Aveiro, Portugal

E-mail addresses: taniameo@ua.pt (T. Melo), montero.bullon@ua.pt (J.-F. Montero-Bullón), p.domingues@ua.pt (P. Domingues), mrd@ua.pt (M.R. Domingues).




Fig. 1. Schematic representation of nitrated lipids pathways: from their generation to their biological effect.

2. Endogenous nitro-fatty acids

2.1. Chemistry and analysis

NO₂-FA are endogenous chemical entities generated by the attack of nitric oxide (NO)-derived reactive species, collectively called reactive nitrogen species (RNS), with unsaturated fatty acids. Nitrogen dioxide (NO₂), nitronium cation (NO₂⁺), and peroxyxynitrite/peroxyxynitrous acid, whose decomposition yields •NO₂ and hydroxyl radical (•OH), were reported as RNS that most frequently initiate nitration or nitroxidation reactions in biomolecules, including lipids. The prevalence and the yield of one process of these processes over the others are dependent on the oxygen levels, concentration of ROS versus RNS, the presence of secondary target molecules (scavengers, thiols, and transition metals), pH, and the partition between hydrophilic and hydrophobic milieu in cellular compartments [50]. The mechanism of FA nitration and nitroxidation in biological systems is not yet wholly undisclosed, and there are some alternative routes to explain the generation of NO₂-FA (Fig. 2). The free radical-induced nitration of FA mediated by •NO₂ is one of the most prominent reaction *in vivo* as a source of NO₂-FA [51]. The endogenous formation of NO₂-FA during free radical-mediated nitration reactions occurs in several biological processes such as digestion [52], metabolic stress, and inflammatory conditions [53]. Thus NO₂-FA were already identified in human red blood cells [8,9], plasma [6,8–10,12], urine [6,7], and tissues [1–5] at concentrations ranging from picomolar [12] to micromolar [6]. Dietary sources of nitrite can also lead to the generation of NO₂-FA via acid-catalyzed nitration reactions [52,54–56]. Recently, NO₂-FA were also reported in plants, fresh olives, and in extra virgin olive oil [56,57], which are considered external sources of NO₂-FA and can contribute to rising the endogenous levels of NO₂-FA [58].

The most common NO₂-FA identified *in vivo* were the nitrated forms of the nitro-oleic acid (NO₂-OA), nitro-linoleic acid (NO₂-LA), and nitro-conjugated linoleic acid (NO₂-CLA) [6,8,9]. However, the reaction of RNS with fatty acids can lead to the generation of several nitroalkene derivatives of other fatty acids, such as the nitro-palmitoleic acid (NO₂-POA), nitro linolenic acid (NO₂-LNA), nitro-arachidonic acid (NO₂-AA), nitro eicosapentaenoic acid (NO₂-EPA), and nitro-docosahexaenoic acid (NO₂-Dha) [6,8,38,51]. Different stereo or positional isomers of NO₂-FA were detected *in vitro* and in biological samples [6,8,38], as represented in Figs. 3–5 and Table 1.

Nevertheless, the recovery of NO₂-FA from biological samples, together with their detection and accurate quantification is a challenge

due to their low concentration, stability issues, metabolism (β-oxidation and saturation/desaturation reactions) [59], reactivity with proteins [59] and esterification [39–41], and different distribution among tissues and biofluids [6,12]. In line with these limitations, there has been an effort for the development of specific, standardized and reproducible methodologies of sample preparation and sensitive analytical approaches. The advent of more sensitive and sophisticated instruments, allied with the possibility of high-throughput analysis prompted by MS-based approaches, combined or not with liquid chromatography (LC-MS), has been the selected tool for the identification, structural characterization and quantification of free NO₂-FA. Indeed, the detection of these lipids is an indication to disclose the bioactive properties of these nitrated derivatives. The progress in the field of MS-based approaches enabled the discovery of NO₂-FA and contributed to the knowledge of NO₂-FA biological roles giving information on the structure-function relationships [60]. The development of improved sample preparation techniques, chromatographic separations, high-resolution instruments with great sensitivity, and innovative tools raised the possibility of detection, structural characterization and quantification of nitro lipids in human samples and animal models both under physiological and pathological conditions [1,3,4,8,59,61,62], and also in plants [56] as summarized in Table 1. The identification of NO₂-FA by MS is based on the detection of specific mass shifts compared to non-modified fatty acid (FA + 45 Da). Using MS-based approaches, NO₂-FA are preferentially analyzed in negative-ion mode as [M-H]⁻ ions [3,63]. However, positive-ion mode ionization can also occur, and NO₂-FA can also be identified as [M+H]⁺ [26], [M+Li]⁺ [51], [M+NH₄]⁺ [9,41], and [M+Na]⁺ ions [9]. Tandem mass spectra acquired both in positive- and negative-ion mode provides information that allows the structural characterization of NO₂-FA [8,38,51,63,64]. The fragmentation pattern of NO₂-FA under tandem MS (MS/MS) conditions includes the typical neutral losses of 47 Da (HNO₂) and product ions formed by cleavage of the hydrocarbon chain in the vicinity of the NO₂ group that allow assigning this modified FA. The differentiation of isomers can be addressed by the identification of reporter fragment ions that are formed by cyclization, followed by heterolytic carbon chain fragmentation, which allows pinpointing the correct position of the NO₂ group [2,63]. These product ions have been used as diagnostic ions broadly employed for targeted analysis and quantitation of specific NO₂-FA using reversed phase LC-MS/MS approaches, in biomimetic systems and in cells, tissues and biofluids [3,6,8–10,12,17,38,59,65,66]. Structural information gathered by using MS studies can be further confirmed by infrared and nuclear




Fig. 2. Representative mechanisms of nitro-fatty acid ($\text{NO}_2\text{-FA}$) formation. Radical-induced nitration of unsaturated fatty acids by nitrogen dioxide (NO_2) yields a β -nitroalkyl radical that can further react with other NO_2 generating the nitronitrite intermediates. Further loss of nitrous acid (HNO_2) leads to the generation of the nitroalkene derivatives also called $\text{NO}_2\text{-FA}$. Electrophilic substitution at the double bond mediated by nitronium cation (NO_2^+) also yields $\text{NO}_2\text{-FA}$.

magnetic resonance analysis for the confirmation of the functional groups and the final structure [6,9,10,30,31,51].

The generation of $\text{NO}_2\text{-FA}$ can be considered as the first step of nitration reactions. These species can be precursors of other nitrated and nitroxidized species because $\text{NO}_2\text{-FA}$ can undergo additional reaction with ROS and RNS to be further nitrated, leading to the formation of nitroso, dinitroso, nitronitroso, di and trinitro species, or oxidized generating the assorted nitroxidized species as nitrohydroxy, nitrohydroperoxy, nitro-epoxy and nitro-keto (Table 1) [3,6,8,26,54,64]. All of these derivatives were already identified by (LC)-MS and characterized by (LC)-MS/MS [1,6,9,38]. In fact, the great sensitivity of MS-based approaches allowed to identify both nitro and nitrohydroxy derivatives of palmitoleic, oleic, linoleic, linolenic, arachidonic and eicosapentaenoic acids in human plasma and urine [6]. However by far the $\text{NO}_2\text{-FA}$ are the most studied mostly because, in opposition to other nitrated and nitroxidized FA, they are electrophilic molecules with great capability to react with protein with the formation of lipoxidation adducts.

2.2. Biological roles of nitro-fatty acids as new metabolic mediators, signaling molecules, and new therapeutic drugs candidates

$\text{NO}_2\text{-FA}$ have raised the interest of the scientific community in last years, mainly because of their biological roles as key mediators in physiological and pathophysiological processes, as demonstrated in a variety of preclinical animal models of disease and in plants [2,5,13,15,20,28,32,34,45,56]. They were assigned as biologically relevant and putative signaling molecules in cardiovascular disease [28,33], myocardial ischemia/reperfusion and ischemia preconditioning [1,2,24], nephropathy [24], renal ischemia/reperfusion

[24], diabetes and metabolic syndrome [14], pulmonary inflammation [15,67] and chronic inflammatory disease [65]. $\text{NO}_2\text{-FA}$ reach endogenous levels that allow them to mediate pivotal signaling actions as cytoprotective and pro-survival players [2,34–37], and based on their pleiotropic actions, $\text{NO}_2\text{-FA}$ has emerged as potential therapeutic agents with high potential for therapeutic use (Table 2). In fact, $\text{NO}_2\text{-FA}$ already undergo clinical trials [68]. The 10- $\text{NO}_2\text{-OA}$ (CXA-10) demonstrated promising pharmacokinetic and pharmacodynamics characteristics during preclinical experiments [61,68,69]. CXA-10 is currently in phase II clinical trials for the treatment of chronic inflammatory and metabolic-related diseases, namely focal segmental glomerulosclerosis and pulmonary arterial hypertension, since it demonstrated beneficial effects when administered via intravenous injections or through ingestion [61,68,69].

The biological actions of $\text{NO}_2\text{-FA}$ are mediated via a) decay reactions and transduction of nitric oxide (NO) signaling actions [29,70], since $\text{NO}_2\text{-FA}$ can be considered NO donor; b) via receptor-dependent and c) via electrophilic adduction reactions to proteins [42], with formation of lipoxidation adducts. All these processes mediate important and specific signaling roles. These signaling actions are summarized in Table 2. Nitric oxide release by $\text{NO}_2\text{-FA}$ has been associated with potential antioxidant properties through inhibition of lipid peroxidation process [71]. Additionally, the release of NO by both $\text{NO}_2\text{-FA}$ and nitrohydroxy FA derivatives has also been related with vasorelaxation properties of these nitrated lipid [26,29–31,51]. The nitro derivatives of arachidonic acid, $\text{NO}_2\text{-AA}$ and nitrohydroxy-AA, were also reported to be able to release NO and thus to induce cGMP-dependent vasorelaxation in rat aortic ring in an endothelium-independent manner [26,31,51]. $\text{NO}_2\text{-LA}$, $\text{NO}_2\text{-cLA}$ and nitrohydroxy-LA promoted vessel relaxation via cGMP-dependent and endothelial-independent manner in




Fig. 3. Proposed structures for nitro-oleic (NO₂-OA) and nitro-linoleic acids (NO₂-LA), with assignment of their different positional isomers, which were previously detected in in vitro studies and/or biological samples.

pre-constricted rat aortic rings [29,30]. Nevertheless, the ¹NO release by nitro lipids remains a controversial issue, and at some level, considered of minor relevance at endogenous levels [28,29,51,70,72]. Actually, ¹NO signaling actions mediated by NO₂-FA mainly occurs via cGMP-independent mechanisms. NO₂-FA modulates endothelial (eNOS) and inducible nitric oxide synthase (iNOS) gene expression and activity and consequently the eNOS- and iNOS-mediated ¹NO generation and reactions. Also, NO₂-FA modulates a broad array of signaling pathways that culminates with the downstream activation or inactivation of ¹NO signaling [67,73,74].

The covalent adduction to key proteins propelled by the electrophilic character of NO₂-FA seems to be the most prominent mechanism by which these nitro lipids spread their modulatory and protective actions. The identification and characterization of these NO₂-FA-protein adducts in distinct biological conditions have been achieved by reversed phase LC-MS-based proteomics approaches [28,42,56,59,75,76]. This topic will be discussed in more detail in the next section.

As endogenous molecules, NO₂-FA undergo a series of metabolic, trafficking and clearance pathways that influences the regulation of activity, half-life and levels of free NO₂-FA. Protein adduction and esterification in complex lipids [70,77,78] are considered as reservoirs of NO₂-FA, allowing to regulate their endogenous levels [70,77,78]. NO₂-FA-protein adducts are reversible in biological systems [59,73,79] and NO₂-FA esterified forms can be hydrolyzed and mobilized by esterases and lipases, allowing NO₂-FA to return to free active forms [70,80]. NO₂-FA can be metabolized via β -oxidation that mediates the formation of shorter and more polar electrophilic species [59] that retains the electrophilic power, but also to inactive nitroalkane species [7,59]. In fact, in humans and rodents, the bio-elimination pathways of 10-NO₂-OA involves the generation of a series of shorter metabolites that were detected in urine using C18-HPLC-ESI-MS and MS/MS using both LTQ Velos Orbitrap and API 5000 triple quadrupole instruments [61].

However, the electrophilic functionality of NO₂-FA is irreversibly inactivated after reduction and conversion to the correspondent nitroalkane derivative by the nitroalkene saturase prostaglandin reductase-1 [81]. Both saturation and desaturation of the double bond of NO₂-FA are related with the generation of non-electrophilic NO₂-FA [59], which are nitro derivatives without signaling abilities. Adduction to peptides or proteins seems to have other proposes, such as the case of conjugation with GSH, which increases the urinary excretion rate of NO₂-FA excreted in urine [82]. Incorporation of NO₂-FA into lipoproteins is another way for NO₂-FA to enter in circulation and to be systemically distributed among tissues. The modulation of all of these diverse pathways will impact the potential reactivity, the efficacy of signaling actions and behavior of these nitration products.

The signaling actions of NO₂-FA are also mediated through the modulation of the structure and regulation of the expression and activity of anti- and pro-inflammatory proteins, heat shock proteins and phase II antioxidant response proteins. The capability of NO₂-FA to react with specific peptides and proteins determines the role of this nitrated lipids in redox regulation with consequence in cell signaling, as will be described in the following section.

3. Nitro-fatty acids and protein lipoxidation adducts

3.1. Main target and biological significance of PTM by nitro-fatty acids

NO₂-FA are electrophilic compounds, able to react via reversible Michael addition with nucleophiles within key proteins, leading to the formation of lipid-protein adducts (lipoxidation) in a process generally denominated nitroalkylation [83,84]. The target nucleophiles in peptides and proteins include the deprotonated thiolate group of cysteine and the nucleophilic amino group of the imidazole moiety of histidine or the amino groups of lysine and arginine [83–85]. The high electro-negative olefinic NO₂ group facilitate the addition to the double bond of

Nitrated Arachidonic Acid (NO₂-AA) isomers




Fig. 4. Proposed structures for nitro arachidonic acid (NO₂-AA), with assignment of its different positional isomers, which were previously detected in in vitro studies and/or biological samples.

Nitrated Docosahexaenoic Acid (NO₂-DHA) isomers




Fig. 5. Proposed structures for nitro-docosahexaenoic acid (NO₂-DHA), with assignment of its different positional isomers, which were previously detected in in vitro studies and/or biological samples.

Table 1
Nitro-fatty acids identified in biological samples and *in vitro* mimetic model systems.

<i>In vitro</i> mimetic model systems				
NO ₂ -FA	Isomer	Experimental model	Method	Ref.
Nitro-oleic acid (NO₂-OA)				
NO ₂ -OA	9-NO ₂ -OA 10-NO ₂ -OA	Gastric juice artificial + NO ₂ ⁻	C18-HPLC-ESI-MS and MS/MS in a API 4000 triple quadrupole and LTQ Orbitrap Velos	[56]
NO ₂ -OA		Pancreatic lipase-digested EVOO MPO + H ₂ O ₂ + NO ₂ ⁻ ONOO ⁻ NO ₂ ⁻ in acidic conditions	C18-HPLC-ESI-MS and MS/MS in a hybrid triple quadrupole-linear ion trap (4000 Q-Trap)	[6]
NO ₂ -OA	9-NO ₂ -OA 10-NO ₂ -OA	●NO ₂	C18-HPLC-ESI-MS and MS/MS in API 2000 triple quadrupole	[80]
Nitro-linoleic acid (NO₂-LA)				
NO ₂ -LA	9-NO ₂ -LA 10-NO ₂ -LA 12-NO ₂ -LA 13-NO ₂ -LA	Gastric juice artificial + NO ₂ ⁻	C18-HPLC-ESI-MS and MS/MS in an API 4000 triple quadrupole and LTQ Orbitrap Velos	[56]
NO ₂ -LA		Pancreatic lipase-digested EVOO		
NO ₂ -LA		NO ₂ ⁻ in acidic conditions	C18-HPLC-ESI-MS and MS/MS in a Quattro triple quadrupole	[10]
Nitro-conjugated linoleic acid (NO₂-cLA)				
NO ₂ -cLA	8-NO ₂ -cLA 9-NO ₂ -cLA 11-NO ₂ -cLA 12-NO ₂ -cLA	Gastric juice artificial + NO ₂ ⁻	C18-HPLC-ESI-MS and MS/MS in an API 4000 triple quadrupole and LTQ Orbitrap Velos	[56]
NO ₂ -cLA	9-NO ₂ -cLA	Pancreatic lipase-digested EVOO		
NO ₂ -cLA	9-NO ₂ -cLA	MPO + H ₂ O ₂ + NO ₂ ⁻ ONOO ⁻ ●NO ₂	C18-HPLC-ESI-MS and MS/MS in an API 5000 triple quadrupole, API Q-Trap 4000, and Velos Orbitrap	[3]
NO ₂ -cLA	12-NO ₂ -cLA NO ₂ -cLA	Photocontrollable peroxyxynitrite donor 2,3,5,6-tetramethyl-4-(methylnitrosoamino)phenol (P-NAP)	ESI-MS and MS/MS in a hybrid triple quadrupole-linear ion trap (4000 Q-Trap)	[38]
Cholesteryl-nitro linoleic acid (Chol-NO₂-LA)				
Chol-NO ₂ -LA		NO ₂ ⁻ in acidic conditions	C18-HPLC-ESI-MS/MS in a Quattro II triple quadrupole	[9]
Chol-NO ₂ -LA		NO ₂ ⁻ in acidic conditions	ESI-MS and MS/MS in a 2000 Q-Trap C18-HPLC-ESI-MS and MS/MS in a 2000 Q-Trap	[66]
Nitro-arachidonic acid (NO₂-AA)				
NO ₂ -AA		●NO ₂	C18-HPLC-ESI-MS and MS/MS in an Esquire ion trap	[26]
NO ₂ -AA	9-NO ₂ -AA 12-NO ₂ -AA 14-NO ₂ -AA 15-NO ₂ -AA	NO ₂ ⁻ in acidic conditions	C18-HPLC-ESI-MS and MS/MS in a hybrid quadrupole-linear ion trap	[51]
Biological samples				
Nitro-palmitoleic acid (NO₂-POA)				
NO ₂ -POA		Human plasma and urine	C18-HPLC-ESI-MS and MS/MS in a hybrid triple Q-linear ion trap (4000 Q-Trap)	[6]
Nitrohydroxy-palmitoleic acid (NO₂OH-POA)				
NO ₂ OH-POA		Human plasma and urine	C18-HPLC-ESI-MS and MS/MS in a hybrid triple Q-linear ion trap (4000 Q-Trap)	[6]
Nitro-oleic acid (NO₂-OA)				
NO ₂ -OA		Human red cells, plasma and urine	C18-HPLC-ESI-MS and MS/MS in a hybrid triple Q-linear ion trap (4000 Q-Trap)	[6]
NO ₂ -OA	9-NO ₂ -OA 10-NO ₂ -OA	Myocardial heart tissue from a murine model of focal myocardial ischemia/reperfusion	C18-HPLC-ESI MS/MS	[1]
NO ₂ -OA	NO ₂ -OA and β-oxidation metabolites	NO ₂ -OA acute intravenous treatment of mice with LPS-induced inflammation	C18-HPLC-ESI-MS/MS in an API 5000 triple quadrupole	[107]
NO ₂ -OA	NO ₂ -OA and its metabolic derivatives	Human and rat urine after intravenous administration of NO ₂ -OA	C18-HPLC-ESI-MS and MS/MS in a LTQ Velos Orbitrap and API 5000 triple quadrupole	[61]
NO ₂ -OA	NO ₂ -OA and its metabolic derivatives	Mitochondrial extracts from rat hearts after ischemia-reperfusion	BME trans-nitroalkylation + C18-HPLC-ESI-MS and MS/MS in a 4000 Q trap hybrid triple quadrupole-linear ion trap	[62]
Dinitro-OA		Rat cardiomyocytes treated with peroxyxynitrite donor 3-morpholinosydnonimine (SIN-1)	C18-HPLC-ESI-MS and MS/MS in a hybrid triple quadrupole-linear ion trap (4000 Q-Trap)	[38]
NO ₂ -OA	NO ₂ -OA and its Saturation, Desaturation β-oxidation metabolic derivatives	Plasma from NO ₂ -OA-treated mice	C18-HPLC-ESI MS/MS coupled to an API 4000 hybrid triple quadrupole or API 5000 triple quadrupole	[59]
NO ₂ -OA	NO ₂ -OA saturation derivatives	NO ₂ -OA-treated BAEC cells	C18-HPLC-ESI MS/MS coupled to an API 4000 hybrid triple quadrupole or API 5000 triple quadrupole	[59]
NO ₂ -OA	NO ₂ -OA and its derivatives	Liver lipid extracts from NO ₂ -OA-treated mice	C18-HPLC-ESI MS/MS coupled to an API 4000 hybrid triple quadrupole or API 5000 triple quadrupole	[59]
Nitrohydroxy-oleic acid (NO₂OH-OA)				
NO ₂ OH-OA		Human red cells, plasma and urine	C18-HPLC-ESI-MS and MS/MS in a hybrid triple Q-linear ion trap (4000 Q-Trap)	[6]

(continued on next page)

Table 1 (continued)

<i>In vitro</i> mimetic model systems				
NO ₂ -FA	Isomer	Experimental model	Method	Ref.
Nitro-linoleic acid (NO₂-LA)				
NO ₂ -LA		Myocardial heart tissue from a murine model of focal myocardial ischemia/reperfusion	C18-HPLC-ESI MS and MS/MS	[1]
NO ₂ -LA		Human plasma and urine	C18-HPLC-ESI-MS and MS/MS in a hybrid triple Q-linear ion trap (4000 Q-Trap)	[6]
NO ₂ -LA		Human blood plasma	C18-HPLC-ESI-MS and MS/MS in a Quattro triple quadrupole	[10]
NO ₂ -LA	9-NO ₂ -LA 12-NO ₂ -LA	Human red cell membranes and plasma	C18-HPLC-ESI-MS and MS/MS in a hybrid triple Q-linear ion trap	[8]
NO ₂ -LA	NO ₂ -LA and its metabolic derivatives	Mitochondrial extracts from rat hearts after ischemia-reperfusion	BME trans-nitroalkylation + C18-HPLC-ESI-MS and MS/MS in a 4000 Q trap hybrid triple quadrupole-linear ion trap	[62]
Nitrohydroxy-linoleic acid (NO₂OH-LA)				
NO ₂ OH-LA		Myocardial heart tissue from a murine model of focal myocardial ischemia/reperfusion	C18-HPLC-ESI MS and MS/MS	[1]
NO ₂ OH-LA		Human plasma and urine	C18-HPLC-ESI-MS and MS/MS in a hybrid triple Q-linear ion trap (4000 Q-Trap)	[6]
Nitro-oxo-linoleic acid (NO₂-oxo-LA)				
NO ₂ -oxo-LA		Myocardial heart tissue from a murine model of focal myocardial ischemia/reperfusion	C18-HPLC-ESI MS/MS	[1]
Nitro-conjugated linoleic acid (NO₂-cLA)				
NO ₂ -cLA		Plasma and vaginal lavages after cLA inoculation in the vaginal lumen from mice infected intravaginally with HSV-2	C18-HPLC-MS/MS in a 6500+ Q-trap or a API 5000	[16]
Nitro-linolenic acid (NO₂OH-LNA)				
NO ₂ -LNA		Human plasma and urine	C18-HPLC-ESI-MS and MS/MS in a hybrid triple Q-linear ion trap (4000 Q-Trap)	[6]
Nitrohydroxy-linolenic acid (NO₂OH-LNA)				
NO ₂ OH-LNA		Human plasma and urine	C18-HPLC-ESI-MS and MS/MS in a hybrid triple Q-linear ion trap (4000 Q-Trap)	[6]
Nitro-arachidonic acid (NO₂-AA)				
NO ₂ -AA		Human plasma and urine	C18-HPLC-ESI-MS and MS/MS into a hybrid triple Q-linear ion trap (4000 Q-Trap)	[6]
NO ₂ -AA		Rat cardiomyocytes treated with peroxynitrite donor 3-morpholinosydnonimine (SIN-1)	C18-HPLC-ESI-MS and MS/MS in a hybrid triple quadrupole-linear ion trap (4000 Q-Trap)	[38]
Nitrohydroxy-arachidonic acid (NO₂OH-AA)				
NO ₂ OH-AA		Human plasma and urine	C18-HPLC-ESI-MS and MS/MS in a hybrid triple Q-linear ion trap (4000 Q-Trap)	[6]
Nitro-Eicosapentaenoic acid (NO₂-EPA)				
NO ₂ -EPA		Human plasma and urine	C18-HPLC-ESI-MS and MS/MS in a hybrid triple Q-linear ion trap (4000 Q-Trap)	[6]
Nitrohydroxy-Eicosapentaenoic acid (NO₂OH-EPA)				
NO ₂ OH-EPA		Human plasma and urine	C18-HPLC-ESI-MS and MS/MS in a hybrid triple Q-linear ion trap (4000 Q-Trap)	[6]
Nitro-Docosahexaenoic acid (NO₂-DHA)				
NO ₂ -DHA and dinitro-DHA		Rat cardiomyocytes treated with peroxynitrite donor 3-morpholinosydnonimine (SIN-1)	C18-HPLC-ESI-MS and MS/MS in a hybrid triple quadrupole-linear ion trap (4000 Q-Trap)	[38]
Nitrohydroxy-Docosahexaenoic acid (NO₂OH-DHA)				
NO ₂ OH-DHA		Rat cardiomyocytes treated with peroxynitrite donor 3-morpholinosydnonimine (SIN-1)	C18-HPLC-ESI-MS and MS/MS in a hybrid triple quadrupole-linear ion trap (4000 Q-Trap)	[38]
Nitrohydroxy-Docosapentaenoic acid (NO₂OH-DPA)				
NO ₂ OH-DPA		Rat cardiomyocytes treated with peroxynitrite donor 3-Morpholinosydnonimine (SIN-1)	C18-HPLC-ESI-MS and MS/MS in a hybrid triple quadrupole-linear ion trap (4000 Q-Trap)	[38]
Nitro-conjugated linoleic acid (NO₂-cLA)				
NO ₂ -cLA	9-NO ₂ -cLA 12-NO ₂ -cLA	Pancreatic lipase-digested EVOO	C18-HPLC-ESI-MS and MS/MS in an API 4000 triple quadrupole and LTQ Orbitrap Velos	[56]
NO ₂ -cLA	9-NO ₂ -cLA 12-NO ₂ -cLA β-oxidation-metabolic derivatives of NO ₂ -cLA	Urine of healthy humans	C18-HPLC-ESI-MS and MS/MS in a LTQ Velos Orbitrap and AB 5000 or API4000 Q-trap triple quadrupole	[7]
NO ₂ -cLA	9-NO ₂ -cLA	Rodents urine, plasma, and tissues (stomach, small intestine, colon, liver) after supplementation with cLA + NO ₂ ⁻ and gastric acidification	C18-HPLC-ESI-MS and MS/MS in an API 5000 triple quadrupole, API Q-Trap 4000, and Velos Orbitrap	[3]
	12-NO ₂ -cLA	Rodents liver and cardiac mitochondria incubated with NO ₂ ⁻ in acidic conditions Rodents cardiac tissue under ischemia-reperfusion Raw 264.7 macrophages stimulated with LPS/IFN _γ Healthy human plasma		
NO ₂ -cLA	9-NO ₂ -cLA 12-NO ₂ -cLA Reduction and β-oxidation-metabolic derivatives	RAW264.7 macrophages stimulated with LPS/IFN _γ and M1, M2 and M0 polarized bone marrow-derived macrophages (BMDM) treated with cLA Mice Peritoneal exudates after zymosan-A induced peritonitis and cLA supplementation	C18-HPLC-ESI-MS and MS/MS in an API 5000 or a Q-Trap 6500+ and LTQ Velos Orbitrap	[13]

(continued on next page)

Table 1 (continued)

<i>In vitro</i> mimetic model systems				
NO ₂ -FA	Isomer	Experimental model	Method	Ref.
NO ₂ -cLA	NO ₂ -cLA and β-oxidation-metabolic derivatives	Urine and plasma healthy humans after ingestion of nitrite, nitrate and cLA	C18-HPLC-ESI-MS and MS/MS in a 5000 triple quadrupole	[58]
Cholesteryl-nitro linoleic acid (Chol-NO₂-LA)				
Chol-NO ₂ -LA		Human blood plasma and lipoproteins from normolipidemic/healthy subjects	C18-HPLC-ESI/MS/MS in a Quattro II triple quadrupole	[9]
Chol-NO ₂ -LA		J774.1 macrophages stimulated with LPS/IFN γ	C18-HPLC-ESI-MS and MS/MS in a 2000 Q-Trap	[66]

Table 2

Modulation of target signaling pathways by NO₂-FA and related biological properties.

Anti-inflammatory
- NF- κ B ↓ [1,13,15,21,23,46,60,66,86,90,108]
- TL4R signaling ↓ [107]
- PG H synthase ↓ [86,109]
- 5-LOX ↓ [45]
- STAT1 ↓ [87,92]
- Cytokine production ↓ [1,5,7,13,20,21,23,24,66,86,110–112]
- MPO ↓ [24]
- Leukocyte recruitment, adhesion and infiltration ↓ [13,23,24,87,108,112]
- iNOS ↓ [13,24,36,51,60,66,86,112]
- COX-2 ↓ [112]
- PGE ₂ ↓ [112]
- Leukocytes number and activity ↓ [5,13,18,108]
- TNF- α ↓ [23]
- Xanthine oxidoreductase (XOR) ↓ [73]
- MKP-1 ↑ [92]
- PPAR γ ↑ [6,15,21,65,100]
- TRPA-1 ↑ [98]
- CD36 expression ↑ [100]
- Heme oxygenase 1 (HO-1) ↑ [17,19,22,23,60,66,67,89,93]
- STING ↓ [16]
- PDI ↓ [95]
- NADPH oxidase (NOX 2) ↓ [24,74,113]
Vasorelaxation
- Nitric oxide ↑ [29–31,51]
- eNOS ↑ [114]
- Ang II-induced vasoconstriction ↓ [28]
Antioxidant
- Lipid peroxidation ↓ [71,86]
- Nrf2 ↑ [13,15,22,46,60,88]
- HO-1 ↑ [17,19,22,60,66,89]
- Reduction of protein nitration and oxidation [87]
- Heat shock response (HSP) proteins [22]
- XOR ↓ [73]
- NOX 2 ↓ [24,74,113]
- O ₂ [•] , ONOO ⁻ and NO production ↓ [36,48,86,108,111,113]
Anti-hypertensive
- AT1R ↓ [28]
- sEH ↓ [32]
- Smooth muscle cell proliferation ↓ [111]
- Prostaglandin F ₂ ↓ [111]
- Differentiation of myofibroblast through Smad2 ↓ [113]
Anti-hyperglycemic
- PPAR γ ↑ [6,28,100]
- Glucose uptake ↑ [110]
- Insulin sensitivity ↑ [6,28,100]
Anti-thrombotic
- Platelets activation and clotting ↓ [18,31,33]
- Platelet production of thromboxane ↓ [109]
- Thrombin-induced aggregation ↓ [33]
- PGHS ↓ [109,115,116]
Cytoprotective
- ANT1 ↑ [2,34]
- UCP-2 ↑ [2]
- Mitochondrial dysfunction ↓ [36,37]
- Mitochondrial respiratory complex ↑ [48]
- Metabolic shift ↑ [48]
Anti-tumorigenic
- Tumor cell proliferation, migration and invasion ↓ [90]

the unsaturated hydrocarbon chain of NO₂-FA. This addition generates an important positive density of charge in the methylenic β-carbon adjacent to the nitration site. The oxygens of the NO₂ group withdraw electrons and the double bond is rearranged over the C–N bond, generating a carbocation. This conjugation makes the β-carbon adjacent to the NO₂ group electron poor and with potential reactivity. The NO₂-FA-protein covalent adducts generated during the nitroalkylation process are reversible, which seems to be related with the possibility of redox regulation [59,73,79] and thus can be associated with the apparent lack of toxicity of these modified lipids. All of the aforementioned characteristics make NO₂-FA as promising pharmacological compounds. In fact, pre-clinical and human trials has demonstrated the NO₂-FA favorable pharmacokinetics and safety.

The formation of NO₂-FA adducts with proteins is considered a key PTM of proteins. This modification of functionally-relevant proteins can modulate the patterns of gene expression programs, transcription factors function, enzyme function and activity, metabolic and inflammatory responses, and cell signaling networks [50,59,73,84]. This lead to a series of downstream signaling events that are intrinsically related to the biological signaling roles of NO₂-FA [2,6,13–37,61,74,75,86–90] (Table 2). The activation of several of these pathways are considered essential for restoring the homeostasis and the redox balance and makes NO₂-FA promising pharmacological compounds [91].

There are several proteins reported to be targets of NO₂-FA electrophilic reactivity, for example, the p65 subunit of NF- κ B [1,23,92], heme oxygenase-1 (HO-1) [17,19,22,67,89,93], mitogen-activated protein kinase (MAPK) phosphatase 1 (MPK-1) [92], Kelch-like ECH-associated protein 1 (Keap 1) [17,22,46,88], metalloproteinases (MMP-7 and MMP-9) [75], glyceraldehyde-3-phosphate dehydrogenase (GAPDH) [42,94], protein disulfide isomerase (PDI) [95], and transient receptor potential (TRP) channels [96–99] (Table 3). NO₂-FA can also conduct their biological signaling roles by a receptor-dependent signaling action and peroxisome proliferator-activated receptor gamma (PPAR γ) is one of the main targets, which is a significant route for the anti-inflammatory effect associated with NO₂-FA derivatives [6,23,47,65,93,100–102].

The nitro lipoxidation PTM of the proteins shown in the Table 3 have been correlated with specific biological effects. For example, nitroalkylation of the p65 subunit of NF- κ B [23], induction of HO-1 expression [93], PPAR γ modulation [100], inhibition of the correct assembly of the active NADPH oxidase (NOX2) [74], and inhibition of both reductase and chaperone activities of PDI and possible prevention of NOX2 activation [95] have been associated with the anti-inflammatory properties of NO₂-FA. Another important anti-inflammatory action of NO₂-FA is attributed to their capability to induce PTM of 5-Lipoxygenase (5-LOX) limiting the inflammation induced by the 5-LOX-dependent leukotriene synthesis. This point deserves to be further explored as a potential therapeutic/pharmacological strategy due to the physiological relevance of 5-LOX, namely in inflammation [45]. Induction of HO-1 and activation of Nrf2 have been correlated with protection against oxidative stress and antioxidant actions of NO₂-FA [93]. Activation of PPAR γ by NO₂-FA has also been associated with glucose uptake and anti-hyperglycemic effects [100]. Inhibition of the

Table 3
Nitro-fatty acids lipoxidation adducts identified in biological samples and *in vitro* mimetic model systems by using mass spectrometry-based approaches.

Protein/peptide	Model system	Method	Molecular mechanism signaling action	Biological role	Ref.
Nitro-oleic acid (NO₂-OA)					
Cysteine	Incubation of NO ₂ -OA and cysteine Incubation of NO ₂ -OA and cysteine Whole olives, mesocarp and peel	C18-HPLC-MS and MS/MS in a triple quadrupole API 4000 C18-HPLC-MS and MS/MS in a triple quadrupole C18-HPLC-MS and MS/MS in a triple quadrupole API 4000			[56] [43] [56] [62]
GSH	NO ₂ -OA-Cys adduct generation after incubation between NO ₂ -OA and GSH NO ₂ -OA-Cys adduct generation after incubation between NO ₂ -OA and GSH	BME trans-nitroalkylation + C18-HPLC-ESI-MS and MS/MS in a Q trap 4000 (BME trans-nitroalkylation +) C18-HPLC-ESI-MS and MS/MS in a hybrid triple quadrupole API 4000 or API 5000 triple quadrupole			[59]
	Plasma from NO ₂ -OA-treated mice	(BME trans-nitroalkylation) C18-HPLC-ESI-MS and MS/MS in a hybrid triple quadrupole API 4000 or API 5000 triple quadrupole			[59]
	NO ₂ -OA-Cys adduct generation after incubation between NO ₂ -OA and GSH	ESI-MS in LCQ ion trap Tryptic digestion + C18-HPLC-ESI-MS and MS/MS in an ESI-LCQ ion trap			[42]
	Red blood cells obtained from healthy humans	C18-HPLC-ESI-MS/MS in a Q-Trap 4000 C18-HPLC-ESI-MS/MS in a Q-Trap 4000	Translocation to membrane	Regulation of enzyme function, cell signaling, and protein trafficking	[42]
GAPDH	Cytosolic and membrane-associated protein fractions from red blood cells obtained from healthy humans	SDS-PAGE under non-reducing and denaturing conditions + Tryptic digestion + C18-nanospray LC-MS and MS/MS in a LTQ ion trap	Translocation to membrane	Regulation of enzyme function, cell signaling, and protein trafficking	[42]
Cys416	Incubation of NO ₂ -OA and GAPDH	MALDI-TOF MS (Voyager DE PRO system) C18-HPLC-MS in a LTQ ion trap			[42]
Cys418		Tryptic digestion and MALDI-TOF MS (Voyager DE PRO system)			
His125		Tryptic digestion + C18-HPLC-ESI-MS in an ESI-LCQ ion trap			
His360	Incubation of NO ₂ -OA and GAPDH	Tryptic digestion + C18-nanospray LC-MS and MS/MS in a LTQ ion trap			[62]
His362		Electrophoresis under reducing conditions + BME trans-nitroalkylation + C18-HPLC-ESI-MS and MS/MS in a 4000 Q trap hybrid triple quadrupole-linear ion trap			
His367	Incubation of NO ₂ -OA and human recombinant 5-LOX	Tryptic digestion and C18-nanoHPLC-ESI-MS and MS in an Orbitrap XL	Irreversible inhibition of 5-LOX activity and	Anti-inflammatory	[45]
His372					
His432					
Keap1	Incubation of NO ₂ -OA and recombinant Keap1	Tryptic digestion C18-HPLC-MS and MS/MS in a LTQ	Prevention of lung injury and systemic immune responses		[46]
Cys38					
Cys151					
Cys226					
Cys273	Human embryonic kidney (HEK)-293T cells transfected with recombinant Keap1 and treated with NO ₂ -OA		Release of Nrf2 transcription factor to the nucleus for induction of expression of antioxidant phase II enzymes	Antioxidant	[46]
Cys288					
Cys257					
Cys489					

(continued on next page)

Table 3 (continued)

Protein/peptide	Model system	Method	Molecular mechanism signaling action	Biological role	Ref.
Cathepsin S (Cat S)	Incubation of NO ₂ -OA with a synthetic Cat S peptide (Cat S23–29)	C18-HPLC-MS and MS/MS in a Q Exactive Hybrid Orbitrap	Downregulation of Cat S expression and activity	Tissue Protection Anti-inflammatory	[5]
Fp subunit of mitochondrial complex II	Incubation of NO ₂ -OA with recombinant human complex II Fp subunit	Tryptic digestion and C18-HPLC-MS and MS/MS in a LTQ-XL	Inhibition of mitochondrial respiration complex II and O ₂ ^{•-} production	Cytoprotective	[48]
His2					
His5					
His6	Rat heart mitochondria treated with OA-NO ₂	Blue native electrophoresis, BME trans-nitroalkylation, C18-HPLC-MS and MS/MS in a hybrid triple-quadrupole linear ion trap mass spectrometer (4000 Q trap)	Promotion of glycolysis	Antioxidant	
Cys9					
Cys14					
AT ₁ R	HEK293 cells overexpressing AT ₁ R treated with NO ₂ -OA	Immunoprecipitation of AT ₁ R from cell lysates, BME trans-nitroalkylation reaction of AT ₁ R-bound NO ₂ -OA, and C18-HPLC-MS and MS/MS in a 4000 Q-Trap triple quadrupole	Inhibits AT ₁ R-dependent vasoconstriction by reduction of heterotrimeric G-protein coupling and inhibition of IP ₃ and calcium mobilization	Anti-hypertensive	[28]
MMP-7	Incubation of NO ₂ -OA with recombinant human proMMP-7 and proMMP-9	C18-HPLC-ESI-MS and MS/MS in a LTQ-XL	Modulation of proteolytic activity	Anti-inflammatory	[75]
Cys70					
MMP-9					
Cys100					
Albumin	Plasma from intraperitoneal NO ₂ -OA-treated mice	Electrophoresis under reducing conditions + BME trans-nitroalkylation + C18-HPLC-ESI-MS and MS/MS in a 4000 Q trap hybrid triple quadrupole-linear ion trap	Decrease of enzyme expression		[62]
	Plasma from NO ₂ -OA-treated mice	Electrophoresis + BME trans-nitroalkylation + C18-HPLC-ESI-MS and MS/MS in a hybrid triple quadrupole API 4000 or API 5000 triple quadrupole			[59]
PPAR γ	Incubation of NO ₂ -OA with human recombinant PPAR γ LBD	Tryptic digestion and C18-HPLC-MS and MS/MS in a LTQ	Activation of PPAR γ -related gene expression for glucose regulation and adipogenesis	Anti-hyperglycemic	[47]
Cys285	HEK 293 T cells were transfected with PPAR γ and treated with NO ₂ -OA	Immunoprecipitation, gel electrophoresis, BME- trans-nitroalkylation and ESI-MS and MS/MS in a hybrid triple quadrupole-linear ion trap mass spectrometer (4000 Q Trap,	Decrease in adipogenesis Increase glucose uptake	Anti-adipogenic effect	
His266					
His323					
His425					
His449					
STING	Incubation human STING-transfected HEK293T cells with 10-NO ₂ -OA	Purification with magnetic beads, tryptic digestion and MALDI LTQ Orbitrap XL	Deregulation of STING palmitoylation Inhibition of STING signaling Inhibition the release of type I IFN	Anti-inflammatory	[16]
Cys88					
Cys91					
His16					
Nitro-linoleic acid (NO ₂ -LA)					
cysteine	NO ₂ -LA-Cys adduct generation after incubation between NO ₂ -LA and cysteine	(C18-HPLC)-ESI-MS and MS/MS in a triple quadrupole			[107]
GSH	NO ₂ -LA-Cys adduct generation after incubation between NO ₂ -LA and GSH	ESI-MS in LCQ ion trap			[42]
	NO ₂ -LA-Cys adduct generation after incubation between NO ₂ -LA and GSH	C18-HPLC-ESI-MS/MS in a Q-Trap 4000			[62]
	NO ₂ -LA-Cys adduct	BME trans-nitroalkylation + C18-HPLC-ESI-MS and MS/MS in a Q trap 4000			[76]
	Generation after incubation between NO ₂ -LA and GSH	C18-HPLC/ESI/MS in Micromass Quattro II triple quadrupole			
	Red blood cells obtained from healthy humans	C18-HPLC-ESI-MS/MS in Q-Trap 4000			[42]
	MCF7/WT and MCF7/MRP1–10 cells treated with NO ₂ -LA	C18-HPLC/ESI/MS in Micromass Quattro II triple quadrupole			[76]
ANT1	NO ₂ -LA-treated intact perfused hearts	Immunoprecipitation + SDS-PAGE + in-gel digestion tryptic digestion + ABSciex 5800 MALDI-TOF MS and MS/MS	Mitochondrial uncoupling	Cytoprotective	[34]
Cys57					

(continued on next page)

Table 3 (continued)

Protein/peptide	Model system	Method	Molecular mechanism signaling action	Biological role	Ref.
Nitro-conjugated linoleic acid (NO ₂ -cLA) cysteine	NO ₂ -cLA-Cys adduct generation after incubation between NO ₂ -cLA and cysteine Urine from healthy humans	C18-HPLC-MS and MS/MS in LTQ Velos Orbitrap and AB 5000 or API4000 Q-trap triple quadrupole C18-HPLC-MS and MS/MS in LTQ Velos Orbitrap and AB 5000 or API 4000 Q-trap triple quadrupole C18-HPLC-MS and MS/MS in API 5000 triple quadrupole			[7] [7] [44]
	NO ₂ -cLA-Cys adduct generation after incubation between NO ₂ -cLA and cysteine Urine from healthy humans	C18-HPLC-MS and MS/MS in API 5000 triple quadrupole			[44]
Nitro-arachidonic acid (NO ₂ -AA) PDI	Incubation of human recombinat PDI with NO ₂ -AA	C4-HPLC-MS of intact protein in a hybrid triple quadrupole/linear ion trap mass spectrometer (Q-trap 4500) digestion tryptic digestion and C18-HPLC-MS and MS/MS	Inhibition of reductase and chaperone activity of PDI	Anti-inflammatory	[95]
	Cys397 Cys400				

catalytic activity of sHE was associated with anti-hypertensive properties of NO₂-FA [32]. Finally, neuroprotective effects associated with the decrease of protein aggregation were related with PTMs of α -synuclein by NO₂-OA [35].


3.2. Identification of protein-nitro-fatty acids adducts: tools and challenges

Identification of protein nitroalkylation by NO₂-FA has been disclosed by using different experimental approaches, as crystallographic analysis [100,101], western immunoblot-based assay [2,23,32,34,87], spectrophotometry [7,43,44,94] and MS-based approaches [4,7,34,42–44]. However, spectrophotometry and immunoassays do not give detailed structural information and crystallography requires pure proteins, being difficult to be used in the analysis of complex biological samples.

Mass spectrometry, namely using matrix assisted laser desorption/ionization (MALDI) or electrospray (ESI) MS-based proteomics approaches, often coupled to reverse phase (RP) liquid chromatography (LC-MS), are the most suitable methods for detection and characterization of adducts formed between NO₂-FA and proteins. *In vitro* generation of NO₂-FA-protein adducts, in biomimetic systems, between standards of NO₂-FA and candidate peptides or proteins has been used as strategy for the initial identification by (LC)-MS and further characterization of these adducts by MS/MS. Data obtained using these biomimetic approaches using controlled reaction conditions are more straightforward and relatively easy to analyze. This, in turn, allows to obtain knowledge on the reactivity of each individual NO₂-FA and the typical fragmentation pathways under MS/MS needed to identify these adducts. The information gathered by tandem mass experiments concerning the typical fragmentation pathways and reporter ions can be used to identify these lipoxidation products in complex biological samples by using MS-based proteomics approaches and to develop MS target analysis, namely multiple reaction monitoring (MRM) analysis. This has contributed to achieve the ultimate goal that consists of the identification of the NO₂-FA-protein adducts in complex biological samples as cells, tissues, biological fluids, which requires specific and targeted approaches. Bottom-up proteomics approaches are usually performed. Through these analytical approaches, it is possible to unequivocally identify the modified peptides after enzymatic digestion of NO₂-FA-protein adduct, usually using trypsin, followed by the analysis of the tryptic peptides by reverse-phase (RP)-LC-MS and MS/MS. The addition of the NO₂-FA moiety increases the retention time of the modified peptides [42], which are identified on the mass spectra as singly, $[M + H]^+$ ions, or multiple charged ions, $[M + nH]^{n+}$, based on the mass shift against the unmodified peptide. This gives information on the nature of NO₂-FA covalently attached to the protein. The observed mass shift in the mass spectra for the Michael adducts will be equal to the molecular weight of the NO₂-FA. Thus, a mass shift of + 327 Da and + 325 Da corresponds to the addition of NO₂-OA and NO₂-LA, respectively [42]. MS/MS data allows to confirm the nature of the modification and provides information on the fragmentation pattern of NO₂-FA-peptide adducts. These data further allows to pinpoint the location of the modification site and thus the targeted residue in the peptide backbone [103,104]. Detailed information to identify the sites of adduction is revealed by a mass shift of the typical *b* and *y* product ions of the adducted peptide, when compared with the non-modified one. The modified immonium ions are also useful to confirm the presence of a modified amino acid residue within the adducted peptide.

RP-LC-ESI-MS and MS/MS were used to detect lipoxidation adducts formed between NO₂-OA or NO₂-LA and GAPDH and GSH *in vivo* in healthy human red cells [42]. This methodology was also applied to confirm the post-translational modifications of matrix metalloproteinase by NO₂-OA [75], and for the identification of reversible Michael adducts of NO₂-OA and thiols of proteins and GSH in liver and plasma of NO₂-OA-treated mice [59]. Significant levels of protein cysteine adducts of NO₂-OA were also observed in fresh olives, especially in the

A) Mono charged $[M+H]^+$ ions



B) Double charged $[M+2H]^{2+}$ ions




Fig. 6. ESI-MS spectra of mono charged $[M+H]^+$ (A) and double charged $[M+2H]^{2+}$ ions (B) of $\text{NO}_2\text{POPC-GSH}$ adducts, acquired in Q-Exactive Orbitrap, with identification of major fragmentation pathways. (Reprinted with permission from [49], copyright 2018 [John Wiley & Sons]).

peel [56]. $\text{AT}_1\text{-R}$ adducts with $\text{NO}_2\text{-OA}$ were quantified by HPLC-MS/MS using MRM scan mode in the negative-ion mode as BME adducts (BME- $\text{NO}_2\text{-OA}$ adducts) after a nucleophilic exchange of $\text{NO}_2\text{-OA}$ from $\text{AT}_1\text{-R}$ to BME. The presence of exchangeable $\text{NO}_2\text{-OA}$ demonstrated the direct adduction of $\text{AT}_1\text{-R}$ by $\text{NO}_2\text{-OA}$, and therefore that $\text{AT}_1\text{-R}$ is a relevant cellular target for $\text{NO}_2\text{-OA}$ alkylation [28]. RP-LC-MRM scan in the positive-ion mode ($[M+H]^+$ ions) was applied for the characterization of $\text{NO}_2\text{-LA-GSH}$ adducts in vitro and further identification in MCF7 cells treated with $\text{NO}_2\text{-LA}$ [76]. Nitroalkylation of albumin by $\text{NO}_2\text{-OA}$ and $\text{NO}_2\text{-LA}$ have been found in the plasma of mice gavaged with these fatty acids [62]. Nitroalkylation of p65 subunit of nuclear factor κB (NF- κB) was observed in vivo in myocardial tissue of a murine model of ischemia-reperfusion with intravenous supplementation of OA and LA [1]. One study also reported the direct analysis by MALDI-TOF-TOF MS and MS/MS, in positive-ion mode, of adenine nucleotide translocase 1 (ATN 1) adducts after $\text{NO}_2\text{-LA}$ infusion into intact perfused hearts allowing to pinpoint that the nitroalkylation of ANT1 by $\text{NO}_2\text{-LA}$ occurred on Cys57 [34]. Adduction of $\text{NO}_2\text{-OA}$ to PPAR- γ [47],

and to Keap1 [46] are also examples of biological detection and characterization of $\text{NO}_2\text{-OA}$ -protein adducts by MS.

The Michael addition reactions between $\text{NO}_2\text{-FA}$ and proteins is remarkably selective and depends on the nature and structural features of the $\text{NO}_2\text{-FA}$. The fatty acyl chain length and the position of the electrophilic carbon, i.e., the position of the nitroalkene group, has a pivotal effect on the reactivity of $\text{NO}_2\text{-FA}$ [102]. Therefore both factors regulate the formation of $\text{NO}_2\text{-FA}$ -protein adducts and the biological activity of the $\text{NO}_2\text{-FA}$ [22,42,65,73,100,101]. In spite of its four possible isomers (at C9, C10, C12 or C13), only the $\text{NO}_2\text{-LA}$ isomers bearing the NO_2 at C10 and C12 were reported to selectively bind to cysteine 285 (Cys285) in the ligand-binding domain and activate PPAR γ [101]. The C10 isomer of $\text{NO}_2\text{-OA}$ is more reactive toward to Cys285 in the ligand binding domain of PPAR γ than the C9 isomer [47]. On the other hand, Keap1 is easily activated by the C9 isomer via nitroalkylation of Cys273 and Cys288 [22,46]. Xanthine oxidoreductase activity is preferentially inhibited by the C9 isomer of $\text{NO}_2\text{-OA}$ or a mixture of both C9 and C10 isomers [73]. It has been reported that $\text{NO}_2\text{-FA}$ with shorter

acyl chains interact stronger with Nrf2 and NF- κ B [60].

Overall, the identification of NO₂-FA-protein adducts is important, because it may give information, as shown in several examples reported earlier, on the potential protein targets whose modulation by NO₂-FA can have potential therapeutic interest.

4. Esterified nitro-fatty acids

4.1. Nitrated phospholipids and their lipoxidation adducts

In spite of their free forms, NO₂-FA can be stabilized by esterification in more complex lipids in hydrophobic compartments, as the biological membranes. Nitrated derivatives of phospholipids were identified in biomimetic model systems and also in vivo [39,40]. In mimetic model studies, nitrated PLs were generated after in vitro incubation of PL standards (phosphatidylcholines, PCs and phosphatidylethanolamines, PEs) and NO₂BF₄, and its characterization was performed using C5-LC-MS and MS/MS in a Linear ion trap [39,40]. Nitrated PCs and nitrated PEs were detected by HILIC-LC-MS and MS/MS-based lipidomic approaches in cardiac mitochondria from diabetic rats [39] and cardiomyoblasts subjected to starvation [40]. Nitrated 1-palmitoyl-2-oleyl-*sn*-glycero-3-phosphocholine (nitrated POPC) was reported to have antioxidant properties as scavenging agent, mediated by its anti-radical potential and ability to inhibit lipid peroxidation. Anti-inflammatory properties of nitrated POPC, related with its ability to inhibit iNOS expression in LPS-activated macrophages, were also reported [105].

NO₂-FA incorporation in PLs was also reported by using C18-HPLC-ESI-MS and MS/MS in API 4000 Q-trap triple quadrupole in adipocytes supplemented with NO₂-SA, NO₂-OA, NO₂-cLA, and NO₂-LA, before and after acidic hydrolysis. The incorporation yield and profile was specific for each supplemented NO₂-FA and PL class, being PC the PL class with highest levels of incorporation of NO₂-FA [106].

Nitrated POPC was also found to have the capability to form adducts with peptides. The identification of the covalent adducts of NO₂POPC with GSH was characterized by tandem MS in different instruments and the typical fragmentation pathways were disclosed for the first time. In this study, the NO₂POPC-GSH adducts were generated under biomimetic conditions and characterized by direct infusion MS and MS/MS using different instrumental platforms including LXQ linear ion trap, Q-TOF 2, and Q-Exactive Hybrid Orbitrap. The observed fragmentation pattern of NO₂POPC-GSH adducts included product ions that confirmed the presence of the phosphatidylcholine moiety (m/z 184.074 and neutral loss of 183 Da), the nitro group (neutral loss of HNO₂), and ^{*}y₂, ^{*}b₂ and ^{*}C₁ fragment ions of the modified peptide. All of these product ions pinpointing that NO₂POPC was linked to a cysteine residue of GSH (Fig. 6) and can be used as reporter ions applied in the search of these lipoxidation adducts in biological samples [49].

4.2. Nitrated triacylglycerides

Nitrated triacylglycerides (NO₂-FA-TAG) have been reported in rat plasma after oral administration of NO₂-OA, together with β -oxidation and dehydrogenation derivatives of NO₂-FA-TAG in adipocytes supplemented with NO₂-OA. These studies were performed by C18-HPLC-ESI-MS and MS/MS in API4000 Q-trap triple quadrupole and LTQ Velos Orbitrap instruments [41]. Another study reported the differential esterification profile of NO₂-FA and their metabolites in TAGs in adipose tissue of rats fed with 10-NO₂-OA. By using C18-HPLC-ESI-MS and MS/MS in API 4000 Q-trap triple quadrupole, the NO₂-FA were observed to be preferentially incorporated in monoacyl- and diacylglycerides. This was found to be in opposite to its reduced metabolites, which were favorably incorporated in TAGs. These observations were corroborated by the analysis of the lipid polar and neutral fractions from adipocytes supplemented with NO₂-SA (nitro-stearic acid), NO₂-OA, NO₂-cLA, and NO₂-LA, after acidic hydrolysis [106].

The occurrence of nitrated phospholipids and triacylglycerides can be of high relevance at biological level. The NO₂-FA-containing phospholipids and triacylglycerides can act as a reservoir of NO₂-FA. Additionally, these esterified NO₂-FA can be further mobilized by lipases in turn to exert their adaptive and anti-inflammatory signaling actions. In the case of NO₂-FA-containing phospholipids, the NO₂-FA moiety seems to be able to retain the electrophilic character, and thus the ability to undergo reversible reactions via Michael addition with key proteins. Also, these phospholipid-esterified NO₂-FA can have an impact as anti-inflammatory and cytoprotective species. The nitration of esterified NO₂-FA or its incorporation into more complex lipids, together with the occurrence of lipoxidation products of NO₂-FA-containing phospholipids, and perhaps NO₂-FA-TAGs, can also contribute to the systemic distribution and metabolism of NO₂-FA.

5. Conclusion and future perspectives

NO₂-FA own important physiological functions that are mediated via formation of lipoxidation adducts and associated regulation of protein function. Several signaling proteins, with key roles in anti-inflammatory, anti-hypertensive, anti-hyperglycemic, and cytoprotective pathways, are targets of NO₂-FA adduction. This points to potential for new therapeutic strategies in important non-communicable diseases as cardiovascular, renal, pulmonary, and metabolic diseases. Mass spectrometry is a promising analytical tool in the detection of NO₂-FA-protein adducts. Nevertheless, there is a need for new methodological developments to improve the detection of these elusive lipoxidation adducts, and to obtain more insights regarding the protein targets of NO₂-FA and its roles in biological signaling pathways.

Acknowledgments

We acknowledge funding from the European Commission's Horizon 2020 research and innovation programme for the Marie Skłodowska-Curie grant agreement number 675132 (MSCA-ITN-ETN MASSTRPLAN) to University of Aveiro, Portugal. Thanks are due to the University of Aveiro, FCT/MEC, European Union, QREN, COMPETE for the financial support to the QOPNA (FCT UID/QUI/00062/2013) and CESAM (UID/AMB/50017 - POCI-01-0145-FEDER-007638), through national funds and where applicable co-financed by the FEDER, within the PT2020 Partnership Agreement, to the Portuguese Mass Spectrometry Network, RNEM, Portugal (LISBOA-01-0145-FEDER-402-022125). Tânia Melo (BPD/UI51/5388/2017) is grateful to FCT for her grant.

References

- [1] V. Rudolph, T.K. Rudolph, F.J. Schopfer, G. Bonacci, S.R. Woodcock, M.P. Cole, P.R.S. Baker, R. Ramani, B.A. Freeman, Endogenous generation and protective effects of nitro-fatty acids in a murine model of focal cardiac ischaemia and reperfusion, *Cardiovasc. Res.* 85 (2010) 155–166, <https://doi.org/10.1093/cvr/cvp275>.
- [2] S.M. Nadtochiy, P.R.S. Baker, B.A. Freeman, P.S. Brookes, Mitochondrial nitroalkene formation and mild uncoupling in ischaemic preconditioning: implications for cardioprotection, *Cardiovasc. Res.* 82 (2009) 333–340, <https://doi.org/10.1093/cvr/cvn323>.
- [3] G. Bonacci, P.R.S. Baker, S.R. Salvatore, D. Shores, N.K.H. Khoo, J.R. Koenitzer, D.A. Vitturi, S.R. Woodcock, F. Golin-Bisello, M.P. Cole, S. Watkins, C. Croix St, C.I. Batthyany, B.A. Freeman, F.J. Schopfer, Conjugated linoleic acid is a preferential substrate for fatty acid nitration, *J. Biol. Chem.* 287 (2012) 44071–44082, <https://doi.org/10.1074/jbc.M112.401356>.
- [4] J.C. Schoeman, A.C. Harms, M. van Weeghel, R. Berger, R.J. Vreeken, T. Hankemeier, Development and application of a UHPLC-MS/MS metabolomics based comprehensive systemic and tissue-specific screening method for inflammatory, oxidative and nitrosative stress, *Anal. Bioanal. Chem.* 410 (2018) 2551–2568, <https://doi.org/10.1007/s00216-018-0912-2>.
- [5] A.T. Reddy, S.P. Lakshmi, R.R. Muchumarri, R.C. Reddy, Nitrated fatty acids reverse cigarette smoke-induced alveolar macrophage activation and inhibit protease activity via electrophilic S-alkylation, *PLoS One* 11 (2016) e0153336, <https://doi.org/10.1371/journal.pone.0153336>.
- [6] P.R.S. Baker, Y. Lin, F.J. Schopfer, S.R. Woodcock, A.L. Groeger, C. Batthyany,

- S. Sweeney, M.H. Long, K.E. Iles, L.M.S. Baker, B.P. Branchaud, Y.E. Chen, B.A. Freeman, Fatty acid transduction of nitric oxide signaling: multiple nitrated unsaturated fatty acid derivatives exist in human blood and urine and serve as endogenous peroxisome proliferator-activated receptor ligands, *J. Biol. Chem.* 280 (2005) 42464–42675, <https://doi.org/10.1074/jbc.M504212200>.
- [7] S.R. Salvatore, D.A. Vitturi, P.R.S. Baker, G. Bonacci, J.R. Koenitzer, S.R. Woodcock, B.A. Freeman, F.J. Schopfer, Characterization and quantification of endogenous fatty acid nitroalkene metabolites in human urine, *J. Lipid Res.* 54 (2013) 1998–2009, <https://doi.org/10.1194/jlr.M037804>.
- [8] P.R. Baker, F.J. Schopfer, S. Sweeney, B.A. Freeman, Red cell membrane and plasma linoleic acid nitration products: synthesis, clinical identification, and quantitation, *Proc. Natl. Acad. Sci. USA* 101 (2004) 11577–11582, <https://doi.org/10.1073/pnas.0402587101>.
- [9] E.S. Lima, P. Di Mascio, D.S.P. Abdalla, Cholesteryl nitrooleate, a nitrated lipid present in human blood plasma and lipoproteins, *J. Lipid Res.* 44 (2003) 1660–1666, <https://doi.org/10.1194/jlr.M200467-JLR200>.
- [10] É.S. Lima, P. Di Mascio, H. Rubbo, D.S.P. Abdalla, Characterization of linoleic acid nitration in human blood plasma by mass spectrometry, *Biochemistry* 41 (2002) 10717–10722, <https://doi.org/10.1021/bi025504j>.
- [11] D. Tsikas, A.A. Zoerner, J. Jordan, Oxidized and nitrated oleic acid in biological systems: analysis by GC-MS/MS and LC-MS/MS, and biological significance, *Biochim. Biophys. Acta - Mol. Cell Biol. Lipids* (2011) 694–705, <https://doi.org/10.1016/j.bbalip.2011.06.015>.
- [12] D. Tsikas, A.A. Zoerner, A. Mitschke, F.M. Gutzki, Nitro-fatty acids occur in human plasma in the picomolar range: a targeted nitro-lipidomics GC-MS/MS study, *Lipids* 44 (2009) 855–865, <https://doi.org/10.1007/s11745-009-3332-4>.
- [13] L. Villacorta, L. Minarrieta, S.R. Salvatore, N.K. Khoo, O. Rom, Z. Gao, R.C. Berman, S. Jobbagy, L. Li, S.R. Woodcock, Y.E. Chen, B.A. Freeman, A.M. Ferreira, F.J. Schopfer, D.A. Vitturi, In situ generation, metabolism and immunomodulatory signaling actions of nitro-conjugated linoleic acid in a murine model of inflammation, *Redox Biol.* 15 (2018) 522–531, <https://doi.org/10.1016/j.redox.2018.01.005>.
- [14] T. Yang, H. Wang, H. Liu, Z. Jia, G. Guan, Effects of endogenous PPAR agonist nitro-oleic acid on metabolic syndrome in obese Zucker rats, *PPAR Res.* (2010), <https://doi.org/10.1155/2010/601562>.
- [15] A.T. Reddy, S.P. Lakshmi, R.C. Reddy, The nitrated fatty acid 10-nitro-oleate diminishes severity of lps-induced acute lung injury in mice, *PPAR Res.* (2012), <https://doi.org/10.1155/2012/617063>.
- [16] A.L. Hansen, G.J. Buchan, M. Rühl, K. Mukai, S.R. Salvatore, E. Ogawa, S.D. Andersen, M.B. Iversen, A.L. Thielke, C. Gunderstofte, M. Motwani, C.T. Møller, A.S. Jakobsen, K.A. Fitzgerald, J. Roos, R. Lin, T.J. Maier, R. Goldbach-Mansky, C.A. Miner, W. Qian, J.J. Miner, R.E. Rigby, J. Rehwinkel, M.R. Jakobsen, H. Arai, T. Taguchi, F.J. Schopfer, D. Olgner, C.K. Holm, Nitro-fatty acids are formed in response to virus infection and are potent inhibitors of STING palmitoylation and signaling, *Proc. Natl. Acad. Sci. USA* 115 (2018) E7768–E7775, <https://doi.org/10.1073/pnas.1806239115>.
- [17] K.E. Iles, M.M. Wright, M.P. Cole, N.E. Welty, L.B. Ware, M.A. Matthey, F.J. Schopfer, P.R.S. Baker, A. Agarwal, B.A. Freeman, Fatty acid transduction of nitric oxide signaling: nitrooleic acid mediates protective effects through regulation of the ERK pathway, *Free Radic. Biol. Med.* 46 (2009) 866–875, <https://doi.org/10.1016/j.freeradbiomed.2008.12.001>.
- [18] B. Coles, A. Bloodsworth, S.R. Clark, M.J. Lewis, A.R. Cross, B.A. Freeman, V.B. O'Donnell, Nitrooleate inhibits superoxide generation, degranulation, and integrin expression by human neutrophils: novel antiinflammatory properties of nitric oxide-derived reactive species in vascular cells, *Circ. Res.* 91 (2002) 375–381, <https://doi.org/10.1161/01.RES.0000032114.68919.EF>.
- [19] M.P. Cole, T.K. Rudolph, N.K.H. Khoo, U.N. Motanya, F. Golin-Bisello, J.W. Wertz, F.J. Schopfer, V. Rudolph, S.R. Woodcock, S. Bolisetty, M.S. Ali, J. Zhang, Y.E. Chen, A. Agarwal, B.A. Freeman, P.M. Bauer, Nitro-fatty acid inhibition of neointima formation after endoluminal vessel injury, *Circ. Res.* 105 (2009) 965–972, <https://doi.org/10.1161/CIRCRESAHA.109.199075>.
- [20] A.R. Mathers, C.D. Carey, M.E. Killen, J.A. Diaz-Perez, S.R. Salvatore, F.J. Schopfer, B.A. Freeman, L.D. Faló, Electrophilic nitro-fatty acids suppress allergic contact dermatitis in mice, *Allergy Eur. J. Allergy Clin. Immunol.* 72 (2017) 656–664, <https://doi.org/10.1111/all.13067>.
- [21] J. Hwang, K.E. Lee, J.Y. Lim, S.I. Park, Nitrated fatty acids prevent TNF α -stimulated inflammatory and atherogenic responses in endothelial cells, *Biochem. Biophys. Res. Commun.* 387 (2009) 633–640, <https://doi.org/10.1016/j.bbrc.2009.07.030>.
- [22] E. Kansanen, H.K. Jyrkkänen, O.L. Volger, H. Leinonen, A.M. Kivelä, S.K. Häkkinen, S.R. Woodcock, F.J. Schopfer, A.J. Horrovoets, S. Ylä-Herttua, B.A. Freeman, A.L. Levenon, Nrf2-dependent and -independent responses to nitro-fatty acids in human endothelial cells: identification of heat shock response as the major pathway activated by nitro-oleic acid, *J. Biol. Chem.* 284 (2009) 33233–33241, <https://doi.org/10.1074/jbc.M109.064873>.
- [23] T. Cui, F.J. Schopfer, J. Zhang, K. Chen, T. Ichikawa, P.R.S. Baker, C. Batthyany, B.K. Chacko, X. Feng, R.P. Patel, A. Agarwal, B.A. Freeman, Y.E. Chen, Nitrated fatty acids: endogenous anti-inflammatory signaling mediators, *J. Biol. Chem.* 281 (2006) 35686–35698, <https://doi.org/10.1074/jbc.M603357200>.
- [24] H. Liu, Z. Jia, S. Soodvilai, G. Guan, M.-H. Wang, Z. Dong, J.D. Symons, T. Yang, Nitro-oleic acid protects the mouse kidney from ischemia and reperfusion injury, *Am. J. Physiol. Physiol.* 295 (2008) F942–F949, <https://doi.org/10.1152/ajprenal.90236.2008>.
- [25] E. Kansanen, S.M. Kuosmanen, A.-K. Ruotsalainen, H. Hynynen, A.-L. Levenon, Nitro-oleic acid regulates endothelin signaling in human endothelial cells, *Mol. Pharmacol.* 92 (2017) 481–490, <https://doi.org/10.1124/mol.117.109751>.
- [26] M. Balazy, T. Iesaki, J.L.L. Park, H. Jiang, P.M.M. Kaminski, M.S.S. Wolin, Vicinal nitrohydroxyeicosatrienoic acids: vasodilator lipids formed by reaction of nitrogen dioxide with arachidonic acid, *J. Pharmacol. Exp. Ther.* 299 (2001) 611–619.
- [27] M. Rudnicki, L.A. Faine, N. Dehne, D. Namgaladze, S. Ferderbar, R. Weinlich, G.P. Amarante-Mendes, C.Y.I. Yan, J.E. Krieger, B. Brüne, D.S.P. Abdalla, Hypoxia inducible factor-dependent regulation of angiogenesis by nitro-fatty acids, *Arterioscler. Thromb. Vasc. Biol.* 31 (2011) 1360–1367, <https://doi.org/10.1161/ATVBAHA.111.224626>.
- [28] J. Zhang, L. Villacorta, L. Chang, Z. Fan, M. Hamblin, T. Zhu, C.S. Chen, M.P. Cole, F.J. Schopfer, C.X. Deng, M.T. Garcia-Barrio, Y.H. Feng, B.A. Freeman, Y.E. Chen, Nitro-oleic acid inhibits angiotensin II-induced hypertension, *Circ. Res.* 107 (2010) 540–548, <https://doi.org/10.1161/CIRCRESAHA.110.218404>.
- [29] É.S. Lima, M.G. Bonini, O. Augusto, H.V. Barbeiro, H.P. Souza, D.S.P. Abdalla, Nitrated lipids decompose to nitric oxide and lipid radicals and cause vasorelaxation, *Free Radic. Biol. Med.* 39 (2005) 532–539, <https://doi.org/10.1016/j.freeradbiomed.2005.04.005>.
- [30] D.G. Lim, S. Sweeney, A. Bloodsworth, C.R. White, P.H. Chumley, N.R. Krishna, F. Schopfer, V.B. O'Donnell, J.P. Eiserich, B.A. Freeman, Nitrooleate, a nitric oxide-derived mediator of cell function: synthesis, characterization, and vasomotor activity, *Proc. Natl. Acad. Sci. USA* 99 (2002) 15941–15946, <https://doi.org/10.1073/pnas.232409599>.
- [31] F. Blanco, A.M. Ferreira, G.V. López, L. Bonilla, M. González, H. Cerecetto, A. Trostchansky, H. Rubbo, 6-Methylnitroarachidonate: a novel esterified nitroalkene that potently inhibits platelet aggregation and exerts cGMP-mediated vascular relaxation, *Free Radic. Biol. Med.* 50 (2011) 411–418, <https://doi.org/10.1016/j.freeradbiomed.2010.11.031>.
- [32] R.L. Charles, O. Rudyk, O. Prisyazhna, A. Kamynina, J. Yang, C. Morisseau, B.D. Hammock, B.A. Freeman, P. Eaton, Protection from hypertension in mice by the Mediterranean diet is mediated by nitro fatty acid inhibition of soluble epoxide hydrolase, *Proc. Natl. Acad. Sci. USA* 111 (2014) 8167–8172, <https://doi.org/10.1073/pnas.1402965111>.
- [33] B. Coles, A. Bloodsworth, J.P. Eiserich, M.J. Coffey, R.M. McLoughlin, J.C. Giddings, M.J. Lewis, R.J. Haslam, B.A. Freeman, V.B. O'Donnell, Nitrooleate inhibits platelet activation by attenuating calcium mobilization and inducing phosphorylation of vasodilator-stimulated phosphoprotein through elevation of cAMP, *J. Biol. Chem.* 277 (2002) 5832–5840, <https://doi.org/10.1074/jbc.M105209200>.
- [34] S.M. Nadtochiy, Q. Zhu, W. Urciuoli, R. Rafikov, S.M. Black, P.S. Brookes, Nitroalkenes confer acute cardioprotection via adenine nucleotide translocase, *J. Biol. Chem.* 287 (2012) 3573–3580, <https://doi.org/10.1074/jbc.M111.298406>.
- [35] J.M. Souza, A. Trostchansky, C. Batthyany, R. Durán, B.A. Freeman, H. Rubbo, Posttranslational modification of human alpha-synuclein by nitro-oleic acid, *Free Radic. Biol. Med.* 49 (2010) S158, <https://doi.org/10.1016/j.freeradbiomed.2010.10.445>.
- [36] B. Sánchez-Calvo, A. Cassina, N. Rios, G. Peluffo, J. Boggia, R. Radi, H. Rubbo, A. Trostchansky, Nitro-arachidonic acid prevents angiotensin II-induced mitochondrial dysfunction in a cell line of kidney proximal tubular cells, *PLoS One* 11 (2016) e0150459, <https://doi.org/10.1371/journal.pone.0150459>.
- [37] P.M. Van Hoose, N.Q. Kelm, K.M. Piell, M.P. Cole, Conjugated linoleic acid and nitrite attenuate mitochondrial dysfunction during myocardial ischemia, *J. Nutr. Biochem.* 34 (2016) 8–16, <https://doi.org/10.1016/j.jnutbio.2016.04.001>.
- [38] I. Milic, E. Griesser, V. Vemula, N. Ieda, H. Nakagawa, N. Miyata, J.M. Galano, C. Oger, T. Durand, M. Fedorova, Profiling and relative quantification of multiply nitrated and oxidized fatty acids, *Anal. Bioanal. Chem.* 407 (2015) 5587–5602, <https://doi.org/10.1007/s00216-015-8766-3>.
- [39] T. Melo, P. Domingues, R. Ferreira, I. Milic, M. Fedorova, S.M. Santos, M.A. Segundo, M.R.M. Domingues, Recent advances on mass spectrometry analysis of nitrated phospholipids, *Anal. Chem.* 88 (2016) 2622–2629, <https://doi.org/10.1021/acs.analchem.5b03407>.
- [40] T. Melo, P. Domingues, T.M. Ribeiro-Rodrigues, H. Girão, M.A. Segundo, M.R.M. Domingues, Characterization of phospholipid nitroxidation by LC-MS in biomimetic models and in H9c2 Myoblast using a lipidomic approach, *Free Radic. Biol. Med.* 106 (2017) 219–227, <https://doi.org/10.1016/j.freeradbiomed.2017.02.033>.
- [41] M. Fazzari, N. Khoo, S.R. Woodcock, L. Li, B.A. Freeman, F.J. Schopfer, Generation and esterification of electrophilic fatty acid nitroalkenes in triacylglycerides, *Free Radic. Biol. Med.* 87 (2015) 113–124, <https://doi.org/10.1016/j.freeradbiomed.2015.05.033>.
- [42] C. Batthyany, F.J. Schopfer, P.R.S. Baker, R. Durán, L.M.S. Baker, Y. Huang, C. Cerveñansky, B.P. Branchaud, B.A. Freeman, Reversible post-translational modification of proteins by nitrated fatty acids in vivo, *J. Biol. Chem.* 281 (2006) 20450–20563, <https://doi.org/10.1074/jbc.M602814200>.
- [43] M.N. Padilla, C. Mata-Pérez, M. Melguizo, J.B. Barroso, In vitro nitro-fatty acid release from Cys-NO2-fatty acid adducts under nitro-oxidative conditions, *Nitric Oxide - Biol. Chem.* 68 (2017) 14–22, <https://doi.org/10.1016/j.niox.2016.12.009>.
- [44] L. Turell, D.A. Vitturi, E.L. Coitino, L. Lebrato, M.N. Möller, C. Sagasti, S.R. Salvatore, S.R. Woodcock, B. Alvarez, F.J. Schopfer, The chemical basis of thiol addition to nitro-conjugated linoleic acid, a protective cell-signaling lipid, *J. Biol. Chem.* 292 (2017) 1145–1159, <https://doi.org/10.1074/jbc.M116.756288>.
- [45] K. Awwad, S.D. Steinbrink, T. Frömel, N. Lill, J. Isaak, A.-K. Häfner, J. Roos, B. Hofmann, H. Heide, G. Geisslinger, D. Steinhilber, B.A. Freeman, T.J. Maier, I. Fleming, Electrophilic fatty acid species inhibit 5-lipoxygenase and attenuate sepsis-induced pulmonary inflammation, *Antioxid. Redox Signal.* 20 (2014) 2667–2680, <https://doi.org/10.1089/ars.2013.5473>.
- [46] E. Kansanen, G. Bonacci, F.J. Schopfer, S.M. Kuosmanen, K.I. Tong, H. Leinonen,

- S.R. Woodcock, M. Yamamoto, C. Carlberg, S. Ylä-Herttuala, B.A. Freeman, A.L. Levonen, Electrophilic nitro-fatty acids activate Nrf2 by a Keap1 cysteine 151-independent mechanism, *J. Biol. Chem.* 286 (2011) 14019–14027, <https://doi.org/10.1074/jbc.M110.190710>.
- [47] F.J. Schopfer, M.P. Cole, A.L. Groeger, C.S. Chen, N.K.H. Khoo, S.R. Woodcock, F. Golin-Bisello, U. Nkiru Motanya, Y. Li, J. Zhang, M.T. Garcia-Barrio, T.K. Rudolph, V. Rudolph, G. Bonacci, P.R.S. Baker, H.E. Xu, C.I. Batthyany, Y.E. Chen, T.M. Hallis, B.A. Freeman, Covalent peroxisome proliferator-activated receptor γ adduction by nitro-fatty acids: selective ligand activity and anti-diabetic signaling actions, *J. Biol. Chem.* 285 (2010) 12321–12333, <https://doi.org/10.1074/jbc.M109.091512>.
- [48] J.R. Koenitzer, G. Bonacci, S.R. Woodcock, C.S. Chen, N. Cantu-Medellin, E.E. Kelley, F.J. Schopfer, Fatty acid nitroalkenes induce resistance to ischemic cardiac injury by modulating mitochondrial respiration at complex II, *Redox Biol.* 8 (2016) 1–10, <https://doi.org/10.1016/j.redox.2015.11.002>.
- [49] J.-F. Montero-Bullon, T. Melo, M.R. Domingues, P. Domingues, Characterization of nitrophospholipid-peptide covalent adducts by electrospray tandem mass spectrometry: a first screening analysis using different instrumental platforms, *Eur. J. Lipid Sci. Technol.* 0 (2018) 1800101, <https://doi.org/10.1002/ejlt.201800101>.
- [50] B.A. Freeman, P.R.S. Baker, F.J. Schopfer, S.R. Woodcock, A. Napolitano, M. D'Ischia, Nitro-fatty acid formation and signaling, *J. Biol. Chem.* (2008) 15515–15519, <https://doi.org/10.1074/jbc.R800004200>.
- [51] A. Trostchansky, J.M. Souza, A. Ferreira, M. Ferrari, F. Blanco, M. Trujillo, D. Castro, H. Cerecetto, P.R.S. Baker, V.B. O'Donnell, H. Rubbo, Synthesis, isomer characterization, and anti-inflammatory properties of nitroarachidonate, *Biochemistry* (2007), <https://doi.org/10.1021/bi602652j>.
- [52] A. Napolitano, L. Panzella, M. Savarese, R. Sacchi, I. Giudicianni, L. Paolillo, M. D'Ischia, Acid-induced structural modifications of unsaturated fatty acids and phenolic olive oil constituents by nitrite ions: a chemical assessment, *Chem. Res. Toxicol.* 17 (2004) 1329–1337, <https://doi.org/10.1021/tx049880b>.
- [53] C. Pereira, N.R. Ferreira, B.S. Rocha, R.M. Barbosa, J. Laranjinha, The redox interplay between nitrite and nitric oxide: from the gut to the brain, *Redox Biol.* 1 (2013) 276–284, <https://doi.org/10.1016/j.redox.2013.04.004>.
- [54] A. Napolitano, E. Camera, M. Picardo, M. D'Ischia, Acid-promoted reactions of ethyl linoleate with nitrite ions: formation and structural characterization of isomeric nitroalkene, nitrohydroxy, and novel 3-nitro-1,5-hexadiene and 1,5-dinitro-1,3-pentadiene products, *J. Org. Chem.* 65 (2000) 4853–4860, <https://doi.org/10.1021/jo000090q>.
- [55] E. Weitzberg, J.O. Lundberg, Novel aspects of dietary nitrate and human health, *Annu. Rev. Nutr.* 33 (2013) 129–159, <https://doi.org/10.1146/annurev-nutr-071812-161159>.
- [56] M. Fazzari, A. Trostchansky, F.J. Schopfer, S.R. Salvatore, B. Sánchez-Calvo, D. Vitturi, R. Valderrama, J.B. Barroso, R. Radi, B.A. Freeman, H. Rubbo, Olives and olive oil are sources of electrophilic fatty acid nitroalkenes, *PLoS One* 9 (2014) e84884, <https://doi.org/10.1371/journal.pone.0084884>.
- [57] C. Mata-Pérez, B. Sánchez-Calvo, M.N. Padilla, J.C. Begara-Morales, R. Valderrama, F.J. Corpas, J.B. Barroso, Nitro-fatty acids in plant signaling: new key mediators of nitric oxide metabolism, *Redox Biol.* (2017), <https://doi.org/10.1016/j.redox.2017.01.002>.
- [58] M. Delmastro-Greenwood, K.S. Hughan, D.A. Vitturi, S.R. Salvatore, G. Grimes, G. Potti, S. Shiva, F.J. Schopfer, M.T. Gladwin, B.A. Freeman, S. Gelhaus Wendell, Nitrite and nitrate-dependent generation of anti-inflammatory fatty acid nitroalkenes, *Free Radic. Biol. Med.* 89 (2015) 333–341, <https://doi.org/10.1016/j.freeradbiomed.2015.07.149>.
- [59] V. Rudolph, F.J. Schopfer, N.K.H. Khoo, T.K. Rudolph, M.P. Cole, S.R. Woodcock, G. Bonacci, A.L. Groeger, F. Golin-Bisello, C.S. Chen, P.R.S. Baker, B.A. Freeman, Nitro-fatty acid metabolome: saturation, desaturation, β -oxidation, and protein adduction, *J. Biol. Chem.* 284 (2009) 1461–1473, <https://doi.org/10.1074/jbc.M802298200>.
- [60] N.K.H. Khoo, L. Li, S.R. Salvatore, F.J. Schopfer, B.A. Freeman, Electrophilic fatty acid nitroalkenes regulate Nrf2 and NF- κ B signaling: a medicinal chemistry investigation of structure-function relationships, *Sci. Rep.* 8 (2018) 2295, <https://doi.org/10.1038/s41598-018-20460-8>.
- [61] S.R. Salvatore, D.A. Vitturi, M. Fazzari, D.K. Jorkasky, F.J. Schopfer, Evaluation of 10-nitro oleic acid bio-elimination in rats and humans, *Sci. Rep.* 7 (2017) 39900, <https://doi.org/10.1038/srep39900>.
- [62] F.J. Schopfer, C. Batthyany, P.R.S. Baker, G. Bonacci, M.P. Cole, V. Rudolph, A.L. Groeger, T.K. Rudolph, S. Nadtchivy, P.S. Brookes, B.A. Freeman, Detection and quantification of protein adduction by electrophilic fatty acids: mitochondrial generation of fatty acid nitroalkene derivatives, *Free Radic. Biol. Med.* 46 (2009) 1250–1259, <https://doi.org/10.1016/j.freeradbiomed.2008.12.025>.
- [63] G. Bonacci, E.K. Asciutto, S.R. Woodcock, S.R. Salvatore, B.A. Freeman, F.J. Schopfer, Gas-phase fragmentation analysis of nitro-fatty acids, *J. Am. Soc. Mass Spectrom.* 22 (2011) 1534–1551, <https://doi.org/10.1007/s13361-011-0185-x>.
- [64] S.V.S. Chakravartula, M. Balazy, Characterization of nitro arachidonic acid and nitro linoleic acid by mass spectrometry, *Anal. Lett.* 45 (2012), <https://doi.org/10.1080/00032719.2012.693558> (2412–242).
- [65] S. Borniquel, E.A. Jansson, M.P. Cole, B.A. Freeman, J.O. Lundberg, Nitrated oleic acid up-regulates PPAR γ and attenuates experimental inflammatory bowel disease, *Free Radic. Biol. Med.* 48 (2010) 499–505, <https://doi.org/10.1016/j.freeradbiomed.2009.11.014>.
- [66] A.M. Ferreira, M.I. Ferrari, A. Trostchansky, C. Batthyany, J.M. Souza, M.N. Alvarez, G.V. López, P.R.S. Baker, F.J. Schopfer, V. O'Donnell, B.A. Freeman, H. Rubbo, Macrophage activation induces formation of the anti-inflammatory lipid cholesteryl-nitrolinoleate, *Biochem. J.* 417 (2009) 223–234, <https://doi.org/10.1042/BJ20080701>.
- [67] N.K.H. Khoo, V. Rudolph, M.P. Cole, F. Golin-Bisello, F.J. Schopfer, S.R. Woodcock, C. Batthyany, B.A. Freeman, Activation of vascular endothelial nitric oxide synthase and heme oxygenase-1 expression by electrophilic nitro-fatty acids, *Free Radic. Biol. Med.* 48 (2010) 230–239, <https://doi.org/10.1016/j.freeradbiomed.2009.10.046>.
- [68] F.J. Schopfer, D.A. Vitturi, D.K. Jorkasky, B.A. Freeman, Nitro-fatty acids: new drug candidates for chronic inflammatory and fibrotic diseases, *Nitric Oxide* 79 (2018) 31–37, <https://doi.org/10.1016/j.niox.2018.06.006>.
- [69] W. Wang, C. Li, T. Yang, Protection of nitro-fatty acid against kidney diseases, *Am. J. Physiol. Ren. Physiol.* 310 (2016) F697–F704, <https://doi.org/10.1152/ajprenal.00321.2015>.
- [70] F.J. Schopfer, P.R.S. Baker, G. Giles, P. Chumley, C. Batthyany, J. Crawford, R.P. Patel, N. Hogg, B.P. Branchaud, J.R. Lancaster, B.A. Freeman, Fatty acid transduction of nitric oxide signaling: nitrolinoleic acid is a hydrophobically stabilized nitric oxide donor, *J. Biol. Chem.* 280 (2005) 19289–19297, <https://doi.org/10.1074/jbc.M414689200>.
- [71] H. Rubbo, S. Parthasarathy, S. Barnes, M. Kirk, B. Kalyanaraman, B.A. Freeman, Nitric oxide inhibition of lipoygenase-dependent liposome and low-density lipoprotein oxidation: termination of radical chain propagation reactions and formation of nitrogen-containing oxidized lipid derivatives, *Arch. Biochem. Biophys.* 324 (1995) 15–25, <https://doi.org/10.1006/abbi.1995.9935>.
- [72] M.J. Gorczynski, J. Huang, H. Lee, S.B. King, Evaluation of nitroalkenes as nitric oxide donors, *Bioorg. Med. Chem. Lett.* 17 (2007) 2013–2017, <https://doi.org/10.1016/j.bmcl.2007.01.016>.
- [73] E.E. Kelley, C.I. Batthyany, N.J. Hundley, S.R. Woodcock, G. Bonacci, J.M. Del Rio, F.J. Schopfer, J.R. Lancaster Jr, B.A. Freeman, M.M. Tarpey, Nitro-oleic acid, a novel and irreversible inhibitor of xanthine oxidoreductase, *J. Biol. Chem.* 283 (2008) 36176–36184, <https://doi.org/10.1074/jbc.M802402200>.
- [74] L. González-Perilli, M.N. Álvarez, C. Prolo, R. Radi, H. Rubbo, A. Trostchansky, Nitroarachidonic acid prevents NADPH oxidase assembly and superoxide radical production in activated macrophages, *Free Radic. Biol. Med.* 58 (2013) 126–133, <https://doi.org/10.1016/j.freeradbiomed.2012.12.020>.
- [75] G. Bonacci, F.J. Schopfer, C.I. Batthyany, T.K. Rudolph, V. Rudolph, N.K.H. Khoo, E.E. Kelley, B.A. Freeman, Electrophilic fatty acids regulate matrix metalloproteinase activity and expression, *J. Biol. Chem.* 286 (2011) 16074–16081, <https://doi.org/10.1074/jbc.M111.225029>.
- [76] R.L. Alexander, D.J.P. Bates, M.W. Wright, S.B. King, C.S. Morrow, Modulation of nitrated lipid signaling by multidrug resistance protein 1 (MRP1): glutathione conjugation and MRP1-mediated efflux inhibit nitrolinoleic acid-induced, PPAR γ -dependent transcription activation, *Biochemistry* 45 (2006) 7889–7896, <https://doi.org/10.1021/bi0605639>.
- [77] T.K. Rudolph, B.A. Freeman, Transduction of redox signaling by electrophile-protein reactions, *Sci. Signal.* 2 (2009), <https://doi.org/10.1126/scisignal.290re7> (re7–re7).
- [78] A. Trostchansky, H. Rubbo, Nitrated fatty acids: mechanisms of formation, chemical characterization, and biological properties, *Free Radic. Biol. Med.* 44 (2008) 1887–1896, <https://doi.org/10.1016/j.freeradbiomed.2008.03.006>.
- [79] D. Lin, S. Saleh, D.C. Liebler, Reversibility of covalent electrophile-protein adducts and chemical toxicity, *Chem. Res. Toxicol.* 21 (2008) 2361–2369, <https://www.ncbi.nlm.nih.gov/pubmed/19548357>.
- [80] K. Jain, A. Siddam, A. Marathi, U. Roy, J.R. Falck, M. Balazy, The mechanism of oleic acid nitration by \cdot NO $_2$, *Free Radic. Biol. Med.* 45 (2008) 269–283, <https://doi.org/10.1016/j.freeradbiomed.2008.04.015>.
- [81] D.A. Vitturi, C.-S. Chen, S.R. Woodcock, S.R. Salvatore, G. Bonacci, J.R. Koenitzer, N.A. Stewart, N. Wakabayashi, T.W. Kensler, B.A. Freeman, F.J. Schopfer, Modulation of nitro-fatty acid signaling: prostaglandin reductase-1 is a nitroalkene reductase, *J. Biol. Chem.* 288 (2013) 25626–25637, <https://doi.org/10.1074/jbc.M113.486282>.
- [82] B. Alvarez, L. Turell, D.A. Vitturi, E.L. Coitiño, L. Lebrato, M.N. Moller, C. Sagasti, S.R. Salvatore, S.R. Woodcock, F.J. Schopfer, Thiol addition to conjugated nitro linoleic acid, *FASEB J.* 31 (2017), https://doi.org/10.1096/fasebj.31.1_supplement.605.1 (605.1–605.1).
- [83] H. Rubbo, Nitro-fatty acids: novel anti-inflammatory lipid mediators, *Braz. J. Med. Biol. Res.* 46 (2013) 728–734, <https://doi.org/10.1590/1414-431X20133202>.
- [84] A.C. Geisler, T.K. Rudolph, Nitroalkylation — a redox sensitive signaling pathway, *Biochim. Biophys. Acta - Gen. Subj.* 1820 (2012) 777–784, <https://doi.org/10.1016/j.bbagen.2011.06.014>.
- [85] G. Aldini, M.R. Domingues, C.M. Spickett, P. Domingues, A. Altomare, F.J. Sánchez-Gómez, C.L. Oeste, D. Pérez-Sala, Protein lipoxidation: detection strategies and challenges, *Redox Biol.* 5 (2015) 253–266, <https://doi.org/10.1016/j.redox.2015.05.003>.
- [86] G. Ambrozova, H. Martiskova, A. Koudelka, T. Ravekes, T.K. Rudolph, A. Klinke, V. Rudolph, B.A. Freeman, S.R. Woodcock, L. Kubala, M. Pekarova, Nitro-oleic acid modulates classical and regulatory activation of macrophages and their involvement in pro-fibrotic responses, *Free Radic. Biol. Med.* 90 (2016) 252–260, <https://doi.org/10.1016/j.freeradbiomed.2015.11.026>.
- [87] T.K. Rudolph, V. Rudolph, M.M. Edreira, M.P. Cole, G. Bonacci, F.J. Schopfer, S.R. Woodcock, A. Franek, M. Pekarova, N.K.H. Khoo, A.H. Hasty, S. Baldus, B.A. Freeman, Nitro-fatty acids reduce atherosclerosis in apolipoprotein E-deficient mice, *Arterioscler. Thromb. Vasc. Biol.* 30 (2010) 938–945, <https://doi.org/10.1161/ATVBAHA.109.201582>.
- [88] L. Villacorta, J. Zhang, M.T. Garcia-Barrio, X. Chen, B.A. Freeman, Y.E. Chen, T. Cui, Nitro-linoleic acid inhibits vascular smooth muscle cell proliferation via the Keap1/Nrf2 signaling pathway, *Am. J. Physiol. Heart Circ. Physiol.* 293 (2007) H770–H776, <https://doi.org/10.1152/ajpheart.00261.2007>.

- [89] M.M. Wright, J. Kim, T.D. Hock, N. Leitinger, B.A. Freeman, A. Agarwal, Human haem oxygenase-1 induction by nitro-linoleic acid is mediated by cAMP, AP-1 and E-box response element interactions, *Biochem. J.* 422 (2009) 353–361, <https://doi.org/10.1042/BJ20090339>.
- [90] C.-S.C. Woodcock, Y. Huang, S.R. Woodcock, S.R. Salvatore, B. Singh, F. Golin-Bisello, N.E. Davidson, C.A. Neumann, B.A. Freeman, S.G. Wendell, Nitro-fatty acid inhibition of triple-negative breast cancer cell viability, migration, invasion, and tumor growth, *J. Biol. Chem.* 293 (2018) 1120–1137, <https://doi.org/10.1074/jbc.M117.814368>.
- [91] A. Trostchansky, M. Mastrogianni, E. Miquel, S. Rodríguez-Bottero, L. Martínez-Palma, P. Cassina, H. Rubbo, Profile of arachidonic acid-derived inflammatory markers and its modulation by nitro-oleic acid in an inherited model of amyotrophic lateral sclerosis, *Front. Mol. Neurosci.* 11 (2018) 131, <https://doi.org/10.3389/fnmol.2018.00131>.
- [92] T. Ichikawa, J. Zhang, K. Chen, Y. Liu, F.J. Schopfer, P.R.S. Baker, B.A. Freeman, Y.E. Chen, T. Cui, Nitroalkenes suppress lipopolysaccharide-induced signal transducer and activator of transcription signaling in macrophages: a critical role of mitogen-activated protein kinase phosphatase 1, *Endocrinology* 149 (2008) 4086–4094, <https://doi.org/10.1210/en.2007-1639>.
- [93] M.M. Wright, F.J. Schopfer, P.R.S. Baker, V. Vidyasagar, P. Powell, P. Chumley, K.E. Iles, B.A. Freeman, A. Agarwal, Fatty acid transduction of nitric oxide signaling: nitro-linoleic acid potentially activates endothelial heme oxygenase 1 expression, *Proc. Natl. Acad. Sci. USA* 103 (2006) 4299–4304, <https://doi.org/10.1073/pnas.0506541103>.
- [94] L.M.S. Baker, P.R.S. Baker, F. Golin-Bisello, F.J. Schopfer, M. Fink, S.R. Woodcock, B.P. Branchaud, R. Radi, B.A. Freeman, Nitro-fatty acid reaction with glutathione and cysteine. Kinetic analysis of thiol alkylation by a Michael addition reaction, *J. Biol. Chem.* 282 (2007) 31085–31093, <https://doi.org/10.1074/jbc.M704085200>.
- [95] L. González-Perilli, M. Mastrogianni, D. de Castro Fernandes, H. Rubbo, F. Laurindo, A. Trostchansky, Nitroarachidonic acid (NO₂AA) inhibits protein disulfide isomerase (PDI) through reversible covalent adduct formation with critical cysteines, *Biochim. Biophys. Acta - Gen. Subj.* (2017), <https://doi.org/10.1016/j.bbagen.2017.02.013>.
- [96] D.E. Artim, F. Bazely, S.L. Daugherty, A. Sculporeanu, K.B. Koronowski, F.J. Schopfer, S.R. Woodcock, B.A. Freeman, W.C. de Groat, Nitro-oleic acid targets transient receptor potential (TRP) channels in capsaicin sensitive afferent nerves of rat urinary bladder, *Exp. Neurol.* 232 (2011) 90–99, <https://doi.org/10.1016/j.expneurol.2011.08.007>.
- [97] X. Zhang, K.B. Koronowski, L. Li, B.A. Freeman, S. Woodcock, W.C. de Groat, Nitro-oleic acid desensitizes TRPA1 and TRPV1 agonist responses in adult rat DRG neurons, *Exp. Neurol.* 251 (2014) 12–21, <https://doi.org/10.1016/j.expneurol.2013.10.020>.
- [98] A. Sculporeanu, F.A. Kullmann, D.E. Artim, F.A. Bazely, F. Schopfer, S. Woodcock, B.A. Freeman, W.C. de Groat, Nitro-oleic acid inhibits firing and activates TRPV1- and TRPA1-mediated inward currents in dorsal root ganglion neurons from adult male rats, *J. Pharmacol. Exp. Ther.* 333 (2010) 883–895, <https://doi.org/10.1124/jpet.109.163154>.
- [99] X. Zhang, J.M. Beckel, S.L. Daugherty, T. Wang, S.R. Woodcock, B.A. Freeman, W.C. de Groat, Activation of TRPC channels contributes to OA-NO₂-induced responses in guinea-pig dorsal root ganglion neurons, *J. Physiol.* 592 (2014) 4297–4312, <https://doi.org/10.1113/jphysiol.2014.271783>.
- [100] F.J. Schopfer, Y. Lin, P.R.S. Baker, T. Cui, M. Garcia-Barrio, J. Zhang, K. Chen, Y.E. Chen, B.A. Freeman, Nitro-linoleic acid: an endogenous peroxisome proliferator-activated receptor gamma ligand, *Proc. Natl. Acad. Sci. USA* 102 (2005) 2340–2345, <https://doi.org/10.1073/pnas.0408384102>.
- [101] Y. Li, J. Zhang, F.J. Schopfer, D. Martynowski, M.T. Garcia-Barrio, A. Kovach, K. Suino-Powell, P.R.S. Baker, B.A. Freeman, Y.E. Chen, H.E. Xu, Molecular recognition of nitrated fatty acids by PPAR gamma, *Nat. Struct. Mol. Biol.* 15 (2008) 865–867, <https://doi.org/10.1038/nsmb.1447>.
- [102] M.J. Gorczynski, P.K. Smitherman, T.E. Akiyama, H.B. Wood, J.P. Berger, S.B. King, C.S. Morrow, Activation of peroxisome proliferator-activated receptor gamma (PPARgamma) by nitroalkene fatty acids: importance of nitration position and degree of unsaturation, *J. Med. Chem.* 52 (2009) 4631–4639, <https://doi.org/10.1021/jm900326c>.
- [103] Y.V. Vasil'ev, S.-C. Tzeng, L. Huang, C.S. Maier, Protein modifications by electrophilic lipoxidation products: adduct formation, chemical strategies and tandem mass spectrometry for their detection and identification, *Mass Spectrom. Rev.* 33 (2014) 157–182 <https://www.ncbi.nlm.nih.gov/pubmed/24818247>.
- [104] M. Fedorova, R.C. Bollineni, R. Hoffmann, Protein carbonylation as a major hallmark of oxidative damage: update of analytical strategies, *Mass Spectrom. Rev.* 33 (2013) 79–97, <https://doi.org/10.1002/mas.21381>.
- [105] T. Melo, S.S. Marques, I. Ferreira, M.T. Cruz, P. Domingues, M.A. Segundo, M.R.M. Domingues, New insights into the anti-inflammatory and antioxidant properties of nitrated phospholipids, *Lipids* 53 (2018) 117–131, <https://doi.org/10.1002/lipd.12007>.
- [106] M. Fazzari, N.K.H. Khoo, S.R. Woodcock, D.K. Jorkasky, L. Li, F.J. Schopfer, B.A. Freeman, Nitro-fatty acid pharmacokinetics in the adipose tissue compartment, *J. Lipid Res.* 58 (2017) 375–385, <https://doi.org/10.1194/jlr.M072058>.
- [107] L. Villacorta, L. Chang, S.R. Salvatore, T. Ichikawa, J. Zhang, D. Petrovic-Djergovic, L. Jia, H. Carlsen, F.J. Schopfer, B.A. Freeman, Y.E. Chen, Electrophilic nitro-fatty acids inhibit vascular inflammation by disrupting LPS-dependent TLR4 signalling in lipid rafts, *Cardiovasc. Res.* 98 (2013) 116–124, <https://doi.org/10.1093/cvr/cvt002>.
- [108] A.T. Reddy, S.P. Lakshmi, S. Dornadula, S. Pinni, D.R. Rampa, R.C. Reddy, The Nitrated Fatty Acid 10-Nitro-Oleate Attenuates Allergic Airway Disease, *J. Immunol.* (2013), <https://doi.org/10.4049/jimmunol.1300730>.
- [109] A. Trostchansky, L. Bonilla, C.P. Thomas, V.B. O'Donnell, L.J. Marnett, R. Radi, H. Rubbo, Nitroarachidonic acid, a novel peroxidase inhibitor of prostaglandin endoperoxide H synthases 1 and 2, *J. Biol. Chem.* 286 (2011) 12891–12900, <https://doi.org/10.1074/jbc.M110.154518>.
- [110] E.E. Kelley, J. Baust, G. Bonacci, F. Golin-Bisello, J.E. Devlin, C.M. Croix St, S.C. Watkins, S. Gor, N. Cantu-Medellin, E.R. Weidert, J.C. Frisbee, M.T. Gladwin, H.C. Champion, B.A. Freeman, N.K.H. Khoo, Fatty acid nitroalkenes ameliorate glucose intolerance and pulmonary hypertension in high-fat diet-induced obesity, *Cardiovasc. Res.* 101 (2014) 352–363, <https://doi.org/10.1093/cvr/cvt341>.
- [111] A. Klinke, A. Möller, M. Pekarova, T. Ravekes, K. Friedrichs, M. Berlin, K.M. Scheu, L. Kubala, H. Kolarova, G. Ambrozova, R.T. Schermuly, S.R. Woodcock, B.A. Freeman, S. Rosenkranz, S. Baldus, V. Rudolph, T.K. Rudolph, Protective effects of 10-nitro-oleic acid in a hypoxia-induced murine model of pulmonary hypertension, *Am. J. Respir. Cell Mol. Biol.* 51 (2014) 155–162, <https://doi.org/10.1165/rncmb.2013-00630C>.
- [112] H. Wang, H. Liu, Z. Jia, C. Olsen, S. Litwin, G. Guan, T. Yang, Nitro-oleic acid protects against endotoxin-induced endotoxemia and multiorgan injury in mice, *Am. J. Physiol. Ren. Physiol.* 298 (2010) F754–F762, <https://doi.org/10.1152/ajprenal.00439.2009>.
- [113] T.K. Rudolph, T. Ravekes, A. Klinke, K. Friedrichs, M. Mollenhauer, M. Pekarova, G. Ambrozova, H. Martiskova, J.-J. Kaur, B. Matthes, A. Schwoerer, S.R. Woodcock, L. Kubala, B.A. Freeman, S. Baldus, V. Rudolph, Nitrated fatty acids suppress angiotensin II-mediated fibrotic remodelling and atrial fibrillation, *Cardiovasc. Res.* 109 (2016) 174–184, <https://doi.org/10.1093/cvr/cvz254>.
- [114] E. Shin, E. Yeo, J. Lim, Y.H. Chang, H. Park, E. Shim, H. Chung, H.J. Hwang, J. Chun, J. Hwang, Nitrooleate mediates nitric oxide synthase activation in endothelial cells, *Lipids* 49 (2014) 457–466, <https://doi.org/10.1007/s11745-014-3893-8>.
- [115] L. Bonilla, V.B. O'Donnell, S.R. Clark, H. Rubbo, A. Trostchansky, Regulation of protein kinase C by nitroarachidonic acid: impact on human platelet activation, *Arch. Biochem. Biophys.* 533 (2013) 55–61, <https://doi.org/10.1016/j.abb.2013.03.001>.
- [116] T. Uchiyama, H. Atsuta, T. Utsugi, M. Oguri, A. Hasegawa, T. Nakamura, A. Nakai, M. Nakata, I. Maruyama, H. Tomura, F. Okajima, S. Tomono, S. Kawazu, R. Nagai, M. Kurabayashi, HSF1 and constitutively active HSF1 improve vascular endothelial function (heat shock proteins improve vascular endothelial function), *Atherosclerosis* 190 (2007) 321–329, <https://doi.org/10.1016/j.atherosclerosis.2006.03.037>.