



## ORIGINAL ARTICLE

# Clinical characteristic of a Haitian stroke cohort and a scoping review of the literature of stroke among the Haitian population

Axler Jean Paul<sup>1\*</sup>, Jude Hassan Charles<sup>2</sup>, Gandhi Marius Edwitch Gedner<sup>1</sup>, Richardson Roche<sup>1</sup>, Wislet Andre<sup>1</sup>, Garly Rushler Saint Croix<sup>3</sup>, Gillian Gordon Perue<sup>2</sup>

<sup>1</sup>Department of Internal Medicine, State University of Haiti, Port-au-Prince, West, Haiti, <sup>2</sup>Department of Neurology, University of Miami Hospital/Jackson Health System, Miami, Florida, United States of America, <sup>3</sup>Interventional Cardiology, Einstein College of Medicine, Montefiore Medical Center, Bronx, New York, United States of America

## ARTICLE INFO

*Article history:*

Received: December 19, 2022

Revised: April 4, 2023

Accepted: April 17, 2023

Published: May 12, 2023

*Keywords:*

Cerebrovascular disease

Haiti

Low- and middle-income countries

Stroke

*\*Corresponding author:*

Axler Jean Paul

Department of internal medicine, State

University of Haiti, 79, Impasse Dady,

Juvénat, Petion-ville, Haiti.

Email: [jeanpaulaxler@hotmail.com](mailto:jeanpaulaxler@hotmail.com)

© 2023 Author(s). This is an Open-Access article distributed under the terms of the Creative Commons Attribution-Noncommercial License, permitting all non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

## ABSTRACT

**Background and Aim:** There are significant disparities in stroke care and outcomes between low- and middle-income countries compared to high-income countries. Haiti, a lower-middle-income country, suffers from a lack of resources for acute stroke management. This study is the first to report the epidemiological profile of the Haitian population presenting with stroke symptoms at the largest academic hospital in the nation.

**Methods:** This is an observational study conducted over a period of 5 months from April 2021 to August 2021 in the Internal Medicine Department of the State University Hospital of Haiti. There were 51 included patients who were suspected to have had an acute stroke. A descriptive statistical analysis was conducted. A scoping review of the literature was also conducted.

**Results:** Over 50% of included patients were between 19 and 65 years old. The mean age at presentation was 61 years, and patients were predominantly female (64.7%). The prevalence of severe motor deficits was over 96%. The mean National Institutes of Health Stroke Scale was 12. Only 15.7% of patients (8/51) had a computed tomography (CT) scan during their hospitalization. The median time to CT scan was 84 h after symptom onset. About 80% of those with complications took more than 24 h to arrive at the hospital after the onset of symptoms. Eleven percent of patients had complications, and the mortality rate was 3.9%. There was a significant association between the Modified Rankin Scale and the occurrence of complications ( $p = 0.016$ ). National Institutes of Health Stroke Scale (NIHSS) score had a significant association with the Glasgow score ( $F = 6.3; p < 0.001$ ) where an inversely proportional correlation was observed between them ( $r = -0.7; p < 0.001$ ) and a proportional correlation with the Rankin prediction score and the NIHSS ( $r = 0.3, p = 0.04$ ). Little is known about the epidemiology of stroke patients in Haiti, and this limits the ability to develop targeted interventions to improve outcomes. In our scoping review, only three pertinent studies were identified over a 25-year period, this leads to a lack of data in regard to stroke care in Haiti mainly due to the absence of trained personnel.

**Conclusion:** In our cohort, stroke is mainly affecting female patients. The majority of stroke patients have moderate to severe motor deficits and took more than 24 h to arrive at the hospital. Urgent assistance is needed to strengthen personnel and infrastructure dedicated to stroke. Neurological assessment based on NIHSS and Rankin score should be systematic in stroke evaluation in Haiti.

**Relevance for Patients:** This study is relevant for patients because it emphasizes the challenges of stroke management in Haiti due to the non-availability of reference drugs, the time to arrive at the hospital to start treatment, as well as the means of diagnosis which are limited, like the CT scan. While stroke prevalence is on the rise in the country, it is the highest in the Caribbean and Latin America region.

## 1. Introduction

Haiti has the highest incidence of stroke in the Caribbean and Latin American region with 176 cases/100.000 population, this is 2 times higher than the neighboring country of the Dominican Republic [1], representing the most frequent cause of admission to the emergency room [2,3]. However, little is known about the epidemiology of stroke in Haiti. In the Global Stroke Statistics 2019 report [4], Haiti was noted to have a low stroke mortality rate, but the authors acknowledge that the data are more than 18 years old. In a report from the World Health Organization (WHO) in 2021 [5], 31% of mortality in Haiti can be accounted for by cardiovascular events. Acute stroke treatment is time-dependent [6], and it includes intravenous thrombolysis, mechanical thrombectomy, or anticoagulation reversal depending on the stroke subtype. Early management has been shown to promote better outcomes. In Haiti, there are no data available on the time to hospital presentation for stroke patients, and there is no systematic use of the National Institutes of Health Stroke Scale (NIHSS) score to measure stroke severity, and unfortunately, thrombolysis and thrombectomy are not yet practical in Haiti, due to lack of availability and cost. Our study aims to describe the clinical characteristics of stroke patients in the largest hospital in Haiti and to highlight challenges associated with stroke management, including delays in presentation and challenges to imaging acquisition and treatment. We also conducted a scoping review of the literature about stroke in the Haitian population.

## 2. Methodology

This is an observational prospective cohort pilot study and a scoping review of stroke in Haiti. The cohort study was conducted over 5 months from April 2021 to August 2021 in the emergency department of the internal medicine of the State University Hospital of Haiti (HUEH), the main teaching hospital of the Faculty of Medicine of the Haitian State University. Patients referred to us often did not have these parameters, as neurological evaluation with Rankin score and NIHSS is not systematic in other internal medicine departments in the country, and those patients were not included.

All stroke cases received and admitted to the emergency department of the internal medicine were included, sample was not considered for the study. We included stroke patients with complete neurological assessment with Glasgow, NIHSS, and Rankin score. Patients referred to us did not often have these parameters, as neurological evaluation with Rankin score and NIHSS is not systematic in other internal medicine departments in the country, are not included without them. During this period, we identified 51 cases with a clinical diagnosis of stroke that was admitted to HEUH with neurological assessment. All stroke cases admitted were considered without specific precision on stroke subtype. Patient data was extracted from the medical record using a standardized collection form. Data extraction spanned from the time of admission discharge or death. Anonymity and confidentiality of patients were protected, and oral consent was obtained from relatives or responsible parents. The data collection form included details on demographic data such as age, sex,

address, occupation, level of education and marital status; clinical data such as chief complaint, history and/or associated defects, date and time of symptoms onset, admission date, vital signs, time to computed tomography (CT) scan, length of stay, referral or not; and neurological deficits including language, motor and sensory deficits, NIHSS score on admission, modified ranking scale (mRS), Glasgow score, state of consciousness, and outcomes such as complications, death or recovery. The NIHSS score was used to categorize stroke into mild (1 – 4), moderate (5 – 15), severe (16 – 20), and grave (>20). The mRS was used, at the first clinical evaluation at admission, to categorize stroke outcome as favorable (0 – 1 point) or unfavorable (2 – 5 points) depending on the newly considered mRS dichotomization [7]. Arrival time was considered as the time of arrival to hospital after stroke onset. This was classified into five categories: <3 h; 3 – 6 h; 6 – 12 h; 12 – 24 h; and >24 h after symptom onset.

The data were then entered into Excel and exported to SPSS 26 for processing and analysis. A descriptive analysis of demographic and clinical parameters was performed and presented in table and graph form. Correlation analyses were performed between the NIHSS score and the variables age, time of arrival, SBP, DBP, Glasgow score, ranking score, and length of stay in hospital. Multiple regression analysis was used to identify significant correlations. An ANOVA test was used to verify associations, Student's *t* tests were performed for comparison of means, and the Levene test was used for homogeneity of variances. The significance level was set at  $p < 0.05$ .

A scoping review of the literature was conducted (Table 1). The authors developed a search in PubMed/Medline, EMBASE, Scopus, clinical trials.gov, and the Cochrane Library along with several grey literature sources. The searches were conducted from circa January 1997 to January 2022. The search was limited to English and French. Both subject headings and article text were searched for the following terms: “ischemic stroke,” “cerebrovascular disease/stroke,” “hemorrhagic stroke,” “brain ischemia,” “acute stroke,” “acute ischemic stroke,” “intracranial occlusion,” or “arterial occlusion,” “subarachnoid hemorrhage” AND “Haiti,” “low- to middle-income countries,” and “Haitian populations.” Only three pertinent studies were identified over a 25-year period.

## 3. Results


### 3.1. Demographics

There were 51 included patients who were admitted for stroke based on clinical evaluation to the internal medicine department at HUEH from April 2021 to August 2021. Nearly two-thirds (64.7%) of our patients were women, and the mean age was  $61.60 \pm 16.13$  years. More than 50% of our patients were between 19 and 65-years-old, and there was a predominance of women in all age groups. However, males had a slightly higher mean age than females (62.94 vs. 60.88 years, respectively) (Figure 1).

The districts of Port-au-Prince and Carrefour (the closest to the HUEH) were the most represented, constituting 47% and 25% of included patients, respectively. The most common occupation was retailer (49%).

**Table 1.** Scoping review of the literature

Study	Year	Mean age/% sex	Mean NIHSS reported	Arrival time	Key finding
Stroke epidemiology among Haitians living in Miami (PMID 16103730)	2005 [29]	60 years/48% female	7	N/A	HTN was the most common stroke risk factor at 87%
Head CT findings at public hospital in rural Haiti	2017 [30]	60.8±17 years/N/A	N/A	N/A	36% of stroke were hemorrhagic. AMS and focal deficit most common presentation
Protocolized emergency department observation care improves quality of ischemic stroke care in Haiti	2020 [31]	62±14 years/59% Female	N/A	4.9±7.1 days	Reported mean numbers of comorbidities 0.7±0.6

**Figure 1.** Bivariate analysis between age and sex for our 51 patients. (A) A box-and-whisker plot of patient age stratified by sex. (B) A histogram showing number of patients by age category and sex. (C) A histogram showing number of patients above and below 35 years of age by sex.

### 3.2. Clinical presentations

The three main chief complaints were as follows: altered mental status (18 patients, 35.29%), hemiplegia (ten patients, 19.60%), and aphasia (13 patients, 25.5%). The findings of neurological examinations are described in Table 2.

Aphasia was present in 25.5% of the cases, dysarthria in 29.4%, facial paralysis in 66.7%, a motor deficit in 96.1% of the cases with the right-sided deficit in 41.1% and left sided 49.1%, and a sensory deficit in 23.5% of the cases. About 10% were referred from small urban clinics. Nearly two-thirds of the patients (65%) were known to have hypertension and 4% were diabetic. The mean time to arrival at the hospital after symptom onset was 75.14 h.

The NIHSS score was only valuable for 36/51 patients with a mean 12.58 points (Table 2). We found that 7.8% of patients had mild stroke, 41.2% had moderate stroke, and 13.7% had severe stroke. About 74.5% of the patients had a baseline mRS between 0 and 2 (Table 2).

### 3.3. Outcomes

Only 15% of patients (8/51) had a CT scan during their hospitalization. The majority were delayed with a median time to CT of 84 h after symptoms onset. The most common finding ischemia in one of the middle cerebral artery territories. Patients spent about 10 days in hospital after admission. Six complications were from aspiration pneumonia, and two in-hospital deaths were

recorded (3.9%). The patients with complications were generally over 65 years of age (83.3%) with a mean age of  $69.33 \pm 25.48$  compared with  $60.5 \pm 14.58$  for uncomplicated cases. This was not a statistically significant difference ( $t = 1.25$ ;  $p = 0.21$ ). We observed that 80% of those with complications took more than 24 h to arrive at the hospital after the onset of symptoms, on average  $129.33 \pm 105.09$  h compared with  $66.11 \pm 106.63$  h for uncomplicated cases. This also was not a statistically significant

difference ( $t = 1.34$ ;  $p = 0.18$ ). All cases of complications were in the category of moderate stroke, with mean NIHSS scores of  $17.33 \pm 14.46$  point for complications versus  $12.15 \pm 5.85$  for patients with no complications ( $t = 1.28$ ;  $p = 0.20$ ), and the variance was homogeneous according to Levene's test  $F = 9.80$ ;  $p = 0.004$ . However, no association was found between complication and NIHSS score ( $F = 1.65$ ;  $p = 0.20$ ). Two-thirds (66.7%) of patients had a moderate Glasgow score (9 – 12) (Table 2). There was a significant association between the mRS and the occurrence of a complication ( $F = 6.33$ ;  $p = 0.016$ ). The mRS was higher in patients with complications with a score  $>3$  versus (0 – 2) in the patients without complications, and the difference was significant ( $t = 2.51$ ;  $p = 0.016$ ). It should be noted that the variance in this group is not homogeneous when considering the Levene test  $F(5) = 5.98$ ;  $p = 0.019$ . A mRS of 0.5 (cut off point) has a sensitivity of 60% and a specificity of 86% to predict the occurrence of a complication with an accuracy of 76% (AUC = 0.76) (Figure 2).

**Table 2.** Clinical characteristics of patients presents with acute stroke symptoms to HEUH


Demographics	Findings (n; %)
Age (mean±SD)	61.60±16.13
<19 years	1; 2%
19–35 years	1; 2%
35–65 years	26; 51%
>65 years	23; 45%
Sex (% female)	33; 64.74%
Chief complaints	
Aphasia	13; 25.50%
Dysarthria	15; 29.4%
Facial drop	34; 66.6%
AMS	20; 39.2%
Hemiparesis	15; 19.4%
Limb ataxia	1; 2%
Hemiplegia	28; 54.9%
Monoplegia	3; 5.8%
Paraparesis	1; 2%
Hypoesthesia	12; 23.5%
Comorbid medical illnesses	
Hypertension	33; 65%
Diabetes	2; 4%
Heart disease	1; 2%
Epilepsy	1; 2%
None	14; 27%
Average arrival time (mean±SD)	75±108 h
NIHSS (mean±SD)	12±6 points
Minor	4; 7.8%
Moderate	21; 41.2%
Severe	7; 13.7%
Grave	4; 7.8%
Time to CT (mean±SD)	84±44 h
GCS (mean±SD)	11.47±3.38
Severe	10; 19.6%
Moderate	19; 37.3%
Low	22; 43.1%
Average length of stay (mean±SD)	10±7 days
Outcome	
mRS 0 – 2	38; 74.50%
mRS 3 – 5	11; 21.60%
mRS 6	2; 3.90%

CT: Computed tomography; NIHSS: National Institutes of Health Stroke Scale; GCS: Glasgow coma scale; SD: Standard deviation

#### 4. Discussion

Stroke is one of the leading causes of death and disability in the world [8]. In Central America and the Caribbean, Haiti has the highest mortality rate from stroke [1]. To the best of our knowledge, this is the first study describing sociodemographic and clinical characteristics of stroke patients in Haiti as well as the scarcity of human and material resources in terms of stroke care.

For the 51 patients of our study, a predominance of women was observed which is similar to international studies [9,10] where 55,000 more annual cases of stroke were in women [11]. However, in our study, men were slightly older than women (62.94 vs. 60.88 years). This is different from international observations where women with stroke are often older than men. [11-13]. The mean age of our patients is 61.60 years compared with the global average age of stroke patients of 70 years. Strokes among Haitian patients are occurring an average of 10 years earlier than those in developed countries [1]. For instance, Abreu *et al.* reported an



**Figure 2.** Receiver operating characteristic curve analysis to assess the sensitivity and specificity of the Rankin score in the risk of complications.



average age of 70 years [12], and Soto-Cámara *et al.* reported an average age of 75.39 years [14]. The average age of our study is close to some developing countries such as Jordan where Al-Qawasmeh *et al.* reported an average age of 66.5 years, and in Tanzania where Matuja *et al.* reported an average age of 57.9 years [13,15]. Regarding occupation, almost half of included patients (49%) were retail merchant operating in the informal market. This demonstrates that stroke is affecting patients in their economic prime, which imposes a significant economic burden to the society. Moreover, 65% of included patients had hypertension which is a treatable risk factor for stroke. It has been shown that the incidence of stroke is increasing in low-income countries compared with high-income countries [16,17]. One explanation for this is the impact of social determinants such as poverty, education, lifestyle, and stress on the cardiovascular system, including hypertension [14,18,19]. A 2021 report by the WHO reported that about one-third of mortality in Haiti is related to a cardiovascular event.

There is evidence that delay in seeking care compromises management, diagnosis, and outcomes in stroke patients [20,21]. Factors such as symptom onset, history, gender, and stroke severity are known to influence the time arrival at the hospital [21]. The mean time to hospital after symptom onset in our study was 75.14 h, or 3.13 days. Approximately 39.2% of patients arrived after 24 h, and more than half or 51% of patients arrived after 12 h. These are much longer delays than reported elsewhere in the literature, for example, publications have reported a mean time to arrival at the hospital of 21.96 h in China and 10.73 h in the USA [17]. The recommended ideal time is <4.5 h to allow for the use of intravenous thrombolysis and between six and 24 h for endovascular thrombectomy [21,22], both treatments that are not available in Haiti. We observed wide variability in time to arrival based on sex (99.16 h for women vs. 7.45 h for men), a difference was significant ( $t = -2.59$ ,  $p = 0.013$ ). Similar findings are described in a study by Le *et al.*, in which women faced greater delays than men [21].

Immediate assessment, including neurological evaluation, is critical in the management of stroke patients. Several scores are used to assess the neurological status of stroke patients, including the Scandinavian Stroke Scale (SSS), the Canadian Neurological Score (CNS), the modified Rankin Scale (mRS), and the NIHSS score [23]. The NIHSS score is the most widely used because it allows estimation of the neurological impact, severity, complications, and prognosis of stroke with simplicity, rapidity, and good sensitivity and reproducibility [24,25]. In our study, the NIHSS score and the Glasgow score were used to assess and categorize stroke severity, and the modified Rankin score was used to establish pre-stroke functional level. In 80% of cases, patients who developed complications arrived after 24 h from the onset of symptoms ( $129.33 \pm 105.09$  h,  $p = 0.18$ ). All of these had a moderate stroke according to the NIHSS score ( $17.33$  vs.  $\pm 14.46$ ,  $p = 0.20$ ). Similarly, Schlegel *et al.* observed an association between NIHSS score and complications ( $p < 0.001$ ). Garavelli *et al.* reported the same finding ( $p = 0.048$ ), and Bovim *et al.* observed that 85.4% of the cases of complications had a delay

of more than 24 h after the onset of symptoms [25-27]. The mRS score was higher in patients with complications ( $1.20 \pm 1.64$  vs.  $0.17 \pm 0.70$ ), with a statistically significant difference ( $t = 2.51$ ;  $p = 0.016$ ). The Rankin score was significantly associated with the occurrence of complications in patients with stroke ( $p = 0.016$ ), such as aspiration pneumonia, with a sensitivity of 60% and a specificity of 86% (AUC = 0.76) (Figure 2). Thus, the Rankin score can be used to predict the occurrence of complications in stroke patients. This is valuable because studies have shown that the occurrence of complications after stroke is a poor prognostic factor [26]. We observed that the NIHSS score has a very highly significant association with the Glasgow score ( $p < 0.001$ ) where an inverse proportional correlation was observed ( $r = -0.7$ ;  $p < 0.001$ ) and a proportional correlation with the Rankin prediction score ( $r = 0.3$ ,  $p = 0.04$ ). This means that in the initial clinical assessment of stroke patients, the NIHSS score and the Rankin score can help predict which patients may develop complications. This has been proven by several previous studies [23,27,28].

Little is known about the epidemiology of stroke patients in Haiti, and this limits the ability to develop targeted interventions to improve outcomes. In our scoping review, only three pertinent studies were identified over a 25-year period (Table 1).

The first study was done by Koch *et al.* on the Haitian population living in Miami and does not provide information on the local resources, stroke treatments, and outcomes in Haiti [29]. The other two studies were conducted in Haiti. Like our study, the mean age of stroke among these Haitian populations is 10 years younger than cohorts from developed countries, suggesting that stroke affects Haitians at their economic prime [30,31]. Rouhani *et al.* also document delays in care including delays to arrival time and CT imaging [31]. Alongside our present study, this literature demonstrates the need for ongoing research and stroke interventions to save lives and prevent stroke-related disability among the Haitian population. The utilization of stroke-related metric such the use of NIHSS, modified Rankin scores, and time to intervention is not systematic even in the largest academic center in the metropolitan area. This leads to a lack of data in regard of stroke care in Haiti mainly due to the absence of trained personals.

#### 4.1. Limitations

Our study has a small cohort of 51 patients. This may limit the generalizability of our findings. Unfortunately, due to challenges associated with socioeconomic barriers to care and limited stroke health-care systems, none of the patients attended follow-up appointment. Some potential bias might be present in the evaluation due to delay to care and lack of standard management regarding the length of stay, findings, and complication. Those limit the evaluation of long-term outcomes. However, this study will serve as a foundation for future observational and interventional studies on stroke in Haiti.

## 5. Conclusion

In our cohort, stroke is mainly affecting female patient. The majority of stroke patient have moderate cerebral deficit based on

the NIHSS score and took more than 24 h to arrive at hospital. Time to have a CT scan is 84 h after stroke events, and only 15% of patient had a CT scan during their hospitalization. There is a salient lack of equipped health-care facilities and appropriate treatment for stroke management. Factors such as low income, limited education, and long arrival time after onset of symptoms are associated with poor outcomes and are found across the board in our population. Neurological assessment based on NIHSS and Rankin score should be systematic in stroke evaluation. There is an urgent need for assistance in both human and infrastructural resources dedicated to stroke unit care, specialized training program for providers in Haiti. Further study is necessary to assess the stroke issues in Haiti.

### Acknowledgments

Special thanks to Dr. Rolvix Patterson, Dr. Raema Mimrod Jean, and Dr. Ansly Jefferson Desravannes.

### Conflicts of Interest

The authors declare that there are no conflicts of interest.

### Ethics Approval and Consent to Participate

This study has been approved for publication by the head directory of the Internal Medicine of the State University Hospital of Haiti as it is a teaching hospital. Verbal consent was obtained from patients or responsible person of patients.

### Consent for Publication

Verbal informed consent of patients was obtained.

### References

- [1] Lookens J, Tymejczyk O, Rouzier V, Smith C, Preval F, Joseph I, *et al.* The Haiti Cardiovascular Disease Cohort : Study Protocol for a Population-based Longitudinal Cohort. *BMC Public Health* 2020;20:1633.
- [2] Barthe EJ, Ernst S, Osborn I, Germano IM. A Prospective Emergency Department-based Study of Pattern and Outcome of Neurologic and Neurosurgical Diseases in Haiti. *World Neurosurg* 2014;82:948-53.
- [3] Lavados PM, Hennis AJ, Fernandes JG, Medina MT, Legetic B, Hoppe A, *et al.* Stroke Epidemiology, Prevention, and Management Strategies at a Regional Level : Latin America and the Caribbean. *Lancet Neurol* 2027;6:362-72.
- [4] Kim J, Thayabaranathan T, Donnan GA, Howard G, Howard VJ, Rothwell PM, *et al.* Global Stroke Statistics 2019. *Int J Stroke* 2020;15:819-38.
- [5] WHO, World Health Organization (WHO). World Health Statistics 2019: Moanitoring Health for the SDGs, Sustainable Development Goals. Vol. 8. Geneva: World Health Organization; 2016. p. 5.
- [6] Powers WJ, Rabinstein AA, Ackerson T, Adeoye OM, Bambakidis NC, Becker K, *et al.* Guidelines for the Early Management of Patients with Acute Ischemic Stroke: 2019 Update to the 2018 Guidelines for the Early Management of Acute Ischemic Stroke a Guideline for Healthcare Professionals from the American Heart Association. *Stroke* 2019;50:e344-418.
- [7] Dijkland SA, Voormolen DC, Venema E, Roozenbeek B, Polinder S, Haagsma JA, *et al.* Utility-Weighted Modified Rankin Scale as Primary Outcome in Stroke Trials A Simulation Study. *Stroke* 2018;49:965-71.
- [8] Wang YX, Wei WB, Xu L, Jonas JB. Prevalence, Risk Factors and Associated Ocular Diseases of Cerebral Stroke: The Population-based Beijing Eye Study. *BMJ Open* 2020;10:e024646.
- [9] Renoux C, Coulombe J, Li L, Ganesh A, Silver L, Rothwell PM. Confounding by Pre-morbid Functional Status in Studies of Apparent Sex Differences in Severity and Outcome of Stroke. *Stroke* 2017;48:2731-8.
- [10] Haast RA, Gustafson DR, Kiliaan AJ. Sex Differences in Stroke. *J Cereb Blood Flow Metab* 2012;32:2100-7.
- [11] Zrelak PA. Sex-based Differences in Symptom Perception and Care-Seeking Behavior in Acute Stroke. *Perm J* 2018;22:18-042.
- [12] Abreu P, Magalhães R, Baptista D, Azevedo E, Silva MC, Correia M. Readmissions and Mortality during the First Year after Stroke-Data from a Population-based Incidence Study. *Front Neurol* 2020;11:636.
- [13] Al Qawasmeh M, Aldabbour B, Momani A, Obiedat D, Alhayek K, Kofahi R, *et al.* Epidemiology, Risk Factors, and Predictors of Disability in a Cohort of Jordanian Patients with the First Ischemic Stroke. *Stroke Res Treat* 2020;2020:1920583.
- [14] Soto-Cámara R, González-Bernal JJ, González-Santos J, Aguilar-Parra JM, Trigueros R, López-Liria R. Knowledge on Signs and Risk Factors in Stroke Patients. *J Clin Med*. 2020;9:2557.
- [15] Matuja SS, Khanbhai K, Mahawish KM, Munseri P. Stroke Mimics in Patients Clinically Diagnosed with Stroke at a Tertiary Teaching Hospital in Tanzania: A Prospective Cohort Study. *BMC Neurol* 2020;20:270.
- [16] Mathur S, Walter S, Grunwald IQ, Helwig SA, Lesmeister M, Fassbender K. Improving Prehospital Stroke Services in Rural and Underserved Settings with Mobile Stroke Units. *Front Neurol* 2019;10:159.
- [17] Wangqin R, Laskowitz DT, Wang Y, Li Z, Wang Y, Liu L, *et al.* International Comparison of Patient Characteristics and Quality of Care for Ischemic Stroke: Analysis of the China National Stroke Registry and the American Heart Association Get with the Guidelines-stroke Program. *J Am Heart Assoc* 2018;7:20.
- [18] Havranek EP, Mujahid MS, Barr DA, Blair IV, Cohen MS, Cruz-Flores S, *et al.* Social Determinants of Risk and Outcomes for Cardiovascular Disease A

- Scientific Statement From the American Heart Association. *Circulation* 2015;132:873-98.
- [19] Molokhia M, Yousif S, Durbaba S, Ashworth M, Harding S. Social Determinants of Diabetes, Hypertension, Stroke, and Coronary Heart Disease in Black Caribbean and Black African Women Aged 40 Years or Older in South London : Findings from 70 582 Primary Care Records from 2000-18. *Lancet* 2021;398:S96.
- [20] Jaffe TA, Goldstein JN, Yun BJ, Etherton M, Leslie-Mazwi T, Schwamm LH, *et al.* Impact of Emergency Department Crowding on Delays in Acute Stroke Care. *West J Emerg Med* 2020;21:891-8.
- [21] Le SM, Copeland LA, Zeber JE, Bemge JF, Allen L, Cho J, *et al.* Factors Affecting Time between Symptom Onset and Emergency Department Arrival in Stroke Patients. *eNeurologicalSci* 2020;21:100285.
- [22] Nguyen TT, Kruyt ND, Pierik JG, Doggen CJ, Van der Lugt P, Ramessersing SA, *et al.* Stroke Patient's Alarm Choice: General Practitioner or Emergency Medical Services. *Acta Neurol Scand* 2021;143:164-70.
- [23] Zöllner JP, Misselwitz B, Kaps M, Stein M, Konzalla J, Roth C, *et al.* National Institutes of Health Stroke Scale (NIHSS) on Admission Predicts Acute Symptomatic Seizure Risk in Ischemic Stroke : A Population-based Study Involving 135, 117 Cases. *Sci Rep* 2020;10:3779.
- [24] Marsh EB, Lawrence E, Gottesman RF, Llinas RH. The NIH Stroke Scale Has Limited Utility in Accurate Daily Monitoring of Neurologic Status. *Neurohospitalist* 2016;6:97-101.
- [25] Garavelli F, Ghelfi AM, Kilstein JG. Usefulness of NIHSS Score as a Predictor of Non-Neurological in-Hospital Complications in Stroke. *Med Clin (Barc)* 2021;157:434-7.
- [26] Bovim MR, Askim T, Lydersen S, Fjærtøft H, Indredavik B. Complications in the First Week after Stroke : A 10-Year Comparison. *BMC Neurol* 2016;16:133.
- [27] Schlegel D, Kolb SJ, Luciano JM, Tovar JM, Cucchiara BL, Liebeskind DS, *et al.* Utility of the NIH Stroke Scale as a Predictor of Hospital Disposition. *Stroke* 2003;34:134-7.
- [28] Aoki J, Kimura K, Koga M, Kario K, Nakagawara J, Furui E. NIHSS-Time Score Easily Predicts Outcomes in rt-PA Patients: The SAMURAI rt-PA Registry. *J Neurol Sci* 2013;327:6-11.
- [29] Koch S, Pabon D, Rabinstein AA, Chirinos J, Romano JG, Forteza A. Stroke Etiology among Haitians Living in Miami. *Neuroepidemiology* 2015;25:192-5.
- [30] Valtis YK, Cochran MF, Martineau L, Mendel JB, Berkowitz AL. Head CT Findings at a Public Hospital in Rural Haiti. *J Neurol Sci* 2017;379:327-30.
- [31] Rouhani SA, Marsh RH, Rimpel L, Anderson K, Outhay M, Edmond MC, *et al.* Protocolized Emergency Department Observation Care Improves Quality of Ischemic Stroke Care in Haiti. *Afr J Emerg Med* 2020;10:145-51.

#### Publisher's note

Whioce Publishing remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.