

Meeting abstract

Open Access

Decreased blood-brain barrier P-glycoprotein function with aging

Martin Bauer¹, Rudolf Karch², Aiman Abraham^{1,3}, Claudia C Wagner¹, Kurt Kletter⁴, Markus Müller¹ and Oliver Langer*^{1,3}

Address: ¹Department of Clinical Pharmacology, Medical University of Vienna, 1090 Vienna, Austria, ²Department of Medical Computer Sciences, Medical University of Vienna, 1090 Vienna, Austria, ³Department of Radiopharmaceuticals, Austrian Research Centers GmbH – ARC, 2444 Seibersdorf, Austria and ⁴Department of Nuclear Medicine, Medical University of Vienna, 1090 Vienna, Austria

Email: Oliver Langer* - oliver.langer@meduniwien.ac.at

* Corresponding author

from 14th Scientific Symposium of the Austrian Pharmacological Society (APHAR) Innsbruck, Austria. 21–22 November 2008

Published: 5 November 2008

BMC Pharmacology 2008, 8(Suppl 1):A48 doi:10.1186/1471-2210-8-S1-A48

This abstract is available from: <http://www.biomedcentral.com/1471-2210/8/S1/A48>

© 2008 Bauer et al; licensee BioMed Central Ltd.

Introduction

P-glycoprotein (P-gp) acts at the blood-brain barrier (BBB) as an active cell membrane efflux pump for several endogenous and exogenous compounds. The P-gp substrate (R)-[¹¹C]verapamil (VPM) can be used to measure P-gp-mediated transport at the BBB *in vivo* with positron emission tomography (PET). The distribution volume (DV) of VPM has been shown to inversely reflect P-gp function in the BBB [1].

Materials and methods

A young (n = 7, mean age: 28.0 ± 3.8 years) and an aged group (n = 6, mean age: 69.4 ± 8.5 years) of healthy volunteers underwent dynamic VPM PET scans and arterial blood sampling. Radiolabelled metabolites of VPM were quantified by a previously described combined solid-phase extraction/HPLC protocol [1]. A whole-brain grey matter region was defined by using the Hammersmith n20r49 brain atlas [2]. The DV of VPM was estimated by using a 2-rate-constant-1-tissue-compartment model [1].

Results

Mean DVs (± standard deviation) of VPM were 0.50 ± 0.08 for the young and 0.63 ± 0.13 for the aged group (+27% for the aged group, p = 0.04, 2-tailed *t*-test). There was no significant difference in VPM metabolism between the young and the aged group (area under the curve of the fraction of polar [¹¹C]metabolites of VPM versus time in

arterial plasma: 12.7 ± 2.4 and 14.1 ± 3.6 for the young and the aged group, respectively, p = 0.19, 2-tailed *t*-test).

Conclusion

Our data confirm previous results that older subjects show significantly decreased P-gp function in the BBB [1,3]. Decreased P-gp function can lead to increased accumulation of toxins and drugs in the aging brain and could thus be a risk factor for the development of neurodegenerative disease.

References

1. Toornvliet R, van Berckel BN, Luurtsema G, Lubberink M, Geldof AA, Bosch TM, Oerlemans R, Lammertsma AA, Franssen EJ: **Effect of age on functional P-glycoprotein in the blood-brain barrier measured by use of (R)-[¹¹C]verapamil and positron emission tomography.** *Clin Pharmacol Ther* 2006, **79**:540-548.
2. Hammers A, Allom R, Koeppe MJ, Free SL, Myers R, Lemieux L, Mitchell TN, Brooks DJ, Duncan JS: **Three-dimensional maximum probability atlas of the human brain, with particular reference to the temporal lobe.** *Hum Brain Mapp* 2003, **19**:224-247.
3. Bartels AL, Kortekaas R, Bart J, Willemsen AT, de Klerk OL, de Vries JJ, van Oostrom JC, Leenders KL: **Blood-brain barrier P-glycoprotein function decreases in specific brain regions with aging: A possible role in progressive neurodegeneration.** *Neurobiol Aging* in press.