

# Effectiveness of median nerve neural mobilization versus oral ibuprofen treatment in subjects who suffer from cervicobrachial pain: a randomized clinical trial

David Rodríguez Sanz<sup>1</sup>, Francisco Unda Solano<sup>2</sup>, Daniel López López<sup>3</sup>, Irene Sanz Corbalán<sup>4</sup>, Carlos Romero Morales<sup>1</sup>, Cesar Calvo Lobo<sup>5</sup>

<sup>1</sup>Department of Physical Therapy and Podiatry, Physical Therapy and Health Sciences Research Group Universidad Europea de Madrid, Madrid, Spain

<sup>2</sup>Interuniversity Degree in Physiotherapy UB-UdG/Grau en Fisioteràpia EUSES-UdG, Barcelona, Spain

<sup>3</sup>Research, Health and Podiatry Unit, Department of Health Sciences, Faculty of Nursing and Podiatry, Universidade da Coruña, Spain

<sup>4</sup>Department of Nursing, Physical Therapy and Podiatry, Universidad Complutense de Madrid, Madrid, Spain

<sup>5</sup>Nursing and Physical Therapy Department, Institute of Biomedicine (IBIOMED), Universidad de Leon, Leon, Spain

## Corresponding author:

Dr. Daniel López López PhD  
Faculty of Nursing  
And Podiatry  
University of A Coruña  
Campus de Esteiro  
Ferrol, 15403  
Coruna, Spain  
E-mail: daniellopez@udc.es@

**Submitted:** 20 July 2017

**Accepted:** 16 August 2017

Arch Med Sci 2018; 14, 4: 871–879

DOI: <https://doi.org/10.5114/aoms.2017.70328>

Copyright © 2017 Termedia & Banach

## Abstract

**Introduction:** Oral ibuprofen (OI) and median nerve neural mobilization (MNNM) are first line treatments for patients who suffer cervicobrachial pain (CP). OI may produce side effects which are not tolerated by all subjects who suffer CP, whereas MNNM has no known side effects. Therefore, the aim of this study was to assess the effectiveness of both treatments (OI vs. MNNM) in CP.

**Material and methods:** This investigation was a blinded parallel randomized clinical trial (NCT02593721). Sixty-two participants diagnosed with CP were recruited and randomly assigned to 2 groups ( $n = 31$ ), which received MNNM or 1200 mg/day OI treatment for 6 weeks. The numeric rating scale for pain intensity was the primary outcome. The cervical rotation range of motion (CROM) and the upper limb function were the secondary outcomes.

**Results:** The results showed that OI treatment ( $\eta^2 = 0.612-0.755$ ) was clearly superior to MNNM ( $\eta^2 = 0.816-0.821$ ) in all assessments ( $p < 0.05$ ) except for the CROM device results, which were equivalent to those of the MNNM group ( $p > 0.05$ ). Three subjects were discharged because of OI side effects.

**Conclusions:** Oral ibuprofen may be superior to MNNM for pain reduction and upper limb function increase of subjects with CP. Nevertheless, both treatments were effective. Median nerve neural mobilization may be considered an effective non-pharmaceutical treatment option in subjects with CP. Regarding OI adverse effects, our findings challenge the role of pharmacologic versus manual therapy as possible treatments that may improve pain intensity and upper limb functionality in subjects with CP.

**Key words:** non-steroidal anti-inflammatory agents, musculoskeletal manipulations, rehabilitation, upper extremity.

## Introduction

Cervicobrachialgia, also known as cervicobrachial pain (CP), is a high incidence and high prevalent well-described disabling medical condition

that affects 83 per every 100,000 people in the course of their lives [1–3]. It is described in the scientific literature as the presence of pain in the neck that radiates to or is associated with the arm [4, 5], and has a gold standard diagnosis through correlative pathological findings in a magnetic resonance imaging (MRI) [6–8]. Other helpful tools in the proper diagnosis of CP may be the presence of positive outcomes in the Spurling, upper limb and distraction orthopedic tests, as well as altered results in nerve conduction assessment [9–11]. The classical pattern of pain symptoms related to this condition are caused by the existence of musculoskeletal damage and neuropathic irradiation of pain due to underlying neural tissue injuries of the cervicobrachial anatomical region [12–14]. There is current evidence of pathologic nerve trunk mechano-sensibility alterations, central sensitizing and visco-elastic distortion of the cervicobrachial neural tissue during the onset of CP symptoms; this constitutes a key point in the proper treatment selection for CP and its adequate implementation [12–17].

Traditionally, first line treatment of CP involves the use of non-steroidal anti-inflammatory drugs (NSAIDs) and physical therapy techniques such as neural tissue mobilization of the median nerve. The principal NSAIDs prescribed worldwide to treat pain and CP include oral ibuprofen (OI) [18–25]. Both median nerve neural mobilization (MNNM) and OI relieve CP, but through different mechanisms and with a different onset of side effects. MNNM is a specific physical therapy technique for the treatment of CP described by Butler, Butler and Coppieters, and Elvey-Hall [14, 26–31] that achieves pain relief through mechanical stimulation of the median nerve and the brachial plexus. The proper mechanical stimulation of the nerve and its surrounding neural tissue induces a wide variety of physiologic responses that may reduce cervicobrachial pain including the activation of descending nervous system pain modulation mechanisms, although the entire set of underlying reasons for this pain reduction is not completely understood. The effects caused by the neural tissue mobilization that produce pain relief include intraneural edema reduction, changes in the intraneural nerve pressure, dispersion of pro-inflammatory substances and an increase in nerve mobility [31–33].

OI is a nonsteroidal anti-inflammatory drug used worldwide to control pain, fever and inflammation [34, 35]. Oral intake of ibuprofen, like most NSAIDs, produces a hypoalgesic effect due to biochemical inhibition of the COX enzymes, which convert arachidonic acid to prostaglandin H<sub>2</sub> (PGH<sub>2</sub>). PGH<sub>2</sub> is converted by other enzymes to several types of prostaglandins and thrombox-

anes (which are mediators of pain and inflammation) [36–38]. Side effects derived from the oral intake of ibuprofen are related to its systemic action and can be severe in some patients. This issue suggests that OI and other NSAIDs with similar mechanisms of action may not be suitable to be administered in all types of patients suffering from CP, whereas MNNM has little to no known side effects when applied properly. This reveals a point of interest in determining comparative effectiveness between these two first line therapeutic options of CP [18, 35, 37, 38].

Currently, there is a lack of single blinded randomized controlled clinical trials (RCT) regarding the comparison between MNNM and OI in CP, which by itself is an additional stand-alone problem for the practicing clinician who desires high quality evidence on the level of effectiveness of MNNM when compared to common over-the-counter pharmaceutical treatment for CP [36, 39–47]. Therefore, the purpose of the present study was to compare the effectiveness for pain intensity, range of movement and functionality between MNNM and OI in treating patients who suffer from CP.

## Material and methods

### Study design

This investigation was an interventional phase 2/phase 3 parallel (2 arm) single blind (outcome assessor) randomized clinical trial with an endpoint classification of efficacy study 1 : 1 allocation ratio conducted at only 1 clinical center. No changes were made in the proposed methodology because the original protocol was not altered. Furthermore, the Consolidated Statement for Reporting Trials (CONSORT) statement and checklist were considered [48].

### Ethical considerations and trial registry

The present study was designed and applied following the ethical principles for medical research involving human subjects established in the Declaration of Helsinki. The study protocol was approved by the Ethics Committee from La Viña Medical Center, Valencia, Venezuela (code CE0072015) and registered at ClinicalTrials (NCT02593721; October 26, 2015). Informed consent of all participants was obtained and the rights of subjects were protected.

### Randomization and blinding

Subjects were assigned to 1 of the 2 groups using restricted block randomization through block computerized randomization software ([www.randomizer.com](http://www.randomizer.com)) assuring that each block had a size

of 31 participants. Randomization and allocation to the trial group was carried out by computer software randomized printed cards contained in consecutively numbered opaque sealed envelopes that were handed to the participant by the physiatrist medical doctor (PMD). The outcome assessor (OA) was blinded to the randomization of group allocation. The principal investigator did not participate in the application of treatment or data collection.

### Participants

Subjects were recruited from a group of voluntary consecutive individuals seeking treatment or assessment related to CP in the established medical center that was selected to apply the study, or by referral of 5 other private and public medical centers. A total of 133 individuals were screened from July to August 2015. MRI corroboration and other necessary medical assessment including the need for diagnosis were performed by a specialized PMD. A total of 62 participants were recruited and divided into 2 groups of 31 subjects by the PMD in concordance with the randomization schedule that the data analyst (DA) generated. Group A contained MNNM participants and group B contained OI participants.

All eligible participants were adults who belonged to both genders ranging from 18 to 45 years of age with a clinical diagnosis of cervicobrachial pain confirmed by magnetic MRI and presence of unilateral symptoms of arm pain, paresthesia or numbness in the upper limb and positive results in all of the following tests: Spurling, distraction, and upper limb during at least 3 continuous months previous to enrollment. Exclusion criteria of participants were contraindication for intake of NSAIDs, the use of any type of treatment (therapy, procedure or drug) to relieve pain, and presence in participants of vertebral instability, vertebral osteoporosis, vertebral or spine infection and neurologic diseases of genetic, infectious or neoplastic origin, cervical stenosis myelopathy, pregnancy, kinesiphobia, endocrine disorders and menopause, history of spine surgery, intellectual disability, severe mental illness, intoxication, severe sleep deprivation and Alzheimer's disease. Furthermore, patients with any neurological or neuromuscular disorders (different or not related to CP) diagnosed by the PMD were excluded.

### Interventions

Eligible subjects who consented to participate were randomly allocated to receive 1 of the 2 proposed treatments. The first group was named group A and received a non-surgical non-invasive MNNM procedure. MNNM was applied con-

tinuously for 2 min on 5 different occasions with 1 min of rest between each 2-minute application of the MNNM procedure by a PT with at least 2 years of experience in manual therapy. The intervention was applied during a period of 6 weeks (from Monday to Friday). The maximum level of elbow extension movement without the reproduction of symptoms during the application of the MNNM treatment was determined through the baseline use of a universal hand held goniometer device. The MNNM intervention was applied according to the previously described principles of neural tissue mobilization for treating CP by Butler, Butler and Coppieters, Coppieters and Butler, and Elvey-Hall [14, 26–30].

The MNNM was performed by PT in the affected upper limb and consisted of initial supine neutral positioning of the patient, shoulder girdle depression, glenohumeral 90° abduction with a lateral rotation component, supination of the forearm, elbow and wrist flexion, thumb and finger extension followed by an immediate second movement of elbow extension with wrist, thumb, and finger flexion, while maintaining the initial shoulder girdle, glenohumeral and forearm positioning [12]. This passive movement sequence of upper limb flexion and extension was done repetitively at an approximate speed of 1 complete repetition of upper limb flexion and extension movement every 2 s without the reproduction of symptoms.

The second group of individuals was named group B and received an oral ibuprofen tablet treatment that was prescribed by the PMD who is familiar with the proper use and side effects of OI in CP. The PMD was responsible for modulating the OI dose to the subject's tolerance whilst trying to achieve the maximum hypoalgesic desired effect. The starting dose on the first day of treatment was a single dose of 400 mg. Doses were increased in case of subjects' adequate tolerance to a maximum dose of 1200 mg/day, divided into 3 doses every 8 h. Patients could leave the present study when they wanted to or if the PMD considered that symptoms worsened significantly; therefore at all times the free will of participants to continue or abandon the experiment was fully respected.

### Outcome measurements

Both groups, MNNM and OI, were assessed in the same time frames. The primary outcome measure was change from baseline using the numeric rating scale for pain (NRSP). The time frame was at baseline and 1 h after the application of treatment, corresponding to intervention sessions 1, 15 and 30 (baseline, 4 weeks and 6 weeks, respectively). The NRSP is an 11-point scale for patient self-reporting of pain [49–51]. It was employed to

evaluate the presence and relief of cervicobrachial pain symptoms by a blinded PT outcomes assessor (PTOA). Secondary outcomes were related to individual physical function; the first secondary outcome measure was change from baseline of the physical function involving the affected upper limb using the QuickDASH (Disabilities of the Arm, Shoulder and Hand) scale. The time frame corresponded to intervention sessions 1 and 30 (baseline and 6 weeks, respectively). The QuickDASH test is a self-report short questionnaire designed to measure physical function and symptoms in people with any of several musculoskeletal disorders of the upper limb [52, 53]. The last secondary outcome measure was the change from baseline cervical rotation range of motion at 1 h. This outcome measure time frame was at baseline and 1 h after the application of treatment, corresponding to intervention sessions 1 and 30 (baseline and 6 weeks, respectively). Cervical rotation was assessed in units of rotation degrees, using a cervical range-of-motion device (CROM) [54, 55]. All secondary outcomes were assessed by the PTOA. No changes were made in the proposed outcomes because there was no deviation from the original protocol.

### Sample size calculation

G\*Power software 3.1.9.2 was used to set the sample size. For sample size determination the NRSP was used as the primary reference measurement. The effect size for the NRSP was estimated as 0.509. Assuming at least six measurements,

correction of sphericity was determined as 0.5 with 0.95 power and a 0.05  $\alpha$  level; 20% of the sample was considered lost. Consequently a total sample size of 54 subjects was estimated to be randomly divided into two groups of 27 subjects. Criteria for participant removal were the presence of any exclusion criterion at the time of measurement or diagnosis and/or during the study period or monitoring process, failure to attend 2 sessions of treatment or evaluation, voluntary abandonment, death or removal by decision of the present study holders.


### Statistical analysis

Demographic characteristics including age and gender of participants were analyzed. The Kolmogorov-Smirnov test was used to confirm a normal distribution of the sample. NRSP, CROM and QuickDASH mean, between-session variance percentages and confidence interval according to each session and type of treatment were performed. Student's *t* test was employed for between-group mean difference comparisons as well as for intragroup analyses. In addition, eta square values ( $\eta^2$ ) of subjects' intereffect testing were added in both treatment groups. Box-plot graphics were developed in order to clarify the data. Finally, all data analyses were performed with commercially available software (SPSS 22.0, SPSS Inc, Chicago, IL) and a confidence interval (CI) of 95% ( $p$ -value < 0.05).

### Results

Of the original 62 participants enrolled in the present study only 50 completed the trial. Of the 31 subjects who belonged to the MNNM group (group A) 24 subjects completed the trial: 6 subjects abandoned participation claiming personal reasons and 1 participant was removed by the study holders because of anticonvulsant intake concealment. In the OI group (group B) 5 participants of the original 31 abandoned the study; 2 participants were removed from the study because they missed evaluation sessions and a total of 3 participants were removed by the PMD because they showed mild signs of gastric side effects associated with OI. No important side effects were reported related to MNNM treatment (Figure 1).

There were no significant differences related to the size, age and sex of both groups' participants; therefore demographic values did not influence the primary and secondary outcomes of the present study (Table I). NRSP results of the MNNM group show significantly ( $p < 0.05$ ) inferior mean values to those obtained by the OI group in all of the performed assessments. The MNNM



**Figure 1.** Participant flow diagram of voluntary participants through different study phases

*n* – number of subjects, MNNM – median nerve neural mobilization, OI – oral ibuprofen.

group had a constant decrease of NRSP scores that ranged between 6.9% and 15%, whereas the OI group did not exhibit a constant decrease in its NRSP values. It was observed that OI NRSP scores decreased dramatically 1 h after the drug was administered but OI NRSP scores had a tendency to increase back again at the baseline measure of the next session.

Because of this series of results we can clearly state that according to the applied protocol OI is more effective in reducing pain in terms of mean values than MNNM even though a tendency of constant pain reduction was observed in MNNM. The results of this investigation reveal that both MNNM and OI proved to be effective in treating CP. All ranges calculated for the confidence interval were contained within the lower and upper limits (between 1.13 and 1.45), implying that 95% differences in similar investigations would not exceed a higher value than 2 (Table II).

CROM outcome values showed an increase in the ipsilateral rotation range of cervical mo-

**Table I.** Demographic data of participants according to type of treatment

Demographic data	MNNM (n = 24)	OI (n = 26)	Statistical significance
Age [years]	32.3 ±3.7	30.8 ±4.3	$p < 0.195$
Sex:			
Female	13 (54)	19 (73)	$p < 0.27$
Male	11 (46)	7 (27)	

\* $p < 0.05$  was considered as statistically significant. Age – age in years of participants, Sex – biological sex of participants, MNNM – median nerve neural mobilization, n – number of participants, f – frequency, OI – oral ibuprofen.

tion (IRRCM) in both treatment groups, but did not show a significant ( $p > 0.05$ ) difference in between-group analysis. CROM device results reveal that MNNM and OI were equally effective in increasing IRRCM (Table III). Responses to the QuickDASH instrument in each study group expressed highly significant values ( $p < 0.0004$  and

**Table II.** NRSP mean, between-session variance percentages and confidence interval according to each session and type of treatment

Session	MNNM (n = 24)	%	OI (n = 26)	%	MNNM – OI difference 95% CI (range IL – SL)
One	6.5 (0.9)	–	5.9 (1.0)	–	0.6 ( $p < 0.033$ )* 0.05–1.18 (1.13)
+ 1 h	5.9 (1.1)	–9.2	3.7 (1.4)	–37.3	2.2 ( $p < 0.0001$ )* 1.45–2.90 (1.45)
Two	4.9 (1.4)	–6.9	3.9 (0.9)	+5.4	1.0 ( $p < 0.004$ )* 0.36–1.77 (1.41)
+ 1 h	4.5 (1.4)	–8.2	2.1 (0.9)	–46.2	2.4 ( $p < 0.0001$ )* 1.75–3.17 (1.42)
Three	3.8 (1.3)	–15.5	2.9 (0.8)	+38.1	0.9 ( $p < 0.009$ )* 0.22–1.50 (1.28)
+ 1 h	3.5 (1.4)	–8.5	1.7 (0.7)	–41.2	1.8 ( $p < 0.0001$ )* 1.12–2.42 (1.30)

\* $p < 0.05$  was considered as statistically significant. CI – 95% confidence interval, MNNM – median nerve neural mobilization, NRSP – numeric rating scale for pain, Session – time lapse when treatment was applied, OI – oral ibuprofen, One – time lapse corresponding to baseline assessment, (range IL – SL) – range lower limit minus range upper limit, +1 h – time lapse corresponding to 1 h after baseline application of treatment assessment, n – number of participants, % – between-session variance percentage.

**Table III.** CROM device mean, between-session variance percentages and confidence interval according to each session and type of treatment

Session	MNNM (n = 24)	%	OI (n = 26)	%	MNNM – OI difference 95% CI (range IL – SL)
One	60.4 (7.0)	–	57.7 (5.6)	–	2.7 ( $p < 0.143$ ) –0.94 – 6.31 (7.25)
+ 1 h	63.8 (7.7)	5.6	66.0 (7.4)	14.3	–2.2 ( $p < 0.317$ ) –6.47 – 2.14 (8.61)
Two	65.0 (5.9)	1.8	66.1 (9.0)	1.5	–1.1 ( $p < 0.489$ ) –5.94 – 2.88 (8.82)
+ 1 h	69.3 (7.4)	6.6	71.1 (4.5)	7.5	–0.84 ( $p < 0.160$ ) –6.07 – 1.03 (7.10)

\* $p < 0.05$  was considered as statistically significant. CI – 95% confidence interval, CROM – cervical range of motion, MNNM – median nerve neural mobilization, Session – time lapse when treatment was applied, OI – oral ibuprofen, One – time lapse corresponding to baseline assessment, (range IL – SL) – range lower limit minus range upper limit, +1 h – time lapse corresponding to 1 h after baseline application of treatment assessment, n – number of participants, % – between-session variance percentage.



**Table IV.** QuickDASH mean, between-session variance percentages and confidence interval according to each session and type of treatment

Session	MNNM (n = 24)	%	OI (n = 26)	%	MNNM – OI difference 95% CI (range IL – SL)
One	60.8 (10.0)	–	52.2 (10.2)	–	8.6 ( $p < 0.004$ )* 2.85–14.43 (11.58)
Two	32.2 (12.6)	47.0	17.8 (7.6)	65.9	14.4 ( $p < 0.0001$ )* 8.32–20.39 (12.07)

\* $p < 0.05$  was considered as statistically significant. CI – 95% confidence interval, DASH – The Disabilities of the Arm, Shoulder and Hand questionnaire, MNNM – median nerve neural mobilization, Session – time lapse when treatment was applied, OI – oral ibuprofen, One – time lapse corresponding to baseline assessment, (range IL – SL) – range lower limit minus range upper limit, n – number of participants, % – between-session variance percentage.

$p < 0.0001$ ) in each of the QuickDASH assessment sessions. The improvement was greater in the OI treatment group, with a difference of 14.4 points and a 65.9% decrease compared to session one, while the MNNM group showed a 47.0% QuickDASH score decrease. Both treatments are capable of improving the physical condition of subjects who suffer CP according to QuickDASH instrument criteria (Table IV).

Eta squared values, expressed in percentages, show that the variations observed in the lower average values of the NRSP scale and increased degrees of CROM were the result of the applied treatments. In this regard, MNNM group average value explained variation of NRSP was 75.5%, whereas in the OI group the percentage was 81.6%; percentages for the CROM device were 61.2% and 82.1% respectively. These percentages reflect the percentage of variance in NRSP and CROM that was explained by the administration of treatments for each group (Table V).

## Discussion

Therefore MNNM was not superior or equal to the OI treatment in reducing pain perception in subjects who suffer CP according to the applied

protocol. Effectiveness values regarding NRSP and QuickDASH of MNNM proved to be significantly inferior ( $p < 0.05$ ) than values obtained through the application of an OI treatment in achieving analgesia and improving function of the affected upper limb during CP.

The MNNM and OI outcomes of this study are believed to be of clinical importance according to Farrar *et al.* [56] and are in line with the pain reduction values (> 2 points) achieved by Cleland *et al.*, Allison *et al.* and Savva *et al.* [57–59]. Because of the novel nature of the applied protocol there was no available literature which could be used to compare specific parameters. All the available studies that applied MNNM had a mixed combination of MNNM with other types of treatments in the same group of subjects. The exclusive application and assessment of passive MNNM performed in this study may explain the smaller pain reduction effect of MNNM when compared to the results reported by De la Llave-Rincon *et al.*, Nee *et al.* and Savva *et al.* [12, 36, 59]. Pain reduction, function improvement and increase in IRRCM were continuous in the MNNM group mean while the OI group outcomes related to pain had a tendency to increase again after a time lapse. This event was linked to the extensively described pharmacokinetics and pharmacodynamics of OI [24].

CROM device outcomes revealed that both treatments applied in this investigation were significantly effective in increasing IRRCM in subjects who suffer CP, as seen in research by Cowell and Phillips [4]. Comparative analysis between the two treatment groups of the present study did not reveal a significant difference of effectiveness in increasing the IRRCM. This reveals that MNNM and OI are equally effective ( $p < 0.05$ ) in increasing IRRCM according to the applied protocol. There was no evidence of side effects or an important exacerbation of symptoms related to MNNM resulting from the application of the protocol [36].

A discrepancy of results was found related to OI effectiveness in CP [60]. According to Sheather-Reid and Cohen [20] OI did not exert any analgesic effect in a group of subjects who suffered

**Table V.** Eta square values ( $\eta^2$ ) of subjects intereffect testing\*

Instruments and treatments	$\eta^2$	%
NRSP:		
MNNM	0.755	75.5
OI	0.816	81.6
CROM device:		
MNNM	0.612	61.2
OI	0.821	82.1

\*NRSP scores and CROM device degrees results of MNNM and OI treatment sessions processed through  $\eta^2$  ( $\eta^2$ ) subject intereffect testing related to repeated measure analysis of the general linear model. CROM – cervical range of motion, NRSP – numeric rating scale for pain, MNNM – median nerve neural mobilization, OI – oral ibuprofen, % –  $\eta^2$  percentage value possibly related to treatment effect.

CP [60–63]. It is important to state that although OI is recommended worldwide to treat CP, the research performed by Sheather-Reid and Cohen [20] in 1998 constitutes the only available study of OI effectiveness in CP. We believe that the reason for the discrepancy of results lies in the fact that the OI dose used to treat subjects in the Sheather-Reid and Cohen [20] investigation was only 800 mg/day, which, according to Rainsford and Portenoy and Kanner [35], is considered an inferior dosage to the recommended 1200 mg/day dose to properly achieve analgesia and reduce inflammation. This explanation is supported by a study performed by McQuay and Moore [63] which concludes that OI is a dose-dependent drug that may vary in its pain reduction effect according to the size of the dose administered to the subject. Another possible reason for the discrepancy of the results with the mentioned Sheather-Reid and Cohen [20] study was the small sample size (only 4 subjects completed the trial) and the inclusion criteria for participants.

The beneficial results of OI and MNNM obtained in the present study support the theory suggested by Allison *et al.* [58] that there is an inextricable link between articular, soft tissue and neural structures of the cervicobrachial anatomic region. The authors of the present study believe that CP is a multimodal condition where joint and neurogenic pain may coexist and overlap as proposed by Elvery and Hall [30]. This strong interrelation among joint and nervous tissue could explain the dual effectiveness of both OI and MNNM treatments in CP. We believe that OI exerted a systemic anti-inflammatory action on joints and soft tissue that could have reached a ceiling effect because the neural components were not specifically treated, whereas the MNNM treatment had a direct impact on CP by reducing peripheral and central neurogenic pain, which is widely known to be a source of local inflammation and nociception. This conjecture is consistent with the histological and Western blot findings of Santos *et al.* [64] which suggested that neurogenic and peripheral inflammation activated through dorsal horn sensitizing of glial cells is reversible by the application of neural mobilization and anti-inflammatory treatment.

A constant limitation faced during the present study was the lack of prior high quality research related to MNNM and OI in CP. Other limitations and potential sources of bias were found in the primary outcome and the participant blinding method. We believe that the NRSP results of this study cannot be used to determine whether pain was of neurogenic or musculoskeletal inflammatory origin. Nevertheless, special care was taken in the participants' inclusion criteria to ensure that pain of neural origin was also present. Fur-

thermore, the time frames of the outcome measurements may have been influenced by the drug half-life and confounded the results between the groups. Blinding of participants was an additional challenge; in this regard, the study hypothesis, control group existence and group allocation blinding could have been the method of choice for subject masking, which is considered according to Boutron *et al.* [42] an effective blinding procedure. Furthermore, other conditions such as arthrogryposis [65] or coronary heart disease [66], which may influence CP, were not excluded. New approaches testing the feasibility and safety in orthopedic rehabilitation should be considered in future studies [67]. Due to the lack of dipper subgroup analysis and additional reproductions of the applied protocol, the authors considered that generalization of our study results was not possible.

In conclusion, OI may be superior to MNNM for pain reduction and upper limb function increase of subjects with CP according to the applied protocol in the present investigation. Nevertheless, it is important to acknowledge that both treatments may be effective in treating CP. MNNM may be considered an effective non-pharmaceutical treatment option in specific cases of CP.

#### Conflict of interest

The authors declare no conflict of interest.

#### References

1. Huisstede BM, Bierma-Zeinstra SM, Koes BW, Verhaar JA. Incidence and prevalence of upper-extremity musculoskeletal disorders. A systematic appraisal of the literature. *BMC Musculoskel Disord* 2006; 7: 7.
2. Radhakrishnan K, Litchy WJ, O'Fallon WM, Kurland LT. Epidemiology of cervical radiculopathy. A population-based study from Rochester, Minnesota, 1976 through 1990. *Brain* 1994; 117: 325-35.
3. Daffner SD, Hilibrand AS, Hanscom BS, Brislin BT, Vaccaro AR, Albert TJ. Impact of neck and arm pain on overall health status. *Spine (Phila Pa 1976)* 2003; 28: 2030-5.
4. Cowell IM, Phillips DR. Effectiveness of manipulative physiotherapy for the treatment of a neurogenic cervicobrachial pain syndrome: a single case study – experimental design. *Manip Ther* 2002; 7: 31-8.
5. Salt E, Wright C, Kelly S, Dean A. A systematic literature review on the effectiveness of non-invasive therapy for cervicobrachial pain. *Manip Ther* 2011; 16: 53-65.
6. Eubanks JD. Cervical radiculopathy: nonoperative management of neck pain and radicular symptoms. *Am Fam Phys* 2010; 81: 33-40.
7. Peolsson A, Söderlund A, Engquist M. Physical function outcome in cervical radiculopathy patients after physiotherapy alone compared with anterior surgery followed by physiotherapy: a prospective randomized study with a 2-year follow-up. *Spine (Phila Pa 1976)* 2013; 38: 300-7.
8. Caridi JM, Pumberger M, Hughes AP. Cervical radiculopathy: a review. *HSS J* 2011; 7: 265-72.
9. Dillingham TR. Evaluating the patient with suspected radiculopathy. *Phys Med Rehabil* 2003; 5: 41-9.

10. Boyles R, Toy P, Mellon J, Hayes M, Hammer B. Effectiveness of manual physical therapy in the treatment of cervical radiculopathy: a systematic review. *J Man Manip Ther* 2011; 19: 135-42.
11. Young IA, Michener LA, Cleland JA, Aguilera AJ, Snyder AR. Manual therapy, exercise, and traction for patients with cervical radiculopathy: a randomized clinical trial. *Phys Ther* 2009; 89: 632-42.
12. De-la-Llave-Rincón A, Puentedura EJ, Fernández-de-Las-Peñas C. Clinical presentation and manual therapy for upper quadrant musculoskeletal conditions. *J Man Manip Ther* 2011; 19: 201-11.
13. Asghar W, Jamali F. The effect of COX-2-selective meloxicam on the myocardial, vascular and renal risks: a systematic review. *Inflammopharmacology* 2015; 23: 1-16.
14. Clark BD, Barr AE, Safadi FF. Median nerve trauma in a rat model of work-related musculoskeletal disorder. *J Neurotrauma* 2003; 20: 681-95.
15. Bove GM. Epi-perineurial anatomy innervation, and axonal nociceptive mechanisms. *J Bodyw Mov Ther* 2008; 12: 185-90.
16. Elvery R. Brachial plexus tension test and the pathoanatomical origin of the arm pain. In: *Aspects of Manipulative Therapy*. Idezak R (ed.). Lincoln Institute of Health Science, Melbourne 1979; 105-10.
17. Elliot MB, Barr AE, Kietrys DM, AL-Shatti T, Amin M, Barbe MF. Peripheral neuritis and increased spinal cord neurochemicals are induced in a model of repetitive motion injury with low force and repetition exposure. *Brain Res J* 2008; 1218: 103-13.
18. Dabbs V, Lauretti WJ. A risk assessment of cervical manipulation vs. NSAIDs for the treatment of neck pain. *J Manip Physiol Therap* 1995; 18: 530-6.
19. Van der Velde G, Hogg-Johnson S, Bayoumi AM. Identifying the best treatment among common nonsurgical neck pain treatments: a decision analysis. *J Manip Phys Ther* 2009; 32: 209-18.
20. Sheather-Reid RB, Cohen M. Efficacy of analgesics in chronic pain: a series of N-of-1 studies. *J Pain Symptom Manage* 1998; 15: 244-52.
21. Nemes D, Amaricai E, Tanase D, Popa D, Catan L, Andrei D. Physical therapy vs. medical treatment of musculoskeletal disorders in dentistry: a randomized prospective study. *Ann Agricult Environm Med* 2013; 20: 301-6.
22. Borghi B, Aurini L, White PF. Treatment of chronic cervicalbrachial pain with periradicular injection of meloxicam. *Minerva Anesthesiol* 2015 Sep 3 [Epub ahead of print].
23. Cohen ML, Arroyo JF, Champion GD, Browne CD. In search of the pathogenesis of refractory cervicobrachial pain syndrome. A deconstruction of the RSI phenomenon. *Med J Aust* 1992; 156: 432-6.
24. Day RO. Variability in response to NSAIDs: what progress? *Agents Actions Suppl* 1993; 44: 3-6.
25. Maguire TA. Ibuprofen: a model medicine for self-care of common conditions. *Int J Clin Pract Suppl* 2013; 178: 43-6.
26. Butler D. *Mobilisation of the Nervous System*. 1<sup>st</sup> ed. Churchill Livingstone, Melbourne 1991.
27. Butler D. *The Sensitive Nervous System*. 1<sup>st</sup> ed. Noigroup Publication, Adelaide, Australia 2000.
28. Butler D, Coppeters MW. Neurodynamics in a broader perspective. *Man Ther* 2007; 12: 7-8.
29. Coppeters MW, Butler DS. Do 'sliders' slide and 'tensioners' tension? An analysis of neurodynamic techniques and considerations regarding their application. *Manip Ther* 2008; 13: 213-21.
30. Elvery RL, Hall TM. *Neural tissue evaluation and treatment. Physical Therapy of the Shoulder*. Churchill Livingstone, New York 1997.
31. Allison GT, Nagy BM, Hall TA. Randomized clinical trial of manual therapy for cervico-brachial pain syndrome: a pilot study. *Manual Therapy* 2002; 7: 95-102.
32. Beneciuk JM, Bishop MD, George SZ. Effects of upper extremity neural mobilization on thermal pain sensitivity: a sham-controlled study in asymptomatic participants. *J Orthop Sports Phys Ther* 2009; 39: 428-38.
33. Coppeters MW, Stappaerts KH, Wouters LL, Janssens K. The immediate effects of a cervical lateral glide treatment technique in patients with neurogenic cervicobrachial pain. *J Orthop Sports Phys Ther* 2003; 33: 369-78.
34. Kuchari E, Han S, Karłowicz-Bodalska K, Miśkiewicz K, Kutyccka E. Safety of oral ibuprofen: analysis of data from the spontaneous reporting system in Poland. *Acta Poloniae Pharmaceut* 2014; 71: 687-90.
35. Rainsford KD. Ibuprofen: from invention to an OTC therapeutic mainstay. *Int J Clin Pract Suppl* 2013; 178: 9-20.
36. Nee RJ, Vicenzino B, Jull GA, Cleland JA, Coppeters MW. Neural tissue management provides immediate clinically relevant benefits without harmful effects for patients with nerve related neck and arm pain: a randomized trial. *J Physiother* 2012; 58: 23-31.
37. Borenstein D, Wiesel S, Boden S. *Low Back and Neck Pain*. 3<sup>rd</sup> ed. Saunders, Philadelphia 2004; 771-849.
38. Sniezek PJ, Brodland DG, Zitelli JA. A randomized controlled trial comparing acetaminophen, acetaminophen and ibuprofen, and acetaminophen and codeine for postoperative pain relief after Mohs surgery and cutaneous reconstruction. *Dermatol Surg J* 2011; 37: 1007-13.
39. Carlesso LC, Gross AR, Santaguida PL, Burnie S, Voth S, Sadi J. Adverse events associated with the use of cervical manipulation and mobilization for the treatment of neck pain in adults: a systematic review. *Man Ther* 2010; 15: 434-44.
40. Portenoy RK, Kanner RM. Nonopioid and adjuvant analgesics. In: *Pain Management: Theory and Practice*. Portenoy RK, Kanner RM (eds.). FD Davis Company, Philadelphia 1996; 219, 247.
41. Carlesso LC, Macdermid JC, Gross AR, Walton DM, Santaguida PL. Treatment preferences amongst physical therapists and chiropractors for the management of neck pain: results of an international survey. *Chiropr Man Therap* 2014; 22: 11.
42. Boutron I, Dutton S, Ravaud P, Altman D. Reporting and interpretation of randomized controlled trials with statistically nonsignificant results for primary outcomes. *J Am Med Assoc* 2010; 303: 2058-64.
43. Bono CM, Ghiselli G, Gilbert TJ. North American Spine Society. An evidence-based guideline for the diagnosis and treatment of cervical radiculopathy from degenerative disorders. *Spine J* 2009; 11: 64-72.
44. American Cancer Society. (2015). *Complementary and Alternative Methods and Cancer* <http://www.cancer.org/acs/groups/cid/documents/webcontent/acspsc-041660-pdf.pdf>.
45. Cassileth BR, Deng GE, Gomez JE, Johnstone PA, Kumar N, Vickers AJ. American College of Chest Physicians. Complementary therapies and integrative oncology in lung cancer: ACCP evidence-based clinical practice guidelines (2nd edition). *Chest* 2007; 132: 340S-54S.
46. Williams AM, Davies A, Griffiths G. Facilitating comfort for hospitalized patients using non-pharmacological measures: preliminary development of clinical practice guidelines. *Int J Nurs Pract* 2009; 15: 145-55.



47. Weiner DK, Ernst E. Complementary and alternative approaches to the treatment of persistent musculoskeletal pain. *Clin J Pain* 2004; 20: 244-55.
48. Schulz KF, Altman DG, Moher D; the CONSORT Group. CONSORT 2010 Statement: updated guidelines for reporting parallel group randomised trials. *Ann Intern Med* 2010; 152: 726-32.
49. Hjermstad MJ, Fayers PM, Haugen DF. European Palliative Care Research Collaborative (EPCRC). Studies comparing Numerical Rating Scales, Verbal Rating Scales, and Visual Analogue Scales for assessment of pain intensity in adults: a systematic literature review. *J Pain Symptom Manag* 2011; 41: 1073-93.
50. Bahreini M, Jalili M, Moradi-Lakeh MA. Comparison of three self-report pain scales in adults with acute pain. *J Emerg Med* 2015; 48: 10-8.
51. Price DD, Bush FM, Long S, Harkins SW. A comparison of pain measurement characteristics of mechanical visual analogue and simple numerical rating scales. *Pain* 1994; 56: 217-26.
52. Angst F, Schwyzer HK, Aeschlimann A, Simmen BR, Goldhahn J. Measures of adult shoulder function: Disabilities of the Arm, Shoulder, and Hand Questionnaire (DASH) and its short version (QuickDASH), Shoulder Pain and Disability Index (SPADI), American Shoulder and Elbow Surgeons (ASES) Society standardized shoulder assessment form, Constant (Murley) Score (CS), Simple Shoulder Test (SST), Oxford Shoulder Score (OSS), Shoulder Disability Questionnaire (SDQ), and Western Ontario Shoulder Instability Index (WOSI). *Arthritis Care Res (Hoboken)* 2011; 63: 174-88.
53. Dowrick AS, Gabbe BJ, Williamson OD, Cameron PA. Does the disabilities of the arm, shoulder and hand (DASH) scoring system only measure disability due to injuries to the upper limb? *J Bone Joint Surg* 2006; 88: 524-7.
54. Audette I, Dumas JP, Côté JN, De Serres SJ. Validity and between-day reliability of the cervical range of motion (CROM) device. *J Orthop Sports Phys Ther* 2010; 40: 318-23.
55. Fletcher JP, Bandy WD. Intrarater reliability of CROM measurement of cervical spine active range of motion in persons with and without neck pain. *J Orthop Sports Phys Ther* 2008; 38: 640-5.
56. Farrar JT, Young JP, LaMoreaux L, Werth JL, Poole RM. Clinical importance of changes in chronic pain intensity measured on an 11-point numerical pain rating scale. *Pain* 2001; 94: 149-58.
57. Cleland JA, Whitman JM, Fritz JM, Palmer JA. Manual physical therapy, cervical traction, and strengthening exercises in patients with cervical radiculopathy: a case series. *J Orthop Sports Phys Ther* 2005; 35: 802-11.
58. Imani F, Motavaf M, Safari S, Alavian SM. The therapeutic use of analgesics in patients with liver cirrhosis: a literature review and evidence based recommendations. *Hepat Mon* 2014; 14: e23539.
59. Savva C, Giakas G. The effect of cervical traction combined with neural mobilization on pain and disability in cervical radiculopathy. A case report. *Manip Ther* 2013; 18: 443-6.
60. U.S. Food and Drug Administration (FDA). (April 01, 2015). FDA Meeting Topic: NDAC Meeting on Risks of NSAIDs. [http://www.fda.gov/ohrms/dockets/ac/02/briefing/3882B2\\_04\\_Wyeth-Ibuprophen.pdf](http://www.fda.gov/ohrms/dockets/ac/02/briefing/3882B2_04_Wyeth-Ibuprophen.pdf). On March 03, 2016.
61. Unzueta A, Vargas HE. Nonsteroidal anti-inflammatory drug-induced hepatotoxicity. *Clin Liver Dis* 2013; 17: 43-56.
62. Youdas JW, Carey JR, Garrett TR. Reliability of measurements of cervical spine range of motion: comparison of three methods. *Phys Ther* 1991; 71: 98-104.
63. McQuay HJ, Moore RA. Dose-response in direct comparisons of different doses of aspirin, ibuprofen and paracetamol (acetaminophen) in analgesic studies. *Br J Clin Pharmacol* 2007; 63: 271-8.
64. Santos FM, Silva JT, Giardini AC. Neural mobilization reverses behavioral and cellular changes that characterize neuropathic pain in rats. *Molecul Pain* 2012; 8: 57.
65. Kowalczyk B, Feluś J. Arthrogyrosis: an update on clinical aspects, etiology, and treatment strategies. *Arch Med Sci* 2016; 12: 10-24.
66. Kałka D, Karpiński Ł, Gebala J, et al. Sexual health of male cardiac patients – present status and expectations of patients with coronary heart disease. *Arch Med Sci* 2017; 13: 302-10.
67. Ficklscherer A, Stapf J, Meissner KM, et al. Testing the feasibility and safety of the Nintendo Wii gaming console in orthopedic rehabilitation: a pilot randomized controlled study. *Arch Med Sci* 2016; 12: 1273-8.