



CircRNA expression profiles and circRNA-miRNA-mRNA crosstalk in allergic rhinitis

Chang-Yu Qiu^{a,1}, Xin-Yan Cui^{a,1}, Mei-Ping Lu^a, Min Yin^{a,b}, Wan-Yun Xu^a, Xin-Jie Zhu^a, Qing Yang^a and Lei Cheng^{a,b*}

ABSTRACT

Background: Circular RNAs (circRNAs) are involved in inflammation; however, their role in allergic rhinitis (AR) remains unclear. In this study, we analyzed circRNA expression and identified a circRNA-miRNA-mRNA network through which circRNAs regulate AR pathogenesis.

Methods: We analyzed circRNA, miRNA, and mRNA expression profiles in the nasal mucosa by high-throughput sequencing (HTS), using a fold-change >1.5 and *p*-value < 0.05 to pinpoint significantly differentially expressed (DE) circRNAs, miRNAs, and mRNAs in AR. A DEcircRNA-DEmiRNA-DEmRNA crosstalk network was then constructed using bioinformatics and statistical analysis. Gene ontology and Kyoto encyclopedia of genes and genomes pathway analyses were performed to identify the biological terms enriched in the network; whereas RT-PCR and Sanger sequencing were used to confirm the circRNAs.

Results: A total of 264 DEcircRNAs were identified by HTS, including 120 upregulated and 144 downregulated in AR compared to controls. A DEcircRNA-DEmiRNA-DEmRNA crosstalk network was constructed with 17 miRNAs, 11 circRNAs, 29 mRNAs, and 64 interaction pairs. These genes were involved in the Wnt signaling pathway, TNF biosynthesis, inflammatory responses, the PI3K-Akt signaling pathway, and Toll-like receptors. Of the 11 DEcircRNAs, hsa_circ_0008668 and circTRIQQ were upregulated, whereas hsa_circ_0029853 and circRNA_01002 were downregulated in AR tissues. Sanger sequencing confirmed the back-splicing junctions of these circRNAs.

Conclusions: We constructed a novel DEcircRNA-DEmiRNA-DEmRNA network for AR that provides a basis for future studies to investigate its underlying molecular mechanisms.

Keywords: Allergic rhinitis, Circular RNA, microRNA, mRNA, Nasal mucosa

^aDepartment of Otorhinolaryngology & Clinical Allergy Center, The First Affiliated Hospital, Nanjing Medical University, Nanjing, China
^{*}Corresponding author. Department of Otorhinolaryngology & Clinical Allergy Center, The First Affiliated Hospital, Nanjing Medical University, 300 Guangzhou Road, Nanjing, 210029, China E-mails: chenglei@jsph.org.cn; jspent@126.com

¹ Contributed equally to the study.

Full list of author information is available at the end of the article <https://doi.org/10.1016/j.waojou.2021.1000548>

Received 26 January 2021; Received in revised form 22 April 2021; Accepted 26 April 2021

1939-4551/© 2021 The Author(s). Published by Elsevier Inc. on behalf of World Allergy Organization. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

INTRODUCTION

Allergic rhinitis (AR) is a chronic multi-factorial inflammatory disease, with a prevalence rate of 17.6% in adults that is expected to increase yearly in China.^{1,2} Despite the discovery of some genetic factors responsible for AR progression, the underlying molecular mechanisms have not yet been fully elucidated.

Previous studies have demonstrated that microRNAs (miRNAs) and long non-coding RNAs (lncRNAs) play critical roles in AR^{3,4} and circular RNAs (circRNAs), a newly-discovered family of endogenous non-coding RNAs, have been found to exert a variety of functions.⁵ For instance, large-scale studies have confirmed that circRNAs are involved in neuronal function, tumor development, and innate immune responses,⁶ while others are associated with inflammatory diseases.^{7,8} Fang et al⁹ have reported that circANKRD36 expression positively correlates with the level of inflammatory factor IL-6 and may be involved in type 2 diabetes mellitus and inflammation-related pathways. Moreover, Liu et al¹⁰ have found circRNA Ddx17 can exert protective effects by inhibiting miR-17-5p expression in ovalbumin-induced AR mice. Similarly, Zhu et al¹¹ have found circHIPK3 is highly expressed in AR mucosa, and that circHIPK3 can upregulate Th2 cell-specific transcription factor GATA-3 via modulating miR-495. Wang et al¹² have found circARRDC3 promotes the development of AR by regulating the miR-375/KLF4 axis.

Competing endogenous RNA (ceRNA) networks are able to reveal previously unknown functions of the human genome in diseases.¹³ For instance, circRNAs have been shown to regulate mRNAs by sponging miRNA, ie, competitively binding to miRNAs via common miRNA response elements (MREs).¹⁴ Zhang et al⁸ have constructed a network of differentially expressed (DE) circRNAs, miRNAs, and mRNAs, with gene enrichment analysis revealing that they regulate cell adhesion, cell activation, and immune responses in atherosclerosis. CircRNA-related ceRNA networks have also been established for illustrating the pathogenesis of various diseases, including liver cancer,¹⁵ glioma,¹⁶ and osteoarthritis.¹⁷ However, no such networks have been elucidated for AR.

Therefore, in this study, we described circRNA, miRNA, and mRNA expression profiles in the nasal mucosa from patients, with and without AR, using high-throughput sequencing (HTS), and constructed a DEcircRNA-DEmiRNA-DEmRNA cross-talk network to illustrate the potential molecular mechanisms of AR progression.

METHODS

Clinical data and nasal mucosa tissues

The nasal mucosa tissues were collected from a cohort of 20 adult subjects (10 with perennial AR and 10 non-allergic controls) aged 18-53 years who received endoscopic septoplasty from August 2019 to January 2020. Tissue samples were immediately frozen and stored in liquid nitrogen, and then subjected to transcriptomic profiling analysis. AR was diagnosed according to the clinical guidelines.¹ Serum total IgE and allergen-specific IgE were measured by the Automated Immuno-Strip Analysis System (LG Life Sciences Ltd., Korea) in all the subjects. Patients with perennial AR enrolled were sensitized mainly by house dust mites, including *Dermatophagoides pteronyssinus* and/or *Dermatophagoides farinae* (Table S1). The controls had negative IgE test results in sera. All subjects had ceased treatment with glucocorticoids, antihistamines, antileukotrienes, and other immunomodulating drugs for at least 4 weeks prior to surgery. Subjects who had an immune deficiency, upper respiratory tract infection, chronic rhinosinusitis, nasal polyps, sinonasal neoplasms, asthma and other allergic diseases, and a history of nasal surgery or immunotherapy were excluded.

Total RNA isolation, library construction, and sequencing

Total RNA was extracted from nasal mucosa samples using TRIzol reagent (Life Technologies, CA, USA) and its quantity and quality were measured using a NanoDrop ND-1000 (Thermo Scientific, DE, USA). RNA integrity was assessed using an Agilent 2100 Bioanalyzer (Agilent Technologies, CA, USA) and RNAs with an integrity value of ≥ 7 were retained.

Six randomly selected samples (three AR and three controls) from the 20 subjects were used to construct circRNA, miRNA, and mRNA libraries,




Fig. 1 CircRNAs in the nasal mucosa. (A) Box plots showing that circRNA distribution intensities from all normalized datasets were similar between the six samples. (B) A total of 30,936 circRNAs were identified in the nasal mucosa, of which 12,376 were known and 18,560 were novel, by comparing the predicted results with the circBase database (<http://www.circbase.org/>). (C) Chromosomal circRNA expression distribution. (D) Spliced reads per million (RPM) circRNA distribution. circRNA, circular RNA

and the 6 samples and the remaining 14 samples were sent to reverse transcription-polymerase chain reaction (RT-PCR). Approximately 1 μ g of RNA was extracted from each sample, and ribosomal and linear RNAs were depleted from the samples using Ribo-zero (Illumina, CA, USA) and RNase R (Epicenter, CA, USA), respectively, according to the manufacturers' instructions. CircRNA libraries were constructed using a TruSeq Stranded Total RNA HT/LT Sample Prep Kit (Illumina) according to the manufacturer's instructions. Library quality was assessed using an Agilent 2100 system (Agilent Technologies) and the libraries were sequenced using an Illumina sequencing platform (HiSeqTM 2500).

RNA (1 μ g) from each of the 6 samples was used as input material for small RNA library generated using TreSeq Small RNA Sample Prep Kits (Illumina), according to the manufacturer's instructions. Library quality was assessed using an Agilent 2100 system and libraries were sequenced using an Illumina sequencing platform (HiSeqTM 2500). All sequencing procedures and analyses were performed by OEbiotech (Shanghai, China).

Identification of miRNA-target interactions

MiRNA-mRNA and miRNA-circRNA pairs were selected using miRanda, which predicts miRNA targets based on sequence complementarity, conserved target sites, and free formation energy. Next, we filtered out miRNAs with crosstalk with circRNAs and mRNAs and used the 3 types of RNAs to produce a circRNA-miRNA-mRNA network. To identify circRNA-mRNA pairs, we used a ceRNA score that evaluates the significance (p -value) of miRNAs shared by each circRNA and mRNA as the ratio of common miRNA MRE to circRNA-miRNA binding MRE, as follows:

$$p\text{-value} = \frac{\sum_{i=M4}^{\min(M2,M3)} \binom{M3}{M4} \binom{M1-M3}{M2-M4}}{\binom{M1}{M2}}$$

where M1 represents the quantity of all miRNAs, M2 represents the quantity of miRNAs regulating mRNAs, M3 represents the quantity of miRNAs with regulatory effects on the circRNA, and M4 represents the quantity of common miRNAs. p -values < 0.05 were considered statistically significant.




Fig. 2 CircRNA, miRNA and mRNA expression. Clustered heat map showing all differentially expressed circRNAs (A), miRNAs (B) and mRNAs (C), respectively (fold-change > 1.5, $p < 0.05$). Red and blue represent high and low relative expression, respectively. circRNA, circular RNA; miRNA, microRNA

Bioinformatics and data analysis

Gene ontology (GO) analysis was used to annotate gene function. Enrichment scores were calculated as the negative logarithm of the p -value and used to determine the statistical significance of GO term clusters targeted by DE genes. Kyoto encyclopedia of genes and genomes (KEGG) analysis was used to determine the biological pathways involving genes in the network, as quantified by the enrichment score.




Fig. 3 Top 300 predicted DE circRNA-miRNA relationships. Some circRNAs were annotated using circBase. Novel circRNAs were described in terms of their genome location. Green and red nodes represent circRNAs and miRNAs, respectively. circRNA, circular RNA; miRNA, microRNA; DE, differentially expressed

Sanger sequencing and RT-PCR

Sanger sequencing was performed by Tsingke Biotechnology (Nanjing, China) to confirm the back-splicing junctions of each circRNA. Sanger sequencing data that matched the HTS results and displayed a unimodal dissolution curve were retained. To ensure that each PCR amplification fragment contained junction sites, the designed circRNA primers were divergent, instead of the frequently used convergent primers (Table S2). The experiment was repeated 3 times for each sample from the 20 subjects (10 with perennial AR and 10 controls). The RT-PCR dissolution curve was unimodal, proving that each pair of specific reverse primers had good repeatability and a single product.

Statistical analysis

All statistical analyses were carried out using SPSS 20.0 (IBM SPSS, Chicago, IL, USA). Between-group differences were analyzed using Student's t -tests. CircRNA expression was displayed as a fold-change calculated using the $2^{-\Delta\Delta ct}$ method on the RT-PCR data. Data from at least 3 independent experiments were presented as the mean \pm standard deviation (SD). p -values < 0.05 were considered statistically significant.

RESULTS

CircRNA expression and characteristics in the nasal mucosa

Three pairs of nasal mucosa tissues from 10 patients with AR and 10 controls were randomly selected for HTS analysis. The clinical characteristics of the subjects are shown in Table S1. All

CircRNA	Chromosome position	Type	Gene symbol	Expression
hsa_circ_0008668	Chr22:46097926_46109908_+	intergenic	-	up
hsa_circ_0029853	Chr13:28256292_28281379_+	sense-overlapping	PAN3	down
circTRIQK	Chr8:92916929_92930707_-	sense-overlapping	TRIQK	up
circRNA_01002	Chr1:56429304_56430306_-	intergenic	-	down
circCDKAL1	Chr6:20781145_20805719_+	sense-overlapping	CDKAL1	up
circPCMT1	Chr6:149765691_149773169_+	sense-overlapping	PCMT1	up
circARHGEF33	Chr2:38895777_38921423_+	sense-overlapping	ARHGEF33	down
circNCOA1	Chr2:24554389_24593721_+	sense-overlapping	NCOA1	up
circRNA_15993	Chr2:90082996_90159914_+	intergenic	-	up
circRNA_07795	Chr13:40826506_40870597_-	intergenic	-	down
circRNA_07800	Chr13:40908159_40914105_-	intergenic	-	down

Table 1. Details of the 11 circRNAs in the DEcircRNA-DEmiRNA-DEmRNA network in AR AR, allergic rhinitis; circRNA, circular RNA; DE, differentially expressed; miRNA, microRNA

normalized sample datasets were found to display similar circRNA distribution densities (Fig. 1A). CIRI software¹⁸ was used to identify and annotate 30,936 circRNAs from the 2 groups, including 12 376 known and 18 560 novel (Fig. 1B), and widely distributed circRNA transcripts were found on all chromosomes (Fig. 1C). CircRNA transcript expression was quantified using the spliced reads per million (RPM) distribution density, revealing

that almost all circRNA transcripts were expressed at low levels (Fig. 1D).

Among the DEcircRNAs, 1119 were antisense and 29 335 were sense, including 28 694 exonic, 641 intronic, and 1119 intergenic circRNAs. In particular, 93% of the circRNAs were sense exons and most circRNA transcripts carried 1-7 exons, with a length of 200-400bp or >2000bp. Hierarchical clustering revealed DE circRNAs between samples (Fig. 2A) and HTS analysis identified 264 DEcircRNAs, including 120 upregulated and 144 downregulated in AR. Based on the number of possible miRNA-binding sites, structure, energy score and *p*-value, the top 300 relevant circRNA-miRNA pairs were screened and their network illustrated using Cytoscape software (Fig. 3).




Fig. 4 Depiction of the DEcircRNA-DEmRNA network. The network includes 11 circRNAs (green nodes) and 29 mRNAs (yellow nodes). CircRNAs were annotated using circBase. Novel circRNAs were described in terms of their genome location. circRNA, circular RNA; DE, differentially expressed

MiRNA and mRNA expression profiles in the nasal mucosa

Next, we used HTS to analyze the miRNA and mRNA expression profiles in the nasal mucosa tissues from the AR patients and controls. Of the 1784 miRNAs available, 13 were upregulated and 15 were downregulated. On the other hand, of the 2030 mRNAs, 93 were upregulated and 79 were downregulated. Hierarchical clustering revealed that these DEmiRNAs (Fig. 2B) and DEmRNAs

(Fig. 2C) were distinct between samples, confirming that miRNAs and mRNAs were DE in patients with AR, compared to controls.

DEcircRNA-DEmiRNA, DEmiRNA-DEmRNA, and DEcircRNA-DEmRNA interactions

DEmiRNA-DEcircRNA and DEmiRNA-DEmRNA interactions were detected using miRanda software, which defined 911 and 379 pairs, respectively. Next, we calculated the Pearson correlation coefficient (PCC) of DEmiRNA-DEcircRNA and DEmiRNA-DEmRNA pairs according to their expression values, with p -values < 0.05 and an absolute PCC of ≥ 0.7 (PCC ≥ 0.7 indicated positive correlation and PCC ≤ -0.7 indicated negative correlation). A total of 941 DEmiRNA-DEcircRNA and 865 DEmiRNA-DEmRNA pairs with negative expression correlation were screened out, while 116 DEmiRNA-DEcircRNA and 379 DEmiRNA-DEmRNA pairs were screened by combining miRanda and correlation analyses. Since circRNAs act as ceRNAs form RNAs, we constructed a DEcircRNA-DEmRNA network (Fig. 4) based on 11 and 29 positively correlated circRNAs and mRNAs, respectively.

DEcircRNA-DEmiRNA-DEmRNA network and functional annotation

Next, we identified miRNAs interacting with both circRNAs and mRNAs to construct a DEcircRNA-DEmiRNA-DEmRNA network, which

included 17 miRNAs, 11 circRNAs (Table 1), 29 mRNAs, and 64 interaction pairs (Fig. 5).

To determine the biological roles of the DEcircRNAs, we performed GO and KEGG analyses on the DEmRNAs in the DEcircRNA-DEmiRNA-DEmRNA crosstalk network. These genes converged in 159 terms, including 172 biological processes, 47 cell components, and 46 molecular functional components. The GO terms mainly involved the Wnt signaling pathway, lung epithelial cell development, and TNF biosynthesis (biological processes), endoplasmic reticulum lumen, flotillin complex, and intercellular canalculus (cell components), fibronectin binding, collagen binding, and calcium: sodium antiporter activity (molecular functions; Fig. 6A). In addition, these genes were enriched in the inflammatory response and various signaling pathways (such as cell surface receptor signaling pathways), indicating that circRNAs may participate in multiple pathways related to AR pathogenesis. KEGG pathways analysis revealed 24 significantly involved pathways, of which local adhesion, ECM receptor interaction, protein degradation and absorption, and PI3K-Akt signaling pathways were the most enriched (Fig. 6B). However, other pathways were related to AR, such as the Toll-like receptor (TLR) signaling pathway, suggesting that circRNAs may




Fig. 5 Depiction of the DEcircRNA-DEmiRNA-DEmRNA crosstalk network. The network includes 17 miRNAs (red nodes), 11 circRNAs (green nodes), and 29 mRNAs (yellow nodes). CircRNAs and miRNAs were annotated using circBase and miRbase, respectively. Novel circRNAs and miRNAs were described in terms of their genome location. circRNA, circular RNA; miRNA, microRNA; DE, differentially expressed

can also control the cellular levels of specific miRNAs,²³ the majority of which harbor highly conserved sequences and act in multiple cellular processes via post-transcriptional regulation. Thus, disturbed miRNA activity can be partly explained by circRNA involvement.²⁴ With the help of MREs, circRNAs, miRNAs and mRNAs participate in large-scale ceRNA crosstalk regulating post-transcriptional genes in various manners. Previous studies have found that Ddx17, circHIPK3 and circARRDC3 can function as ceRNAs to sponge miRNAs in AR.¹⁰⁻¹² However, each of these studies has just proved one circRNA related to immunity regulation. To the best of our knowledge, our study is the first to investigate circRNA expression profiles and ceRNA network in AR.

We predicted DEmiRNA-DEcircRNA interactions using miRanda and screened 116 pairs with negative expression relationships, including 17

miRNAs (such as hsa-miR-98-5p and hsa-miR-455-5p), while HTS revealed that hsa-miR-98-5p was significantly upregulated. Takuse et al²⁵ have found that hsa-miR-98-5p negatively correlates with peripheral IL-10⁺ cell count and IL-10. Moreover, an increasing number of studies have demonstrated imbalance between CD4⁺ T cells and B cells in peripheral blood plays a role in AR.²⁶ Moreover, studies have reported reduced numbers of regulatory T cells (Treg)²⁷ and regulatory B cells (Breg)^{28,29} in AR, which are major sources of IL-10, an important molecule in allergen tolerance.^{30,31} Hence, hsa-miR-98-5p may affect allergen tolerance in AR by negatively regulating Tregs, Bregs, and IL-10 secretion. Our DEmiRNA-DEcircRNA network revealed that hsa-miR-98-5p competed with several circRNAs, such as hsa_circ_0029853, suggesting that circRNAs may interact with hsa-miR-98-5p during AR immune tolerance.




Fig. 7 Sanger sequencing and RT-PCR of circRNAs. (A-D) Back-splice junction sequences for hsa_circ_0008668, hsa_circ_0029853, circTRIQQ, and circRNA_01002 were confirmed by Sanger sequencing, respectively. Arrows represent divergent primers binding to the genomic region of circRNAs. (E-H) RT-PCR results for circRNAs in the nasal mucosa of ten patients with AR and ten control subjects. Hsa_circ_0008668 (E) and circTRIQQ (G) were upregulated in AR, whereas hsa_circ_0029853 (F) and circRNA_01002 (H) were downregulated, consistent with the HTS data. Data represent the mean \pm SD of at least three independent experiments. * $p < 0.05$, ** $p < 0.01$, **** $p < 0.0001$. circRNA, circular RNA

Similarly, we identified 379 DE miRNA-DE mRNA pairs and many DE genes (such as DOCK8 and MUC19) that were significantly associated with AR. In particular, DOCK8 expression was decreased in the nasal mucosa of patients with AR. DOCK8 deficiency plays a key role in allergic inflammation, susceptibility to infection, autoimmunity, and malignant tumors^{32,33} as it exerts negative effects on the migration, function, and survival of immune cells as well as innate and adaptive immunity.³² For instance, DOCK8 participates in the migration of interstitial dendritic cells³⁴ and maintains protective immunity by repressing the function of ROR γ ^t ILCs cells.³⁵ DOCK8 has also been shown to regulate adaptive immune cells, while Janssen et al³³ have found that DOCK8 is related to Treg cell homeostasis and function and enhances peripheral B cell tolerance. In addition, DOCK8 modulates MyD88-dependent TLR signal transduction and activates the STAT3 transcription factor.³² By examining DE miRNA-DE mRNA relationships, we found that DOCK8 could bind to hsa-miR-548aq-3p, hsa-miR-582-5p and hsa-miR-96-5p, and that its expression correlated negatively with that of these miRNAs. We also found that MUC19, SLC8A3³⁶ and SPP1³⁷ in the ceRNA network were associated with AR, while PTPN22 and FCRL5 expression was downregulated in the nasal mucosa. Since single nucleotide polymorphisms in PTPN22³⁸ and FCRL5³⁹ is associated with asthma and AR in Chinese children, future studies should focus on the mechanisms via which DE genes regulate AR.

Finally, we constructed a novel DE circRNA-DE miRNA-DE mRNA network for AR consisting of 64 ceRNA pairs, including 11 circRNAs bound to 17 miRNAs, such as hsa-miR-98-5p, to regulate the expression of 29 genes. Previous studies have found that AR is regulated by many signaling pathways, including the PI3K-Akt⁴⁰ and TLR⁴¹ signaling pathways. Consistent with these findings, we found that the 29 AR-related genes were involved in various important biological functions and pathways, including the PI3K-Akt and TLR signaling pathways, suggesting that these pathways play key roles in AR development.

The DE circRNA-DE miRNA-DE mRNA network also revealed that hsa_circ_0029853 could competitively bind to hsa-miR-98-5p, hsa-miR-140-5p, and hsa-miR-3065-5p to regulate ERRFI1 and NRCAM (Fig. 5), whose expressions were significantly downregulated in the nasal mucosa of patients with AR. ERRFI1 is also downregulated in human breast, skin, pancreatic, and ovarian cancers.⁴² Further, Izumchenko et al⁴³ have found that ERRFI1 can significantly inhibit LPS-induced TNF- α , IL-1 β , and IL-6 mRNA and protein expression. Moreover, our previous studies revealed that these pro-inflammatory cytokines were upregulated in the nasal mucosa of patients with AR,⁴¹ suggesting that their overexpression may be related to ERRFI1 downregulation in these patients. In addition, ERRFI1 has been implicated in the epidermal growth factor receptor (EGFR) signaling pathway⁴³ and studies have demonstrated an interplay between EGFR and eosinophilic airway inflammation.^{44,45} Moreover, intranasal administration of the EGFR inhibitor AG1478 has been found to significantly suppress GM-CSF, eotaxin-1, and RANTES secretion by nasal mucosal cells.⁴⁵ Since ERRFI1 is a negative regulator of EGFR,⁴³ it may participate in AR pathogenesis by regulating the EGFR signaling pathway in the nasal mucosa. Additionally, ERRFI1 is regulated by multiple miRNAs, including miR-126,⁴⁶ miR-205⁴⁷ and miR-2355-5p⁴³; therefore, hsa_circ_0029853 may sponge hsa-miR-98-5p, hsa-miR-140-5p, and hsa-miR-3065-5p to inhibit ERRFI1 expression and thereby promote AR progression.

There are some limitations in the present study. First, we did not perform a functional analysis for the miRNAs (such as hsa-miR-98-5p, hsa-miR-455-5p) and genes (such as DOCK8, ERRFI1), especially their ceRNA network, to determine their roles in AR pathogenesis. The second is that we did not explore the mechanism of circRNAs in sponging miRNAs to regulating mRNAs, or other functions of circRNAs, such as acting as coding protein.

In conclusion, circRNA expression profiles and the ceRNA network for AR could provide a basis for future studies on the underlying molecular mechanisms of AR.

Abbreviations

AR, allergic rhinitis; miRNA, microRNA; lncRNA, long non-coding RNA; circRNA, circular RNA; ceRNA, competing endogenous RNA; MREs, miRNA response elements; DE, differentially expressed; HTS, high-throughput sequencing; GO, gene ontology; KEGG, Kyoto encyclopedia of genes and genomes; RT-PCR, reverse transcription-polymerase chain reaction; SD, standard deviation; RPM, reads per million; PCC, Pearson correlation coefficient; Treg, regulatory T cells; Breg, regulatory B cells; TLR, Toll-like receptor; EGFR, epidermal growth factor receptor.

Ethics approval and consent to participate

This study was approved by the Ethics Committee of the First Affiliated Hospital of Nanjing Medical University (Approval No. 2019-SR-023), and all the participants have signed written informed consent.

Availability of data and materials

The datasets used and/or analyzed during the current study are available from the corresponding author on reasonable request.

Consent for publication

All authors have approved the manuscript for submission.

Author contributions

C-Y Qiu and X-Y Cui performed data analysis and wrote the draft manuscript. M-P Lu, M Yin, W-Y Xu, and X-J Zhu collected samples and performed specimen quality screening. Q Yang performed RT-PCR. L Cheng designed the study and finalized the manuscript. All authors approved the final version of the manuscript.

Funding

This study was supported by a grant (No. 81970862) from the National Natural Science Foundation of China.

Potential competing interests

The authors report no competing interests.

Confirmation of unpublished work

A part of data and materials in this manuscript has been posted to the AUTHORIA preprint server at <https://authoria.com/doi/full/10.22541/au.160193459.95947821/v1>. The authors confirm that the final version of the manuscript is original, has not been published before, and is not currently being considered for publication elsewhere.

Acknowledgements

This study was supported by a grant (No. 81970862) from the National Natural Science Foundation of China. We thank Associate Professor Yong-Ke Cao at the College of

Foreign Languages of Nanjing Medical University for professional English-language proofreading of the manuscript.

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.waojou.2021.100548>.

Author details

^aDepartment of Otorhinolaryngology & Clinical Allergy Center, The First Affiliated Hospital, Nanjing Medical University, Nanjing, China. ^bInternational Centre for Allergy Research, Nanjing Medical University, Nanjing, China.

REFERENCES

1. Cheng L, Chen J, Fu Q, et al. Chinese Society of Allergy guidelines for diagnosis and treatment of allergic rhinitis. *Allergy Asthma Immunol Res.* 2018;10:300-353.
2. Wang XD, Zheng M, Lou HF, et al. An increased prevalence of self-reported allergic rhinitis in major Chinese cities from 2005 to 2011. *Allergy.* 2016;71:1170-1180.
3. Shaoqing Y, Ruxin Z, Guojun L, et al. Microarray analysis of differentially expressed microRNAs in allergic rhinitis. *Am J Rhinol Allergy.* 2011;25:e242-e246.
4. Ma Z, Teng Y, Liu X, et al. Identification and functional profiling of differentially expressed long non-coding RNAs in nasal mucosa with allergic rhinitis. *Tohoku J Exp Med.* 2017;242:143-150.
5. Barrett SP, Salzman J. Circular RNAs: analysis, expression and potential functions. *Development.* 2016;143:1838-1847.
6. Li X, Yang L, Chen LL. The biogenesis, functions, and challenges of circular RNAs. *Mol Cell.* 2018;71:428-442.
7. Pan Z, Li GF, Sun ML, et al. MicroRNA-1224 splicing circularRNA-Filip11 in an Ago2-dependent manner regulates chronic inflammatory pain via targeting Ubr5. *J Neurosci.* 2019;39:2125-2143.
8. Zhang F, Zhang R, Zhang X, et al. Comprehensive analysis of circRNA expression pattern and circRNA-miRNA-mRNA network in the pathogenesis of atherosclerosis in rabbits. *Aging (Albany NY).* 2018;10:2266-2283.
9. Fang Y, Wang X, Li W, et al. Screening of circular RNAs and validation of circANKRD36 associated with inflammation in patients with type 2 diabetes mellitus. *Int J Mol Med.* 2018;42:1865-1874.
10. Liu J, Cao Z. Protective effect of circular RNA (circRNA) Ddx17 in ovalbumin (OVA)-induced allergic rhinitis (AR) mice. *Med Sci Monit.* 2020;26, e919083.
11. Zhu X, Wang X, Wang Y, Zhao Y. The regulatory network among CircHIPK3, lncGAS5, and miR-495 promotes Th2 differentiation in allergic rhinitis. *Cell Death Dis.* 2020;11:216.
12. Wang T, Wang P, Chen D, Xu Z, Yang L. circARRDC3 contributes to interleukin-13-induced inflammatory cytokine and mucus production in nasal epithelial cells via the miR-375/KLF4 axis. *Mol Med Rep.* 2021;23:141.

13. Salmena L, Poliseno L, Tay Y, Kats L, Pandolfi PP. A ceRNA hypothesis: the Rosetta Stone of a hidden RNA language? *Cell*. 2011;146:353-358.
14. Tay Y, Rinn J, Pandolfi PP. The multilayered complexity of ceRNA crosstalk and competition. *Nature*. 2014;505:344-352.
15. Xu L, Feng X, Hao X, et al. CircSETD3 (Hsa_circ_0000567) acts as a sponge for microRNA-421 inhibiting hepatocellular carcinoma growth. *J Exp Clin Oncol*. 2019;38:98.
16. Shi F, Shi Z, Zhao Y, Tian J. CircRNA hsa-circ-0014359 promotes glioma progression by regulating miR-153/PI3K signaling. *Biochem Biophys Res Commun*. 2019;510:614-620.
17. Zhou ZB, Huang GX, Fu Q, et al. circRNA.33186 contributes to the pathogenesis of osteoarthritis by sponging miR-127-5p. *Mol Ther*. 2019;27:531-541.
18. Gao Y, Wang J, Zhao F. CIRI: an efficient and unbiased algorithm for de novo circular RNA identification. *Genome Biol*. 2015;16:4.
19. Holdt LM, Kohlmaier A, Teupser D. Molecular roles and function of circular RNAs in eukaryotic cells. *Cell Mol Life Sci*. 2018;75:1071-1098.
20. Li Z, Huang C, Bao C, et al. Exon-intron circular RNAs regulate transcription in the nucleus. *Nat Struct Mol Biol*. 2015;22:256-264.
21. You X, Vlatkovic I, Babic A, et al. Neural circular RNAs are derived from synaptic genes and regulated by development and plasticity. *Nat Neurosci*. 2015;18:603-610.
22. Legnini I, Di Timoteo G, Rossi F, et al. Circ-ZNF609 is a circular RNA that can be translated and functions in myogenesis. *Mol Cell*. 2017;66:22-37. e9.
23. Tost J. A translational perspective on epigenetics in allergic diseases. *J Allergy Clin Immunol*. 2018;142:715-726.
24. Verduci L, Strano S, Yarden Y, Blandino G. The circRNA-microRNA code: emerging implications for cancer diagnosis and treatment. *Mol Oncol*. 2019;13:669-680.
25. Takase Y, Watanabe M, Inoue N, et al. Association of IL-10-regulating microRNAs in peripheral blood mononuclear cells with the pathogenesis of autoimmune thyroid disease. *Immunol Invest*. 2017;46:590-602.
26. Shamji MH, Durham SR. Mechanisms of allergen immunotherapy for inhaled allergens and predictive biomarkers. *J Allergy Clin Immunol*. 2017;140:1485-1498.
27. Sogut A, Yilmaz O, Kirmaz C, et al. Regulatory-T, T-helper 1, and T-helper 2 cell differentiation in nasal mucosa of allergic rhinitis with olive pollen sensitivity. *Int Arch Allergy Immunol*. 2012;157:349-353.
28. Luo J, Guo H, Liu Z, et al. Analysis of peripheral B cell subsets in patients with allergic rhinitis. *Allergy Asthma Immunol Res*. 2018;10:236-243.
29. Wang M, Gu Z, Yang J, Zhao H, Cao Z. Changes among TGF- β 1⁺ Breg cells and helper T cell subsets in a murine model of allergic rhinitis with prolonged OVA challenge. *Int Immunopharm*. 2019;69:347-357.
30. Matsuda M, Morie Y, Oze H, et al. Phenotype analyses of IL-10-producing Foxp3⁺ CD4⁺ T cells increased by subcutaneous immunotherapy in allergic airway inflammation. *Int Immunopharm*. 2018;61:297-305.
31. Taitano SH, van der Vlugt LEM, Shea MM, Yang J, Lukacs NW, Lundy SK. Differential influence on regulatory B cells by T_H2 cytokines affects protection in allergic airway disease. *J Immunol*. 2018;201:1865-1874.
32. Biggs CM, Keles S, Chatila TA. DOCK8 deficiency: insights into pathophysiology, clinical features and management. *Clin Immunol*. 2017;181:75-82.
33. Janssen E, Morbach H, Ullas S, et al. Deducator of cytokines 8-deficient patients have a breakdown in peripheral B-cell tolerance and defective regulatory T cells. *J Allergy Clin Immunol*. 2014;134:1365-1374.
34. Harada Y, Tanaka Y, Terasawa M, et al. DOCK8 is a Cdc42 activator critical for interstitial dendritic cell migration during immune responses. *Blood*. 2012;119:4451-4461.
35. Singh AK, Eken A, Fry M, Bettelli E, Oukka M. DOCK8 regulates protective immunity by controlling the function and survival of ROR γ ⁺ ILCs. *Nat Commun*. 2014;5:4603.
36. Shang Y, Das S, Rabold R, Sham JS, Mitzner W, Tang WY. Epigenetic alterations by DNA methylation in house dust mite-induced airway hyperresponsiveness. *Am J Respir Cell Mol Biol*. 2013;49:279-287.
37. Kasetty G, Bhongir R, Papareddy P, et al. Osteopontin protects against pneumococcal infection in a murine model of allergic airway inflammation. *Allergy*. 2019;74:663-674.
38. Ke X, Song S, Wang X, Shen Y, Kang H, Hong S. Associations of single nucleotide polymorphisms of PTPN22 and Ctla4 genes with the risk of allergic rhinitis in a Chinese Han population. *Hum Immunol*. 2017;78:227-231.
39. Gu Z, Shen Y, Tang XY, et al. Genetic risk of FCRL3 and FCRL5 polymorphisms in children with asthma and allergic rhinitis in a Chinese Han population. *Int J Pediatr Otorhinolaryngol*. 2019;120:58-63.
40. Chen X, Li YY, Zhang WQ, Zhang WM, Zhou H. House dust mite extract induces growth factor expression in nasal mucosa by activating the PI3K/Akt/HIF-1 α pathway. *Biochem Biophys Res Commun*. 2016;469:1055-1061.
41. Cui XY, Chen X, Yu CJ, et al. Increased expression of toll-like receptors 2 and 4 and related cytokines in persistent allergic rhinitis. *Otolaryngol Head Neck Surg*. 2015;152:233-238.
42. Zhang YW, Vande Woude GF. Mig-6, signal transduction, stress response and cancer. *Cell Cycle*. 2007;6:507-513.
43. Izumchenko E, Sidransky D. Understanding the MIG6-EGFR signaling axis in lung tumorigenesis. *Canc Discov*. 2015;5:472-474.
44. El-Hashim AZ, Khajah MA, Renno WM, et al. Src-dependent EGFR transactivation regulates lung inflammation via downstream signaling involving ERK1/2, PI3K δ /Akt and NF κ B induction in a murine asthma model. *Sci Rep*. 2017;7:9919.
45. Shimizu S, Takezawa-Yasuoka K, Ogawa T, Tojima I, Kouzaki H, Shimizu T. The epidermal growth factor receptor inhibitor AG1478 inhibits eosinophilic inflammation in upper airways. *Clin Immunol*. 2018;188:1-6.

12 Qiu et al. *World Allergy Organization Journal* (2021) 14:100548
<http://doi.org/10.1016/j.waojou.2021.100548>

46. Wang W, Zheng Y, Wang M, Yan M, Jiang J, Li Z. Exosomes derived miR-126 attenuates oxidative stress and apoptosis from ischemia and reperfusion injury by targeting ERRF1. *Gene*. 2019;690:75-80.
47. Migliore C, Morando E, Ghiso E, et al. miR-205 mediates adaptive resistance to MET inhibition via ERRF1 targeting and raised EGFR signaling. *EMBO Mol Med*. 2018;10, e8746.