



CASE REPORT

Open Access

A patient with features of Albright hereditary osteodystrophy and unusual neuropsychiatric findings without coding Gs α mutations

Shirin Hasani-Ranjbar^{1,2,3*}, Zahra Jouyandeh², Mahsa Mohammad Amoli², Akbar Soltani² and Seyed Masoud Arzaghi²

Abstract

Background: Pseudohypoparathyroidism(PHP) is a heterogeneous group of rare metabolic disorders characterized by hypocalcemia and hyperphosphatemia resulting from PTH resistance. Different forms of PHP have been reported based on biochemical and clinical manifestation and genetic findings. Most of these forms are caused by defects in GNAS, an imprinted gene locus with multiple subunits. We reported a 12-year-old girl with unusual clinical manifestations of Pseudopseudohypoparathyroidism(PPHP).

Methods: After clinical and biochemical evaluations, the patients' genomic DNA was isolated from peripheral blood leukocytes using salting out method. The whole coding sequences of GNAS gene including 13 exons were amplified by PCR. Quantitative PCR reactions were performed too.

Findings: We described a 12-year-old girl with Albright Hereditary osteodystrophy (AHO) phenotype, poor school performance, some abnormal movements, TSH resistance with normal serum calcium and phosphorus levels and normal Gs α bioactivity with no mutation in GNAS exons. Unusual neuropsychiatric findings in this patient were compatible with Asperger syndrome.

Conclusions: According to our findings this patient could not be categorized in any of PHP subgroups. Identifying of such individuals may be useful to discover different genetic patterns in pseudohypoparathyroidism and pseudopseudohypoparathyroidism. It is important to identify patients in whom PHP is caused by novel GNAS mutations, as careful investigations of these findings will likely further our knowledge of this complex and this unique disorder. In addition this case presented with unusual neuropsychiatric findings which has not been reported up to now.

Keywords: Pseudohypoparathyroidism, GNAS mutation, Albright Hereditary osteodystrophy, Pseudopseudohypoparathyroidism

Background

Pseudohypoparathyroidism(PHP) is a heterogeneous group of rare metabolic disorders whose common feature is resistance to the parathyroid hormone which results to hypocalcemia and hyperphosphatemia [1,2]. Different forms of PHP are caused by mutations in those exons of GNAS

encoding the α subunit [3]. Based on biochemical and clinical manifestation and genetic findings, variable forms of this disorder have been described.

Heterogeneous mutations located in GNAS exons which encode Gs α inherited from mother leads to PHP-Ia in which the characteristic phenotype appearance is Albright's hereditary osteodystrophy(AHO), including obesity, short stature, heterotopic ossification, brachydactyly and/or mental retardation which has been described by Albright et al. for the first time in 1942 [4]. This form is also associated with other hormonal abnormalities, including hypothyroidism and hypogonadism [5]. Pseudopseudohypoparathyroidism

* Correspondence: shirinhasanir@yahoo.com

¹Obesity & Eating Habits Research Center, Endocrinology and metabolism Cellular & Molecular Science Institute, Endocrinology & Metabolism research institute, Tehran University of Medical Sciences, Tehran, Iran

²Endocrinology and Metabolism Research center, Endocrinology & Metabolism research institute, Tehran University of Medical Sciences, Tehran, Iran

Full list of author information is available at the end of the article

(PPHP) a subgroup closely related to PHP-Ia have the typical features of AHO without any hormonal resistance [6] caused by heterozygous inactivating mutations in G_sα coding exons of GNAS which is inherited paternally [7]. In Type-Ib the resistance is restricted only to kidney, without AHO and other hormonal abnormalities. These patients have maternal microdeletions of STX16 which is associated with loss of methylation at GNAS exon A/B [1,3]. Another subtype called I-c is the same as type-Ia with no difference in clinical phenotype with in some patients, a mutation that affect the coupling of G protein to the PTH receptor [8,9]. Patients with PHP Type 2 do not have the physical appearance of AHO with a normal cAMP in response to PTH stimulation despite the inherent abnormality in calcium regulation [10,11]. However, recent reports have provided evidence that overlap can exist between the clinical features of different PHP subtypes [12,13]. Different subtypes of pseudohypoparathyroidism are shown in Table 1.

The psychiatric symptoms of pseudohypoparathyroidism have been reported to be related to Intracerebral calcification in basal ganglia [14]. Moderately severe to entirely normal school performance and moderate degrees of mental retardation has been reported in patients with PHP-Ia but not in PHP-Ib [15,16]. Seizure, tetany and laryngeal stridor may also occur in PHP-Ia because of hypocalcemia [17].

Here we present a case with Albright's hereditary osteodystrophy phenotype, with normal calcium, phosphate and PTH levels, in which neurological symptoms, resistance to other hormones and negative GNAS gene mutations were unique findings.

Patients and methods

Mutation analysis of the GNAS gene

The patients' genomic DNA was isolated from peripheral blood leukocytes using salting out method. The whole coding sequences of GNAS gene including 13 exons were amplified by PCR according to the assay described before [18].

Molecular analysis

After collection of heparinized venous blood, lymphocyte separation was performed using density gradient ficoll (1.077) and total RNA was extracted using Tripure Isolation Reagent (Roche Applied Science), according to the manufacturer's protocol. One µg aliquot of total RNA from patients sample and 3 normal control samples were reverse transcribed into single-stranded cDNA using random

hexanucleotides primers and expand reverse transcriptase (Roche Applied Science). cDNA for sample was subjected to quantitative real-time PCR using the following primers for HPRT internal quantitative control forward 5'-CC TGGCGTCGTGATTAGTGAT-3', reverse 5'- AGACGTT CAGTCCTGTCCATAA-3' and GNAS, Forward 5'- TGTA CAAGCAGTTAACACCCACCA -3' and reverse 5'- TCT GTAGGCCGCCTTAAGCTTTC -3' primers.

Quantitative PCR

For evaluation of G_s alpha gene quantitative PCR reactions were performed in 25 µl reaction mixtures containing 250 ng cDNA, 10 µl Takara Real-Time™ SYBR Green/ROX PCR Master, primer pairs and nuclease-free water to 20 µl. Each biological replicate was run in duplicate on an ABI Step One quantitative PCR system. Thermocycling conditions consisted of an initial polymerase activation step at 95°C for 10 min, followed by 40 cycles at 95°C for 5 s and 60°C for 30 s. Afterwards, melting curves were generated to confirm a single gene-specific peak and to detect primer dimmer formation.

Case presentation

Patient

A 12-year-old girl whom parents were distant relatives was referred to our clinic for evaluation of short stature and poor school performance. Her past medical history was positive for post term delivery (2 weeks delay) with cesarean section and meconium aspiration, poor feeding, severe high arched palate in infancy and a history of reddened face while crying, amblyopia at the age of 5 year old, poor school performance and attention deficit (concentration disturbance). She showed some abnormal repetitive verbal and physical movements. Physical examination revealed obesity with short neck, round face and brown spots on her skin. Her height was 127 cm (below the third percentile) and her weight was 50 kg (body mass index > 97th percentile). Mother's height was 158 cm and father's height was 172 cm. Both hands appeared short, with markedly short fourth and fifth metacarpals. Thyroid physical exam showed a 15–20 gr, normal, not firm gland without any palpable nodules. Laboratory data revealed normal serum calcium (9.5 mg/dl), phosphorus (5.6 mg/dl) and intact PTH (10 pg/ml). Serum TSH was 11 µU/ml. Her puberty showed normal timing (menarche at 11 y and 6 months). All laboratory tests are shown in Table 2.

Table 1 Classification of pseudohypoparathyroidism with AHO phenotype

	PTH resistance	Hormone resistance	Typical AHO phenotype	G _s α levels (%)	GNAS defects
PHP-Ia	Yes	Yes	Yes	50	+ G _s α mutations
PPHP	No	No	Yes	50	± G _s α mutations
Case	No	Yes	Yes	100	- G _s α mutations

Table 2 Clinical and laboratory findings of the patient

Variable	Patient	Normal range
Height (cm)	127	Below the 3th percentile
Weight (kg)	50	Above 97 th percentile
Calcium (mg/dl)	9.5	8.5-10.5
Phosphate (mg/dl)	5.6	3.6-5.8
PTH (pg/ml)	10	10-62
TSH (μ U/ml)	11	0.5-4
Anti-TPO Ab	Negative	<15

Radiography of the left hand revealed markedly short fourth and fifth metacarpals (Figure 1). Bone age was compatible with her age. Electromyography (EMG) and Nerve conduction velocity (NCV) tests of four limbs were normal. Echocardiography showed an ejection fraction of 60%, other cardiac components were normal.

Karyotype analysis on peripheral lymphocytes was 46, XX as a normal female chromosomal pattern with no chromosomal defect. We analyzed the 13 GNAS exons encoding Gs α in genomic DNA and no mutations or alternations could be identified. Gs α mRNA level, measured by qRT-PCR using primers at exons 2 and 11, showed that the level



Figure 1 Left hand x-ray of case. Short 4th & 5th metacarpal.

was not different from the controls. The subject and her family gave informed consent to the study.

Neuropsychiatric findings

She had a febrile seizure at 1.5 years old and in early childhood recurrent generalized Tonic-clonic seizures with Abnormal EEG (frequent paroxysmal dysrhythmia) which treated up to now with provision of neurologist. Stereotyped and repetitive motor mannerisms (hand flapping) were present during infancy and early childhood as parents registered. Severe impairment in social interactions as failure to develop age-appropriate peer relationship and severe communication problems with family members especially with mother are present. Beside, the patient had ritualistic behaviors and compulsions (washing and repeating) which significantly impaired her functioning and proposed comorbid obsessive-compulsive disorder. Academic failure was present from elementary school stage and total intellectual score using Wechsler intelligence scale was 68 and suggests mild mental retardation. Episodes of binge eating behavior present and patient was obviously obese. Magnetic Resonance Imaging (MRI) revealed bilateral ventricles enlargement due to Cavum Septum Pellucidum.

Discussion

In this report, we present a patient with short stature and poor school performance, with elevated TSH level diagnosed as hypothyroidism. The patient had normal calcium and phosphate levels with no resistance to PTH hormone, which did not lead to suspicion of PHP-Ia or Ib. We found a short stature male in father's family (his uncle) and a short stature member (grandfather) in mother's family; however we have report of AHO phenotype in 2 cousins of her, which suggest the paternally inherited pattern of the disease.

These data and her laboratorial profile were compatible with PPHP [7,19]. The patient had normal serum concentrations of calcium and phosphate for his age, and just his elevated levels of TSH and his mother's background was compatible with the diagnosis of PHP-Ia [17]. Mariana Tenorio Antunes Reis et al. [20] have also reported an infant boy with AHO phenotype, normal calcium and phosphate levels, whose mother was diagnosed as PHP-Ia. The functional analysis of GNAS exons showed a novel heterozygous missense mutation in exon 5 of GNAS. They suggest that AHO phenotype and normocalcemia does not always mean PPHP, especially when the allele is maternally inherited.

To investigate this possibility, the GNAS coding region of patient was studied. There was no typical mutation in GNAS coding region and the Gs α mRNA level did not appear to be diminished, which is compatible with Gs α pattern in PHP-Ic [9]. It should be noted that our qRT-PCR analysis of Gs-alpha mRNA was performed by generating

an amplicon consisting of the region encoded by exons 2 through 11. Thus, given that certain other GNAS transcripts share this region, the level of Gs-alpha mRNA determined in our experiments may not entirely reflect the expression of Gs-alpha transcript in case of any significant changes being observed in the level of expression. Sussane Thiele et al. described 5 patients diagnosed as PHP-Ic in whom they found TSH and PTH resistance, with normal Gs α activity and inherited maternal inactivating mutations in GNAS exons [21]. Linglart et al. [22] in a study on genotype, phenotype relationship on the maternal transmission of the hormonal resistance did not observe any relationship between the erythrocyte Gs α activity and the phenotype which may justify our findings in this patient.

This patient has high TSH and low anti TPO level which leads to hypothyroidism. Joaquin Lado-Abeal et al. [23] reported a family with congenital hypothyroidism due to maternal inherited loss-of-function mutation of TSH receptor and AHO phenotype in two children of family due to a paternal inherited mutation of GNAS,G alpha protein subunit. This study suggests that our subject may have the same inheritance pattern which leads to TSH resistance and AHO phenotype, although we do not have enough evidence to prove this hypothesis. Also studies have shown that Gs α has a tissue specific manner, as YU S et al. showed a mild PTH resistance despite normal TSH and thyroid hormone levels in a mice because of imprinted gene in renal tubules but not in other tissues [24]. A justification for some unusual and discordant findings in PHP patients which has been confirmed by other studies recently [25,26] may explain the hormonal resistance pattern in this case.

These unusual findings make a difficulty in net diagnosis for this case. This patient did not have PTH resistance which cannot be classified as type Ia or Ib and at the same time GNAS mutation was negative. Israeli et al. also reported a new variant of the syndrome, affecting 5 individuals in a 3 generation family with AHO, normal Gs activity and hypothyroidism [27]. Giovanna Mantovani and colleges in a study on 40 patients have detected an absence of mutations in Gs α -coding GNAS exons although they had the presence of PTH resistance [12]. Due to Gs α inheritance pattern, it is suggested that PHP-Ia results from a heterozygous inactivating mutation at GNAS inherited from maternal allele while PPHP is inherited by paternal alleles [7]. In this case the paternally inheritance pattern, with normal Gs α activity and resistance to TSH hormone make a discordance to categorize this patient as PHP-Ia, Ic or PPHP. Although we did not do genome sequencing to identify specific GNAS mutations in this patient, Linglart et al. suggest that western blot could be used to assess Gs α protein expression. They have report three patients with PHP-Ia phenotype without any GNAS mutations that hypothesize that normal Gs α protein

expression and the absence of a mutation in the coding regions makes the presence of a GNAS1 molecular defect unlikely [22]. Davids et al. proposed that an alternative etiology for PPHP is the deletion of STK25, inhibiting G protein signaling by a mechanism different from mutation of the Gs α gene which may account for the PPHP condition in the small group of patients with normal Gs α who still experience the AHO phenotype [28]. Also, due to normal karyotype in this patient, brachydactyly-mental retardation syndrome with deletion in chromose 2 is not considered. This deletion (MIM #600430) is often reffered to as Albright hereditary osteodystrophy-like syndrome. These patients with distal 2q deletions present with mild to moderate intellectual disabilities, hypotonia, obesity, short stature, and brachydactyly with short phalanges[especially the third to fifth phalanges;brachydactyly type E], as seen in patients with Albright syndrome [29]. No dysmorphic features including thin, highly arched eyebrows, prominent forehead, depressed nasal bridge, hypoplastic alae nasi, prominent nasal septum, thin upper lip, and ear anomalies (which are part of brachydactyly-mental retardation syndrome) were not seen in our patient.

However the normal laboratory tests in our case, decline the diagnosis of PHP-Ia. Analysis of PHP-Ia and PPHP cases shows that individuals with isolated AHO and normal Gs α activity exists [22]. So identifying of such individuals may be useful to discover different genetic patterns in pseudohypoparathyroidism and pseudopseudohypoparathyroidism. Regarding to the reports by Nanclares et al. [5] Mantovani et al. [12] and Sussane Thiele et al. [21], these findings strengthen the need of new classifications of GNAS related disorders due to this discordance reported in this case and other similar cases described before.

The neuropsychiatric findings in our subject were compatible with Asperger syndrome diagnosis. Asperger syndrome is characterized by impairment in social interactions and restrictive-repetitive pattern of behaviors and interest. The syndrome belongs of Pervasive Developmental Disorders (PDD) or Autism spectrum disorders [30]. The aforementioned case with multiple medical complications had prominent neuropsychiatric manifestations such as; generalized Tonic-clonic seizures with Abnormal EEG, Stereotyped and repetitive motor mannerisms, severe impairment in social interactions, severe communication problems with family members, ritualistic behaviors and compulsions (washing and repeating), mild mental retardation. Warwick et al. [31] describes a case of a 23 years old man with obsessive-compulsive behavior and agitation, temporal lobe epilepsy (multiple generalized tonic-colonic seizures), some developmental delay, and markedly impaired social interactions toward family and at work with stereotyped and repetitive movements. He was diagnosed and treated as Asperger syndrome, that his symptoms are compatible with neuropsychiatric findings in our subject. In addition, the

case of co-morbid Pseudohypoparathyroidism and Asperger syndrome manifestations has not been reported to now as we searched.

It will be important to identify patients in whom PHP (or PPHP subtype) is caused by novel GNAS findings or unusual clinical findings, as careful investigations of these findings will likely further our knowledge of this complex and this unique disorder. The most important point is that this case presented with unusual neurologic findings which has not been reported up to now.

Conclusion

According to our findings this patient could not be categorized in any of PHP subgroups. Identifying of such individuals may be useful to discover different genetic patterns in pseudohypoparathyroidism and pseudopseudohypoparathyroidism. It is important to identify patients in whom PHP is caused by novel GNAS mutations, as careful investigations of these findings will likely further our knowledge of this complex and this unique disorder. In addition this case presented with unusual neuropsychiatric findings which has not been reported up to now.

Consent

Written informed consent was obtained from the patient's parents for publication of this Case report.

Competing interest

The authors declare that they have no competing interest.

Authors' contributions

SHR and AS gave the idea and designed the study and do the Follow up of the patient. SHR also edited the article. ZJ does the follow up and drafted the manuscript. SMA investigates and drafted the neuropsychiatric findings' section of the article. MMA does the genetic evaluations. All authors have read and approved content of the article.

Acknowledgements

We gratefully thank the patient and her family for consenting to publication of this study.

Author details

¹Obesity & Eating Habits Research Center, Endocrinology and metabolism Cellular & Molecular Science Institute, Endocrinology & Metabolism research institute, Tehran University of Medical Sciences, Tehran, Iran. ²Endocrinology and Metabolism Research center, Endocrinology & Metabolism research institute, Tehran University of Medical Sciences, Tehran, Iran. ³Endocrinology & Metabolism Research Institute, 5th Floor, Shariati Hospital, North Kargar Ave, Tehran 14114, Iran.

Received: 27 August 2013 Accepted: 10 April 2014

Published: 22 May 2014

References

1. Weinstein LS, Yu S, Warner DR, Liu J: Endocrine manifestations of stimulatory G protein α -subunit mutations and the role of genomic imprinting. *Endocr Rev* 2001, **22**(5):675–705.
2. Mantovani G, Spada A: Mutations in the Gs alpha gene causing hormone resistance. *Best Pract Res Clin Endocrinol Metab* 2006, **20**(4):501–513.
3. Bastepo M, Jüppner H: GNAS locus and pseudohypoparathyroidism. *Horm Res* 2005, **63**(2):65–74.
4. Albright F, Burnett C, Smith PH, Parson W: Pseudohypoparathyroidism: an example of "Seabright-Bantam syndrome". *Endocrinology* 1942, **30**(6):922–932.
5. de Nanclares GP, Fernández-Rebolledo E, Santin I, García-Cuartero B, Gatzambide S, Menéndez E, Morales MJ, Pombo M, Bilbao JR, Barros F, Zazo N, Ahrens W, Jüppner H, Hiort O, Castaño L, Bastepo M: Epigenetic defects of GNAS in patients with pseudohypoparathyroidism and mild features of Albright's hereditary osteodystrophy. *J Clin Endocrinol Metab* 2007, **92**(6):2370–2373.
6. Albright F, Forbes A, Henneman P: Pseudo-pseudohypoparathyroidism. *Trans Assoc Am Phys* 1952, **65**:337.
7. Wilson LC, Oude Luttkhuis ME, Clayton PT, Fraser WD, Trembath RC: Parental origin of Gs alpha gene mutations in Albright's hereditary osteodystrophy. *J Med Genet* 1994, **31**(11):835–839.
8. Farfel Z, Brothers VM, Brickman AS, Conte F, Neer R, Bourne HR: Pseudohypoparathyroidism: inheritance of deficient receptor-cyclase coupling activity. *Proc Natl Acad Sci U S A* 1981, **78**(5):3098–3102.
9. ALDRED MA: Genetics of pseudohypoparathyroidism types Ia and Ic. *J Pediatr Endocrinol Metab* 2006, **19**(Suppl 2):635–640.
10. Dreznar M, Neelon FA, Lebovitz HE: Pseudohypoparathyroidism type IIa: a possible defect in the reception of the cyclic AMP signal. *N Engl J Med* 1973, **289**(20):1056–1060.
11. Rodriguez HJ, Villarreal JRH, Klahr S, Slatopolsky E: Pseudohypoparathyroidism type II: restoration of normal renal responsiveness to parathyroid hormone by calcium administration. *J Clin Endocrinol Metab* 1974, **39**(4):693–701.
12. Mantovani G, de Sanctis L, Barbieri AM, Elli FM, Bollati V, Vaira V, Labarile P, Bondioni S, Peverelli E, Lania AG, Beck-Peccoz P, Spada A: Pseudohypoparathyroidism and GNAS epigenetic defects: clinical evaluation of Albright hereditary osteodystrophy and molecular analysis in 40 patients. *J Clin Endocrinol Metab* 2010, **95**(2):651–658.
13. Lecumberri B, Fernández-Rebolledo E, Sentchordi L, Saavedra P, Bernal-Chico A, Pallardo LF, Bustos JM, Castaño L, de Santiago M, Hiort O, Pérez de Nanclares G, Bastepo M: Coexistence of two different pseudohypoparathyroidism subtypes (Ia and Ib) in the same kindred with independent Gsa coding mutations and GNAS imprinting defects. *J Med Genet* 2010, **47**(4):276–280.
14. Illum F, Dupont E: Prevalences of CT-detected calcification in the basal ganglia in idiopathic hypoparathyroidism and pseudohypoparathyroidism. *Neuroradiology* 1985, **27**(1):32–37.
15. Farfel Z, Friedman E: Mental deficiency in pseudohypoparathyroidism type I is associated with Ns-protein deficiency. *Ann Intern Med* 1986, **105**(2):197–199.
16. Narch H: A girl with poor school performance. *Eur J Pediatr* 2000, **159**(1):121–122.
17. Wilson LC, Trembath RC: Albright's hereditary osteodystrophy. *J Med Genet* 1994, **31**(10):779–784.
18. Mantovani G, Romoli R, Weber G, Brunelli V, De Menis E, Beccio S, Beck-Peccoz P, Spada A: Mutational analysis of GNAS1 in patients with pseudohypoparathyroidism: identification of two novel mutations. *J Clin Endocrinol Metab* 2000, **85**(11):4243–4248.
19. Davies SJ, Hughes HE: Imprinting in Albright's hereditary osteodystrophy. *J Med Genet* 1993, **30**(2):101–103.
20. Reis MT, Cattani A, Mendonca BB, Corrêa PH, Martin RM: A novel GNAS mutation in an infant boy with pseudohypoparathyroidism type Ia and normal serum calcium and phosphate levels. *Arq Bras Endocrinol Metabol* 2010, **54**(8):728–731.
21. Thiele S, de Sanctis L, Werner R, Grötzinger J, Aydin C, Jüppner H, Bastepo M, Hiort O: Functional characterization of GNAS mutations found in patients with pseudohypoparathyroidism type Ic defines a new subgroup of pseudohypoparathyroidism affecting selectively Gsa-receptor interaction. *Hum Mutat* 2011, **32**(6):653–660.
22. Linglart A, Carel JC, Garabédian M, Lé T, Mallet E, Kottler ML: GNAS1 lesions in pseudohypoparathyroidism Ia and Ic: genotype phenotype relationship and evidence of the maternal transmission of the hormonal resistance. *J Clin Endocrinol Metab* 2002, **87**(1):189–197.
23. Lado-Abeal J, Castro-Piedras I, Palos-Paz F, Labarta-Aizpún JL, Albergo-Gamboa R: Family with congenital hypothyroidism caused by a combination of loss-of-function mutations in the thyrotropin receptor and adenylate cyclase-stimulating g alpha-protein subunit genes. *Thyroid* 2011, **21**(2):103–109.
24. Yu S, Gavrilova O, Chen H, Lee R, Liu J, Pacak K, Parlow AF, Quon MJ, Reitman ML, Weinstein LS: Paternal versus maternal transmission of a stimulatory G-protein alpha subunit knockout produces opposite effects on energy metabolism. *J Clin Invest* 2000, **105**(5):615–623.

25. Mantovani G, Ballare E, Giannone E, Beck-Peccoz P, Spada A: The *Gsa* gene: predominant maternal origin of transcription in human thyroid gland and gonads. *J Clin Endocrinol Metab* 2002, **87**(10):4736–4740.
26. Germain-Lee EL, Ding CL, Deng Z, Crane JL, Saji M, Ringel MD, Levine MA: Paternal imprinting of *Gαs* in the human thyroid as the basis of TSH resistance in pseudohypoparathyroidism type 1a. *Biochem Biophys Res Commun* 2002, **296**(1):67–72.
27. Israeli S, Metzker A, Horev G, Karmi D, Merlob P, Farfel Z: Albright hereditary osteodystrophy with hypothyroidism, normocalcemia, and normal Gs protein activity: a family presenting with congenital osteoma cutis. *Am J Med Genet* 1992, **43**(4):764–767.
28. Davids MS, Crawford E, Weremowicz S, Morton CC, Copeland NG, Gilbert DJ, Jenkins NA, Phelan MC, Comb MJ, Melnick MB: STK25 is a candidate gene for pseudopseudohypoparathyroidism. *Genomics* 2001, **77**(1–2):2–4.
29. GeneReviews: 2q37 Deletion Syndrome; <http://www.ncbi.nlm.nih.gov/book/NBK1158>.
30. American Psychiatric Association: *Diagnostic and statistical manual of mental disorders : DSM-IV-TR*. American Psychiatric Association: Washington; 2000.
31. Warwick TC, Griffith J, Reyes B, Legesse B, Evans M: Effects of vagus nerve stimulation in a patient with temporal lobe epilepsy and Asperger syndrome: case report and review of the literature. *Epilepsy Behav* 2007, **10**(2):344–347.

doi:10.1186/2251-6581-13-56

Cite this article as: Hasani-Ranjbar et al.: A patient with features of albright hereditary osteodystrophy and unusual neuropsychiatric findings without coding *Gsalpha* mutations. *Journal of Diabetes & Metabolic Disorders* 2014 13:56.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

