

Liver-specific ROR α deletion does not affect the metabolic susceptibility to western style diet feeding



Antonio Molinaro¹, Robert Caesar¹, Laurent L'homme², Ara Koh¹, Marcus Ståhlman¹, Bart Staels², Fredrik Bäckhed^{1,3,*}

ABSTRACT

Objectives: The nuclear receptor superfamily is a potential target for the development of new treatments for obesity and metabolic diseases. Increasing evidence has pointed towards the retinoic acid-related orphan receptor-alpha (ROR α) as an important nuclear receptor involved in several biological processes. ROR α full body knockout mice display improved metabolic phenotypes on both chow and high fat (60% fat, 20% carbohydrate) diets, but also have severe behavioral abnormalities. Here we investigated the effect of hepatic ROR α by generating mice with liver-specific ROR α deletion to elucidate the role of this nuclear receptor on host metabolism.

Methods: 8 week-old mice with liver-specific ROR α deletion and littermate controls were fed either chow or western-style diets (40% fat, 40% carbohydrate) for 12 weeks. Metabolic phenotyping was performed at the end of the dietary intervention.

Results: Here, we show that hepatic ROR α deletion does not affect the metabolic susceptibility to either chow or western-style diet in terms of glucose metabolism and adiposity.

Conclusions: Our data indicate that liver deletion of ROR α does not have a pivotal role in the regulation of hepatic glucose and lipid metabolism on chow or western-style diet.

© 2019 The Authors. Published by Elsevier GmbH. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Keywords ROR α ; Obesity; Glucose metabolism; Steatosis

1. INTRODUCTION

The incidence of obesity and metabolic associated disorders, such as cardiovascular diseases, type 2 diabetes mellitus and non-alcoholic fatty liver disease (NAFLD), increase worldwide [1–3]. Furthermore, there are no effective long-term non-surgical treatments for obesity and metabolic associated disorders [4]. However, accumulating evidence points towards nuclear receptors (NRs) as potential molecular treatment targets [5,6]. The NRs superfamily comprises 48 transcription factors in humans, which are involved in the regulation of a wide range of physiologic and pathophysiologic processes (i.e. development, circadian rhythm, response to steroid hormones, inflammation, obesity, diabetes, NAFLD, and cancer) [7–10].

The retinoic acid-related orphan receptor α (ROR α ; NR1F1) is a NR that has been implicated in the regulation of a wide variety of metabolic pathways, including lipid, glucose and steroid metabolism [11–13]. The initial indications that ROR α can affect metabolic processes originate mainly from a natural occurring mutant mouse, the Staggerer mouse, which results in a loss of function [14].

Staggerer mice are protected against diet-induced obesity and associated complications (i.e., adipose tissue inflammation, NAFLD, and impaired glucose metabolism), but also have cerebral phenotypes including impaired locomotion [15,16]. However, activation of ROR α with synthetic ligands also improves NAFLD by reducing hepatic oxidative stress and inflammation suggesting a potential beneficial role of ROR α activation on obesity and metabolic diseases [17–19].

In order to elucidate the role of ROR α in obesity and metabolic diseases, tissue specific knockout models have recently been developed. Liver-specific ablation of ROR α deteriorates metabolic profile in terms of adiposity, liver steatosis, inflammation and hepatic insulin resistance, when mice were fed a high fat diet (60% of energy from fat and 20% from carbohydrates) [20,21]. These findings suggest a central role of ROR α in the regulation of hepatic lipid and glucose metabolism. To further extend and validate the role of ROR α on glucose and lipid metabolism, we generated an independent liver-specific ROR α deficient strain that was metabolically phenotyped on both chow and western-style diet (WSD; 40% of energy from fat and 40% from carbohydrate and 20% protein).

¹Wallenberg Laboratory, Department of Molecular and Clinical Medicine, Sahlgrenska Academy, University of Gothenburg, 413 45, Gothenburg, Sweden ²Univ. Lille, Inserm, CHU Lille, Institut Pasteur de Lille, U1011-EGID, F-59000, Lille, France ³Novo Nordisk Foundation Center for Basic Metabolic Research and Section for Metabolic Receptology and Endoendocrinology, Faculty of Health Sciences, University of Copenhagen, Copenhagen, 2200, Denmark

*Corresponding author. Wallenberg Laboratory, Sahlgrenska University Hospital, S-413 45, Gothenburg, Sweden. E-mail: fredrik.backhed@wlab.gu.se (F. Bäckhed).

Received January 9, 2019 • Revision received February 21, 2019 • Accepted February 28, 2019 • Available online 9 March 2019

<https://doi.org/10.1016/j.molmet.2019.02.010>

2. MATERIALS AND METHODS

2.1. Mice experiments

Mice harbouring *loxP* sequences flanking exon 3, the first common exon coding for both isoform 1 and 4, were generated (Figure 1A). Deletion of the 86 bp exon 3 led to a frame shift resulting in a premature stop codon. Potential translated truncated proteins would be composed of 88 and 32 amino acids instead of 523 and 467 for isoform, 1 and 4 respectively. No functional domain would be present on these putative proteins. Generation of the strain was completed by homologous recombination in C57BL/6J embryonic stem cells followed by microinjection into C57BL/6J blastocysts. The resulting new-borns were crossed with C57BL/6 Flp delete mice to allow the excision of the neomycin selection cassette.

Albumin-Cre mice were purchased from The Jackson Laboratory. Cre efficacy and specificity towards hepatocytes was confirmed by breeding the mice with ROSA^{MT/mG} mice (data not shown). 8-weeks old male ROR α ^{LKO} and ROR α ^{fl/fl} littermate were fed a chow diet (Labdiet) or WSD (Harland, TD.96132, 40% fat, 40% sugar, 20% proteins) *ad libitum* for 12 weeks. MRI, insulin, glucose and pyruvate tolerance tests, and measurements of insulin levels were performed as earlier described [22]. Briefly, insulin, glucose and pyruvate tolerance tests were performed by injecting insulin (0.75 U/kg body weight), glucose (2 g/kg body weight) or pyruvate (2 g/kg weight) respectively, intraperitoneally after a 4 (ITT and GTT) or 12 (PTT) h fast. Tail blood samples were collected at 0, 15, 30, 60, 90 and 120 min and blood glucose levels were determined using a glucose meter (Accu-Check Aviva, Roche). Insulin was measured with ELISA (Crystal Chem) according to the manufacturers' protocols. All experiments including tissue harvesting were performed at the same time of the day (12.00 PM). The local ethics committee in Gothenburg approved all animal experiments.

2.2. Analysis of liver lipids content

Lipids were extracted using the BUMÉ method [23]. The cholesteryl esters, triglycerides and free cholesterol were analyzed using straight phase HPLC coupled to evaporative light scattering detection according to previous work [24]. Briefly, the extract was evaporated under nitrogen and reconstituted in 500 μ l heptane:isopropanol [9:1]. Cholesteryl esters, triglycerides, and free cholesterol were separated using straight-phase HPLC (Sunfire 2.1 \times 100 mm, Waters) coupled to an evaporative light-scattering detector (PL-ELS 1000, Polymer Laboratories), as described previously [24]. Quantification was made using external standards (Larodan Fine Chemicals).

2.3. Primary hepatocytes

Primary hepatocytes were isolated from ROR α ^{LKO} and ROR α ^{fl/fl} littermates mice by digesting liver with perfusion of collagenase type IV, as described previously [25]. After perfusion, 16 \times 10⁵ cells were plated on collagen-coated 60 mm dishes in Dulbecco's modified Eagle's medium (DMEM)/F12 (Thermo Fisher) supplemented with fetal bovine serum (Thermo Fisher), penicillin/streptomycin, and 100 nM dexamethasone (Sigma Aldrich). After 4 h, the medium was changed to DMEM/F12 containing penicillin/streptomycin and then cells were harvested after 18h.

2.4. Preparation of protein extracts and immunoblotting

ROR α deletion in the liver was evaluated by extracting proteins from the snap-frozen liver or primary cells with lysis buffer, as described previously [26] and immunoblotting with anti-ROR α (Abcam, ab60134, 1:300 dilution), GAPDH (Cell Signaling #2118, 1:1000 dilution), and anti-actin (Cell Signaling #4970, 1:1000 dilution) antibody.

2.5. Quantitative RT-PCR

Total RNA was isolated using RNeasy kit with DNase treatment (Qiagen) and cDNA templates were generated using the high-capacity




Figure 1: Generation of liver-specific ROR α knock out mice. Schematic representation of the *Rora* gene-targeting strategy (A, see methods section for details). Expression level of *Rora* in the liver, ileum, white adipose tissue (WAT), and brain in ROR α ^{fl/fl} and ROR α ^{LKO} mice (B, 3–8 mice per group). Protein expression level of ROR α and actin in liver extract from ROR α ^{fl/fl} and ROR α ^{LKO} mice (C). Data are plotted as mean \pm SEM. ***p < 0.001.




Figure 2: RORα hepatic deletion does not affect glucose and lipid metabolism in mice fed chow diet. Body weight (A), body composition (B), relative organ weights (C), liver cholesterol, cholesteryl esters, triglycerides levels (D), fasting glucose (E) and insulin levels (F), intraperitoneal glucose (G), insulin (H), and pyruvate (I) tolerance test in RORα^{fl/fl} and RORα^{LKO} male mice fed 12 weeks chow diet (n = 7–8 mice per group). Data are plotted as mean – SEM.

cDNA reverse transcription kit using random hexamers (Applied Biosystems). qRT-PCR assays were performed in 10 μl reactions with SYBR Green Master Mix buffer (Thermo Scientific). Gene expression data were normalized to the ribosomal protein L32. Primer sequences used are listed in [Supplementary Table 1](#) (Sigma–Aldrich).

2.6. Statistical analysis

Data are presented as mean – SEM. Analyses between groups were performed by Student’s t-test using GraphPad Prism 7 software.

3. RESULTS

3.1. Generation of mice with a hepatocyte specific RORα deletion

We generated mice with a hepatocyte specific *Rora* deletion of by introducing *loxP* sites flanking exon 3 of the *Rora* allele and crossing them with mice expressing Cre under the albumin promoter (Figure 1A and [Supplementary Fig. 1A](#)).

Analyses of *Rora* expression by qRT-PCR in the liver, ileum, white adipose tissue (WAT), and brain, as well as RORα protein levels by Western blotting, confirmed a selective and specific liver deletion in RORα^{LKO} mice compared to littermate RORα^{fl/fl} mice (Figure 1B–C). Liver RT-qPCR analysis showed that there was a trend (p = 0.057) towards a compensatory increased expression of *Rorc* by the selective and specific liver RORα deletion ([Supplementary Fig. 1B](#)). *Rorb* was not detectable in the livers (data not shown). Moreover, the expression of *Rev-erba*, one of the known *RORs* target genes, was reduced by the selective and specific liver RORα deletion ([Supplementary Fig. 1B](#)). The expression of other *RORs* target genes (i.e. *Rev-erbb*, *Pparg*, *Cyp7b1*, and *Lxra*) was not affected by the selective and specific liver RORα deletion ([Supplementary Fig. 1B](#)).

To further confirm a specific hepatocyte RORα deletion, we isolated primary hepatocytes from RORα^{LKO} mice and littermate RORα^{fl/fl} mice. Analyses of *Rora* expression by qRT-PCR as well as RORα protein levels by Western blotting, confirmed a specific RORα

hepatocyte deletion in $ROR\alpha^{LKO}$ mice compared to littermate $ROR\alpha^{fl/fl}$ mice (Supplementary Figs. 1C–D). Mice with a specific liver $ROR\alpha$ deletion displayed a normal fertility and fecundity generating an equal number of male and female pups during $ROR\alpha^{LKO}$ and $ROR\alpha^{fl/fl}$ cross-breeding.

3.2. $ROR\alpha$ hepatic deletion does not affect glucose and lipid metabolism in mice fed chow diet

Weight gain was monitored of 8-week-old male $ROR\alpha^{LKO}$ and littermate $ROR\alpha^{fl/fl}$ mice on chow diet for 12 weeks. We did not observe any significant differences in weight gain (Figure 2A), whole body adiposity determined by MRI (Figure 2B) or liver, WAT, spleen and cecum weights (Figure 2C). Liver RT-qPCR analysis as well as liver lipid measurements did not reveal any differences in lipid metabolism genes nor in cholesterol, cholesteryl ester or triglyceride levels between $ROR\alpha^{LKO}$ and $ROR\alpha^{fl/fl}$ littermates (Figure 2D and Supplementary Fig. 2).

Fasting glucose and insulin levels as well as glucose levels during glucose-, insulin-, and pyruvate tolerance tests did not demonstrate any differences between $ROR\alpha^{LKO}$ and $ROR\alpha^{fl/fl}$ littermates (Figure 2E–I). Similar findings were obtained for female mice (see Supplementary Fig. 3).

Taken together, our findings suggest that hepatocyte $ROR\alpha$ deletion does not affect glucose and lipid metabolism on chow diet neither in males nor in females.

3.3. $ROR\alpha$ hepatic deletion does not affect glucose and lipid metabolism in mice fed Western style diet

To further study the effect of $ROR\alpha$ on metabolic associated disorders, we fed 8-week-old $ROR\alpha^{LKO}$ and littermate $ROR\alpha^{fl/fl}$ male mice with a WSD for 12 weeks. We did not observe any significant differences in weight gain (Figure 3A), whole body adiposity measured with MRI (Figure 3B) or liver, WAT, spleen or cecum weights (Figure 3C) suggesting that $ROR\alpha$ does not affect diet-induced obesity. Liver RT-qPCR




Figure 3: $ROR\alpha$ hepatic deletion does not affect glucose and lipid metabolism in mice fed western-style diet (WSD). Body weight (A), body composition (B), relative organ weights (C), liver cholesterol, cholesteryl esters, triglycerides levels (D), fasting glucose (E) and insulin levels (F) intraperitoneal glucose (G) insulin (H) and pyruvate (I) tolerance test in $ROR\alpha^{fl/fl}$ and $ROR\alpha^{LKO}$ male mice fed 12 weeks WSD ($n = 8-12$ mice per group). Data are plotted as mean \pm SEM.

analysis as well as liver lipid measurements did not reveal any differences in lipid metabolism genes nor in cholesterol, cholesteryl esters or triglyceride levels between ROR α ^{LKO} and ROR α ^{fl/fl} littermates (Figure 3D and Supplementary Fig. 4).

Next, we assessed whether ROR α contributes to glucose metabolism during WSD by analyzing fasting glucose and insulin levels as well as performing glucose-, insulin- and pyruvate tolerance tests. We did not observe any differences between ROR α ^{LKO} and ROR α ^{fl/fl} littermates (Figure 2E–I). Similar findings were obtained when we used female mice (see Supplementary Fig. 5).

Taken together, our findings suggest that a hepatocyte ROR α deletion does not affect glucose and lipids metabolism on WSD in both males and females.

4. DISCUSSION

NRs have been shown to play a crucial role in the regulation of glucose and lipid metabolism. Evidences from the ROR α whole body knockout (Staggerer) mice have pointed towards ROR α as a potential molecular target for the treatment of metabolic diseases [14–16]. However, the improved metabolic phenotype of Staggerer mice is highly affected by the staggering phenotype induced by the cerebellum ROR α deletion [14,15], thus it is difficult to evaluate the role of ROR α in metabolic active tissues, such as the liver. Here, by generating a liver specific ROR α knockout mouse model, we did not detect any effects of ROR α liver deletion on glucose or lipids metabolism on chow diet or after a WSD exposure.

Our data are in contrast with recent published data that showed that liver-specific ablation of ROR α dramatically worsens metabolic profiles in mice eg. increased adiposity, liver fat accumulation, inflammation, and hepatic insulin resistance, when mice are fed a high fat diet [20,21]. Several factors should be taken into account when comparing the discrepancy between the distinct phenotype reported by other groups and our present study. First, we targeted ROR α exon 3 while the Cre-lox system used by others targeted exon 4 and 4–5, leaving the exon 3 that encodes the beginning of the DNA binding domain. Although the stability of potentially produced recombinant proteins, which in our experience usually is low, we cannot exclude that the generation of different truncated proteins containing a portion of the DNA binding domain may explain some of the divergent results between the studies. Importantly, we demonstrated an almost complete elimination of ROR α in hepatocytes similar to other groups and also a significant effect on some of the known ROR α target genes [20,21,27,28]. Second, we challenged our mice with WSD while other reports used HFD. Importantly, it has been well-documented that, despite containing fewer calories, WSD induces more pronounced steatosis and liver inflammation compared with HFD [29,30]. Thus, it may be plausible that in our model, the effect of WSD on steatosis and inflammation overwhelms the effect of ROR α liver deletion.

In support to our findings Zhang et al. recently reported that the single knockout of ROR α or ROR γ in hepatocyte has almost no effect on liver gene expression while the double deletion of both ROR α and ROR γ in hepatocytes has a substantial impact on gene expression profiles, increases lipogenesis and therefore predisposes to HFD-induced steatosis [31]. ROR α and ROR γ bind to the same response element and are both highly expressed in liver. Thus, deletion of only one of them may not be sufficient to induce a metabolic phenotype due to their functional redundancy. In our experimental model upon deletion of ROR α , we observed a trend towards increased *Rorc* expression, which may be a compensation for the lack of ROR α activity. Taken

together, it is thus likely that double targeting of ROR α and ROR γ may be required for achieving maximum effect.

Moreover, it is known that ROR α is a clock gene involved in circadian rhythm regulation [31,32]. In an attempt to reduce experimental variation all metabolic phenotyping was performed at the same time of the day. Thus we cannot exclude that the discrepancy between our data and previous published data may be due to differences in experimental set up such as time of analyses.

5. CONCLUSIONS

In conclusion, our data show that the deletion of ROR α in the liver does not affect glucose or lipid metabolism during WSD in mice. Further studies testing the role of liver ROR α or the double ROR α and ROR γ deletion in different dietary settings are needed to further elucidate the role of ROR α on obesity and metabolic associated disorders.

AUTHOR CONTRIBUTIONS

AM, BS, and FB: conceived the project and designed the experiments. AM, RC, AK, LL, and MS: performed and analyzed experiments. AM and FB wrote the paper. All authors commented and approved the paper.

ACKNOWLEDGEMENTS

The authors thank Zacharias Gulic for superb technical assistance and prof. Martin Bergö for kind donation of ROSA^{MT/mG} mice. This study was supported by the Novo Nordisk Foundation, the Swedish Research Council, Swedish Diabetes Foundation, Swedish Heart-Lung Foundation, Göran Gustafsson's Foundation, Inga-Britt och Arne Lundberg's Foundation, Knut and Alice Wallenberg Foundation, the regional agreement on medical training and clinical research (ALF) between Region Västra Götaland and Sahlgrenska University Hospital. FB is Torsten Söderberg Professor in Medicine and recipient of an ERC Consolidator Grant (European Research Council, Consolidator grant 615362 – METABASE). BS holds an ERC Advanced Grant 694717.

CONFLICTS OF INTERESTS

The authors report no conflict of interest.

APPENDIX A. SUPPLEMENTARY DATA

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.molmet.2019.02.010>.

REFERENCES

- [1] Afshin, A., Forouzanfar, M.H., Reitsma, M.B., Sur, P., Estep, K., Lee, A., et al., 2017. Health effects of overweight and obesity in 195 countries over 25 years. *New England Journal of Medicine* 377(1):13–27.
- [2] Finkelstein, E.A., Khavjou, O.A., Thompson, H., Trogdon, J.G., Pan, L., Sherry, B., et al., 2012. Obesity and severe obesity forecasts through 2030. *American Journal of Preventive Medicine* 42(6):563–570.
- [3] Boyle, J.P., Honeycutt, A.A., Narayan, K.M., Hoerger, T.J., Geiss, L.S., Chen, H., et al., 2001. Projection of diabetes burden through 2050: impact of changing demography and disease prevalence in the U.S. *Diabetes Care* 24(11):1936–1940.
- [4] Sjöström, L., 2013. Review of the key results from the Swedish Obese Subjects (SOS) trial - a prospective controlled intervention study of bariatric surgery. *Journal of Internal Medicine* 273(3):219–234.

- [5] Pearen, M.A., Muscat, G.E., 2012. Orphan nuclear receptors and the regulation of nutrient metabolism: understanding obesity. *Physiology (Bethesda)* 27(3): 156–166.
- [6] Schulman, I.G., 2010. Nuclear receptors as drug targets for metabolic disease. *Advanced Drug Delivery Reviews* 62(13):1307–1315.
- [7] Dhiman, V.K., Bolt, M.J., White, K.P., 2018. Nuclear receptors in cancer — uncovering new and evolving roles through genomic analysis. *Nature Reviews Genetics* 19(3):160–174.
- [8] Germain, P., Staels, B., Dacquet, C., Spedding, M., Laudet, V., 2006. Overview of nomenclature of nuclear receptors. *Pharmacological Reviews* 58(4):685–704.
- [9] Tanaka, N., Aoyama, T., Kimura, S., Gonzalez, F.J., 2017. Targeting nuclear receptors for the treatment of fatty liver disease. *Pharmacology & Therapeutics* 179:142–157.
- [10] Swanson, H.I., Wada, T., Xie, W., Renga, B., Zampella, A., Distrutti, E., et al., 2013. Role of nuclear receptors in lipid dysfunction and obesity-related diseases. *Drug Metabolism & Disposition* 41(1):1–11.
- [11] Jetten, A.M., 2009. Retinoid-related orphan receptors (RORs): critical roles in development, immunity, circadian rhythm, and cellular metabolism. *Nuclear Receptor Signaling* 7:e003.
- [12] Fitzsimmons, R.L., Lau, P., Muscat, G.E., 2012. Retinoid-related orphan receptor alpha and the regulation of lipid homeostasis. *The Journal of Steroid Biochemistry and Molecular Biology* 130(3–5):159–168.
- [13] Marciano, D.P., Chang, M.R., Corzo, C.A., Goswami, D., Lam, V.Q., Pascal, B.D., et al., 2014. The therapeutic potential of nuclear receptor modulators for treatment of metabolic disorders: PPAR γ , RORs, and Rev-erbs. *Cell Metabolism* 19(2):193–208.
- [14] Dussault, I., Fawcett, D., Matthyssen, A., Bader, J.A., Giguère, V., 1998. Orphan nuclear receptor ROR alpha-deficient mice display the cerebellar defects of staggerer. *Mechanisms of Development* 70(1–2):147–153.
- [15] Lau, P., Fitzsimmons, R.L., Raichur, S., Wang, S.C., Lechtken, A., Muscat, G.E., 2008. The orphan nuclear receptor, RORalpha, regulates gene expression that controls lipid metabolism: staggerer (SG/SG) mice are resistant to diet-induced obesity. *Journal of Biological Chemistry* 283(26):18411–18421.
- [16] Kadiri, S., Monnier, C., Ganbold, M., Ledent, T., Capeau, J., Antoine, B., 2015. The nuclear retinoid-related orphan receptor- α regulates adipose tissue glyceroneogenesis in addition to hepatic gluconeogenesis. *American Journal of Physiology Endocrinology and Metabolism* 309(2):E105–E114.
- [17] Han, Y.H., Kim, H.J., Na, H., Nam, M.W., Kim, J.Y., Kim, J.S., et al., 2017. ROR α induces KLF4-mediated M2 polarization in the liver macrophages that protect against nonalcoholic steatohepatitis. *Cell Reports* 20(1):124–135.
- [18] Kim, E.J., Yoon, Y.S., Hong, S., Son, H.Y., Na, T.Y., Lee, M.H., et al., 2012. Retinoic acid receptor-related orphan receptor α -induced activation of adenosine monophosphate-activated protein kinase results in attenuation of hepatic steatosis. *Hepatology* 55(5):1379–1388.
- [19] Han, Y.H., Kim, H.J., Kim, E.J., Kim, K.S., Hong, S., Park, H.G., et al., 2014. ROR α decreases oxidative stress through the induction of SOD2 and GPx1 expression and thereby protects against nonalcoholic steatohepatitis in mice. *Antioxidants and Redox Signaling* 21(15):2083–2094.
- [20] Kim, H.J., Han, Y.H., Na, H., Kim, J.Y., Kim, T., Shin, C., et al., 2017. Liver-specific deletion of ROR α aggravates diet-induced nonalcoholic steatohepatitis by inducing mitochondrial dysfunction. *Scientific Reports* 7(1):16041.
- [21] Kim, K., Boo, K., Yu, Y.S., Oh, S.K., Kim, H., Jeon, Y., et al., 2017. ROR α controls hepatic lipid homeostasis via negative regulation of PPAR γ transcriptional network. *Nature Communications* 8(1):162.
- [22] Molinaro, A., Caesar, R., Holm, L.M., Tremaroli, V., Cani, P.D., Bäckhed, F., 2017. Host-microbiota interaction induces bi-phasic inflammation and glucose intolerance in mice. *Molecular Metabolism* 6(11):1371–1380.
- [23] Homan, R., Anderson, M.K., 1998. Rapid separation and quantitation of combined neutral and polar lipid classes by high-performance liquid chromatography and evaporative light-scattering mass detection. *Journal of Chromatography B Biomedical Sciences and Applications* 708(1–2): 21–26.
- [24] Löfgren, L., Forsberg, G.B., Ståhlman, M., 2016. The BUME method: a new rapid and simple chloroform-free method for total lipid extraction of animal tissue. *Scientific Reports* 6:27688.
- [25] Zhang, W., Sargis, R.M., Volden, P.A., Carmean, C.M., Sun, X.J., Brady, M.J., 2012. PCB 126 and other dioxin-like PCBs specifically suppress hepatic PEPCCK expression via the aryl hydrocarbon receptor. *PLoS One* 7(5):e37103.
- [26] Koh, A., Lee, M.N., Yang, Y.R., Jeong, H., Ghim, J., Noh, J., et al., 2013. C1-Ten is a protein tyrosine phosphatase of insulin receptor substrate 1 (IRS-1), regulating IRS-1 stability and muscle atrophy. *Molecular and Cellular Biology* 33(8):1608–1620.
- [27] Wada, T., Kang, H.S., Angers, M., Gong, H., Bhatia, S., Khadem, S., et al., 2008. Identification of oxysterol 7 α -hydroxylase (Cyp7b1) as a novel retinoid-related orphan receptor alpha (RORalpha) (NR1F1) target gene and a functional cross-talk between RORalpha and liver X receptor (NR1H3). *Molecular Pharmacology* 73(3):891–899.
- [28] Delerive, P., Chin, W.W., Suen, C.S., 2002. Identification of Rev-erb(alpha) as a novel ROR(alpha) target gene. *Journal of Biological Chemistry* 277(38): 35013–35018.
- [29] Bortolin, R.C., Vargas, A.R., Gasparotto, J., Chaves, P.R., Schnorr, C.E., Martinello, K.B., et al., 2017. A new animal diet based on human Western diet is a robust diet-induced obesity model: comparison to high-fat and cafeteria diets in term of metabolic and gut microbiota disruption. *International Journal of Obesity (Lond)* 42(3):525–534.
- [30] Fleissner, C.K., Huebel, N., Abd El-Bary, M.M., Loh, G., Klaus, S., Blaut, M., 2010. Absence of intestinal microbiota does not protect mice from diet-induced obesity. *British Journal of Nutrition* 104(6):919–929.
- [31] Zhang, Y., Papazyan, R., Damle, M., Fang, B., Jager, J., Feng, D., et al., 2017. The hepatic circadian clock fine-tunes the lipogenic response to feeding through ROR α/γ . *Genes & Development* 31(12):1202–1211.
- [32] Zhao, X., Cho, H., Yu, R.T., Atkins, A.R., Downes, M., Evans, R.M., 2014. Nuclear receptors rock around the clock. *EMBO Reports* 15(5):518–528.