

Non-contiguous finished genome sequence and description of *Bacillus algeriensis* sp. nov.

Esma Bendjama^{1,2,3}, Lotfi Loucif^{1,2,3}, Seydina M. Diene¹, Caroline Michelle¹, Djamilia Gacemi-Kirane², Jean-Marc Rolain^{1*}

¹ Unité de recherche sur les maladies infectieuses et tropicales émergentes (URMITE), UMR CNRS, IHU Méditerranée Infection, Faculté de Médecine et de Pharmacie, Aix-Marseille-Université, Marseille, France.

² Département de Biochimie, Faculté des Sciences, Université Badji Mokhtar, Annaba, Algérie

³ Département des Sciences Biologiques, Faculté des Sciences, Université El Hadj Lakhdar, Batna, Algérie.

*Correspondence: Jean-Marc Rolain (jean-marc.rolain@univ-amu.fr)

Keywords: *Bacillus algeriensis*, hypersaline environments, sediments, genome, taxonomics

Strain EB01^T sp. nov. is the type strain of *Bacillus algeriensis*, a new species within the genus *Bacillus*. This strain, whose genome is described here, was isolated from sediment sample of the hypersaline lake Ezzemoul sabkha in northeastern Algeria. *B. algeriensis* is a facultative anaerobic Gram-positive bacillus. Here we describe the features of this organism, together with the complete genome sequence and annotation. The 5,269,577 bp long genome contains 5,098 protein-coding and 95 RNA genes, including 12 rRNA genes.

Introduction

Bacillus algeriensis sp. nov. strain EB01^T (= CSUR P857 = DSM 27334) is the type strain of *B. algeriensis* sp. nov. It is a new Gram-positive, facultatively anaerobic, motile, indole-negative, rod shaped bacterium with rounded ends. It was isolated from a sediment sample from the hypersaline lake Ezzemoul sabkha in the Oum-El-Bouaghi region in northeastern Algeria, which is an important wintering and resting site for several species of waterbirds, including the Greater Flamingo. This site is one of the Ramsar convention wetlands (<http://www.ramsar.org>). The genus *Bacillus* was created by Cohn about 142 years ago [1], and mainly comprises Gram-positive, rod-shaped, aerobic or facultatively anaerobic, spore-forming bacteria. The genus includes 279 species and 7 subspecies with validly published names [2]. Members of *Bacillus* genus are ubiquitous in nature, ranging from freshwater to marine sediments and from hot springs and desert sands to Arctic soils; many strains have been isolated from the gastrointestinal tracts of various insects and animals, from vegetation and from food [3]. *Bacillus* strains are biotechnologically priceless be-

cause of their high capacity to produce a wide range of antimicrobial compounds, enzymes and other metabolites that can be used in industry [4,5]. Some species of *Bacillus* are pathogenic, such as *B. anthracis* (responsible for causing anthrax) [6] and *B. cereus* (a major cause of food poisoning) [7]. Others are opportunists in immunocompromised patients, and may also be involved in various human infections, including pneumonia, endocarditis, ocular, cutaneous, bone or central nervous system infections and bacteraemia [8]. The current bacterial taxonomy is based on a combination of various phenotypic and genetic criteria [9,10]. However, the three essential genetic criteria that are used, comprising 16S rRNA gene based phylogeny [11], G+C content, and DNA-DNA hybridization [10,12] exhibit several drawbacks. As a result of the recent decrease in the cost of genomic sequencing, it has been proposed that whole genome sequencing information and MALDI-TOF spectrum [13] be combined with the main phenotypic characteristics as a polyphasic approach strategy (taxonomics) to describe new bacterial taxa [14-26].

Here we present a summary classification and a set of features for *B. algeriensis* sp. nov. strain EB01^T together with the description of the complete genome sequence and annotation. These characteristics support the circumscription of the species *B. algeriensis*.

Classification and features

In July 2012, a sediment sample was aseptically collected in sterile bottles, 15 cm below the evaporite crust of the hypersaline lake Ezzemoul sabkha of Oum-El-Bouaghi region in northeastern Algeria. Samples were transferred in a cooler (4°C) to our lab in Algeria. Samples were processed the same day. Sediments were diluted 1:10 v/v with sterile saline water (0.9% NaCl) and vigorously shaken, tenfold serial dilutions (10⁻¹-10⁻⁵)

of the sediment suspension were plated in Nutrient Agar (NA) medium (meat extract 1 g/l, peptone 5 g/l, yeast extract 2 g/l, sodium chloride 5 g/l, agar 15 g/l) and the plates were incubated at 30°C for 24-72 h. In order to obtain a pure culture, colonies were transferred to fresh NA medium. *Bacillus algeriensis* sp. nov. strain EB01^T (Table 1) was isolated in July 2012 by cultivation under aerobic conditions at 30°C. This strain exhibited a 97.0% 16S rRNA nucleotide sequence similarity with *Bacillus subterraneus* type strain DSM13966^T (Figure 1), the phylogenetically closest validly published *Bacillus* species. These values were lower than the 98.7% 16S rRNA gene sequence threshold recommended by Stackebrandt and Ebers to delineate a new species without carrying DNA DNA hybridization [11].

Table 1. Classification and general features of *Bacillus algeriensis* strain EB01^T

| MIGS ID | Property | Term | Evidence code ^a |
|----------|------------------------|-------------------------------------|----------------------------|
| | | Domain <i>Bacteria</i> | TAS [27] |
| | | Phylum <i>Firmicutes</i> | TAS [28-30] |
| | | Class <i>Bacilli</i> | TAS [31,32] |
| | Current classification | Order <i>Bacillales</i> | TAS [33,34] |
| | | Family <i>Bacillaceae</i> | TAS [33,35] |
| | | Genus <i>Bacillus</i> | TAS [1,33,36] |
| | | Species <i>Bacillus algeriensis</i> | IDA |
| | | Type strain EB01 ^T | IDA |
| | Gram stain | Positive | IDA |
| | Cell shape | Rod-shaped | IDA |
| | Motility | Motile | IDA |
| | Sporulation | Sporulating | IDA |
| | Temperature range | Between 37°C and 55°C | IDA |
| | Optimum temperature | 37°C | IDA |
| | | Growth in LB medium + 0-2.5% | |
| MIGS-6.3 | Salinity | NaCl | IDA |
| MIGS-22 | Oxygen requirement | Facultative anaerobic | IDA |
| | Carbon source | Unknown | NAS |
| | Energy source | Unknown | NAS |
| MIGS-6 | Habitat | Hypersaline sediment sample | IDA |
| MIGS-15 | Biotic relationship | Free living | IDA |
| | Pathogenicity | Unknown | NAS |
| | Biosafety level | 2 | NAS |
| MIGS-14 | Isolation | Sediment of Ezzemoul Sabkha Lake | IDA |
| MIGS-4 | Geographic location | Algeria | IDA |
| | Sample collection | July 2012 | |
| MIGS-5 | time | | IDA |
| MIGS-4.1 | Latitude | 35.88167 | IDA |
| MIGS-4.1 | Longitude | 6.503272 | IDA |
| MIGS-4.3 | Depth | Unknown | NAS |
| MIGS-4.4 | Altitude | 800 m | IDA |

^aEvidence codes - IDA: Inferred from Direct Assay, TAS: Traceable Author Statement (i.e., a direct report exists in the literature), NAS: Non traceable Author Statement (i.e., not directly observed for the living, isolated sample, but based on a generally accepted property for the species, or anecdotal evidence). These evidence codes are from the Gene Ontology project [37]. If the evidence is IDA, then the property was directly observed for a live isolate by one of the authors or an expert mentioned in the acknowledgements.




Figure 1. A consensus phylogenetic tree based on 16S rRNA gene sequence comparisons, highlighting the position of strain EB01^T *Bacillus algeriensis* relative to other type strains within the *Bacillus* genus. GenBank accession numbers are displayed in parentheses. Sequences were aligned using CLUSTALW, and phylogenetic inferences made using the neighbor-joining method [38] within the MEGA 5 software [39]. Numbers above the nodes are percentages of bootstrap values from 1,000 replicates that support the node. *Paenibacillus polymyxa* was used as the outgroup. The scale bar represents 0.01 substitutions per nucleotide position.

Six different growth temperatures (25, 30, 37, 45, 50 and 55°C), nine NaCl concentrations (0, 2.5, 5, 7.5, 10, 15, 20, 25, 30%) and ten pHs (5, 6, 6.5, 7, 7.5, 8, 8.5, 9, 10, 11) were tested. Growth occurred at all tested temperatures, however the optimal growth was observed at 37°C, between 0% and 2.5% NaCl concentration and pH in the range of 6.5-9 (optimum at pH 7). Colony morphology was observed on sheep blood agar (BioMerieux) after 24 h of aerobic incubation under optimal growth conditions, the colonies of strain EB01^T were circular, light yellow, smooth and 2 mm in diameter. Growth of the strain was tested in anaerobic and microaerophilic atmospheres using GasPak EZ Anaerobe Pouch (Becton, Dickinson and Compa-

ny) and CampyGen Compact (Oxoid) systems, respectively, and in aerobic atmosphere, with or without 5% CO₂. Growth was achieved under aerobic (with and without CO₂) and microaerophilic conditions but weak growth was observed under anaerobic conditions. Gram staining showed Gram-positive rods (Figure 2). Cells grown on agar sporulate. A motility test was positive. The size of cells were determined by negative staining transmission electron microscopy on a Technai G² Cryo (FEI) at an operating voltage of 200 kV, the rods have a length ranging from 2.4 μm to 4.9 μm (mean 3.6 μm) and a diameter ranging from 0.7 μm to 1.1 μm (mean 0.8 μm) (Figure 3).




Figure 2. Gram stain of *B. algeriensis* strain EB01^T.




Figure 3. Transmission electron micrograph of *B. algeriensis* strain EB01^T made using a Technai G² Cryo (FEI) at an operating voltage of 200 kV. The scale bar represents 500 nm.

Strain EB01^T exhibited catalase activity but oxidase activity was negative. Using the commercially available API 50CH system (BioMerieux) according to the manufacturer's instructions, a weak positive reaction was observed for D-ribose, D-glucose, D-fructose, methyl α -D-glucopyranoside, N-acetylglucosamine, D-maltose, D-lactose, D-melibiose, D-saccharose, D-trehalose, D-tagatose, and hydrolysis of starch. Other tests were negative. Using the API ZYM system (BioMerieux), positive reactions were observed for alkaline phosphatase, esterase (C4), esterase lipase (C8), leucine arylamidase, α -chymotrypsin, β -glucuronidase, α -glucosidase, N-acetyl-glucosaminidase and a weak positive reaction was observed for acid phosphatase. The nitrate reduction and β -galactosidase reaction was also positive, but urease and indole production were negative. *B. algeriensis* was susceptible to amoxicillin, nitrofurantoin, erythromycin, doxycycline, rifampicin, vancomycin, gentamicin, imipenem, trimethoprim-sulfamethoxazole, ciprofloxacin, ceftriaxone and amoxicillin-clavulanic acid, but resistant to nalidixic acid.

When compared to other *Bacillus* species [40-48], *Bacillus algeriensis* sp. nov. strain EB01^T exhibited the phenotypic differences detailed in (Table 2).

Matrix-assisted laser-desorption/ionization time-of-flight (MALDI-TOF) MS protein analysis was performed as previously described [26,49,50]. Briefly, strain EB01^T was plated on 5% sheep blood-enriched Columbia agar (BioMerieux) and incubated for 24 h at 37°C. Isolated bacterial colo-

nies were picked, and then deposited as a thin film in 12 replicates on a MALDI-TOF steel target plate (Bruker Daltonics, Bremen, Germany). The plates were allowed to dry at room temperature. Each deposit was overlaid with 1.5 μ l of matrix solution containing α -cyano-hydroxycinnamic acid (Sigma, Saint-Quentin Fallavier, France) saturated with 50% acetonitrile, 2.5% trifluoroacetic acid and high-performance liquid chromatography (HPLC)-grade water, and allowed to co-crystallize with the sample. Measurements were conducted using the Microflex LT spectrometer (Bruker Daltonics). Spectra were recorded in the linear positive ion mode over a mass range of 2 to 20 kDa. The acceleration voltage was 20 kV. Spectra were collected as a sum of 240 shots across a spot. The 12 EB01^T spectra were imported into the MALDI BioTyper software (version 3.0, Bruker) and analyzed by standard pattern matching (with default parameter settings) against 6,335 bacterial spectra including 210 spectra from 104 *Bacillus* species, used as reference data, in the BioTyper database. A score enabled the identification, or not, from the tested species: a score > 2 with a validated species enabled the identification at the species level, a score > 1.7 but < 2 enabled the identification at the genus level; and a score < 1.7 did not enable any identification. For strain EB01^T, the scores obtained ranged from 1.15 to 1.60 thus suggesting that our isolate was a new species. We added the spectrum from strain EB01^T (Figure 4) to our database. Spectrum differences with other of *Bacillus* species are shown in (Figure 5).

Table 2. Differential phenotypic characteristics between *B. algeriensis* sp. nov. strain EB01^T and phylogenetically close *Bacillus* species[†].

| Characteristic | <i>B. ma</i> | <i>B. su</i> | <i>B. fo</i> | <i>B. je</i> | <i>B. th</i> | <i>B. bo</i> | <i>B. ba</i> | <i>B. ne</i> | <i>B. kr</i> |
|---------------------------------|-----------------------|-----------------------|--------------|-----------------------|--------------|--------------|-----------------------|-----------------------|--------------|
| Cell-diameter(µm) | 0.7-1.1 | 0.5-0.8 | 1 | 0.8-1.1 | 0.5-0.7 | 0.5-0.9 | 0.7-1.2 | 1 | 1.4-2.0 |
| Oxygen requirement | facultative anaerobic | facultative anaerobic | aerobic | facultative anaerobic | aerobic | na | facultative anaerobic | facultative anaerobic | aerobic |
| Gram strain | + | - | + | V | V | + | + or V | + | + |
| NaCl range (% w/v) | 0-2.5 | 0-9 | 0-3 | 0-13 | 0-5 | 0-7 | na | 0-8 | 0-6 |
| Motility | + | + | na | + | + | + | + | + | + |
| Endospore formation | + | - | - | + | + | + | + | + | + |
| Production of | | | | | | | | | |
| Alkaline phosphatase | + | na | na | na | na | + | na | na | na |
| Acid phosphatase | w | na | na | na | na | na | na | na | na |
| Catalase | + | + | + | + | + | + | na | + | + |
| Oxidase | - | - | + | - | - | + | na | na | - |
| Nitrate reductase | + | + | + | + | + | - | + | - | - |
| Urease | - | - | + | + | - | - | - | - | - |
| α-galactosidase | - | na | na | na | na | na | na | na | na |
| β-galactosidase | - | - | + | na | na | - | na | + | na |
| β-glucuronidase | + | na | na | na | na | na | na | na | + |
| N-acetyl-β-glucosaminidase | + | na | na | na | na | na | na | na | na |
| Indole | - | - | na | - | - | - | - | - | na |
| Esterase | + | na | na | na | na | na | na | na | + |
| Esterase lipase | + | na | na | na | na | w | na | na | + |
| Naphthyl-AS-BI-Phosphohydrolase | - | na | na | na | na | na | na | na | + |
| Leucine arylamidase | + | na | na | na | na | + | na | na | na |

[†](*Bacillus subterraneus* strain COO13B^T, *Bacillus foraminis* strain CV53^T, *Bacillus jeotgali* strain YK1-10^T, *Bacillus thioparans* strain BMP-1^T, *Bacillus boroniphilus* strain T-15Z^T, *Bacillus bataviensis* strain IDA1115^T, *Bacillus nealsonii* strain FO-92^T, *Bacillus kribbensis* strain BT080^T).

B. algeriensis (*B. ma*), *B. subterraneus* (*B. su*), *B. foraminis* (*B. fo*), *B. jeotgali* (*B. je*), *B. thioparans* (*B. th*), *B. boroniphilus* (*B. bo*), *B. bataviensis* (*B. ba*), *B. nealsonii* (*B. ne*) and *B. kribbensis* (*B. kr*).

+: positive result, -: negative result, var: variable, w: weak positive result, na: data not available.

Table 2. (cont.) Differential phenotypic characteristics between *B. algeriensis* sp. nov. strain EB01^T and phylogenetically close *Bacillus* species[†].

| Characteristic | B. ma | B. su | B. fo | B. je | B. th | B. bo | B. ba | B. ne | B. kr |
|---|-------|---|-------------------------------------|----------------------------|--|-------------------|--------------|---|------------|
| Production of (cont.) | | | | | | | | | |
| Cystine arylamidase | - | na | na | na | na | w | na | na | na |
| Valine arylamidase | - | na | na | na | na | w | na | na | + |
| Utilization of | | | | | | | | | |
| D-mannose | - | - | + | - | - | + | + | + | - |
| Amygdalin | - | na | + | - | - | + | w | - | na |
| L-Arabinose | - | - | + | - | na | na | - | + | - |
| Cellobiose | - | - | + | + | na | + | + | + | + |
| Lactose | w | - | + | - | na | na | + | + | + |
| D-xylose | - | + | + | - | na | na | - | + | + |
| D-Glucose | w | + | + | + | + | na | + | + | + |
| Mannitol | - | - | + | - | - | na | + | + | na |
| Arabinose | - | - | + | - | - | na | - | - | na |
| L-Xylose | - | + | + | - | na | na | - | na | + |
| Glycerol | - | + | + | - | na | + | w | + | - |
| D-Galactose | - | - | + | - | na | na | + | + | na |
| Hydrolysis of | | | | | | | | | |
| Starch | w | + | + | + | na | na | v | + | - |
| Gelatin | - | + | + | + | w | - | + | - | + |
| G+C content (mol%) | 42,22 | 43±1 deep subterranean thermal waters | 43,1 alkaline ground water | 41 fermented seafood | 43,8 wastewater treatment culture system | 41,1-42,2 soil | 39,6 soil | na spacecraft- assembly facility | na soil |
| Habitat | | | | | | | | | |
| [†] (<i>Bacillus subterraneus</i> strain COO13B ^T , <i>Bacillus foraminis</i> strain CV53 ^T , <i>Bacillus jeotgali</i> strain YKJ-10 ^T , <i>Bacillus thioparans</i> strain BMP-1 ^T , <i>Bacillus boroniphilus</i> strain T-15Z ^T , <i>Bacillus bataviensis</i> strain IDA1115 ^T , <i>Bacillus nealsonii</i> strain FO-92 ^T , <i>Bacillus kribbensis</i> strain BT080 ^T). | | | | | | | | | |
| <i>B. algeriensis</i> (B. ma), <i>B. subterraneus</i> (B. su), <i>B. foraminis</i> (B. fo), <i>B. jeotgali</i> (B. je), <i>B. thioparans</i> (B. th), <i>B. boroniphilus</i> (B. bo), <i>B. bataviensis</i> (B. ba), <i>B. nealsonii</i> (B. ne) and <i>B. kribbensis</i> (B. kr). | | | | | | | | | |
| +: positive result, -: negative result, var: variable, w: weak positive result, na: data not available. | | | | | | | | | |




Figure 4. Reference mass spectrum from *B. algeriensis* strain EB01^T. Spectra from 12 individual colonies were compared and a reference spectrum was generated.




Figure 5. Gel view comparing *Bacillus algeriensis* EB01^T spectra with other members of the *Bacillus* genus (*B. weihenstephanensis*, *B. vallismortis*, *B. thuringiensis*, *B. thioparans*, *B. subtilis subsp. subtilis*, *B. subterraneus*, *B. shackletonii*, *B. novalis*, *B. niacini*, *B. nealsonii*, *B. jeotgali*, *B. flexus*, *B. circulans*, *B. bataviensis* and *B. asahii*). The Gel View displays the raw spectra of all loaded spectrum files as a pseudo-electrophoretic gel. The x-axis records the m/z value. The left y-axis displays the running spectrum number originating from subsequent spectra loading. The peak intensity is expressed by a grey scale scheme code. The grey scale bar on the right y-axis indicates the relation between the shade of grey a peak is displayed with and the peak intensity in arbitrary units.

Genome sequencing information

Genome project history

The organism was selected for sequencing on the basis of its phylogenetic position and 16S rRNA similarity to other members of the genus *Bacillus*, and is part of a study of *Bacillus* genus diversity in hypersaline lakes of Algeria. It was the 398th ge-

nome of a *Bacillus* species and the first genome of *Bacillus algeriensis* sp. nov. The EMBL accession number is ERP003483 and consists of 46 contigs. Table 3 shows the project information and its association with MIGS version 2.0 compliance [51].

Table 3. Project information

| MIGS ID | Property | Term |
|-----------|----------------------|---|
| MIGS-31 | Finishing quality | High-quality draft |
| MIGS-28 | Libraries used | Nextera XT library |
| MIGS-29 | Sequencing platform | Miseq-Illumina |
| MIGS-31.2 | Sequencing coverage | 34× |
| MIGS-30 | Assemblers | Velvet |
| MIGS-32 | Gene calling method | Prodigal |
| | EMBL Date of Release | January 10, 2014 |
| | EMBL ID | ERP003483 |
| MIGS-13 | Project relevance | Study of the <i>Bacillus</i> genus diversity in hypersaline lakes of northeastern Algeria |

Growth conditions and DNA isolation

Bacillus algeriensis sp. nov. strain EB01^T, was grown aerobically on 5% sheep blood enriched Columbia agar at 37°C. Three Petri dishes were spread and resuspended in a 2 ml sterile Eppendorf tube containing 1ml of TE buffer with acid-washed glass beads (diameter ≤106 μm, Sigma, Saint-Quentin Fallavier, France). Three cycles of shaking were performed using a FastPrep BIO 101 apparatus (Qbiogene, Strasbourg, France) for 15 sec at level 6.5 (full speed). Then, the supernatant was placed in a new tube along with one hundred μl of 10% SDS and 50 μl of Proteinase K (Qiagen GmbH, Hilden, Germany) and incubated over night at 56°C. The digested mixture was used to perform DNA extraction using the classical phenol-chloroform method. The quality of the DNA was checked on an agarose gel (0.8%) stained with SYBR safe.

Genome sequencing

Genomic DNA of *B. algeriensis* sp. nov. strain EB01^T was sequenced on the MiSeq platform (Illumina, Inc, San Diego CA 92121, USA) with a paired end and barcode strategy in order to be mixed with 7 others genomic projects constructed with the Nextera XT library kit (Illumina).

The gDNA was quantified by a Qubit assay with the high sensitivity kit (Life technologies, Carlsbad, CA, USA) to 34.4 ng/μL and dilution was performed to provide 1 ng of each small genome as input. The “tagmentation” step fragmented and tagged the DNA to generate an optimum insert

size of 1.6 kb, validated on a high sensitivity labchip Calliper-Perkin Elmer (Caliper Life Sciences, Inc, Massachusetts, USA). Then limited cycle PCR amplification completed the tags adapters and introduced dual-index barcodes. After purification on Ampure beads (Lifetechnologies, Carlsbad, CA, USA), the libraries were normalized on specific beads according to the Nextera XT protocol (Illumina). Normalized libraries are pooled into a single library for sequencing on the MiSeq. The pooled single strand library was loaded onto the reagent cartridge and then onto the instrument along with the flow cell. Automated cluster generation and paired-end sequencing with dual index reads was performed in a single 39-hour run with a 2×250 bp read length. Within this pooled run, the index representation was determined to 7.1%. Total information of 2.4 G bases was obtained from a 320 K/mm² density with 94.9% (5,757,000 clusters) of the clusters passing quality control (QC) filters. From the genome sequencing process, the 775,420 produced Illumina reads for *B. algeriensis* EB01^T were filtered according to the read qualities and sizes using the fastq-mcf program (Ea-utils: command-line tools for processing biological sequencing data) [52]. 714,540 filtered read sequences were kept for genome assembly. The Velvet assembler was used with different kmer values (from 51 to 95) and the best assembly result with kmer value (n=91) producing 46 contigs with sizes from 872 bp to 409,112 bp, was retained for genome annotation.

Genome annotation

Open Reading Frames (ORFs) were predicted using Prodigal [53] with default parameters. The predicted bacterial protein sequences were searched against the Clusters of Orthologous Groups (COG) database and the GenBank database [54] using BLASTP. Ribosomal RNAs were found by using RNAmmer 2.1 server [55,56] and BLASTn against the GenBank database, whereas the tRNAscanSE tool [57] was used to find tRNA genes. Transmembrane helices and lipoprotein signal peptides were predicted using phobius web server [58]. ORFans were identified if their BLASTP *E*-value was lower than $1e-03$ for alignment length greater than 80 amino acids. If alignment lengths were smaller than 80 amino acids, we used an *E*-value of $1e-05$. Artemis [59] was used for data management and DNA Plotter [60] was used for visualization of genomic features. To estimate the mean level of nucleotide sequence similarity at the genome level between *B. algeriensis* sp nov. strain EB01^T and seven other *Bacillus* species, we use the Average Genomic Identity of Orthologous gene Sequences (AGIOS) in-house software. Briefly, this software combines

the Proteinortho software [61] for pairwise comparison and detection of orthologous proteins between genomes, then retrieves the corresponding genes and determines the mean percentage of nucleotide sequence identity among orthologous ORFs using the Needleman-Wunsch global alignment algorithm.

Genome properties

The genome is 5,269,577 bp long with 42.22% GC content (Figure 6 and Table 4). It is composed of 46 contigs. Of the 5,193 predicted genes, 5,098 were protein-coding genes, and 95 were RNAs (10 genes encode 5S rRNA, 1 gene encodes 16S rRNA, 1 gene encodes 23S rRNA, 83 genes are tRNA genes). A total of 3,217 genes (63.1%) were assigned a putative function (by cogs or by NR blast). 457 genes were identified as ORFans (8.96%). The remaining genes were annotated as hypothetical proteins (1,097 genes, 21.52%). The distribution of genes into COGs functional categories is presented in Table 5. The properties and statistics of the genome are summarized in Tables 4 and 5.




Figure 6. Graphical circular map of the chromosome. From outside to the center: Red and gray bars representing contigs, genes on the forward strand colored by COG categories (only genes assigned to COG), genes on the reverse strand colored by COG categories (only genes assigned to COG), RNA genes (tRNAs green, rRNAs red), GC content. The inner-most circle shows the GC skew, purple and olive indicating negative and positive values, respectively.

Table 4. Nucleotide content and gene count levels of the genome

| Attribute | Value | % of total ^a |
|----------------------------------|-----------|-------------------------|
| Genome size (bp) | 5,269,577 | 100 |
| Coding region (bp) | 4,342,253 | 82.4 |
| G+C content (bp) | 2,224,760 | 42.22 |
| Total genes | 5,193 | 100 |
| RNA genes | 95 | 1.82 |
| Protein-coding genes | 5,098 | 98.17 |
| Genes with function prediction | 3,217 | 63.1 |
| Genes assigned to COGs | 3,041 | 59.65 |
| Genes with peptide signals | 653 | 12.8 |
| Genes with transmembrane helices | 1,297 | 25.44 |

^a The total is based on either the size of the genome in base pairs or the total number of protein coding genes in the annotated genome

Table 5. Number of genes associated with the 25 general COG functional categories

| Code | Value | % age ^a | Description |
|------|-------|--------------------|--|
| J | 176 | 3.45 | Translation, ribosomal structure and biogenesis |
| A | 0 | 0 | RNA processing and modification |
| K | 281 | 5.51 | Transcription |
| L | 165 | 3.23 | Replication, recombination and repair |
| B | 1 | 0.01 | Chromatin structure and dynamics |
| D | 33 | 0.64 | Cell cycle control, mitosis and meiosis |
| Y | 0 | 0 | Nuclear structure |
| V | 63 | 1.23 | Defense mechanisms |
| T | 144 | 2.82 | Signal transduction mechanisms |
| M | 156 | 3.06 | Cell wall/membrane biogenesis |
| N | 48 | 0.94 | Cell motility |
| Z | 0 | 0 | Cytoskeleton |
| W | 0 | 0 | Extracellular structures |
| U | 50 | 0.98 | Intracellular trafficking and secretion |
| O | 110 | 2.15 | Posttranslational modification, protein turnover, chaperones |
| C | 206 | 4.04 | Energy production and conversion |
| G | 277 | 5.43 | Carbohydrate transport and metabolism |
| E | 370 | 7.25 | Amino acid transport and metabolism |
| F | 82 | 1.60 | Nucleotide transport and metabolism |
| H | 109 | 2.13 | Coenzyme transport and metabolism |
| I | 142 | 2.78 | Lipid transport and metabolism |
| P | 213 | 4.17 | Inorganic ion transport and metabolism |
| Q | 86 | 1.68 | Secondary metabolites biosynthesis, transport and catabolism |
| R | 521 | 10.21 | General function prediction only |
| S | 327 | 6.41 | Function unknown |
| - | 2,057 | 40.34 | Not in COGs |

^a The total is based on the total number of protein coding genes in the annotated genome

Comparison with other species *Bacillus* genomes

Here, we compared the genome of *B. algeriensis* strain EB01^T with those of *B. kribbensis* strain DSM 17871, *B. nealsonii* strain AAU1, *B. bataviensis* strain LMG 21833, *B. subtilis subsp. subtilis* strain 168, *B. vallismortis* strain DV1-F-3, *B. thuringiensis* strain BMB171 and *B. weihenstephanensis* strain KBAB4 (Table 6). The draft genome of *B. algeriensis* (5.26Mb) is larger in size than those of *B. kribbensis*, *B. nealsonii*, *B. subtilis subsp. subtilis* and *B. vallismortis* (5.05, 4.98, 4.22 and 3.88 Mb, respectively) but smaller than those of, *B. bataviensis*, *B. thuringiensis* and *B. weihenstephanensis* (5.37, 5.64 and 5.87 Mb, respectively). *B. algeriensis* has a lower G+C content than *B. kribbensis*, *B. subtilis subsp. subtilis* and *B. vallismortis* (42.22% vs 43%, 43.5% and 43.8%, respectively) but higher than *B. nealsonii*, *B. bataviensis*, *B. thuringiensis* and *B. weihenstephanensis* (42.22% vs 35.1%, 39.6%,

35.2% and 35.5%, respectively). *B. algeriensis* has more predicted protein coding genes (5,098) than *B. kribbensis*, *B. nealsonii*, *B. subtilis subsp. subtilis* and *B. vallismortis* (4,918, 4,789, 4,175 and 4,097, respectively) but fewer protein coding genes than *B. bataviensis*, *B. thuringiensis* and *B. weihenstephanensis* (5,207, 5,352 and 5,653, respectively). In addition, *B. algeriensis* shared 1,804, 1,778, 2,017, 1,768, 1,985, 1,541, 1,863 orthologous genes with *B. thuringiensis*, *B. nealsonii*, *B. bataviensis*, *B. subtilis subsp. subtilis*, *B. kribbensis*, *B. vallismortis*, *B. weihenstephanensis* respectively.

The average nucleotide sequence identity of orthologous genes ranges from 64.54 to 91.06% among the 8 *Bacillus* species, and from 64.77 to 69.33% between *Bacillus algeriensis* and the other compared genomes (Table 7), thus confirming its new species status.

Table 6. Genomic comparison of *B. algeriensis* sp. nov. strain EB01^T with seven other *Bacillus* species[†].

| Species | Strain | Genome accession number | Genome size (Mb) | G+C content |
|--|-------------------|-------------------------|------------------|-------------|
| <i>Bacillus algeriensis</i> | EB01 ^T | ERP003483 | 5.26 | 42.22 |
| <i>Bacillus kribbensis</i> | DSM 17871 | AUMQ00000000.1 | 5.05 | 43 |
| <i>Bacillus nealsonii</i> | AAU1 | ASRU00000000.1 | 4.98 | 35.1 |
| <i>Bacillus bataviensis</i> | LMG 21833 | AJLS00000000.1 | 5.37 | 39.6 |
| <i>Bacillus subtilis subsp. subtilis</i> | 168 | NC_000964.3 | 4.22 | 43.5 |
| <i>Bacillus vallismortis</i> | DV1-F-3 | AFSH00000000.1 | 3.88 | 43.8 |
| <i>Bacillus thuringiensis</i> | BMB171 | NC_014171.1 | 5.64 | 35.2 |
| <i>Bacillus weihenstephanensis</i> | KBAB4 | NC_010184.1 | 5.87 | 35.5 |

[†] species and strain names, genome accession numbers, sizes and G+C contents.

Table 7. Genomic comparison of *B. algeriensis* sp. nov. strain EB01^T with seven other *Bacillus* species[†].

| Species | <i>B. th</i> | <i>B. ne</i> | <i>B. ba</i> | <i>B. subt</i> | <i>B. ma</i> | <i>B. kr</i> | <i>B. va</i> | <i>B. we</i> |
|------------------------------------|--------------|--------------|--------------|----------------|--------------|--------------|--------------|--------------|
| <i>B. thuringiensis</i> | 5,352 | 1,642 | 1,825 | 1,786 | 1,804 | 1,783 | 1,548 | 2,369 |
| <i>B. nealsonii</i> | 67.32 | 4,789 | 1,746 | 1,679 | 1,778 | 1,751 | 1,450 | 1,702 |
| <i>B. bataviensis</i> | 66.65 | 68.78 | 5,207 | 1,812 | 2,017 | 1,997 | 1,567 | 1,904 |
| <i>B. subtilis subsp. subtilis</i> | 65.35 | 65.74 | 66.03 | 4,175 | 1,768 | 1,841 | 2,016 | 1,838 |
| <i>B. algeriensis</i> | 64.77 | 66.61 | 69.33 | 65.44 | 5,098 | 1,985 | 1,541 | 1,863 |
| <i>B. kribbensis</i> | 64.54 | 66.05 | 67.20 | 65.86 | 66.92 | 4,918 | 1,604 | 1,864 |
| <i>B. vallismortis</i> | 64.56 | 65.05 | 65.54 | 91.06 | 64.92 | 65.30 | 4,097 | 1,592 |
| <i>B. weihenstephanensis</i> | 89.95 | 67.27 | 66.70 | 65.46 | 64.87 | 64.56 | 64.70 | 5,653 |

[†]Numbers of orthologous protein shared between genomes (above diagonal), average percentage similarity of nucleotides corresponding to orthologous proteins shared between genomes (below diagonal). Bold numbers indicate numbers of proteins per genome.

B. thuringiensis (*B. th*), *B. nealsonii* (*B. ne*), *B. bataviensis* (*B. ba*), *B. subtilis subsp. subtilis* (*B. subt*), *B. algeriensis* (*B. ma*), *B. kribbensis* (*B. kr*), *B. vallismortis* (*B. va*) and *B. weihenstephanensis* (*B. we*).

Conclusion

On the basis of phenotypic (Table 2), phylogenetic and genomic analyses (taxonogenomics) (Table 6), we formally propose the creation of *Bacillus algeriensis* sp. nov. that contains the strain EB01^T. This strain has been found in hypersaline lacustrine sediment sample collected from Algeria.

Description of *Bacillus algeriensis* sp. nov.

Bacillus algeriensis (al.ge.ri.en'sis. NL. masc.adj. *algeriensis*, of or pertaining to Algeria). Strain EB01^T is a facultative anaerobic Gram-positive, endospore-forming, motile and rod shaped bacterium with rounded ends. Growth is achieved aerobically between 30 and 55°C (optimum 37°C), between 0% and 2.5% NaCl concentration and pH in the range of 6.5-9 (optimum at pH 7). Growth is also observed in microaerophilic atmosphere, however, weak growth was observed under anaerobic conditions. After 24h growth on 5% sheep blood-enriched Columbia agar (BioMerieux) at 37°C, bacterial colonies were smooth, light yellow with 2 mm in diameter. Cells have a length ranging from 2.4 µm to 4.9 µm (mean 3.6 µm) and a diameter ranging from 0.7 µm to 1.1 µm (mean 0.8 µm). Catalase positive but oxidase negative. Using the commercially available API 50CH system (BioMerieux) according to the manufacturer's instructions, a weak positive reaction was observed for D-ribose, D-glucose, D-fructose, methyl α-D-glucopyranoside, N-acetylglucosamine, D-maltose,

D-lactose, D-melibiose, D-saccharose, D-trehalose, D-tagatose, and hydrolysis of starch. Other tests were negative. Using the API ZYM system (BioMerieux), positive reactions were observed for alkaline phosphatase, esterase (C4), esterase lipase (C8), leucine arylamidase, α chymotrypsin, β-glucuronidase, α-glucosidase, N-acetyl-glucosaminidase and a weak positive reaction was observed for acid phosphatase. The nitrate reduction and β-galactosidase reaction was also positive, but urease and indole production were negative. *B. algeriensis* was susceptible to amoxicillin, nitrofurantoin, erythromycin, doxycycline, rifampin, vancomycin, gentamycin, imipenem, trimethoprim-sulfamethoxazole, ciprofloxacin, ceftriaxone and amoxicillin/clavulanic acid, but resistant to nalidixic acid.

The G+C content of the genome is 42.22. The 16S rRNA and genome sequences are deposited in GenBank under accession numbers HG315679 and EMBL database under accession number ERP003483, respectively. The type strain EB01^T (= CSUR P857 = DSM 27334) was isolated from sediment sample of the hypersaline lake Ezzemoul sabkha of Oum-El-Bouaghi region in northeastern Algeria.

Acknowledgements

The authors thank Linda Hadjadj for technical assistance and Xegen company for automating the genome annotation process.

References

1. Cohn F. Untersuchungen über Bakterien. *Beitr Biol Pflanz* 1872; **1**:127-224.
2. Parte AC. LPSN-list of prokaryotic names with standing in nomenclature. *Nucleic Acids Res* 2013; **42**:613-616. [PubMed](#) <http://dx.doi.org/10.1093/nar/gkt1111>
3. Nicholson WL. Roles of *Bacillus* endospores in the environment. *Cell Mol Life Sci* 2002; **59**:410-416. [PubMed](#) <http://dx.doi.org/10.1007/s00018-002-8433-7>
4. Moshafi MH, Forootanfar H, Ameri A, Shakibaie M, Noudeh G, Razavi M. Antimicrobial activity of *Bacillus* sp. strain FAS1 isolated from soil. *Pak J Pharm Sci* 2011; **24**:269-275. [PubMed](#)
5. Bumpus SB, Evans BS, Thomas PM, Ntai I, Kelleher NL. A proteomics approach to discovering natural products and their biosynthetic pathways. *Nat Biotechnol* 2009; **27**:951-960. [PubMed](#) <http://dx.doi.org/10.1038/nbt.1565>
6. Jernigan JA, Stephens DS, Ashford DA, Omenaca C, Topiel MS, Galbraith M, Tapper M, Fisk TL, Zaki S, Popovic T, et al. Bioterrorism-related inhalational anthrax: the first 10 cases reported in the United States. *Emerg Infect Dis* 2001; **7**:933-944. [PubMed](#) <http://dx.doi.org/10.3201/eid0706.010604>
7. Bottone EJ. *Bacillus cereus*, a volatile human pathogen. *Clin Microbiol Rev* 2010; **23**:382-398. [PubMed](#) <http://dx.doi.org/10.1128/CMR.00073-09>
8. Mandell GL, Bennett JE, Dolin R. Principles and Practice of Infectious Diseases. Elsevier **2010**, 4320p.
9. Stackebrandt E, Frederiksen W, Garrity GM, Grimont PA, Kämpfer P, Maiden MC, Nesme X, Rosselló-Mora R, Swings J, Trüper HG, et al. Re-

- port of the ad hoc committee for the re-evaluation of the species definition in bacteriology. *Int J Syst Evol Microbiol* 2002; **52**:1043-1047. [PubMed](#) <http://dx.doi.org/10.1099/ijs.0.02360-0>
10. Tindall BJ, Rossello-Mora R, Busse HJ, Ludwig W, Kampfer P. Notes on the characterization of prokaryote strains for taxonomic purposes. *Int J Syst Evol Microbiol* 2010; **60**:249-266. [PubMed](#) <http://dx.doi.org/10.1099/ijs.0.016949-0>
 11. Stackebrandt E, Ebers J. Taxonomic parameters revisited: tarnished gold standards. *Microbial Today* 2006; **33**:152-155.
 12. Rossello-Mora R. DNA-DNA Reassociation Methods Applied to Microbial Taxonomy and Their Critical Evaluation. In: Stackebrandt E (ed), *Molecular Identification, Systematics, and Population Structure of Prokaryotes*. Springer, Berlin **2006**. p. 23-50.
 13. Welker M, Moore ER. Applications of whole-cell matrix-assisted laser-desorption/ionization time-of-flight mass spectrometry in systematic microbiology. *Syst Appl Microbiol* 2011; **34**:2-11. [PubMed](#) <http://dx.doi.org/10.1016/j.syapm.2010.11.013>
 14. Ramasamy D, Mishra AK, Lagier JC, Padhmanabhan R, Rossi M, Sentaosa E, Raoult D, Fournier PE. A polyphasic strategy incorporating genomic data for the taxonomic description of novel bacterial species. *Int J Syst Evol Microbiol* 2014; **64**:384-391.
 15. Ramasamy D, Lagier JC, Gorlas A, Raoult D, Fournier PE. Non contiguous-finished genome sequence and description of *Bacillus massiliosenegalensis* sp. nov. *Stand Genomic Sci* 2013; **8**:264-278. [PubMed](#) <http://dx.doi.org/10.4056/sigs.3496989>
 16. Keita MB, Diene SM, Robert C, Raoult D, Fournier PE, Bittar F. Non-contiguous finished genome sequence and description of *Bacillus massiliogorillae* sp. nov. *Stand Genomic Sci* 2013; **9**:93-105. [PubMed](#) <http://dx.doi.org/10.4056/sigs.4388124>
 17. Mishra AK, Pfliegerer A, Lagier JC, Robert C, Raoult D, Fournier PE. Non-contiguous finished genome sequence and description of *Bacillus massilioanorexius* sp. nov. *Stand Genomic Sci* 2013; **8**:465-479. [PubMed](#) <http://dx.doi.org/10.4056/sigs.4087826>
 18. Kokcha S, Mishra AK, Lagier JC, Million M, Leroy Q, Raoult D, Fournier PE. Non contiguous-finished genome sequence and description of *Bacillus timonensis* sp. nov. *Stand Genomic Sci* 2012; **6**:346-355. [PubMed](#) <http://dx.doi.org/10.4056/sigs.2776064>
 19. Mishra AK, Lagier JC, Rivet O, Raoult D, Fournier PE. Non-contiguous finished genome sequence and description of *Paenibacillus senegalensis* sp. nov. *Stand Genomic Sci* 2012; **7**:70-81. [PubMed](#) <http://dx.doi.org/10.4056/sigs.3056450>
 20. Mishra AK, Lagier JC, Robert C, Raoult D, Fournier PE. Genome sequence and description of *Timonella senegalensis* gen. nov., sp. nov., a new member of the suborder *Micrococccinae*. *Stand Genomic Sci* 2013; **8**:318-335. [PubMed](#) <http://dx.doi.org/10.4056/sigs.3476977>
 21. Ramasamy D, Lagier JC, Nguyen TT, Raoult D, Fournier PE. Non contiguous-finished genome sequence and description of *Dielma fastidiosa* gen. nov., sp. nov., a new member of the Family *Erysipelotrichaceae*. *Stand Genomic Sci* 2013; **8**:336-351. [PubMed](#) <http://dx.doi.org/10.4056/sigs.3567059>
 22. Mishra AK, Hugon P, Lagier JC, Nguyen TT, Couderc C, Raoult D, Fournier PE. Non contiguous-finished genome sequence and description of *Enorma massiliensis* gen. nov., sp. nov., a new member of the Family *Coriobacteriaceae*. *Stand Genomic Sci* 2013; **8**:290-305. [PubMed](#) <http://dx.doi.org/10.4056/sigs.3426906>
 23. Hugon P, Mishra AK, Lagier JC, Nguyen TT, Couderc C, Raoult D, Fournier PE. Non contiguous-finished genome sequence and description of *Brevibacillus massiliensis* sp. nov. *Stand Genomic Sci* 2013; **8**:1-14. [PubMed](#) <http://dx.doi.org/10.4056/sigs.3466975>
 24. Lagier JC, El Karkouri K, Mishra AK, Robert C, Raoult D, Fournier PE. Non-contiguous finished genome sequence and description of *Enterobacter massiliensis* sp. nov. *Stand Genomic Sci* 2013; **7**:399-412. [PubMed](#) <http://dx.doi.org/10.4056/sigs.3396830>
 25. Mishra AK, Lagier JC, Nguyen TT, Raoult D, Fournier PE. Non-contiguous finished genome sequence and description of *Peptoniphilus senegalensis* sp. nov. *Stand Genomic Sci* 2013; **7**:370-381. [PubMed](#) <http://dx.doi.org/10.4056/sigs.3366764>
 26. Roux V, ElKarkouri K, Lagier JC, Robert C, Raoult D. Non-contiguous finished genome sequence and description of *Kurthia massiliensis* sp. nov. *Stand Genomic Sci* 2012; **7**:221-232. [PubMed](#) <http://dx.doi.org/10.4056/sigs.3206554>
 27. Woese CR, Kandler O, Wheelis ML. Towards a natural system of organisms: proposal for the do-

- mains *Archaea*, *Bacteria*, and *Eucarya*. *Proc Natl Acad Sci USA* 1990; **87**:4576-4579. [PubMed](#) <http://dx.doi.org/10.1073/pnas.87.12.4576>
28. Gibbons NE, Murray RGE. Proposals concerning the higher taxa of bacteria. *Int J Syst Bacteriol* 1978; **28**:1-6. <http://dx.doi.org/10.1099/00207713-28-1-1>
29. Murray RGE. The Higher Taxa, or, a Place for Everything...? In: Holt JG (ed). *Bergey's Manual of Systematic Bacteriology*, First Edition, Volume 1, The Williams and Wilkins Co, Baltimore 1984; p. 31-34.
30. Garrity GM, Holt JG. The Road Map to the Manual. In: Garrity GM, Boone DR, Castenholz RW (eds), *Bergey's Manual of Systematic Bacteriology*, Second Edition, Volume 1, Springer, New York, 2001, p. 119-169.
31. Ludwig W, Schleifer KH, Whitman WB. Class I. *Bacilli* class nov. In: De Vos P, Garrity GM, Jones D, N.R. Krieg W, Ludwig W, Rainey EA, Schleifer KH, Withman WB. *Bergey's Manual of Systematic Bacteriology*, second edition, vol 3 (The *Firmicutes*), Springer, Dordrecht, Heidelberg, London, New York **2009**, pp.19-20.
32. List of new names and new combinations previously effectively, but not validly, published. List no. 132. *Int J Syst Evol Microbiol* 2010; **60**:469-472. <http://dx.doi.org/10.1099/ijs.0.022855-0>
33. Skerman VBD, McGowan V, Sneath PHA, eds. Approved Lists of Bacterial Names. *Int J Syst Bacteriol* 1980; **30**:225-420. <http://dx.doi.org/10.1099/00207713-30-1-225>
34. Prévot AR. In: Hauduroy P, Ehringer G, Guillot G, Magrou J, Prévot AR, Rosset. Urbain A (eds) *Dictionnaire des Bactéries Pathogènes*, 2nd ed., Masson, Paris, 1953, pp. 1-692.
35. Fischer A. Untersuchungen über bakterien. *Jahrbücher für Wissenschaftliche Botanik* 1985; **27**:1-163.
36. Gibson T, Gordon RE. Genus I. *Bacillus* Cohn 1872; 174; Nom. gen. cons. Nomencl. Comm. Intern. Soc. Microbiol. 1937, 28; Opin. A. Jud. Comm. 1955, 39. In: Buchanan RE, Gibbons NE (eds), *Bergey's Manual of Determinative Bacteriology*, Eighth Edition, The Williams and Wilkins Co., Baltimore, 1974, p. 529-550.
37. Ashburner M, Ball CA, Blake JA, Botstein D, Butler H, Cherry JM, Davis AP, Dolinski K, Dwight SS, Eppig J T *et al.* Gene ontology: tool for the unification of biology. The Gene Ontology Consortium. *Nat Genet* 2000; **25**:25-29. [PubMed](#) <http://dx.doi.org/10.1038/75556>
38. Saitou N, Nei M. The neighbor-joining method: a new method for reconstructing phylogenetic trees. *Mol Biol Evol* 1987; **4**:406-425. [PubMed](#)
39. Tamura K, Peterson D, Peterson N, Stecher G, Nei M, Kumar S. MEGA5: molecular evolutionary genetics analysis using maximum likelihood, evolutionary distance, and maximum parsimony methods. *Mol Biol Evol* 2011; **28**:2731-2739. [PubMed](#) <http://dx.doi.org/10.1093/molbev/msr121>
40. Kanso S, Greene AC, Patel BKC. *Bacillus subterraneus* sp. nov., an iron- and manganese-reducing bacterium from a deep subsurface Australian thermal aquifer. *Int J Syst Evol Microbiol* 2002; **52**:869-874. [PubMed](#) <http://dx.doi.org/10.1099/ijs.0.01842-0>
41. Tiago I, Pires C, Mendes V, Morais PV, da Costa MS, Verissimo A. *Bacillus foraminis* sp. nov., isolated from a non-saline alkaline groundwater. *Int J Syst Evol Microbiol* 2006; **56**:2571-2574. [PubMed](#) <http://dx.doi.org/10.1099/ijs.0.64281-0>
42. Yoon JH, Kang SS, Lee KC, Kho YH, Choi SH, Kang KH, Park YH. *Bacillus jeotgali* sp. nov., isolated from jeotgal, Korean traditional fermented seafood. *Int J Syst Evol Microbiol* 2001; **51**:1087-1092. [PubMed](#) <http://dx.doi.org/10.1099/00207713-51-3-1087>
43. Euzéby J. Validation List no. 117. List of new names and new combinations previously effectively, but not validly, published. *Int J Syst Evol Microbiol* 2007; **57**:1933-1934. [PubMed](#) <http://dx.doi.org/10.1099/ijs.0.65495-0>
44. Perez-Ibarra BM, Flores ME, Garcia-Varela M. Isolation and characterization of *Bacillus thioparus* sp. nov., chemolithoautotrophic, thio-sulfate-oxidizing bacterium. *FEMS Microbiol Lett* 2007; **271**:289-296. [PubMed](#) <http://dx.doi.org/10.1111/j.1574-6968.2007.00729.x>
45. Ahmed I, Yokota A, Fujiwara T. A novel highly boron tolerant bacterium, *Bacillus boroniphilus* sp. nov., isolated from soil, that requires boron for its growth. *Extremophiles* 2007; **11**:217-224. [PubMed](#) <http://dx.doi.org/10.1007/s00792-006-0027-0>
46. Heyrman J, Vanparys B, Logan NA, Balcaen A, Rodriguez-Diaz M, Felske A, De Vos P. *Bacillus novalis* sp. nov., *Bacillus vireti* sp. nov., *Bacillus soli* sp. nov., *Bacillus bataviensis* sp. nov. and *Bacillus drentensis* sp. nov., from the Drentse A grasslands. *Int J Syst Evol Microbiol* 2004; **54**:47-

57. [PubMed](#)
<http://dx.doi.org/10.1099/ijs.0.02723-0>
47. Venkateswaran K, Kempf M, Chen F, Satomi M, Nicholson W, Kern R. *Bacillus nealsonii* sp. nov., isolated from a spacecraft-assembly facility, whose spores are gamma-radiation resistant. *Int J Syst Evol Microbiol* 2003; **53**:165-172. [PubMed](#)
<http://dx.doi.org/10.1099/ijs.0.02311-0>
48. Lim JM, Jeon CO, Lee JR, Park DJ, Kim CJ. *Bacillus kribbensis* sp. nov., isolated from a soil sample in Jeju, Korea. *Int J Syst Evol Microbiol* 2007; **57**:2912-2916. [PubMed](#)
<http://dx.doi.org/10.1099/ijs.0.65227-0>
49. Seng P, Rolain JM, Fournier PE, La Scola B, Drancourt M, Raoult D. MALDI-TOF-mass spectrometry applications in clinical microbiology. *Future Microbiol* 2010; **5**:1733-1754. [PubMed](#)
<http://dx.doi.org/10.2217/fmb.10.127>
50. Seng P, Drancourt M, Gouriet F, La Scola B, Fournier PE, Rolain JM, Raoult D. Ongoing revolution in bacteriology: routine identification of bacteria by matrix-assisted laser desorption ionization time-of-flight mass spectrometry. *Clin Infect Dis* 2009; **49**:543-551. [PubMed](#)
<http://dx.doi.org/10.1086/600885>
51. Field D, Garrity G, Gray T, Morrison N, Selengut J, Sterk P, Tatusova T, Thomson N, Allen MJ, Angiuoli SV, et al. The minimum information about a genome sequence (MIGS) specification. *Nat Biotechnol* 2008; **26**:541-547. [PubMed](#)
<http://dx.doi.org/10.1038/nbt1360>
52. fastq-mcf program. <http://code.google.com/p/ea-utils>
53. Prodigal. <http://prodigal.ornl.gov>
54. GenBank database.
<http://www.ncbi.nlm.nih.gov/genbank>
55. RNAmmer 1.2 Server.
<http://www.cbs.dtu.dk/services/RNAmmer/>
56. Lagesen K, Hallin P, Rodland EA, Staerfeldt HH, Rognes T, Ussery DW. RNAmmer: consistent and rapid annotation of ribosomal RNA genes. *Nucleic Acids Res* 2007; **35**:3100-3108. [PubMed](#)
<http://dx.doi.org/10.1093/nar/gkm160>
57. Lowe TM, Eddy SR. tRNAscan-SE: a program for improved detection of transfer RNA genes in genomic sequence. *Nucleic Acids Res* 1997; **25**:955-964. [PubMed](#)
<http://dx.doi.org/10.1093/nar/25.5.0955>
58. Kall L, Krogh A, Sonnhammer EL. Advantages of combined transmembrane topology and signal peptide prediction--the Phobius web server. *Nucleic Acids Res* 2007; **35**:W429-W432. [PubMed](#)
<http://dx.doi.org/10.1093/nar/gkm256>
59. Rutherford K, Parkhill J, Crook J, Horsnell T, Rice P, Rajandream MA, Barrell B. Artemis: sequence visualization and annotation. *Bioinformatics* 2000; **16**:944-945. [PubMed](#)
<http://dx.doi.org/10.1093/bioinformatics/16.10.944>
60. Carver T, Thomson N, Bleasby A, Berriman M, Parkhill J. DNAPlotter: circular and linear interactive genome visualization. *Bioinformatics* 2009; **25**:119-120. [PubMed](#)
<http://dx.doi.org/10.1093/bioinformatics/btn578>
61. Lechner M, Findeiss S, Steiner L, Marz M, Stadler PF, Prohaska SJ. Proteinortho: detection of (co-)orthologs in large-scale analysis. *BMC Bioinformatics* 2011; **12**:124-133. [PubMed](#)
<http://dx.doi.org/10.1186/1471-2105-12-124>