

Supplementary Materials

Supplementary Table 1. Baseline characteristics of the WEB and clipping group after propensity score matching. WEB: Woven EndoBridge, SMD: standardized mean difference, SD: standard deviation, BMI: body mass index. Values are presented as number and proportion (%) of the patients not otherwise specified. Asterisk indicates statistical significance ($p < .05$).

	WEB (N = 36)	Clipping (N = 72)	<i>p</i> -value	SMD
Age, yrs	61.5±8.6	61.5±8.9	0.999	0.001
Male (%)	12 (33.3)	20 (27.8)	0.547	0.121
BMI	24.8±3.3	24.6±2.7	0.724	0.071
Comorbidities				
Hypertension	20 (55.6)	38 (52.8)	0.788	0.056
Diabetes mellitus	10 (27.8)	15 (20.8)	0.355	0.162
Hyperlipidemia	17 (47.2)	42 (58.3)	0.288	0.224
Smoker	4 (11.1)	13 (18.1)	0.353	0.198
Alcohol consumption	12 (33.3)	19 (26.4)	0.408	0.152
Aneurysm Morphology				
Width (mm)	4.92±1.75	4.74±2.64	0.676	0.082
Height (mm)	4.64±1.74	4.53±2.38	0.739	0.055
Neck diameter (mm)	3.77±1.30	3.83±1.67	0.837	0.039
Aspect ratio	1.25±0.32	1.20±0.36	0.182	0.144
Dome to neck ratio	1.33±0.23	1.24±0.32	0.035*	0.329

Supplementary Table 2. Clinical and radiologic outcome after propensity score matching. WEB: Woven EndoBridge, OR: odds ratio, CI: confidence interval, SD: standard deviation, mRS: modified Rankin Scale. Values are presented as number and proportion (%) of the patients not otherwise specified. Asterisk indicates statistical significance ($p < .05$).

	WEB (N = 36)	Clipping (N = 72)	OR (95% CI)	<i>p</i>-value
Clinical Follow-up				
Interval (days, Mean±SD)	297.7±92.4	328.2±88.4		0.092
Morbidity (%)	1 (2.8)	1 (1.4)	2.00 (0.13-31.98)	0.624
Major Complications (%)	2 (5.6)	1 (1.4)	4.00 (0.36-44.11)	0.258
Radiologic Follow-up				
Interval (days, Mean±SD)	299.8±105.3	305.9±163.7		0.839
Adequate occlusion (%)	26 (74.3)	71 (98.6)	0.06 (0.01-0.44)	0.006*