



Review Article

Optimization of microbial cell factories for astaxanthin production: Biosynthesis and regulations, engineering strategies and fermentation optimization strategies

Mostafa Basiony^a, Liming Ouyang^{a,*}, Danni Wang^a, Jiaming Yu^a, Liming Zhou^a, Mohan Zhu^a, Xuyuan Wang^a, Jie Feng^a, Jing Dai^a, Yijie Shen^a, Chengguo Zhang^b, Qiang Hua^a, Xiuliang Yang^b, Lixin Zhang^a

^a State Key Laboratory of Bioreactor Engineering, East China University of Science and Technology, Shanghai, 200237, China

^b Shandong Jincheng Bio-Pharmaceutical Co., Ltd., No. 117 Qixing River Road, Zibo, 255130, Shandong, China

ARTICLE INFO

Keywords:

Astaxanthin
Carotenoids
Metabolic engineering
Microbial cell factories
Lipid
Fermentation
Antioxidants

ABSTRACT

The global market demand for natural astaxanthin is rapidly increasing owing to its safety, the potential health benefits, and the diverse applications in food and pharmaceutical industries. The major native producers of natural astaxanthin on industrial scale are the alga *Haematococcus pluvialis* and the yeast *Xanthophyllomyces dendrorhous*. However, the natural production via these native producers is facing challenges of limited yield and high cost of cultivation and extraction. Alternatively, astaxanthin production via metabolically engineered non-native microbial cell factories such as *Escherichia coli*, *Saccharomyces cerevisiae* and *Yarrowia lipolytica* is another promising strategy to overcome these limitations. In this review we summarize the recent scientific and biotechnological progresses on astaxanthin biosynthetic pathways, transcriptional regulations, the interrelation with lipid metabolism, engineering strategies as well as fermentation process control in major native and non-native astaxanthin producers. These progresses illuminate the prospects of producing astaxanthin by microbial cell factories on industrial scale.

1. Introduction

Astaxanthin is a keto-carotenoid with dark reddish color, and is widely distributed in algae, shrimp, crab, shellfish and other organisms [1,2]. Due to its safety and the stable reddish color, astaxanthin became an ideal source for pigmentation in aquaculture and food industry [3]. Additionally, several studies showed potential health benefits associated with astaxanthin supplementation including antioxidant, anti-inflammation, anti-cancer, neuroprotection and immuno-enhancement activities [4]. Owing to these benefits and uses, the demand and market size of astaxanthin have been rapidly increasing. The global astaxanthin market is expected to reach USD 3.5 billion by 2026, from USD 1.37 billion in 2020, at a compound annual growth rate (CAGR) of 16.8% [5]. The animal feed application dominated the market, with a share of 65.5% in 2020, while China dominated the Asia Pacific market for astaxanthin with a share of 40.4% [6].

The chemical structure of astaxanthin (3,3'-dihydroxy-4,4'-dione- β,β' -carotene) resembles that of many other carotenoids. It contains a main chain composed of four isoprene units connected by conjugated double bonds, and one β -ionone ring at each end of the chain. Each ionone ring carries a characteristic 3-hydroxy (OH) and a 4-keto (C=O) groups. These structural arrangements enable the compound to attract free radicals and provide them electrons resulting in termination of the free radical chain reaction, which explains the strong antioxidant activity of astaxanthin [2,7]. Astaxanthin has many optical and geometric isomers [8]. The 3 and 3' hydroxygroup of each ionone ring can exist in the S or R form. Accordingly, three different optical isomers have been reported ((3S, 3'S), (3R, 3'R), and (3R, 3'S)) for astaxanthin. The antioxidant activity is influenced by the form of astaxanthin isomer used, and the (3S, 3'S) form showed stronger antioxidant activity than that of (3R, 3'R) and (3R, 3'S) [9]. Several astaxanthin derivatives have been reported in nature including esterified astaxanthin (mono- and di-ester),

Peer review under responsibility of KeAi Communications Co., Ltd.

* Corresponding author.

E-mail address: ouyanglm@ecust.edu.cn (L. Ouyang).

<https://doi.org/10.1016/j.synbio.2022.01.002>

Received 29 October 2021; Received in revised form 8 December 2021; Accepted 3 January 2022

2405-805X/© 2022 The Authors. Publishing services by Elsevier B.V. on behalf of KeAi Communications Co. Ltd. This is an open access article under the CC

BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

glycosylated astaxanthin (astaxanthin β -D-glucoside, astaxanthin β -D-diglucoside and astaxanthin dirhamnoside) and glycosylated-esterified astaxanthin (astaxanthin diglucoside diester) [10–14]. These modifications take place at the 3 and 3' hydroxygroup of the ionone rings and are believed to influence the polarity, solubility and the biological activity of astaxanthin [4,10,15].

Astaxanthin can be obtained through extraction from natural sources or chemical synthesis [16]. Astaxanthin biosynthesis has been reported in several organisms including microalgae, bacteria, yeast and plants [4, 17]. Although, astaxanthin has been detected in several aquatic animals, these animals cannot synthesize astaxanthin, but they obtain it through feed [16]. Several microalgae have been extensively studied for their ability to synthesize and accumulate astaxanthin naturally when cells are cultured under stress [18]. Amongst these algae, *Haematococcus pluvialis* is recognized as the best natural source of astaxanthin with the highest astaxanthin content, which can reach up to 4% of the dry cell weight under optimal cultivation conditions [18]. Therefore, *H. pluvialis* astaxanthin covers over 50% of the natural astaxanthin market in the global nutraceutical industry [19,20]. In addition to *H. pluvialis*, *Chlorella zofingiensis* is also considered as a promising source of natural astaxanthin production [21]. However, the large scale production using these algae requires high light intensity, large cultivation areas and complex cultivation process which makes the industrial production challenging [22]. Bacterial astaxanthin biosynthesis have been reported in many strains, such as *Brevundimonas* sp., *Sphingomonas* sp. and *Paracoccus* sp [23–25]. The marine bacterium *Paracoccus carotinifaciens* is considered the best among bacterial astaxanthin producers, with a high level at 480 mg/L by fed-batch fermentation [18,26]. One of the major astaxanthin producing yeasts is *Xanthophyllomyces dendrorhous* [27]. *X. dendrorhous* is the earliest strain used in the industrial production of astaxanthin, and contains about 0.2–0.5 mg/g DCW carotenoids, of which 40–95% is astaxanthin [28,29]. Several strategies have been

extensively applied for enhanced production of astaxanthin from *X. dendrorhous*, and the highest yield reported has reached 9.7 mg/g dry cell weight (DCW) [18,30]. In addition to the above mentioned microorganisms, astaxanthin biosynthesis has been detected in some plant species of the genus *Adonis* [17,31].

At present, the human health consumption of astaxanthin is strictly limited to *H. pluvialis* [32]. However, natural astaxanthin cannot meet the market demand due to the low yield and high costs of cultivation and extraction [16]. Therefore, chemically synthesized astaxanthin is the most dominant source for commercial use due to its low cost and higher profitability. Nevertheless, the demand is still in the favor of natural astaxanthin due to safety concerns related to the byproducts formed during the chemical synthesis of astaxanthin [20]. Additionally, natural astaxanthin has higher antioxidant activity, as it mainly exists either in (3S, 3'S) or (3R, 3'R) forms. For instance, more than 95% of astaxanthin from *H. pluvialis* is in (3S, 3'S) form [33], while, the (3R, 3'R) is the dominant form in the red yeast *X. dendrorhous* [34]. However, the chemically synthesized astaxanthin is composed of a mixture of the three isomers at a ratio of 1:2:1 (3R,3'R:3R,3'S:3S,3'S). Thus, to overcome the imperfections of the native producers or chemical synthesis and benefiting from the advances in biotechnology, the engineering of the non-carotenogenic microbes, such as *Escherichia coli*, *Saccharomyces cerevisiae* and *Yarrowia lipolytica*, to produce astaxanthin shows promising results to become better alternative cell factories on industrial scale.

In this review we intend to address the recent advances in microbial astaxanthin production with a main focus on the major native and non-native producers (Table 1) in regards to astaxanthin biosynthesis regulations, engineering strategies and fermentation optimization attempts.

Table 1
Major native and non-native astaxanthin producing microorganisms.

Organism	Highest Yield or Titer	Fermentation scale	Pros	Cons	Ref
Native					
<i>Haematococcus pluvialis</i>	Modified: 87.4 mg/L	500 mL volume glass tubular airlift photobioreactors 8 days	High astaxanthin content; safe for human consumption; environment-friendly; stable non-GMO; well accepted by laws and regulations of different countries	Complex cultivation; slow growth rate; large space required; easy to be contaminated in open culture system; high cost of production; difficult to scale up and extract; limited by light and nutrition; astaxanthin in esterified form	[3,19,220, 239,240]
	Unmodified: 77.2 mg/g DCW	Shake-flask 18 days			
<i>Chlorella zofingiensis</i>	73.3 mg/L	Shake-flask 14 days	High growth rate and high cell density when cultivated in heterotrophic conditions; high lipid content; non-GMO	Complex cultivation; easy to be contaminated; difficult to scale up and extract; slow growth rate and large space required when cultured in photoautotrophic conditions, astaxanthin in esterified form	[3,21,36, 241–243]
	6.8 mg/g DCW	NA			
<i>Xanthophyllomyces dendrorhous</i>	Modified: 9.7 mg/g Unmodified: 420 mg/L	Fermenter 216 h 1 L bioreactor 240 h	Simple requirements for growth; easier to scale up and achieve high biomass; high growth rate; can use a variety of carbon sources; can act as a non-GMO	Low yield; the growth rate of yeast cells is inversely proportional to the accumulation of astaxanthin; low market demand for its astaxanthin configuration; harder to obtain pure free astaxanthin; unknown regulatory pathways and thus harder to be rationally engineered	[3,30,80, 198]
Non-native					
<i>Escherichia coli</i>	18.7 mg/g DCW	1 L bioreactor 85 h	Easier to scale up and extract; high biomass; fastest growth rate; clear genetic background and mature genetic manipulation; low cost of using glucose as a carbon source, simple equipment requirements	Multiple purification steps and high purification cost are required to remove recombinant DNA and endotoxin	[3,124, 138,244]
	1.18 g/L	5 L bioreactor 60 h			
<i>Saccharomyces cerevisiae</i>	404.78 mg/L	5 L bioreactor 180 h	Easier to scale up and extract; high biomass; fast growth rate; GRAS; environment friendly; clear genetic background; mature genetic manipulation	Purification is required to remove recombinant DNA; low yield; complex intermediate metabolites	[3,172, 225,244]
	13.8 mg/g	5 L bioreactor 68 h			
<i>Yarrowia lipolytica</i>	858 mg/L (16.7 mg/g)	Shake Flask (Fed-Batch) 288 h	Easier to scale up and extract; high biomass; fast growth; GRAS; high lipid content	Purification is required to remove recombinant DNA; low yield; complex intermediate metabolites	[3,117, 149,245]

2. Astaxanthin biosynthesis: pathways, regulations and distribution

2.1. Astaxanthin biosynthetic pathways

As a terpenoid, astaxanthin biosynthesis starts from the 5-carbon isomers isopentenyl pyrophosphate (IPP) and dimethylallyl pyrophosphate (DMAPP) (Fig. 1). IPP and DMAPP can be natively synthesized by mevalonic acid (MVA) pathway in eukaryotic microbes using acetyl-CoA as precursor, or 2-methyl-D-erythritol-4-phosphate (MEP) pathway in prokaryotic microbes and plant plastids using glyceraldehyde-3-phosphate and pyruvate as precursors [35,36]. In the MVA pathway, acetyl-CoA acetyltransferase (AACT) condensates two acetyl-CoA molecules to give acetoacetyl-CoA, which is further condensed with additional molecule of acetyl-CoA by hydroxymethylglutaryl-CoA synthase (HMGS) to form 3-hydroxy-3-methylglutaryl-CoA (HMG-CoA) [37]. Then, a reduction step by 3-hydroxy-3-methylglutaryl-CoA reductase (HMGR) results in mevalonate formation. Afterwards, IPP is formed by a three step reactions mediated by mevalonate kinase (MK), phosphomevalonate kinase (PMK), and at last mevalonate diphosphocarbonylase (MVD). The formed IPP is then isomerized to DMAPP by isopentenyl diphosphate isomerase (IDI) [37].

In MEP pathway, the first step starts by condensation of

glyceraldehyde-3-phosphate and pyruvate by 1-deoxy-D-xylulose-5-phosphate synthase (DXS) to form 1-deoxy-D-xylulose-5-phosphate (DXP), followed by a reduction step by 1-deoxy-D-xylulose-5-phosphate reductoisomerase (DXR) to form 2C-methyl-D-erythritol-4-phosphate (MEP) [38]. The formed MEP is then converted into 4-(cytidine 5'-diphospho)-2-C-methyl-D-erythritol (CDP-ME) which is catalyzed by 2-C-methyl-D-erythritol 4-phosphate cytidylyltransferase (MCT). Then 2C-methyl-D-erythritol-2,4-cyclodiphosphate (MECDP) is generated through phosphorylation and cyclization by the activity of CDP-ME kinase (CMK) and MECDP synthase (MDS), respectively. The MECDP is then converted into 1-hydroxy-2-methyl-2-butenyl 4-diphosphate (HMBPP) under the action of 4-hydroxy-3-methylbut-2-enyl-diphosphate synthase (HDS). Then IPP and DMAPP are derived from HMBPP by the activity of 4-hydroxy-3-methylbut-2-enyl diphosphate reductase (HDR) [38]. In addition to these differences between MVA and MEP pathways, the two pathways differ in carbon use efficiency, energy and cofactor requirements, and IPP yields [39,40]. Starting from glucose, the carbon loss in the form of CO₂ Via MVA pathway is higher than that of MEP. In the course of the precursors formation, one molecule of CO₂ is generated to form a single acetyl-CoA molecule as the MVA pathway precursor, while, the formation of the MEP pathway precursors, pyruvate and G3P, does not involve any carbon loss [39]. For each IPP molecule formation in the MVA pathway, a total of 1.5 molecules of




Fig. 1. Scheme for astaxanthin biosynthesis and strategies for enhancing precursors and cofactors supply. Enzymes are as follows: (1) EMP-TCA pathway (purple): HK, Hexokinase; GPD, glyceraldehyde 3-phosphate dehydrogenase; PD, pyruvate dehydrogenase; GltA, citrate synthase; Acon, aconitate hydratase; Icd, aconitate hydratase; SucAB, α -ketoglutarate dehydrogenase; SucCD, succinyl-CoA synthetase; SdhABCD, succinate dehydrogenase; FumAC, fumarate hydratase; Mdh, malate dehydrogenase; ME, malic enzyme. (2) Pentose phosphate pathway (gold): G6PDH, glucose 6-phosphate dehydrogenase; PGL, 6-phosphogluconolactonase; PGDH, 6-phosphogluconate dehydrogenase; Rpe, ribulose 5-phosphate 3-epimerase; Rpi, ribose-5-phosphate isomerase; Tkt A/B, transketolase subunit A/B; Tal, transaldolase. (3) DMAPP/IPP module (green): MVA Pathway: AACT, acetyl-CoA acetyltransferase; HMGS, hydroxymethylglutaryl-CoA synthase; HMGR, 3-hydroxy-3-methylglutaryl-CoA reductase; MK, mevalonate kinase; PMK, phosphomevalonate kinase; MVD, diphosphocarbonylase; IDI, isopentenyl pyrophosphate isomerase; MEP Pathway: DXS, 1-deoxy-D-xylulose-5-phosphate synthase; DXR, 1-deoxy-D-xylulose-5-phosphate reductoisomerase; MCT, 2-C-methyl-D-erythritol 4-phosphate cytidylyltransferase; CMK, CDP-ME kinase; MDS, MECDP synthase; HDS, 4-hydroxy-3-methylbut-2-enyl-diphosphate synthase; HDR, 4-hydroxy-3-methylbut-2-enyl diphosphate reductase; IDI, isopentenyl pyrophosphate isomerase. (4) β -carotene biosynthesis (orange): GPS, geranyl pyrophosphate synthase; FFPS, farnesyl diphosphate synthase; GGPPS, geranylgeranyl pyrophosphate synthase; CrtY(CrtYB), lycopene cyclase. (5) Astaxanthin biosynthesis (red): CrtZ, β -carotene hydroxylase; CrtW, β -carotene ketolase; CBFD, carotenoid beta-ring 4-dehydrogenase; HBFD, 4-hydroxy-beta-ring 4-dehydrogenase; CrtS, cytochrome P450 monooxygenase; CrtR, cytochrome P450 reductase. (6) Electron transport chain (grey). The Solid arrows indicate a single step reaction. The dashed arrows indicate multistep reactions. The dashed purple boxes indicate the engineered steps via overexpression of the corresponding gene in the engineered host. (For interpretation of the references to color in this figure legend, the reader is referred to the Web version of this article.)

glucose, three ATP molecules and two molecules of NADPH are needed [40]. However, the MEP pathway requires 1.25 molecules of glucose, three molecules of NADPH and three ATP molecules (two for the regeneration of CTP, one for CDP-ME conversion to CDP-MEP) (Fig. 1). Thus, the theoretical IPP yield on glucose from MEP pathway is expected to be higher than that of MVA pathway, while more cofactors are needed.

The IPP and DMAPP produced through MEP or MVA pathways are then condensed to give geranyl pyrophosphate (GPP; C10) by the activity of geranyl pyrophosphate synthase (GPS), which is subsequently converted to farnesyl pyrophosphate (FPP; C15) by farnesyl diphosphate synthase (FFPS), and then to geranylgeranyl pyrophosphate (GGPP; C20) by the action of GGPP synthase (GGPPS) [41]. GGPP is then catalyzed by phytoene synthase (PSY or CrtB) to produce the first C40 carotenoid phytoene, which is desaturated and converted into lycopene by the consecutive action of phytoene desaturase (PDS) and zeta-carotene desaturase (ZDS). β -carotene is then formed from lycopene by the activity of lycopene cyclase (LCY, CrtY, CrtYB, CarRRP or LCYB). Afterwards, β -carotene is converted into astaxanthin in enzymatic reactions dependent on the producing organisms [42].

The biosynthesis of astaxanthin in bacteria and algae from β -carotene is mediated by two enzymes, the β -carotene hydroxylase encoded by the *crtZ* (*crtR-B* or *CHYB* in algae) gene, and the β -carotene ketolase encoded by the *crtW* (*bkt* in algae) gene [43–45]. In response to different environmental stress conditions, three different BKTs have been reported to be expressed in *H. pluvialis* [46–48]. The expression level of these proteins varies among each other on a manner dependent on the stress conditions [47]. Luo et al. identified six copies of *bkt* genes (*bkt1a*, *bkt1b*, *bkt1c*, *bkt2a*, *bkt2b* and *bkt3*) in *H. pluvialis* genome, and they are believed to be a result of gene duplications during genome evolution [49]. Similarly, two different β -carotene hydroxylase genes (*crtR-B1* and *crtR-B2*) were also found in *H. pluvialis* genome [50]. In addition to the multi-copies of these genes in *H. pluvialis*, the products of these genes are believed to be localized to different cellular compartments [50–52]. These findings might explain the significant high level of astaxanthin in *H. pluvialis* over other producers. Given that the conversion from β -carotene to astaxanthin involves four reactions that takes place by the addition of two carbonyl groups and two hydroxyl groups at 4, 4'-C and 3, 3'-C positions of the β -ionone ring, respectively (Fig. 1), the hydroxylation and ketolation reactions may occur alternatively dependent on the substrate preferences of the different enzymes which also leads to the accumulation of several intermediates [53].

A bi-functional cytochrome P450 astaxanthin synthase, CrtS, is believed to mediate the conversion of β -carotene to astaxanthin in *X. dendrorhous* (Fig. 1), however the mechanism is not well understood [54]. It should be noted that, the redox partner cytochrome P450 reductase, CrtR, is essential for astaxanthin biosynthesis in this yeast [55]. In this context, the sole expression of *crtS* in β -carotene producing *S. cerevisiae* did not result in accumulation of astaxanthin until *crtR* was co-expressed [54,56].

The biosynthetic pathway of astaxanthin from β -carotene in *Adonis* is different than the above mentioned organisms (Fig. 1). In *Adonis aestivalis*, a carotenoid- β -ring-4-dehydrogenase (CBFD) hydroxylates the fourth carbon of β -ionone ring, then the hydroxylated products are further dehydrogenated into 4-keto group by carotenoid-4-hydroxy- β -ring-4-dehydrogenase (HBFD) [17,57]. The hydroxyl group on the third carbon of β -ionone ring is also introduced by the CBFD enzyme [17].

2.2. Astaxanthin biosynthesis regulation

2.2.1. Transcriptional regulations of astaxanthin biosynthesis

Given their protective rule against stress, carotenoids including astaxanthin are believed to be synthesized in response to different stressors [58,59]. Several studies have been conducted on the natural producers *H. pluvialis*, *C. zofingiensis* and *X. dendrorhous* to reveal the

transcriptional regulations of astaxanthin biosynthesis.

Multiple stressors have been reported to induce astaxanthin synthesis in *H. pluvialis* including high light, nitrogen limitation, and chemical inducers. High Light can cause elevated levels of reactive oxygen species (ROS) which in turn promotes the accumulation of astaxanthin [60]. However, astaxanthin biosynthesis seems to be influenced not only by the light intensity, but the wavelength of light as well. For instance, red light was found to enhance the biomass, while blue light promotes astaxanthin production in a poorly understood mechanism [61,62]. Blue light signaling is known to induce ROS in algae [60]. Thus upon the exposure to the blue light, significant upregulation in the expression level of catalase has been detected in *H. pluvialis* suggesting the exposure of the cells to oxidative stress which in turn led to up-regulation of the genes encoding β -carotene ketolase and hydroxylase [62]. Additionally, changes in the expression levels of the blue light receptors encoded by *CPH1* and *PHOT* genes have been detected and influenced the astaxanthin accumulation, suggesting their role in mediating the signal for astaxanthin biosynthesis [61,62]. For example, upon the upregulation in expression level of *PHOT*, up-regulation of the carotenoid biosynthesis genes *PSY* and *PDS* was detected [61]. Exogenous supplementation with some chemicals can play a synergetic effect in enhancing astaxanthin production as well. Addition of Fe^{2+} under high light can increase astaxanthin levels indirectly through promoting oxidative stress and influencing photosynthesis by up-regulation of photosynthesis-antenna genes including *Lhca1*, *Lhca3*, *Lhca4*, *Lhcb2*, *Lhcb3*, *Lhcb5*, *Lhcb6* and *Lhcb7* [63]. Under high light conditions, γ -aminobutyric acid supplementation resulted in significant improvement of the biomass and subsequently the astaxanthin yields [64]. The γ -Aminobutyric acid addition up-regulated several stress resistance related genes (*PP2C*, *SnRK2*, *CPK*, *HSP90*, *WRKY1*, *PR-1*) and led to increased stress resistance of the algal cells. Additionally, upregulation of *Lhca2* and *PTOX* genes was detected upon the supplementation which resulted in enhancement of light protection and led to improved photosynthetic activity of the cells. The induction of the endogenous methyl jasmonate pathway by ethanol, upregulated *DXS*, the key enzyme in MEP pathway, and carotenogenic genes such as *PSY*, *bkt* and *crtR-B*, which resulted in enhanced astaxanthin accumulation [65].

Although, many transcriptomic analyses revealed the change in the expression level of key genes in astaxanthin biosynthesis upon the exposure to stress in *H. pluvialis*, the regulatory mechanism of these changes is not clear. Transcription factors (TFs) are important regulatory proteins that play important rule in the activation or repression of the expression of their target genes [66]. The bHLH family TFs are reported to be carotenogenesis suppressors in plants [67]. In consistent with these findings, nitrogen limitation in the presence of high light can induce high level of astaxanthin accumulation in *H. pluvialis*, which was accompanied with downregulation of the TFs from bHLH family [68]. On the other hand, C3H, MYB, Nin-like, MTB related and ERF TFs were highly expressed TF families under the effects of salicylic acid in the presence of high light [68,69]. In addition to TFs, some miRNAs also have been identified as responding miRNAs to sodium acetate and high light stress [70]. A total of 83 and 46 miRNAs were considered as light and sodium acetate stress responsive miRNAs, respectively, and 14 miRNAs responded to both stresses. The identified miRNAs were targeting several genes involved in signal transduction, heavy metal stress response, and secondary metabolism. Moreover, four miRNAs (miR482d, PC-3p-859521_20, miR167, and PC-5p-411755_80) involved in regulating astaxanthin synthesis by directly targeting the astaxanthin biosynthetic genes, *lcy-B*, *GGPS*, *PDS* [70].

Many studies have been focused on the genetic control of carotenogenesis in the red yeast *X. dendrorhous*, however the regulations of these processes are not well understood yet [71]. The MVA and sterols biosynthetic pathways are important for astaxanthin biosynthesis as precursors supply and storage medium, respectively. The transcription factor Sre1 has been identified as a regulator that regulates these pathways by directly regulating the *ERG10*, *HMGS*, and *HMGR* genes

whose upregulations is beneficial for astaxanthin synthesis [72,73]. Similar to *H. pluvialis*, the medium components can influence astaxanthin production in this yeast. Catabolic repression by glucose has been reported in *X. dendrorhous* and has been proved to affect astaxanthin production [74,75]. Such repression is mediated by a metabolic suppressor encoded by *MIG1* gene along with the co-repressor complex Cyc8-Tup1. The deletion of *MIG1* upregulated *crtI*, *crtYB* and *crtS* transcripts in presence of glucose [74]. Likewise, deletion of *CYC8* and *TUP1*, upregulated the genes involved in the synthesis of carotenoids precursors including *HMGR*, *IDI* and *FPS* genes [75]. Exogenous supplementation with the phytohormone 6-benzylaminopurine (6-BAP) can induce the astaxanthin synthesis by upregulating the transcription levels of related genes *HMGR*, *IDI* and *crtYB* [76].

2.2.2. Crosstalk between lipid and astaxanthin biosynthesis

The astaxanthin biosynthetic pathway is interfering with that of lipids including fatty acids, TAG and sterols (Fig. 2). On the other hand, due to their hydrophobic nature, carotenoids including astaxanthin are incorporated in the lipid rich cellular compartments including plasma membranes and lipid bodies (LBs), which will be discussed in details in section 2.3.2 [77,78]. Thus, the relation between lipid and astaxanthin biosynthesis is complex and seems to be dependent on the organism and the astaxanthin biosynthetic pathway itself. For instance, in the red yeast *X. dendrorhous*, astaxanthin biosynthesis is competing on the substrate acetyl-CoA which is also the precursor for fatty acids and ergosterol biosynthesis [79]. Miao et al. reported an astaxanthin over-producing mutant where fatty acids production was reduced compared to the wild type, suggesting the inverse correlation between the two pathways in this yeast [80]. Furthermore, the inhibition of the fatty acids biosynthetic pathway in the wild-type strain using triclosan resulted in 2 fold and 1000 fold increase in astaxanthin and ergosterol

content, respectively, which might be attributed to increased acetyl-CoA availability and re-directing the flux toward the mevalonate pathway [81].

Nevertheless, the most popular astaxanthin producing algae, *H. pluvialis* and *C. zofingiensis*, showed different patterns in astaxanthin accumulation in response to fatty acids inhibition [36,51]. These two algae have been reported to accumulate significant amount of TAG and astaxanthin simultaneously in response to stress conditions [82,83]. Additionally, astaxanthin is predominantly accumulated in esterified form in these two algae and stored in TAG filled LBs which acts as a solvent for the synthesized astaxanthin. Because astaxanthin biosynthesis in algae is derived from MEP pathway, and it competes with fatty acid and TAG biosynthesis on pyruvate, there might be a potential regulatory mechanism between both pathways [84,85]. A previous report suggested the presence of a minimal TAG that might be needed to induce the biosynthesis of esterified astaxanthin and accumulation in LBs in *H. pluvialis* [86]. The chemical inhibition of fatty acids biosynthesis using norflurazon and cerulenin in this alga resulted in significant reduction in TAG content, the major fatty acids in astaxanthin esters, such as oleic acid (C18:1) and linolic acid (C18:2), and total astaxanthin [51,87]. Additionally, the cerulenin treated cells showed higher free astaxanthin content in spite of the decrease in the total astaxanthin level suggesting a possible feedback inhibition of the free astaxanthin on its biosynthesis, which is believed to be relieved through esterification. In consistent with this hypothesis, the total astaxanthin level has been restored upon fatty acids supplementation to the cerulenin treated algal cells. On the other hand, cerulenin treated *C. zofingiensis* showed enhanced total astaxanthin level upon the inhibition of fatty acids biosynthesis which is opposite to *H. pluvialis* [36,84,85,88]. Some Studies tried to find explanations for this behavior. For instance, Liu et al. showed that the increase in astaxanthin might be due to enhanced




Fig. 2. The crosstalk between lipid and astaxanthin biosynthesis and the hypothetical distribution of astaxanthin in microorganisms. The Solid arrows indicate a single step reaction. The dashed arrows indicate multistep reactions. The gold arrows represent the lipid biosynthesis pathways. The blue arrows represent the MVA pathway. The green arrows represent the MEP pathway. The red arrows represent the astaxanthin biosynthesis pathway. G3P, glyceraldehyde-3-phosphate; DAG, diacylglycerol; TAG, triacylglycerol; DMAPP, dimethylallyl pyrophosphate; IPP, isopentenyl pyrophosphate; FPP, farnesyl pyrophosphate; PA, phosphatidic acid; LB, lipid body; PM, plasma membrane. (For interpretation of the references to color in this figure legend, the reader is referred to the Web version of this article.)

conversion of other carotenoids into astaxanthin inside the cell, supported by decrease of β -carotene and canthaxanthin in alignment with the increase of astaxanthin and cerulenin concentration [85]. Additionally, cerulenin might induce ROS which might serve as inducer for astaxanthin biosynthesis [36,88]. Another explanation is that, the content of astaxanthin in *C. zofingiensis* is much lower than *H. pluvialis*, however they have almost the same content of total fatty acids. Therefore, in contrast to *H. pluvialis*, the level of TAG might not be a limiting factor for astaxanthin accumulation in *C. zofingiensis* and the fatty acid inhibition might result in enhanced precursor and energy supply for carotenogenesis [88].

Sterols represent a class of lipid that have been found in eukaryotic cells with a great importance in maintaining the membrane integrity and fluidity [89]. Ergosterol is a major lipid component of the fungal cell membranes and helps to maintain the membrane integrity in a manner similar to cholesterol in mammalian cells [90]. In the astaxanthin overproducing *X. dendrorhous* mutant mentioned above, besides the reduced fatty acid content, Miao et al. also reported reduction in the ergosterol content [80]. Similarly, chemical inhibition of ergosterol biosynthesis using fluconazol resulted in 5 fold increase in astaxanthin [81]. Astaxanthin and ergosterol biosynthetic pathways are sharing the substrate FPP, so the flux reduction of the competitive pathway in the mutant is conducive to astaxanthin synthesis, which provides a possible route for metabolic engineering in *X. dendrorhous* [79]. Additionally, the key enzyme HMGR in the MVA pathway have been reported to be negatively regulated by sterols in some fungal species and mammalian cells [89]. The deletion of *CYP61* (*ERG5*), encoding C22-sterol desaturase, which is involved in the late steps of ergosterol biosynthesis, resulted in enhanced HMGR transcription level and 2 fold increase in astaxanthin and carotenoids production in *X. dendrorhous* [91]. Further investigations on this mutant revealed that ergosterol depletion activates the sterol regulatory element-binding protein (SREBP) pathway by the activation of Sre1 protein which is involved in the activation of several genes involved in MVA pathway, sterol biosynthesis and carotenogenesis [72]. However, the deletion of the genes *ERG3* and *ERG4* which are involved in the preceding and the following step to the Erg5 mediated reaction, respectively, had no effect on carotenogenesis, despite the inability to accumulate ergosterol, suggesting the inactivation of SREBP in these mutants [92]. Thus, the activation of SREBP in the Δ erg5 mutant might be dependent on the sterol composition, ergosta-5,7,24(28)-trienol in particular, rather than ergosterol itself. It worth noting that, along with TAG and astaxanthin, an increase in sterols occurred in response to high light stress in *H. pluvialis* [93]. Additionally, sterols biosynthesis is derived from the IPP synthesized in the chloroplast of *H. pluvialis* through MEP pathway which competes with carotenoids biosynthesis [93]. However, the effect of sterol biosynthesis on astaxanthin or carotenoids have not been studied yet in this algae.

2.3. Astaxanthin biosynthesis and storage distribution

The study of the subcellular location of the biosynthesis and storage of astaxanthin is of great importance for engineering astaxanthin overproducers, however only limited studies have been conducted to reveal this mystery in the native astaxanthin producers.

2.3.1. Astaxanthin biosynthesis localization

Given that the biosynthesis of astaxanthin is a multistep reaction with several enzymes, it is possible that the biosynthesis localization involves several subcellular locations. For instance, astaxanthin biosynthesis in *H. pluvialis* is believed to be divided between the chloroplast and the endoplasmic reticulum (ER), where β -carotene is produced in the chloroplast and then transported to the ER by unknown mechanism to be converted to astaxanthin followed by esterification step [1,51,94]. This is consistent with the fact that, carotenogenesis in algae is derived from MEP pathway which is originated in the chloroplast. Additionally, several enzymes involved in β -carotene biosynthesis

were reported to be in the chloroplast [95,96]. At last, *in vitro* assay using several fractions of *H. pluvialis* cell lysates showed that β -carotene conversion to astaxanthin is associated with ER containing fractions [51]. It should be noted that, β -carotene hydroxylase has been detected in LBs and chloroplast membranes [52]. However, ketolase activity was only associated with LBs [52]. LBs are known to be generated from ER in eukaryotes, which might be a possible reason for the detection of astaxanthin biosynthetic activity in the ER containing fractions [51,97].

In contrary to the extensively studied *H. pluvialis*, limited information about the biosynthesis and storage localization are available for the yeast *X. dendrorhous*. Verdoes et al. hypothesized a membrane bound carotenogenic complex for astaxanthin production in this yeast [98]. Additionally, astaxanthin biosynthesis from β -carotene is mediated by a bi-functional p450 monooxygenase (CrtS) which is believed to be localized to the ER [54,99]. Thus, it is possible that carotenogenesis is localized to the ER, which needs further experiments to be confirmed.

2.3.2. Astaxanthin storage localization

As mentioned previously, carotenoids are hydrophobic and they tend to be stored in membranes and LBs due to their high lipid content, however astaxanthin distribution in the cells appears to be influenced by the type of the astaxanthin produced whether it is esterified or in a free form (Fig. 2). In *H. pluvialis*, 95% of astaxanthin is in esterified form and it is believed to be accumulated in the LBs [51]. However, analysis of extracts from thylakoid membranes of *H. pluvialis* showed accumulation of free and esterified astaxanthin in the membranes [100]. Furthermore, at the late stages of the growth period, the ratio of the thylakoid membranes/whole cell free astaxanthin content were much higher than that of the astaxanthin ester and was around 50% of the whole free astaxanthin content, suggesting higher affinity of the free form to the membranes [100]. However, the possibility of contamination of these membranes with LBs during the cell lysates preparation cannot be ruled out [101].

Contrarily, *X. dendrorhous* produces astaxanthin in its free form. Electron paramagnetic resonance spectroscopy experiment on *X. dendrorhous* membranes proved the incorporation of astaxanthin in plasma membrane [102]. In another study, laser confocal fluorescence microscopy analysis (LCFM) for the yeast *X. dendrorhous*, suggested the accumulation of the carotenoids in LBs [103,104]. LBs are composed of a core filled with TAG and sterol esters enclosed in a monolayer of phospholipids [97]. A previous study showed that polar carotenoids (zeaxanthin) are mainly localized to the phospholipid layer of artificial LBs, while apolar carotenoids (β -carotene) were in the core of the LBs [105]. Given the polarity of astaxanthin, due to the two hydroxyl and two keto groups, it is possible that free astaxanthin might be incorporated in the LBs membranes which needs further confirmation. However, the incorporation of carotenoids in membranes is believed to be toxic to the cells, and resulted in several responses included cell membrane destruction, oxidative stress, induction of drug stress responses, in addition to decreasing in the plasma membrane fluidity [106–109]. Carotenoids are embedded in membranes in different orientations dependent on their polarity (Fig. 2), where the apolar carotenoids are perpendicular to the membrane lipid chains and the polar carotenoids are in a parallel orientation [102]. Therefore, the effect of the polar carotenoid on the membranes fluidity is stronger.

3. Strategies for microbial cell factories optimization

There is a growing demand for the sustainable production of astaxanthin in microbial cell factories via metabolic engineering. Several efforts have been taken to enhance the production of astaxanthin in the native producers, however the lack of genetic manipulation tools for these organisms limited their development. Thus, the development of non-native producers for astaxanthin using other microorganisms such as *Escherichia coli*, *Saccharomyces cerevisiae*, *Yarrowia lipolytica* has attracted a great attention (Table 1) [18]. In the past decade, the

biosynthesis of astaxanthin in these non-native microbes has achieved high yields [3]. Compared to the native organisms, these microbes are rapidly growing on low cost substrates and can easily reach high cell density within few days, in addition to the availability of mature genetic manipulation tools for their metabolic engineering. However, the use of *E. coli* in food grade products biosynthesis has several health concerns due to its ability to produce endotoxins. Alternatively, *S. cerevisiae* is a GRAS organism and was proven to be suitable for the production of carotenoids and terpenoid [27]. Additionally, the non-conventional yeast *Y. lipolytica* has attracted a great attention for the production of several hydrophobic compounds, due to its ability to accumulate lipids up to 60% of its dry weight in its LBs [110]. This oleaginous yeast has been successfully engineered for production of lycopene and β -carotene with significant high yields up to 21 mg/g DCW and 90 mg/g DCW, respectively [111–114]. Although, carotenoids biosynthesis in non-native microbes have achieved high yield in several hosts but the transformation from β -carotene to astaxanthin is still low [18,56,115–117]. The possible limitations might include enzymes expression levels and their substrates preferences, the accessibility of enzyme to substrates, precursors and cofactors supply, products and intermediates feedback inhibition, the interference with other biosynthetic pathways, and physiological stress due to the accumulation of astaxanthin or its intermediates. In this section we will review the recent strategies that have been applied to overcome these limitations in the above mentioned microbial cell factories to achieve high yields of astaxanthin.

3.1. Directing the flux toward astaxanthin production

Several strategies have been adopted for directing the flux toward astaxanthin through metabolic engineering by enhancing the precursors supply and optimization of astaxanthin biosynthetic enzymes.

3.1.1. Enhancing precursors and cofactors supply

β -Carotene is the common building block for the biosynthesis of astaxanthin in all the reported pathways. The pathway to produce β -carotene starting from MEP/MVA precursors contains several rate limiting steps that influence its accumulation. Accordingly, and in order to achieve high levels of astaxanthin, an optimization to the β -Carotene biosynthesis is needed.

IPP and DMAPP are essential for isoprenoid production including carotenoids. Therefore, adequate supply of these substrates plays a critical role in enhanced isoprenoid production. MVA or MEP pathways are the main source for the native supply of these key precursors. Thus, upregulation of these two pathways is believed to enhance the IPP and DMAPP supply and isoprenoid production, which can be achieved by optimizing the rate limiting steps or the heterologous expression/overexpression of the whole pathway. In the MVA pathway, HMGR (3-hydroxy-3-methylglutaryl coenzyme A reductase) is a major rate-limiting enzyme which is regulated at translational and post-translational levels by a feedback inhibition system [118]. HMGR is regulated at the translation level through feedback inhibition that is believed to be mediated by mevalonate [119]. Additionally, the enzyme degradation is modulated through a feedback inhibition mediated by sterols, which can be relieved by the deletion of its N-terminus transmembrane domain [37]. The overexpression of this enzyme or its truncated form showed enhancement in the production of isoprenoids including astaxanthin in *S. cerevisiae* and *Y. lipolytica* [116,120–122]. Similar to HMGR, the rate limiting enzyme 1-deoxy-D-xylulose-5-phosphate synthase (DXS) in the MEP pathway is feedback inhibited by IPP and DMAPP [123]. The overexpression of DXS resulted in significant enhancement in β -carotene which was optimized for astaxanthin production in *E. coli* [124,125]. The overexpression of the MEP pathway enzymes IspD (MCT) and IspF (MDS) led to 71% increase in astaxanthin [126]. Although, the overexpression of genes involved in MEP or/and MVA pathways is beneficial for isoprenoid production, the toxicity of the intermediates limits the enhancement of the IPP supply and the subsequent increase in

isoprenoid [40]. For instance, the introduction of MVA pathway in addition to the native MEP pathway in *E. coli* has been found to be toxic due to the accumulation of HMG-CoA [127]. Thus, balancing the expression level of the genes involved in the MVA pathway was needed for enhanced isoprenoid production including astaxanthin [127–129]. Additionally, the overexpression of IspG (HDS) enzyme from the MEP pathway results in accumulation of HMBPP and reduced growth, which requires the co-expression of IspH (HDR) enzyme to alleviate its toxicity and enhance astaxanthin production [124,130]. Some toxic effects have been detected upon the accumulation of IPP in the cells, thus the overexpression of *idi* to convert IPP to DMAPP is always adopted to reduce its toxicity [124,125,131].

Another important intermediate in β -carotene biosynthetic pathway is FPP. As mentioned previously, carotenoids biosynthesis is competing with sterol biosynthesis on FPP [79]. However, the complete in-activation of sterol biosynthesis is not possible due to its deleterious effects on the cells, fungal hosts in particular. Thus the downregulation of the sterol biosynthetic pathway might be more feasible. In this context, the downregulation of squalene synthase encoded by *SQS1* from *Y. lipolytica* by promoter truncation resulted in enhanced β -carotene which further promoted astaxanthin production [116,117]. Alternatively, similar effect can be achieved by directing the flux toward GGPP formation by overexpressing highly active GGPPs like CrTE03 M mutant [122], GGPPs7 from *Synechococcus* sp. [117], or GGPP synthase from *Archaeoglobus fulgidus* [131]. Direct synthesis of GGPP from IPP was achieved through *in vivo* enzyme assembly of GGPP synthase (CrtE) and Idi by fusing a pair of short peptide RIDD and RIAD to their C-termini, respectively, which guided the construction of multienzyme complex to prevent intermediate diffusion and resulted in 2.7 fold increase in astaxanthin level in *E. coli* [128].

The optimization of the two rate limiting steps of phytoene synthesis and lycopene cyclization catalyzed by phytoene synthase (PSY or CrtB) and lycopene cyclase (CrtY), respectively, is critical for enhanced carotenoids production [132]. Thus for effective β -carotene and astaxanthin production, the bifunctional phytoene synthase and lycopene cyclase CrtYB enzyme from *X. dendrorhous* has been overexpressed in several host for pathway optimization [116,117,122].

Aside from substrates, the cofactors NADPH and ATP are essential for carotenoids biosynthesis. Hence, in order to boost astaxanthin production, it is necessary to balance the supply of the cofactors. The central carbon metabolism plays a vital role in supplying these cofactors, NADPH is mainly generated through the pentose phosphate pathway (PPP) and malic enzyme reaction, while ATP is mainly generated through electron transport chain starting from NADH formed in the TCA cycle [133]. Based on this and in order to optimize the β -carotene biosynthesis in astaxanthin producing *E. coli*, combined upregulation of TCA cycle and PPP was adopted by overexpressing *sucAB*, *sdh* and *talB* [37,38,41,124].

3.1.2. Optimization of astaxanthin biosynthetic enzymes

3.1.2.1. Balancing the levels and activities of β -carotene hydroxylases and ketolases.

One of the bottlenecks of astaxanthin biosynthesis in engineered microorganisms, is the accumulation of several intermediates during the conversion of β -carotene into astaxanthin (Fig. 1). The astaxanthin biosynthetic enzymes from different sources have different substrate preferences. For example, the CrtW from *Paracoccus* sp. N81106 shows a strong substrate preference for carotenoids with non hydroxylated β -ionone rings [53]. On the other hand, zeaxanthin (hydroxylated) is used as main substrate by CrtW from *Brevundimonas* sp. strain SD212 [134]. The CrtZ from *Erwinia uredovora* showed substrate preference for β -carotene, while the CrtZ from *Alcaligenes* sp. PC-1 showed higher activity toward canthaxanthin [135]. Accordingly, to efficiently convert β -carotene into astaxanthin, selecting the appropriate combination of enzymes with desired substrate preference is necessary.

A screening of nine CrtZ and eight CrtW from different organisms was conducted to efficiently produce astaxanthin in *S. cerevisiae* [136]. As a result of the screening, a CrtW from *B. vesicularis* DC263 and CrtZ from *Alcaligenes* sp. strain PC-1 were verified to be the best combination for highest astaxanthin yield (3.1 mg/g DCW) with less intermediates accumulation.

In addition to the substrate preferences, the expression level of CrtZ and CrtW is critical for intermediates accumulation control, which can be adjusted through optimization of the copy number of each gene, promoter strength or ribosome binding site engineering [18,27]. For instance, replacement of the CrtZ gene promoter with a stronger one increased the enzyme level ratio of CrtZ to CrtW and yielded 30.4% increase in astaxanthin [136]. In order to finely optimize the ratio the two enzymes, screening of several RBS libraries obtained through introducing six random nucleotides to the RBS region of these genes resulted in optimized cassette with a titer of 3.46 mg/g DCW and 99% astaxanthin [137]. Further increase by 67.6% was obtained by multiple integration of the optimized cassette. In addition, since the expression level of a the heterogeneous gene is generally proportional to its copy number, adjustment of the copy number of CrtZ and CrtW have been successfully applied to enhance astaxanthin biosynthesis in several hosts [116,117,122,138]. For example, introducing additional copy of CrtZ from *P. ananatis* resulted in 68.5% increase in the ratio of astaxanthin to total carotenoid in *E. coli* [138].

Besides the above mentioned strategies to balance the activity of the two enzymes, ensuring the proper folding of the enzymes is critical for obtaining their maximum activity. Thus, to optimize the folding of CrtZ and CrtW, the GroES-GroEL molecular chaperone were overexpressed in *E. coli* which resulted in 1.45 fold increase in astaxanthin with a yield of 1.18 g/L [124].

3.1.2.2. Enzyme engineering of β -carotene hydroxylases and ketolases. As stated previously, astaxanthin biosynthesis from β -carotene is mediated by the activity of CrtW and CrtZ in algae and bacterial cells. The two enzymes are iron (Fe^{2+}) and oxygen dependent oxygenases, which was supported by several *in vitro* assays [53,135,139]. Additionally, the conserved iron binding motifs exist among all sequences of different CrtZ and CrtW strongly supports that the Fe^{2+} is essential for the reaction [135]. The lack of information about the structure of these enzymes up to now is limiting their rational engineering. Thus, the majority of the studies on characterizing and enhancing the properties of these enzymes are mainly focused on alanine-scanning mutagenesis, directed evolution, fusion proteins constructs through artificial linkers, and directed localization of the enzymes through signal peptides.

In order to characterize the amino acid residues which are critical for the enzymes activity, site directed mutagenesis in the conserved amino acid residues among the protein families is of a great help. Similar to other oxygen-dependent and iron-containing integral membrane enzymes, CrtZ and CrtW contain putative iron-binding His motifs and have multiple transmembrane domains with several conserved amino acid residues [18,140]. Therefore, alanine-scanning mutagenesis of these residues at the His motifs and the transmembrane domains was performed to further study the CrtW from *Paracoccus* sp. strain N81106 [140]. Among the conserved amino acids at the three His motifs of CrtW, six His residues (H69, H103, H106, H107, H221 and H222) were essential for its activity, while partial activity was retained upon on the alanine substitution of the three His residues (H65A, H218A, and H219A). Additionally, alanine substitution of the Y134A, F176A, F118A, P116A, W126A, W229A and L232A residues in the transmembrane domains resulted in decreased activity. Site directed mutagenesis to the sumoylation site (K90R) of a β -carotene hydroxylase from *H. pluvialis* resulted in 1.34 fold enhancement in astaxanthin production, which is believed to be due to enhancement on the protein stability [141].

Directed evolution via error-prone PCR has been successfully

employed for enhancing the activity of CrtW from different sources. Six mutations (H96L,R203W, A205V, A208V, F213L and A215T) in the CrtW from *Sphingomonas* sp. DC18 showed improved activity toward hydroxylated carotenoids which resulted in enhanced astaxanthin production in *E. coli* [142]. Enhancement of astaxanthin levels was achieved upon three mutations in L175 M, M99V, and M99I residues of the CrtW from *Paracoccus* sp. N81106 [140]. The mutations increased the affinity of CrtW toward adonixanthin without any significant effect on β -carotene conversion rate. A triple mutant (H165R/V264D/F298Y) of β -carotene ketolase from *H. pluvialis* showed enhanced β -carotene conversion rate which resulted in 2.4 fold increase in canthaxanthin level and subsequent increase in astaxanthin level by 93% in *S. cerevisiae* [143]. Additional mutation at the (M1T/N188D/L271R) residues further enhanced increased the canthaxanthin yield by 51% and decreased the β -carotene accumulation by 44% [144]. In the same study, *H. pluvialis* β -carotene hydroxylase mutant at L288R residue showed 33% enhanced astaxanthin production compared to the wild type. Furthermore, the co-expression of the these two mutants resulted in 39% astaxanthin increase [144]. Another triple mutant (A6T, T105A and L239 M) of CrtW from *Brevundimonas* sp. SD212 enhanced astaxanthin level by 5.35 fold in *E. coli* [137]. The mutant showed enhanced conversion of the intermediates and astaxanthin reached 92% of the total carotenoids produced by this strain.

Creating artificial scaffolds of the key enzymes of a synthetic pathway has been proved as effective strategy for enhancing the efficiency of the pathway by reducing the undesirable intermediates diffusion [145]. In this context, a fusion construct of CrtZ from plant and CrtW from *Brevundimonas* sp. strain SD212 using flexible linker resulted in 1.4 fold increase in astaxanthin level when expressed in *E. coli* [146]. β -carotene has been reported to accumulate in the plasma membrane of engineered *E. coli*, thus bringing the enzymes to close proximity to their substrate might be beneficial to enhance the productivity [147]. Accordingly, and regardless of their predicted transmembrane domains, the directed localization of a fusion protein of CrtW and CrtZ from *Brevundimonas* sp. strain SD212 and *Pantoea agglomerans*, respectively, by linking them to the membrane protein GlpF led to 215.4% improved astaxanthin levels [148]. Likewise, the N-terminus fusion of a signal peptide of the outer membrane protein ompF to truncated BKT from *Chlamydomonas reinhardtii* led to 31% increase in astaxanthin in *E. coli* [126]. Further increase by 34% was obtained upon enhancing the stability of the protein by C-terminus fusion with *E. coli* thioredoxin (*trxA*) as molecular chaperone. In a recent study on astaxanthin production using *Y. lipolytica*, directed localization of a fusion construct of CrtZ from *H. pluvialis* and CrtW from *Paracoccus* sp to the ER, β -carotene biosynthesis location; LBs and peroxisomes, hypothetical β -carotene storage location; resulted in 4.8 fold (139.4 mg/L) increase in astaxanthin level [149]. Further optimization of the culture using fed-batch shake flask fermentation led to a yield of 858 mg/L which is the highest reported in yeast.

3.2. Membrane stress management strategies

The accumulation of astaxanthin and carotenoids in membranes induces toxic effect to the cells, which subsequently limits their yield. Directing the storage to the LBs has been an effective strategy to increase carotenoids production and relieve their toxicity in *S. cerevisiae* and *Y. lipolytica* [114,150,151]. Therefore, directing the storage of astaxanthin to the LBs is promising for optimizing astaxanthin accumulation in engineered microorganisms. However, free astaxanthin is highly probable to be stored in membranes due to its structure and polarity. Astaxanthin in *H. pluvialis* and *C. zofingiensis* is mainly in esterified form and accumulates in LBs, yet the mechanism and the enzymes involved in esterification are not clearly identified [51,84,152]. Astaxanthin esterification in *H. pluvialis* and *C. zofingiensis* is speculated to be mediated by diacylglycerol acyltransferases (DGATs) and the long-chain-alcohol O-fatty-acyltransferase (AAT), respectively, [51,152]. However, in

in vitro assay using microsomal fraction of *S. cerevisiae* expressing the *C. zofingensis* AAT did not show esterification activity toward astaxanthin [153]. Another approach can be adopted by enhancing the water solubility of astaxanthin. Carotenoids glycosylation is known to enhance their solubility [154,155]. For instance, zeaxanthin di-glycosylation mediated by zeaxanthin glycosyltransferase from *Erwinia herbicola* enhanced its water solubility from 12.6 ppm to 800 ppm [154]. Glycosylated astaxanthin has been reported in some microorganisms such as *Paracoccus* sp. N81106 (*Agrobacterium aurantiacum*) and *Sphingomonas astaxanthinifaciens* and showed higher solubility in water based solvents compared to free astaxanthin [11,13]. Few attempts have been made to produce glycosylated astaxanthin in *E. coli* through the heterologous expression of zeaxanthin glycosyltransferase (CrtX) from *E. uredoovora* [12,155]. However, several glycosylated intermediates have been detected which might be due to the non-specificity of the enzyme toward astaxanthin [155]. Furthermore, the total carotenoids level has been decreased accompanied with increase in the accumulation of un-converted β -carotene suggesting an interruption of the astaxanthin biosynthetic pathway upon the expression of CrtX. It should be noted that, optimization of the glycosyl donor supply should be considered for efficient production of glycosylated astaxanthin.

Alternatively, enhancing the capacity of the membranes for storage via membrane engineering strategies can be an effective tool for enhanced astaxanthin production. The knock-out of the membrane morphology genes, *lpp*, *bambB*, and *rodZ*, in *E. coli* resulted in increase in the astaxanthin content by 40% [156]. Upon the knock-out, the *E. coli* cells exhibited elongated morphology which in turn increased the membrane surface area for the storage of astaxanthin. Another strategy has been adopted to increase the membrane surface area by introducing inner and outer membrane vesicles in *E. coli* [157]. Inner membrane vesicles were induced by the heterologous expression of the caveolin protein (Cav1) which is involved in the formation of the cell membranes invaginations (caveolae), while the outer membrane vesicles were formed by the repression of *rffD* and *rfaD* genes which are involved in the outer membrane integrity, and an increase in astaxanthin titer by 50% and 62% was achieved, respectively [157].

The orientation of astaxanthin in the plasma membrane enables its diffusion through the membrane [129]. Therefore, the secretion of astaxanthin to the medium can reduce the membrane stress, the potential astaxanthin mediated feedback inhibition, and the extraction costs. However due to its hydrophobicity, the amount of secreted astaxanthin in the culture medium is very limited. Accordingly, the use of biocompatible organic solvents can facilitate the secretion of astaxanthin in the medium [115].

3.3. Dynamic control systems: a promising strategy for astaxanthin production

The manipulation of cellular metabolism through metabolic engineering strategies including tuning gene expression, gene deletions, as well as protein engineering has enabled improvement in the production levels of many valuable products in microbial cell factories. However directing the flux toward non-native biosynthetic pathways creates burden to the cells due to the competition over the resources and the energy supplies with the native metabolic pathways, in addition to the possible toxicity of the products [158]. Such burden can influence the cell growth and the yield of the desired product. Thus, to overcome these restrictions, decoupling of the cell growth and the biosynthesis of the product of interest is needed, which can be achieved through dynamic control system [158]. Several studies have been conducted to control the production of several products via dynamic control systems, however few examples are available for astaxanthin [159,160]. A temperature dependent dynamic control system has been developed using the temperature-responsive Gal4M9-system in *S. cerevisiae* which resulted in 235 mg/L astaxanthin through fed-batch fermentation [144]. Similarly, decoupling the growth and the production in *E. coli* using the inducible

IPTG promoter resulted in 64% increase in astaxanthin [129]. Although dynamic control systems are promising in optimizing the production of several products, but several limitations still exist including the lack of known sensors for the biosynthetic/toxic intermediates and the limited information about native sensor-response systems in some promising microbial cell factories [161,162].

3.4. Directed evolution of microbial cell factories

The development of ideal microbial cell factories is challenging even with the advances in molecular biology and biotechnology, due to the complexity of intracellular reaction networks in living cells and the limited knowledge about their regulation, which limit the appropriate rational design. Therefore, directed evolution of the whole cell factory is another effective strategy to improve its performance. The majority of the directed evolution research has been made on the native producers *X. dendrorhous* and *H. pluvialis* and little has been done on the non-native producers. In this section we summarize the directed evolution strategies used in astaxanthin producing strains including mutagenesis and adaptive laboratory evolution (ALE) (Tables 2 and 3).

3.4.1. Chemical and physical mutagenesis

Chemical mutagenesis, physical mutagenesis (UV and Gamma irradiation), and atmospheric and room temperature plasma (ARTP) are the most commonly used mutagenesis strategies for microbial cell factories directed evolution [163]. Gamma rays mediated mutagenesis have been employed to increase the production of astaxanthin in *X. dendrorhous* and led to 1.77 fold increase in astaxanthin [164]. ARTP mutagenesis for astaxanthin producing *S. cerevisiae* promoted astaxanthin level by 83% [165]. Treatment of *H. pluvialis* cells with ethyl methane sulfonate (EMS) and UV increased astaxanthin by 2.38 fold and 2.17 fold, respectively [166]. In order to obtain a better mutagenic effect, a combination of physical and chemical mutagenesis is frequently used. A three-stage mutagenesis to *H. pluvialis* has been conducted using UV mutagenesis followed by EMS and screening based on the resistance to the inhibitor diphenylamine (DPA), resulted in selection of a mutant with 1.7 fold enhancement in astaxanthin production compared to the wild type cells [167].

3.4.2. Adaptive laboratory evolution

Over the last few years, adaptive laboratory evolution (ALE) has received a significant attention for the development and optimization of many microbial cell factories [168]. ALE is conducted under laboratory conditions where the cells are subjected to controlled culture conditions until a desired mutant is obtained upon the exposure to several stressors. However, to obtain the desired mutant, a proper selection of the stressors is required. Due to their antioxidant activities, carotenoids producing organisms including astaxanthin producers are believed to be more fit to oxidative stress [32,169]. Accordingly, adaptive evolution of *X. dendrorhous* upon the exposure to six oxidizing agents (ionone, diphenylamine, NaCl, TiO₂, H₂O₂ and NaClO) have been conducted to get an astaxanthin overproducing strain [170]. As a result, an evolved mutant with 48.2% enhancement in astaxanthin was obtained in response to the oxidative stress caused by TiO₂. In another study, combined ARTP mutagenesis and ALE using H₂O₂ led to 4 fold increase in astaxanthin in *S. cerevisiae* [171].

3.4.3. Screening methods

For an ideal directed evolution process at the enzymes or whole cell level, an efficient rapid screening method is needed to select the desired mutant. Owing to its red color, visual color screening depending on the change in color intensity is the most common and easiest method for the selection of astaxanthin overproducers [144,172]. However, the background due to interference of other intermediates (i.e. canthaxanthin) and the color saturation make it challenging [144]. The use of astaxanthin biosynthesis inhibitors diphenylamine (DPA) and nicotine, or the

Table 2
Chemical and physical mutagenesis strategies for astaxanthin production.

Hosts	Mutagenesis methods	Mutagenic conditions	Fermentation scale	Titer (mg/L)	Yield (mg/g DCW)	Fold change	Ref
<i>H. pluvialis</i>	UV + EMS	15min /0.12% (w/w)	–	–	–	3.2	[166]
	UV + EMS + DES	11min/2.0% (V/V) /0.1% (V/V)	shake-flask 15 days	90	47.21	1.7	[167]
<i>X. dendrorhous</i>	γ -irradiation	5.0 kGy	shake-flask 8 days	2.6	15.9	1.77	[164]
	ARTP + UV	50s/40s	shake-flask 96 h	–	–	1.2	[246]
<i>S. cerevisiae</i>	ARTP	30s or 40s	5-L bioreactor 84 h	217.9	13.8	3.26	[165]

Table 3
ALE strategies for astaxanthin production.

Hosts	ALE stress agents	ALE conditions	ALE rounds	Fermentation scale	Titer (mg/L)	Yield (mg/g DCW)	Fold change	Ref
<i>X. dendrorhous</i>	H ₂ O ₂	10 mmol/L	1	shake-flask 120 h	10.4	1.3	1.83	[247]
	TiO ₂	300 mg/L	150	shake-flask 48 h	1.22	–	1.48	[170]
	TiO ₂	500 mg/L	5	shake-flask 72 h	14.74	–	2	[248]
<i>S. cerevisiae</i>	H ₂ O ₂ +ARTP	10 L/min/30%	1	5-L fermenter 210 h	65.9	–	4	[171]

expression of astaxanthin biosynthetic genes under weak promoters can help to overcome these limitations and facilitate the screening [144,166,167]. Other screening methods using flow-cytometry and near-infrared spectroscopy have been employed to screen astaxanthin hyper-producers [173–175]. A high-throughput screening using an Raman-Activated Droplet Sorting (RADS) showed promising results for screening astaxanthin hyper-producers [176]. Using astaxanthin producing *H. pluvialis* variants as a model, a sorting accuracy of 98.3%, enrichment rate of 8 folds, and throughput of ~260 cells/min was achieved.

4. Astaxanthin fermentation optimization

Besides the improvement of the microbial cell factories performance through the above mentioned strategies, the appropriate optimization of the fermentation conditions contributes greatly to boost their productivity and reduce the production cost. Therefore, for optimum and cost efficient industrial scale production of astaxanthin using these organisms, native producers in particular, adequate knowledge of their fermentation process is crucial. As a secondary metabolite, the biosynthesis of astaxanthin is not required for the growth of the cells. Additionally, several reports indicated the dependence of astaxanthin biosynthesis on several environmental stress conditions, which are not the optimum conditions for the cell growth. Thus, the fermentation strategy adopted for astaxanthin production is a two-stage fermentation, where the cell growth is promoted at the first stage, followed by astaxanthin biosynthesis stimulation stage. In this section we will discuss the most significant factors that stimulate astaxanthin biosynthesis including stress conditions, chemical supplementation, pH, temperature and dissolved oxygen (DO).

4.1. Stress and cultivation strategies

High light intensity, nutrients restriction and high salinity are the major stress conditions that trigger astaxanthin biosynthesis in *H. pluvialis* [177]. While, the use of high light intensities requires high amount of energy, nutrients restriction is more convenient in terms of cost and implementation for industrial scale production of astaxanthin. Nitrogen starvation is believed to induce astaxanthin production effectively [178–181]. A recent study showed 25% increase in astaxanthin

using a sequential stress strategy involved extended nitrogen starvation followed by moderate light intensity exposure at the late palmella stage [177]. Moderate sodium chloride concentration promotes astaxanthin production, while high concentration has an inhibitory effect [182,183]. In *C. zofingiensis* separate application of high light (HL) or salinity stress (SS) can moderately enhance astaxanthin accumulation, while the combinatorial HL and SS can significantly enhance the production [83]. The current large-scale cultivation of *H. pluvialis* for astaxanthin production is a complex two-stage culture system, the green stage and red stage. While the green stage is for obtaining high biomass, and the red stage is characterized by the synthesis and accumulation of astaxanthin in response to inducers. At present, studies on the production of astaxanthin from *H. pluvialis* are mainly focused on optimizing the culture and induction conditions [184]. Several fermentation strategies have been developed to save the cost and increase the productivity of astaxanthin using this microalgae including: sequential heterotrophic-phototrophic culture, one-step continuous culture, indoor artificial light culture, biofilm cultivation and sequential mixotrophic culture [183,185–196].

Light can induce carotenogenesis *X. dendrorhous*, however high light intensities are lethal to the cells [197]. Irradiation with white and ultraviolet light resulted in increase of astaxanthin yield in *X. dendrorhous* by 85% and 97%, respectively in a shake flask [198]. Furthermore, large scale fermentation using 10 L and 800 L fermenters under white light irradiation led to astaxanthin yield of 420 mg/L (4.7 mg/g) and 350 mg/L (4.1 mg/g), respectively [198]. High C/N is believed to promote astaxanthin production in *X. dendrorhous*, however high glucose could inhibit the cell growth in. Accordingly, Yamane et al. developed a two-stage fed-batch culture: where the first stage was controlled at low C/N to promote the cell growth, followed by a production stage at high C/N which significantly enhanced astaxanthin production [199]. Similarly, a two stage fed-batch fermentation was conducted with a controlled glucose concentration of 25 g/L at the lag and the early log phases, and 5 g/L at the late log and the stationary phases and 109% increase in astaxanthin was obtained [200]. Sonication has a positive effect on enzyme activity and microbial processes [201]. In this context, exposure of *X. dendrorhous* culture to ultrasonic stimulation led to 26% enhancement of astaxanthin [201].

4.2. Chemical supplements

In addition to light and nutrient deficiency, addition of chemical supplements to the culture media have a significant influence on astaxanthin production. Exogenous addition of acetate to the culture medium of *H. pluvialis* led to enhancement in astaxanthin by 2 fold compared to the control group [202]. Further increase can be obtained through combined oxidative stress with acetate supplementation by addition of ferrous ions or active oxygen species (singlet oxygen, superoxide anion radical, hydrogen peroxide and peroxy radical) [203]. Addition of the cationic polymer polyethyleneimine led to enhanced oxidative stress which was accompanied with 5 fold increase in astaxanthin [204]. Phytohormones such as methyl jasmonate, gibberellin, salicylic acid have synergetic effect on astaxanthin accumulation in *H. pluvialis* [48,205]. Other studies indicated that the addition of the antioxidants butylated hydroxyanisole and butylated hydroxytoluene could significantly induce astaxanthin synthesis and led to 2.03 and 1.66 fold increase, respectively, however the mechanism of induction is not clear [206,207]. Astaxanthin content in *X. dendrorhous* was enhanced by 40.7% upon glutamate feeding [208]. Ethanol and acetic acid feeding after glucose consumption increased astaxanthin by 31% and 26%, respectively, in *X. dendrorhous* culture [209]. Other studies showed that plant and fungal extracts are beneficial for boosting astaxanthin production using *X. dendrorhous* [210–212].

4.3. pH

The pH of the fermentation broth has a significant influence on the metabolic activities of the microorganisms. *H. pluvialis* can grow on a wide range of pH with an optimal growth pH of 7.0–7.5, while a pH higher than 8.25 promotes astaxanthin biosynthesis [182,213,214]. The parasitic fungus *Paraphysoderma sedebokerensis* is one of the most serious contaminants of *H. pluvialis* cultures with a significant influence on the growth and subsequently the astaxanthin level [215]. An acidic cultivation strategy was developed to prevent infection of *H. pluvialis* with the fungus in culture for astaxanthin production [216]. The strategy involved cultivation at a pH 4 for the growth and astaxanthin biosynthesis, while to relief the pH dependent reduction in astaxanthin, gradual light irradiation in addition to nitrogen deficiency were employed at the induction stage which resulted in 141 fold increase in astaxanthin levels compared to the control.

The optimal pH for astaxanthin biosynthesis in *X. dendrorhous* appears to be dependent on the strain. For instance, the earliest reports showed that the optimal pH for *X. dendrorhous* growth and astaxanthin production was 4.5 [217]. A mutant strain showed optimal astaxanthin production pH of 5.0 while the optimum growth was achieved at pH 6.0 [218]. However, the change in the pH had a great influence on growth, but little effect on astaxanthin production. Contrarily, culture of the *X. dendrorhous* DSMZ5626 under uncontrolled pH condition showed significant inhibition of astaxanthin below 5.5, while the cell growth was significantly inhibited when pH dropped to 4.2 [219]. These findings suggest that the genetic background of the strain influences their response to the pH.

4.4. Temperature

Temperature is another critical factor that affects the growth of the microbial cells. The change in the temperature significantly influence the enzymatic activities of the metabolic pathways enzymes and subsequently the cell growth and astaxanthin production. The optimal temperature for astaxanthin accumulation in *H. pluvialis* is 27–28 °C; and the growth and production rate would decrease significantly at temperatures lower than 15 °C or higher than 30 °C [220–222]. While, the optimum temperature for *X. dendrorhous* is 20–22 °C with a significant growth and production inhibition at temperatures higher than 22 °C [217]. In engineered non-native carotenoid producers, lower

temperatures were favored for carotenoids biosynthesis. For instance, In *S. cerevisiae*, low temperatures were thought to benefit the accumulation of crocein, zeaxanthin, and β -carotene [223,224]. Similarly, a 78.96% increase in astaxanthin and 2 fold increase in the total carotenoids was achieved upon the cultivation of a high-yield astaxanthin producing *S. cerevisiae* strain AX15 at 20 °C [225]. Park et al. studied the effect of different temperature (26, 28 and 30 °C) on astaxanthin accumulation in *E. coli*, and 30 °C was the selected as the optimum temperature for astaxanthin production [226].

4.5. Dissolved oxygen (DO)

DO is a key parameter in astaxanthin fermentation process, due to its direct effect on the energy metabolism of the cells, and the astaxanthin biosynthetic enzymes activity. Yamane et al. found that astaxanthin accumulation in *X. dendrorhous* is directly proportional to the oxygen supply [199]. Further kinetic analysis showed that respiration rate was positively correlated with astaxanthin production and negatively correlated with ethanol production [199]. Wang et al. found that *X. dendrorhous* growth and astaxanthin production would be significantly inhibited when the dissolved oxygen tension was adjusted at 20% saturation [227]. Another study investigated the effect of different agitation speeds (250, 400, and 600 rpm) on *X. dendrorhous* DSMZ5626 in a 2 L bioreactor, and found that DO, biomass and astaxanthin production reached the highest when the speed was the highest, and *vice versa* [219]. Similarly, enhancement of the oxygen level in shake flasks using the biocompatible organic oxygen carrier *n*-hexadecane resulted in 58% increase in the carotenoid yield of *X. dendrorhous* [228]. DO is not critical for astaxanthin biosynthesis by *H. pluvialis*. On the other hand, adequate supply of carbon dioxide is required for the photosynthetic activities, and the oxygen generated through photosynthesis is partially consumed by the astaxanthin biosynthetic pathway [229]. Moderate increase in carbon dioxide supply can lead to increase in astaxanthin production [230]. Furthermore, the introduced CO₂ can alter C/N ration which creates nutrients deficiency condition that act as astaxanthin inducer.

5. Conclusion and outlook

Up to date, the highest yield of astaxanthin was achieved in recombinant *E. coli* at 1.18 g/L by employment of a combination of different engineering strategies, which is much higher than the most popular native producer *H. pluvialis* and *X. dendrorhous* [124]. In contrast to this non-native producer, the optimization of astaxanthin in the native producers *H. pluvialis* and *X. dendrorhous* is mainly through directed evolution and fermentation optimization through chemicals supplementation for the induction of astaxanthin biosynthesis. However, these methods have been successfully implied to increase astaxanthin levels but they are laborious and not cost efficient. This illuminate the power of synthetic biology in optimization of microbial cell factories. However, compared to some high terpenoid producing strains, for example, farnesene high-producing *S. cerevisiae*, there still exist huge room for astaxanthin producing strains for production optimization and enhancement [231]. In addition to the metabolic engineering strategies mentioned above, introduction of non-native pathways to enhance the precursors supply might be of great influence on increasing astaxanthin production. For example, IPP enhancement has been achieved in several organisms via introducing exogenous isoprenol utilizing pathway (IUP), which led to enhanced isoprenoids production [232–234]. Additionally, other promising membrane stress management strategies have to be adopted to relief the burden of astaxanthin accumulation on the cells. Enhancing the membrane fluidity can be achieved through increasing the unsaturated fatty acids content of the membranes [235]. The *OLE1* gene encodes a $\Delta 9$ -fatty acid desaturase which is catalyzing the dehydrogenation of the 9-position in saturated fatty acids forming the corresponding unsaturated fatty acid [236]. The overexpression of this gene

has been successfully employed to increase triterpenoids and β -carotene production in yeast by reducing the membrane stress [237,238]. Furthermore, deep knowledge about gene expression and regulation is needed to improve these microbial cell factories, such as epigenetic modifications, non-coding RNAs and post-translational regulations etc., which are still mysterious. Additionally, highly efficient genetic manipulation tools and techniques are also in urgent demand for native microbial producers and non-conventional but promising yeast such as *Y. lipolytica*. It worth noting, the design of microbial cell factory needs to be closely connected to the industrial process and market needs and applications. For example, to increase the stability and bioavailability of astaxanthin, different forms of astaxanthin obtained through esterification, covalent binding to proteins, or PEGylation need to be investigated. And considering the huge differences between flask cultivation and bioreactor fermentation, the evaluation or screening of engineered strains is preferred to be based on industrially used culture mediums and parallel-bioreactors.

At the end, the market application of natural astaxanthin transformed by synthetic biotechnology have to face supervisory regulations on market access by different countries. The removal of regulatory obstacles is also the key to unlocking attractive market prospects of astaxanthin.

Author statement

Mostafa Basiony: Writing - original draft; Writing - review & editing. Visualization; **Liming Ouyang:** Project administration; Resources; Supervision; Validation; Writing - original draft; Writing - review & editing. **Danni Wang:** partial writing; **Jiaming Yu:** partial writing; **Liming Zhou:** partial writing, Visualization; **Mohan Zhu:** partial writing; **Xuyuan Wang:** partial writing; **Jie Feng:** partial writing; **Jing Dai:** partial writing; **Yijie Shen:** partial writing; **Chengguo Zhang:** partial writing; **Qiang Hua:** Investigation, Supervision; **Xiuliang Yang:** Funding acquisition, Investigation, Resources; **Lixin Zhang:** Funding acquisition, Investigation, Resources.

Declaration of competing interest

Nothing declared.

Acknowledgements

We gratefully acknowledge the financial support from the National Key Research and Development Program of China (2020YFA0907800) and Shandong Jincheng Bio-Pharmaceutical Co., Ltd.

References

- [1] Han DX, Li YT, Hu Q. Astaxanthin in microalgae: pathways, functions and biotechnological implications. *Algae* 2013;28(2):131–47.
- [2] Schmidt I, Schewe H, Gassel S, Jin C, Buckingham J, Hübelin M, et al. Biotechnological production of astaxanthin with *Phaffia rhodozyma*/*Xanthophyllomyces dendrorhous*. *Appl Microbiol Biotechnol* 2011;89(3):555–71.
- [3] Zhang C, Chen X, Too H. Microbial astaxanthin biosynthesis: recent achievements, challenges, and commercialization outlook. *Appl Microbiol Biotechnol* 2020;104(13):5725–37.
- [4] Ambati R, Phang S, Ravi S, Aswathanarayana R. Astaxanthin: sources, extraction, stability, biological activities and its commercial applications—a review. *Mar Drugs* 2014;12(1):128–52.
- [5] Grand View Research Inc.. Astaxanthin market size, share & trends analysis report by product (oil, softgel, liquid), by source (natural, synthetic), by application (aquaculture & animal feed, nutraceuticals), by region, and segment forecasts, 2021–2028. Grand View Research; 2021.
- [6] MarketsandMarkets. Market by source (natural, synthetic), form (dry, liquid), method of production (microalgae cultivation, chemical synthesis, fermentation), application (dietary supplements, food & beverages, cosmetics), and region - global forecast to 2026. MarketsandMarkets; 2021.
- [7] Rao A, Sindhuja H, Dharmesh S, Sankar K, Sarada R, Ravishankar G. Effective inhibition of skin cancer, tyrosinase, and antioxidative properties by astaxanthin and astaxanthin esters from the green alga *Haematococcus pluvialis*. *J Agric Food Chem* 2013;61(16):3842–51.
- [8] Qiu D, Wu Y, Zhu W, Yin H, Yi L. Identification of geometrical isomers and comparison of different isomeric samples of astaxanthin. *J Food Sci* 2012;77(9):C934–40.
- [9] Lorenz RT, Cysewski GR. Commercial potential for *Haematococcus microalgae* as a natural source of astaxanthin. *Trends Biotechnol* 2000;18(4):160–7.
- [10] Asker D, Awad TS, Beppu T, Ueda K. Purification and identification of astaxanthin and its novel derivative produced by radio-tolerant *Sphingomonas astaxanthinifaciens*. *Methods Mol Biol* 2018;1852:171–92.
- [11] Yokoyama A, Adachi K, Shizuri Y. New carotenoid glucosides, astaxanthin glucoside and adonixanthin glucoside, isolated from the astaxanthin-producing marine bacterium, *agrobacterium aurantiacum*. *J Nat Prod* 1995;58(12):1929–33.
- [12] Yokoyama A, Shizuri Y, Misawa NJTL. Production of new carotenoids, astaxanthin glucosides, by *Escherichia coli* transformants carrying carotenoid biosynthesis genes 1998;39:3709–12.
- [13] Asker D, Amano S-i, Morita K, Tamura K, Sakuda S, Kikuchi N, et al. Astaxanthin dirhamnoside, a new astaxanthin derivative produced by a radio-tolerant bacterium, *Sphingomonas astaxanthinifaciens*. *J Antibiot* 2009;62(7):397–9.
- [14] Rezanek T, Nedbalová L, Sigler K, Cepák V. Identification of astaxanthin diglucoside diesters from snow alga *Chlamydomonas nivalis* by liquid chromatography–atmospheric pressure chemical ionization mass spectrometry. *Phytochemistry* 2008;69(2):479–90.
- [15] Matsushita Y, Suzuki R, Kotake-Nara E, Yokoyama A, Miyashita K. Antioxidant activity of polar carotenoids including astaxanthin- β -glucoside from marine bacterium on PC liposomes. *Fish Sci* 2008;66:980–5.
- [16] Lim KC, Yusoff FM, Shariff M, Kamarudin MS. Astaxanthin as feed supplement in aquatic animals, vol. 10; 2018. p. 738–73.
- [17] Cunningham Jr FX, Gantt E. Elucidation of the pathway to astaxanthin in the flowers of *Adonis aestivalis*. *Plant Cell* 2011;23(8):3055–69.
- [18] Wan X, Zhou X, Moncalian G, Su L, Chen W, Zhu H, et al. Reprogramming microorganisms for the biosynthesis of astaxanthin via metabolic engineering. *Prog Lipid Res* 2021;81:101083.
- [19] Khoo KS, Lee SY, Ooi CW, Fu X, Miao X, Ling TC, et al. Recent advances in biorefinery of astaxanthin from *Haematococcus pluvialis*. *Bioresour Technol* 2019;288:121606.
- [20] Shah MMR, Liang Y, Cheng JJ, Daroch M. Astaxanthin-producing green microalga *Haematococcus pluvialis*: from single cell to high value commercial products. *Front Plant Sci* 2016:531.
- [21] Zhang Z, Huang J, Sun D, Lee Y, Chen F. Two-step cultivation for production of astaxanthin in *Chlorella zofingiensis* using a patented energy-free rotating floating photobioreactor (RFP). *Bioresour Technol* 2017;224:515–22.
- [22] Panis G, Carreon JR. Commercial astaxanthin production derived by green alga *Haematococcus pluvialis*: a microalgae process model and a techno-economic assessment all through production line. *Algal Res* 2016;18:175–90.
- [23] Tao L, Rouvière PE, Cheng Q. A carotenoid synthesis gene cluster from a non-marine *Brevundimonas* that synthesizes hydroxylated astaxanthin. *Gene* 2006;379:101–8.
- [24] Asker D, Beppu T, Ueda K. *Sphingomonas astaxanthinifaciens* sp. nov., a novel astaxanthin-producing bacterium of the family *Sphingomonadaceae* isolated from Misasa, Tottori, Japan. *FEMS (Fed Eur Microbiol Soc) Microbiol Lett* 2007;273(2):140–8.
- [25] Lee JH, Kim YS, Choi T-J, Lee WJ, Kim YT. *Paracoccus haeundaensis* sp. nov., a Gram-negative, halophilic, astaxanthin-producing bacterium 2004;54(5):1699–702.
- [26] Ide T, Hoya M, Tanaka T, Harayama S. Enhanced production of astaxanthin in *Paracoccus* sp. strain N-81106 by using random mutagenesis and genetic engineering. *Biochem Eng J* 2012;65:37–43.
- [27] Li C, Swofford CA, Sinskey AJ. Modular engineering for microbial production of carotenoids. *Metabolic Engineering Communications* 2020;10:e00118.
- [28] Johnson E. *Phaffia rhodozyma*: colorful odyssey. *Int Microbiol : the official journal of the Spanish Society for Microbiology* 2003;6(3):169–74.
- [29] Calo P, González T. The yeast *Phaffia rhodozyma* as an industrial source of astaxanthin. *Microbiologia* 1995;11(3):386–8.
- [30] Gassel S, Schewe H, Schmidt I, Schrader J, Sandmann G. Multiple improvement of astaxanthin biosynthesis in *Xanthophyllomyces dendrorhous* by a combination of conventional mutagenesis and metabolic pathway engineering. *Biotechnol Lett* 2013;35(4):565–9.
- [31] Li Y, Gong F, Guo S, Yu W, Liu J. *Adonis amurensis* is a promising alternative to as a resource for natural esterified (3,3')-astaxanthin production. *Plants* 2021;10(6):1059.
- [32] Guerin M, Huntley ME, Olaizola M. *Haematococcus astaxanthin*: applications for human health and nutrition. *Trends Biotechnol* 2003;21(5):210–6.
- [33] Koller M, Muhr A, Braunnegg G. Microalgae as versatile cellular factories for valued products. *Algal Research-Biomass Biofuels and Bioproducts* 2014;6:52–63.
- [34] Andrewes AGSM. 3R,3'R)-Astaxanthin from the yeast *Phaffia rhodozyma*. *Phytochemistry* 1976;15(6):3.
- [35] Eisenreich W, Bacher A, Arigoni D, Rohdich F. Biosynthesis of isoprenoids via the non-mevalonate pathway. *Cell Mol Life Sci* 2004;61(12):1401–26.
- [36] Liu J, Mao X, Zhou W, Guarnieri MT. Simultaneous production of triacylglycerol and high-value carotenoids by the astaxanthin-producing oleaginous green microalga *Chlorella zofingiensis*. *Bioresour Technol* 2016;214:319–27.
- [37] Wang Q, Quan S, Xiao H. Towards efficient terpenoid biosynthesis: manipulating IPP and DMAPP supply. *Bioresources and Bioprocessing* 2019;6(1):6.
- [38] Moise AR, Al-Babili S, Wurtzel ET. Mechanistic aspects of carotenoid biosynthesis. *Chem Rev* 2014;114(1):164–93.

- [39] Gruchattka E, Hädicke O, Klamt S, Schütz V, Kayser O. In silico profiling of *Escherichia coli* and *Saccharomyces cerevisiae* as terpenoid factories. *Microb Cell Factories* 2013;12:84.
- [40] Li M, Hou F, Wu T, Jiang X, Li F, Liu H, et al. Recent advances of metabolic engineering strategies in natural isoprenoid production using cell factories. *Nat Prod Rep* 2020;37(1):80–99.
- [41] Liao P, Hemmerlin A, Bach TJ, Chye M-L. The potential of the mevalonate pathway for enhanced isoprenoid production. *Biotechnol Adv* 2016;34(5):697–713.
- [42] Martin JF, Gudina E, Barredo JL. Conversion of beta-carotene into astaxanthin: two separate enzymes or a bifunctional hydroxylase-ketolase protein? *Microb Cell Factories* 2008;7:3.
- [43] Lee JH, Kim YT. Cloning and characterization of the astaxanthin biosynthesis gene cluster from the marine bacterium *Paracoccus haemundensis*. *Gene* 2006;370:86–95.
- [44] Kakizono SKT, Kondo TSK, Nishio TON, Misawa SNN. Isolation and function identification of a novel cDNA for astaxanthin biosynthesis from *Haematococcus pluvialis* and astaxanthin synthesis in *Escherichia coli*. *Plant Mol Biol* 1995;29:343–52.
- [45] Linden H. Carotenoid hydroxylase from *Haematococcus pluvialis* cDNA sequence, regulation and functional complementation. *Biochim Biophys Acta* 1999;1446:203–12.
- [46] Lotan T, Hirschberg J. Cloning and expression in *Escherichia coli* of the gene encoding β -C-4-oxygenase, that converts fl-carotene to the ketocarotenoid canthaxanthin in *Haematococcus pluvialis*. *FEBS (Fed Eur Biochem Soc) Lett* 1995;364:125–8.
- [47] Huang JC, Chen F, Sandmann G. Stress-related differential expression of multiple beta-carotene ketolase genes in the unicellular green alga *Haematococcus pluvialis*. *J Biotechnol* 2006;122(2):176–85.
- [48] Lu Y, Jiang P, Liu S, Gan Q, Cui H, Qin S. Methyl jasmonate- or gibberellins A3-induced astaxanthin accumulation is associated with up-regulation of transcription of beta-carotene ketolase genes (bkts) in microalga *Haematococcus pluvialis*. *Bioresour Technol* 2010;101(16):6468–74.
- [49] Luo Q, Bian C, Tao M, Huang Y, Zheng Y, Lv Y, et al. Genome and transcriptome sequencing of the astaxanthin-producing green microalga, *Haematococcus pluvialis*. *Genome biology and evolution* 2019;11(1):166–73.
- [50] Huang D, Liu W, Li A, Hu Z, Wang J, Wang C. Cloning and identification of a novel β -carotene hydroxylase gene from *Haematococcus pluvialis* and its function in *Escherichia coli*. *Algal Res* 2021;55:102245.
- [51] Chen G, Wang B, Han D, Sommerfeld M, Lu Y, Chen F, et al. Molecular mechanisms of the coordination between astaxanthin and fatty acid biosynthesis in *Haematococcus pluvialis* (Chlorophyceae). *Plant J : for cell and molecular biology* 2015;81(1):95–107.
- [52] Grünewald K, Hirschberg J, Hagen C. Ketocarotenoid biosynthesis outside of plastids in the unicellular green alga *Haematococcus pluvialis*. *J Biol Chem* 2001;276(8):6023–9.
- [53] Fraser PD, Shimada H, Misawa N. Enzymatic confirmation of reactions involved in routes to astaxanthin formation, elucidated using a direct substrate in vitro assay. *Eur J Biochem* 1998;252:229–36.
- [54] Alcaíno J, Fuentealba M, Cabrera R, Baeza M, Cifuentes V. Modeling the interfacial interactions between CrtS and CrtR from *Xanthophyllomyces dendrorhous*, a P450 system involved in astaxanthin production. *J Agric Food Chem* 2012;60(35):8640–7.
- [55] Alcaíno J, Barahona S, Carmona M, Lozano C, Marcoleta A, Niklitschek M, et al. Cloning of the cytochrome p450 reductase (crtR) gene and its involvement in the astaxanthin biosynthesis of *Xanthophyllomyces dendrorhous*. *BMC Microbiol* 2008;8(1):169.
- [56] Ukibe K, Hashida K, Yoshida N, Takagi H. Metabolic engineering of *Saccharomyces cerevisiae* for astaxanthin production and oxidative stress tolerance. *Appl Environ Microbiol* 2009;75(22):7205–11.
- [57] Huang J, Zhong Y, Sandmann G, Liu J, Chen F. Cloning and selection of carotenoid ketolase genes for the engineering of high-yield astaxanthin in plants. *Planta* 2012 Aug;236(2):691–9.
- [58] Fang N, Wang C, Liu X, Zhao X, Liu Y, Liu X, et al. De novo synthesis of astaxanthin: from organisms to genes. *Trends Food Sci Technol* 2019;92:162–71.
- [59] Avalos J, Carmen Limón M. Biological roles of fungal carotenoids. *Curr Genet* 2015;61(3):309–24.
- [60] Fu W, Guðmundsson Ó, Paglia G, Herjólfsson G, Andrésón ÓS, Pálsson BØ, et al. Enhancement of carotenoid biosynthesis in the green microalga *Dunaliella salina* with light-emitting diodes and adaptive laboratory evolution. *Appl Microbiol Biotechnol* 2012;97(6):2395–403.
- [61] Ma R, Thomas-Hall SR, Chua ET, Eltanahy E, Netzel ME, Netzel G, et al. Blue light enhances astaxanthin biosynthesis metabolism and extraction efficiency in *Haematococcus pluvialis* by inducing haematocyst germination. *Algal Res* 2018;35:215–22.
- [62] Lee C, Ahn JW, Kim JB, Kim JY, Choi YE. Comparative transcriptome analysis of *Haematococcus pluvialis* on astaxanthin biosynthesis in response to irradiation with red or blue LED wavelength. *World J Microbiol Biotechnol* 2018;34(7):96.
- [63] He B, Hou L, Dong M, Shi J, Huang X, Ding Y, et al. Transcriptome analysis in *Haematococcus pluvialis*: astaxanthin induction by high light with acetate and Fe²⁺. *Int J Mol Sci* 2018;19(1):175.
- [64] Li L, Chen Z, Huang Q. Exogenous γ -aminobutyric acid promotes biomass and astaxanthin production in *Haematococcus pluvialis*. *Algal Res* 2020;52:102089.
- [65] Liu Y-H, Alimujiang A, Wang X, Luo S-W, Balamurugan S, Yang W-D, et al. Ethanol induced jasmonate pathway promotes astaxanthin hyperaccumulation in *Haematococcus pluvialis*. *Bioresour Technol* 2019:289.
- [66] Spitz F, Furlong EE. Transcription factors: from enhancer binding to developmental control. *Nat Rev Genet* 2012;13(9):613–26.
- [67] Toledo-Ortiz G, Johansson H, Lee KP, Bou-Torrent J, Stewart K, Steel G, et al. The HY5-PIF regulatory module coordinates light and temperature control of photosynthetic gene transcription. *PLoS Genet* 2014;10(6):e1004416.
- [68] Hoys C, Romero-Losada AB, del Río E, Guerrero MG, Romero-Campero FJ, García-González M. Unveiling the underlying molecular basis of astaxanthin accumulation in *Haematococcus* through integrative metabolomic-transcriptomic analysis. *Bioresour Technol* 2021;332:125150.
- [69] Hu Q, Huang D, Li A, Hu Z, Gao Z, Yang Y, et al. Transcriptome-based analysis of the effects of salicylic acid and high light on lipid and astaxanthin accumulation in *Haematococcus pluvialis*. *Biotechnol Biofuels* 2021;14(1):1–20. 2021/4//;14(1).
- [70] Wang X, Miao X, Chen G, Cui Y, Sun F, Fan J, et al. Identification of microRNAs involved in astaxanthin accumulation responding to high light and high sodium acetate (NaAC) stresses in *Haematococcus pluvialis*. *Algal Res* 2021;54:102179.
- [71] Bellora N, Moline M, David-Palma M, Coelho MA, Hittinger CT, Sampaio JP, et al. Comparative genomics provides new insights into the diversity, physiology, and sexuality of the only industrially exploited tremellomycete: *phaffia rhodozyma*. *BMC Genom* 2016;17(1):901.
- [72] Gutiérrez MS, Campusano S, González AM, Gómez M, Barahona S, Sepúlveda D, et al. Sterol regulatory element-binding protein (Sre1) promotes the synthesis of carotenoids and sterols in *xanthophyllomyces dendrorhous*. *Front Microbiol* 2019;10:586.
- [73] Gómez M, Campusano S, Gutiérrez MS, Sepúlveda D, Barahona S, Baeza M, et al. Sterol regulatory element-binding protein Sre1 regulates carotenogenesis in the red yeast *Xanthophyllomyces dendrorhous*. *J Lipid Res* 2020;61(12):1658–74.
- [74] Alcaíno J, Bravo N, Córdova P, Marcoleta AE, Contreras G, Barahona S, et al. The involvement of Mig1 from *xanthophyllomyces dendrorhous* in catabolic repression: an Active mechanism contributing to the regulation of carotenoid production. *PLoS One* 2016;11(9). e0162838-e.
- [75] Córdova P, Alcaíno J, Bravo N, Barahona S, Sepúlveda D, Fernández-Lobato M, et al. Regulation of carotenogenesis in the red yeast *Xanthophyllomyces dendrorhous*: the role of the transcriptional co-repressor complex Cyc8–Tup1 involved in catabolic repression. *Microb Cell Factories* 2016;15(1):1–19.
- [76] Pan X, Wang B, Duan R, Jia J, Li J, Xiong W, et al. Enhancing astaxanthin accumulation in *Xanthophyllomyces dendrorhous* by a phytohormone: metabolomic and gene expression profiles. *Microbial Biotechnology* 2020;13(5):1446–60.
- [77] Gruszecki WI. Carotenoids in membranes. In: Frank HA, Young AJ, Britton G, Cogdell RJ, editors. *The photochemistry of carotenoids*. Dordrecht: Springer Netherlands; 1999. p. 363–79.
- [78] Begum H, Yusoff FMD, Banerjee S, Khatoon H, Shariff M. Availability and utilization of pigments from microalgae. *Crit Rev Food Sci Nutr* 2016;56(13):2209–22.
- [79] Visser H, van Ooyen AJ, Verdoes JC. Metabolic engineering of the astaxanthin-biosynthetic pathway of *Xanthophyllomyces dendrorhous*. *FEMS Yeast Res* 2003;4(3):221–31.
- [80] Miao L, Wang Y, Chi S, Yan J, Guan G, Hui B, et al. Reduction of fatty acid flux results in enhancement of astaxanthin synthesis in a mutant strain of *Phaffia rhodozyma*. *J Ind Microbiol Biotechnol* 2010;37(6):595–602.
- [81] Miao L, Chi S, Tang Y, Su Z, Yin T, Guan G, et al. Astaxanthin biosynthesis is enhanced by high carotenogenic gene expression and decrease of fatty acids and ergosterol in a *Phaffia rhodozyma* mutant strain. *FEMS Yeast Res* 2011;11(2):192–201.
- [82] Gwak Y, Hwang Y-s, Wang B, Kim M, Jeong J, Lee C-G, et al. Comparative analyses of lipidomes and transcriptomes reveal a concerted action of multiple defensive systems against photooxidative stress in *Haematococcus pluvialis*. *J Exp Bot* 2014;65(15):4317–34.
- [83] Kou Y, Liu M, Sun P, Dong Z, Liu J. High light boosts salinity stress-induced biosynthesis of astaxanthin and lipids in the green alga *Chromochloris zofingiensis*. *Algal Res* 2020;50:101976.
- [84] Zhang Y, Ye Y, Bai F, Liu J. The oleaginous astaxanthin-producing alga *Chromochloris zofingiensis*: potential from production to an emerging model for studying lipid metabolism and carotenogenesis. *Biotechnol Biofuels* 2021;14(1):119.
- [85] Zhang Y, Ye Y, Ding W, Mao X, Li Y, Gerken H, et al. Astaxanthin is ketolated from zeaxanthin independent of fatty acid synthesis in *chromochloris zofingiensis*. *Plant Physiol* 2020;183(3):883–97.
- [86] Zhekisheva M, Zarka A, Khozin-Goldberg I, Cohen Z, Boussiba S. Inhibition OF astaxanthin synthesis under high irradiance does not abolish triacylglycerol accumulation IN the green alga *haematococcus PLUVIALIS*. *CHLOROPHYCEAE* 2005;41(4):819–26.
- [87] Holtin K, Kuehnle M, Rehbein J, Schuler P, Nicholson G, Albert K. Determination of astaxanthin and astaxanthin esters in the microalgae *Haematococcus pluvialis* by LC-(APCI)MS and characterization of predominant carotenoid isomers by NMR spectroscopy. *Anal Bioanal Chem* 2009;395(6):1613.
- [88] Zhang Z, Sun D, Mao X, Liu J, Chen F. The crosstalk between astaxanthin, fatty acids and reactive oxygen species in heterotrophic *Chlorella zofingiensis*. *Algal Res* 2016;19:178–83.
- [89] Espenshade PJ, Hughes AL. Regulation of sterol synthesis in eukaryotes. *Annu Rev* 2007;41:401–27.
- [90] Zhang Y, Rao R. Beyond ergosterol: linking pH to antifungal mechanisms. *Virulence* 2010;1(6):551–4.

- [91] Loto I, Gutiérrez MS, Barahona S, Sepúlveda D, Martínez-Moya P, Baeza M, et al. Enhancement of carotenoid production by disrupting the C22-sterol desaturase gene (CYP61) in Xanthophyllomyces dendrorhous. *BMC Microbiol* 2012;12:235.
- [92] Venegas M, Barahona S, González AM, Sepúlveda D, Zúñiga GE, Baeza M, et al. Phenotypic analysis of mutants of ergosterol biosynthesis genes (ERG3 and ERG4) in the red yeast xanthophyllomyces dendrorhous. *Front Microbiol* 2020;11:1312.
- [93] Scodelaro Bilbao PG, Garelli A, Díaz M, Salvador GA, Leonardi PI. Crosstalk between sterol and neutral lipid metabolism in the alga *Haematococcus pluvialis* exposed to light stress. *Biochim Biophys Acta Mol Cell Biol Lipids* 2020;1865(10):158767.
- [94] Grünwald K, Hagen C. β -carotene is the intermediate exported from the chloroplast during accumulation of secondary carotenoids in *Haematococcus pluvialis*. *J Appl Phycol* 2001;13(1):89–93.
- [95] Grünwald K, Eckert M, Hirschberg J, Hagen C. Phytoene desaturase is localized exclusively in the chloroplast and up-regulated at the mRNA level during accumulation of secondary carotenoids in *Haematococcus pluvialis* (Volvocales, chlorophyceae). *Plant Physiol* 2000;122(4):1261–8.
- [96] Steinbrenner J, Linden H. Regulation of two carotenoid biosynthesis genes coding for phytoene synthase and carotenoid hydroxylase during stress-induced astaxanthin formation in the green alga *Haematococcus pluvialis*. *Plant Physiol* 2001;125(2):810–7.
- [97] Walther TC, Chung J, Farese Jr RV. Lipid droplet biogenesis. *Annu Rev Cell Dev Biol* 2017;33:491–510.
- [98] Verdoes JC, Sandmann G, Visser H, Diaz M, van Mossel M, van Ooyen AJJ. Metabolic engineering of the carotenoid biosynthetic pathway in the yeast *Xanthophyllomyces dendrorhous* (*Phaffia rhodozyma*). *Appl Environ Microbiol* 2003;69(7):3728–38.
- [99] Ojima K, Breitenbach J, Visser H, Setoguchi Y, Tabata K, Hoshino T, et al. Cloning of the astaxanthin synthase gene from *Xanthophyllomyces dendrorhous* (*Phaffia rhodozyma*) and its assignment as a beta-carotene 3-hydroxylase/4-ketolase. *Mol Genet Genom* : MGG 2006;275(2):148–58.
- [100] Mascia F, Girolomoni L, Alcocer MJP, Bargigia I, Perozeni F, Cazzaniga S, et al. Functional analysis of photosynthetic pigment binding complexes in the green alga *Haematococcus pluvialis* reveals distribution of astaxanthin in Photosystems. *Sci Rep* 2017;7(1):16319.
- [101] Ota S, Morita A, Ohnuki S, Hirata A, Sekida S, Okuda K, et al. Carotenoid dynamics and lipid droplet containing astaxanthin in response to light in the green alga *Haematococcus pluvialis*. *Sci Rep* 2018;8(1):5617.
- [102] Blasko A, Belagyi J, Dergez T, Deli J, Papp G, Papp T, et al. Effect of polar and non-polar carotenoids on Xanthophyllomyces dendrorhous membranes by EPR. *Eur Biophys J : EBJ (Eur Biophys J)* 2008;37(7):1097–104.
- [103] An GH, Bielich J, Auerbach R, Johnson EA. Isolation and characterization of carotenoid hyperproducing mutants of yeast by flow cytometry and cell sorting. *Biotechnology* 1991;9(1):70–3.
- [104] Johnson EA, An G-H. Astaxanthin from microbial sources. *Crit Rev Biotechnol* 1991;11(4):297–326.
- [105] Borel P, Grolrier P, Armand M, Partier A, Lafont H, Lairon D, et al. Carotenoids in biological emulsions: solubility, surface-to-core distribution, and release from lipid droplets. *J Lipid Res* 1996 Feb;37(2):250–61.
- [106] Sung WS, Lee IS, Lee DG. Damage to the cytoplasmic membrane and cell death caused by lycopene in *Candida albicans*. *J Microbiol Biotechnol* 2007 Nov;17(11):1797–804.
- [107] Siems W, Sommerburg O, Schild L, Augustin W, Langhans CD, Wiswedel I. Beta-carotene cleavage products induce oxidative stress in vitro by impairing mitochondrial respiration. *Faseb J : official publication of the Federation of American Societies for Experimental Biology* 2002 Aug;16(10):1289–91.
- [108] Liu P, Sun L, Sun Y, Shang F, Yan G. Decreased fluidity of cell membranes causes a metal ion deficiency in recombinant *Saccharomyces cerevisiae* producing carotenoids. *J Ind Microbiol Biotechnol* 2016 Apr;43(4):525–35.
- [109] Verwaal R, Jiang Y, Wang J, Daran JM, Sandmann G, van den Berg JA, et al. Heterologous carotenoid production in *Saccharomyces cerevisiae* induces the pleiotropic drug resistance stress response. *Yeast* 2010 Dec;27(12):983–98.
- [110] Qiao K, Wasylenko TM, Zhou K, Xu P, Stephanopoulos G. Lipid production in *Yarrowia lipolytica* is maximized by engineering cytosolic redox metabolism. *Nat Biotechnol* 2017;35(2):173–7.
- [111] Schwartz C, Frogue K, Misa J, Wheeldon I. Host and pathway engineering for enhanced lycopene biosynthesis in *Yarrowia lipolytica* 2017;8(2233).
- [112] Matthäus F, Ketelhot M, Gatter M, Barth G. Production of lycopene in the non-carotenoid-producing yeast *Yarrowia lipolytica*. *Appl Environ Microbiol* 2014;80(5):1660–9.
- [113] Zhang XK, Wang DN, Chen J, Liu ZJ, Wei LJ, Hua Q. Metabolic engineering of β -carotene biosynthesis in *Yarrowia lipolytica*. *Biotechnol Lett* 2020;42(6):945–56.
- [114] Larroude M, Celinska E, Back A, Thomas S, Nicaud JM, Ledesma-Amaro R. A synthetic biology approach to transform *Yarrowia lipolytica* into a competitive biotechnological producer of β -carotene. *Biotechnol Bioeng* 2018;115(2):464–72.
- [115] Li N, Han Z, O'Donnell T, Kurasaki R, Kajihara L, Williams P, et al. Production and excretion of astaxanthin by engineered *Yarrowia lipolytica* using plant oil as both the carbon source and the biocompatible extractant. *Appl Microbiol Biotechnol* 2020;104(16):6977–89.
- [116] Kildegaard KR, Adiego-Perez B, Domenech Belda D, Khangura JK, Holkenbrink C, Borodina I. Engineering of *Yarrowia lipolytica* for production of astaxanthin. *Synth Syst Biotechnol* 2017;2(4):287–94.
- [117] Tramontin LRR, Kildegaard KR, Sudarsan S, Borodina I. Enhancement of astaxanthin biosynthesis in oleaginous yeast *Yarrowia lipolytica* via microalgal pathway. *Microorganisms* 2019;7(10).
- [118] DeBose-Boyd RA. Feedback regulation of cholesterol synthesis: sterol-accelerated ubiquitination and degradation of HMG CoA reductase. *Cell Res* 2008;18(6):609–21.
- [119] Burg JS, Espenshade PJ. Regulation of HMG-CoA reductase in mammals and yeast. *Prog Lipid Res* 2011;50(4):403–10.
- [120] Yan GI, Wen KR, Duan CQ. Enhancement of β -carotene production by over-expression of HMG-CoA reductase coupled with addition of ergosterol biosynthesis inhibitors in recombinant *Saccharomyces cerevisiae*. *Curr Microbiol* 2012;64(2):159–63.
- [121] Polakowski T, Stahl U, Lang C. Overexpression of a cytosolic hydroxymethylglutaryl-CoA reductase leads to squalene accumulation in yeast. *Appl Microbiol Biotechnol* 1998;49(1):66–71.
- [122] Zhou P, Xie W, Li A, Wang F, Yao Z, Bian Q, et al. Alleviation of metabolic bottleneck by combinatorial engineering enhanced astaxanthin synthesis in *Saccharomyces cerevisiae*. *Enzym Microb Technol* 2017;100:28–36.
- [123] Rinaldi MA, Ferraz CA, Scrutton NS. Alternative metabolic pathways and strategies to high-titre terpenoid production in *Escherichia coli*. *Nat Prod Rep* 2021;39:90–118.
- [124] Gong Z, Wang H, Tang J, Bi C, Li Q, Zhang X. Coordinated expression of astaxanthin biosynthesis genes for improved astaxanthin production in *Escherichia coli*. *J Agric Food Chem* 2020;68(50):14917–27.
- [125] Lemuth K, Steuer K, Albermann C. Engineering of a plasmid-free *Escherichia coli* strain for improved in vivo biosynthesis of astaxanthin. *Microb Cell Factories* 2011;10(1):29.
- [126] Park SY, Binkley RM, Kim WJ, Lee MH, Lee SY. Metabolic engineering of *Escherichia coli* for high-level astaxanthin production with high productivity. *Metab Eng* 2018;49:105–15.
- [127] Pitera DJ, Paddon CJ, Newman JD, Keasling JD. Balancing a heterologous mevalonate pathway for improved isoprenoid production in *Escherichia coli*. *Metab Eng* 2007;9(2):193–207.
- [128] Kang W, Ma T, Liu M, Qu J, Liu Z, Zhang H, et al. Modular enzyme assembly for enhanced cascade biocatalysis and metabolic flux. *Nat Commun* 2019;10(1):4248.
- [129] Zhang C, Seow VY, Chen X, Too H-P. Multidimensional heuristic process for high-yield production of astaxanthin and fragrance molecules in *Escherichia coli*. *Nat Commun* 2018;9(1):1858.
- [130] Li Q, Fan F, Gao X, Yang C, Bi C, Tang J, et al. Balanced activation of IspG and IspH to eliminate MEP intermediate accumulation and improve isoprenoids production in *Escherichia coli*. *Metab Eng* 2017;44:13–21.
- [131] Wang C-W, Oh M-K, Liao JC. Engineered isoprenoid pathway enhances astaxanthin production in *Escherichia coli*. *Biotechnol Bioeng* 1999;62(2):235–41.
- [132] Tian L. Chapter 32 - carotenoids, genetically modified foods, and vitamin A nutrition. In: Watson RR, Preedy VR, editors. *Genetically modified organisms in food*. San Diego: Academic Press; 2016. p. 353–60.
- [133] Sauer U, Canonaco F, Heri S, Perrenoud A, Fischer E. The soluble and membrane-bound transhydrogenases UdhA and PntAB have divergent functions in NADPH metabolism of *Escherichia coli*. *J Biol Chem* 2004;279(8):6613–9.
- [134] Choi SK, Nishida Y, Matsuda S, Adachi K, Kasai H, Peng X, et al. Characterization of beta-carotene ketolases, CrtW, from marine bacteria by complementation analysis in *Escherichia coli*. *Mar Biotechnol* 2005;7(5):515–22.
- [135] Fraser PD, Miura Y, Misawa N. In vitro characterization of astaxanthin biosynthetic enzymes. *J Biol Chem* 1997;272(10):6128–35.
- [136] Wang R, Gu X, Yao M, Pan C, Liu H, Xiao W, et al. Engineering of β -carotene hydroxylase and ketolase for astaxanthin overproduction in *Saccharomyces cerevisiae*. *Front Chem Sci Eng* 2017;11(1):89–99.
- [137] Li D, Li Y, Xu JY, Li QY, Tang JL, Jia SR, et al. Engineering CrtW and CrtZ for improved biosynthesis of astaxanthin in *Escherichia coli*. *Chin J Nat Med* 2020;18(9):666–76.
- [138] Lu Q, Bu Y-F, Liu J-Z. Metabolic engineering of *Escherichia coli* for producing astaxanthin as the predominant carotenoid, vol. 15; 2017. p. 296.
- [139] Hundle BS, O'Brien DA, Beyer P, Kleinig H, Hearst JE. In vitro expression and activity of lycopene cyclase and beta-carotene hydroxylase from *Erwinia herbicola*. *FEBS Lett* 1993 Jan 11;315(3):329–34.
- [140] Ye RW, Stead KJ, Yao H, He H. Mutational and functional analysis of the beta-carotene ketolase involved in the production of canthaxanthin and astaxanthin. *Appl Environ Microbiol* 2006 Sep;72(9):5829–37.
- [141] Lin YJ, Chang JJ, Lin HY, Thia C, Kao YY, Huang CC, et al. Metabolic engineering a yeast to produce astaxanthin. *Bioresour Technol* 2017 Dec;245(Pt A):899–905.
- [142] Tao L, Wilczek J, Odom JM, Cheng Q. Engineering a β -carotene ketolase for astaxanthin production. *Metab Eng* 2006;8(6):523–31.
- [143] Zhou P, Xie W, Li A, Wang F, Yao Z, Bian Q, et al. Alleviation of metabolic bottleneck by combinatorial engineering enhanced astaxanthin synthesis in *Saccharomyces cerevisiae*. *Enzym Microb Technol* 2017 May;100:28–36.
- [144] Zhou P, Li M, Shen B, Yao Z, Bian Q, Ye L, et al. Directed coevolution of beta-carotene ketolase and hydroxylase and its application in temperature-regulated biosynthesis of astaxanthin. *J Agric Food Chem* 2019 Jan 30;67(4):1072–80.
- [145] Siu K-H, Chen RP, Sun Q, Chen L, Tsai S-L, Chen W. Synthetic scaffolds for pathway enhancement. *Curr Opin Biotechnol* 2015;36:98–106.
- [146] Nogueira M, Effissi EMA, Welsch R, Beyer P, Zurbriggen MD, Fraser PD. Construction of a fusion enzyme for astaxanthin formation and its characterisation in microbial and plant hosts: a new tool for engineering ketocarotenoids. *Metab Eng* 2019 Mar;52:243–52.
- [147] Wu T, Ye L, Zhao D, Li S, Li Q, Zhang B, et al. Membrane engineering - a novel strategy to enhance the production and accumulation of β -carotene in *Escherichia coli*. *Metab Eng* 2017;43:85–91.

- [148] Ye L, Zhu X, Wu T, Wang W, Zhao D, Bi C, et al. Optimizing the localization of astaxanthin enzymes for improved productivity. *Biotechnol Biofuels* 2018;11:278.
- [149] Ma Y, Li J, Huang S, Stephanopoulos G. Targeting pathway expression to subcellular organelles improves astaxanthin synthesis in *Yarrowia lipolytica*. *Metab Eng* 2021;68:152–61.
- [150] Ma T, Shi B, Ye Z, Li X, Liu M, Chen Y, et al. Lipid engineering combined with systematic metabolic engineering of *Saccharomyces cerevisiae* for high-yield production of lycopene. *Metab Eng* 2019;52:134–42.
- [151] Groenewald M, Boekhout T, Neuveglise C, Gaillardin C, van Dijck PW, Wyss M. *Yarrowia lipolytica*: safety assessment of an oleaginous yeast with a great industrial potential. *Crit Rev Microbiol* 2014 Aug;40(3):187–206.
- [152] Zhang Y, Shi M, Mao X, Kou Y, Liu J. Time-resolved carotenoid profiling and transcriptomic analysis reveal mechanism of carotenogenesis for astaxanthin synthesis in the oleaginous green alga *Chromochloris zofingiensis*. *Biotechnol Biofuels* 2019;12:287.
- [153] Xu Y, Pan X, Lu J, Wang J, Shan Q, Stout J, et al. Evolutionary and biochemical characterization of a *Chromochloris zofingiensis* MBOAT with wax synthase and diacylglycerol acyltransferase activity. *J Exp Bot* 2021;72(15):5584–98.
- [154] Hundle BS, O'Brien DA, Alberti M, Beyer P, Hearst JE. Functional expression of zeaxanthin glucosyltransferase from *Erwinia herbicola* and a proposed uridine diphosphate binding site, vol. 89; 1992. p. 9321–5.
- [155] Chen X, Lim X, Bouin A, Lautier T, Zhang C. High-level de novo biosynthesis of glycosylated zeaxanthin and astaxanthin in *Escherichia coli*. *Bioresources and Bioprocessing* 2021;8(1):67.
- [156] Lu Q, Liu J-Z. Enhanced astaxanthin production in *Escherichia coli* via morphology and oxidative stress engineering. *J Agric Food Chem* 2019;67(42):11703–9.
- [157] Yang D, Park SY, Lee SY. Production of Rainbow Colorants by Metabolically Engineered *Escherichia coli* 2021;8:2100743.
- [158] Venayak N, Anesiadis N, Cluett WR, Mahadevan R. Engineering metabolism through dynamic control. *Curr Opin Biotechnol* 2015;34:142–52.
- [159] Shen HJ, Cheng BY, Zhang YM, Tang L, Li Z, Bu YF, et al. Dynamic control of the mevalonate pathway expression for improved zeaxanthin production in *Escherichia coli* and comparative proteome analysis. *Metab Eng* 2016 Nov;38:180–90.
- [160] Xu P. Production of chemicals using dynamic control of metabolic fluxes. *Curr Opin Biotechnol* 2018;53:12–9.
- [161] Madzak C. *Yarrowia lipolytica* strains and their biotechnological applications: how natural biodiversity and metabolic engineering could contribute to cell factories improvement. *J Fungi (Basel)*. 2021;7(7):548.
- [162] Lee H-M, Vo PNL, Na D. Advancement of Metabolic Engineering Assisted by Synthetic Biology 2018;8(12):619.
- [163] Yu Q, Li Y, Wu B, Hu W, He M, Hu G. Novel mutagenesis and screening technologies for food microorganisms: advances and prospects. *Appl Microbiol Biotechnol* 2020 Feb;104(4):1517–31.
- [164] Najafi N, Ahmadi AR, Hosseini R, Golkhoo S. Gamma irradiation as a useful tool for the isolation of astaxanthin-overproducing mutant strains of *Phaffia rhodozyma*. *Can J Microbiol* 2011 Sep;57(9):730–4.
- [165] Jin J, Wang Y, Yao M, Gu X, Li B, Liu H, et al. Astaxanthin overproduction in yeast by strain engineering and new gene target uncovering. *Biotechnol Biofuels* 2018;11:230.
- [166] Tripathi U, Venkateshwaran G, Sarada R, Gokare R. Studies on *Haematococcus pluvialis* for improved production of astaxanthin by mutagenesis. *World J Microbiol Biotechnol* 2001;17:143–8.
- [167] Wang N, Guan B, Kong Q, Sun H, Geng Z, Duan L. Enhancement of astaxanthin production from *Haematococcus pluvialis* mutants by three-stage mutagenesis breeding. *J Biotechnol* 2016 Oct 20;236:71–7.
- [168] Dragosits M, Mattanovich D. Adaptive laboratory evolution – principles and applications for biotechnology. *Microb Cell Factories* 2013;12(1):64.
- [169] Reyes LH, Gomez JM, Kao KC. Improving carotenoids production in yeast via adaptive laboratory evolution. *Metab Eng* 2014 Jan;21:26–33.
- [170] Gong J, Duan N, Zhao X. Evolutionary engineering of *Phaffia rhodozyma* for astaxanthin-overproducing strain. *Front Chem Sci Eng* 2012;6(2):174–8.
- [171] Jiang G, Yang Z, Wang Y, Yao M, Chen Y, Xiao W, et al. Enhanced astaxanthin production in yeast via combined mutagenesis and evolution. *Biochem Eng J* 2020;156:107519.
- [172] Jin J, Wang Y, Yao M, Gu X, Li B, Liu H, et al. Astaxanthin overproduction in yeast by strain engineering and new gene target uncovering. *Biotechnol Biofuels* 2018;11(1):1–15.
- [173] Chen J, Wei D, Pohnert G. Rapid estimation of astaxanthin and the carotenoid-to-chlorophyll ratio in the green microalga *chromochloris zofingiensis* using flow cytometry. *Mar Drugs* 2017 Jul 19;15(7):231.
- [174] Ukibe K, Katsuragi T, Tani Y, Takagi H. Efficient screening for astaxanthin-overproducing mutants of the yeast *Xanthophyllomyces dendrorhous* by flow cytometry. *FEMS (Fed Eur Microbiol Soc) Microbiol Lett* 2008;286(2):241–8.
- [175] Liu JH, Song L, Huang Q. Rapid screening astaxanthin-hyperproducing *Haematococcus pluvialis* mutants through near-infrared spectroscopy 2016;62:185–91.
- [176] Wang X, Ren L, Su Y, Ji Y, Liu Y, Li C, et al. Raman-activated droplet sorting (RADS) for label-free high-throughput screening of microalgal single-cells. *Anal Chem* 2017 Nov 21;89(22):12569–77.
- [177] Niizawa I, Espinaco BY, Leonardi JR, Heinrich JM, Sihufe GA. Enhancement of astaxanthin production from *Haematococcus pluvialis* under autotrophic growth conditions by a sequential stress strategy. *Prep Biochem Biotechnol* 2018;48(6):1–7.
- [178] Han D, Li Y, Qiang H. *Biology and commercial aspects of Haematococcus pluvialis: handbook of microalgal culture: applied phycology and biotechnology*. second ed. 2013.
- [179] Solovchenko Alexei E. Recent breakthroughs in the biology of astaxanthin accumulation by microalgal cell. *Photosynth Res* 2015;125(3):437–49.
- [180] Boussiba S, Bing W, Yuan JP, Zarka A, Chen F. Changes in pigments profile in the green alga *Haematococcus pluvialis* exposed to environmental stresses. *Biotechnol Lett* 1999;21(7):601–4.
- [181] Scibilia L, Girolomoni L, Berteotti S, Alboresi A, Ballottari M. Photosynthetic response to nitrogen starvation and high light in *Haematococcus pluvialis*. *Algal Res* 2015;12:170–81.
- [182] Sarada R, Tripathi U, Ravishankar GA. Influence of stress on astaxanthin production in *Haematococcus pluvialis* grown under different culture conditions. *Process Biochem* 2002;37(6):623–7.
- [183] Ad A, Spa B, Ao A. Does *Haematococcus pluvialis* need to sleep? *Algal Res* 2019;44:101722.
- [184] Cho S, Sung Y, Lee J, Yu B, Sim S. Robust cyst germination induction in *Haematococcus pluvialis* to enhance astaxanthin productivity in a semi-continuous outdoor culture system using power plant flue gas. *Bioresour Technol* 2021;338:125533.
- [185] Wan M, Zhang Z, Wang J, Huang J, Fan J, Yu A, et al. Sequential heterotrophy–dilution–photoinduction cultivation of *haematococcus pluvialis* for efficient production of astaxanthin. *Bioresour Technol* 2015;198:557–63.
- [186] Hata N, Ogonna JC, Hasegawa Y, Taroda H, Tanaka H. Production of astaxanthin by *Haematococcus pluvialis* in a sequential heterotrophic-photoautotrophic culture. *J Appl Phycol* 2001;13(5):395–402.
- [187] Río Esperanza Del, Acien F Gabriel, García-Malea M Carmen, Rivas Joaquín, Molina-Grima Emilio, Guerrero Miguel G. Efficient one-step production of astaxanthin by the microalga *Haematococcus pluvialis* in continuous culture. *Biotechnol Bioeng* 2005;91(7):808–15.
- [188] Río E, Acien F, García-Malea M, Rivas J, Molina-Grima E, Guerrero MG. Efficiency assessment of the one-step production of astaxanthin by the microalga *Haematococcus pluvialis*. *Biotechnol Bioeng* 2008;100(2):397–402.
- [189] Schulze P, Barreira LA, Pereira H, Perales JA, Varela JCS. Light emitting diodes (LEDs) applied to microalgal production. *Trends Biotechnol* 2014;32(8):422–30.
- [190] Liu T, Wang J, Hu Q, Cheng P, Bei J, Liu J, et al. Attached cultivation technology of microalgae for efficient biomass feedstock production. *Bioresour Technol* 2013;127C:216–22.
- [191] Berner F, Heilmann K, Sheehan M. Microalgal biofilms for biomass production. *J Appl Phycol* 2015;27:1793–804.
- [192] Schultze L, Simon MV, Li T, Langenbach D, Podola B, Melkonian M. High light and carbon dioxide optimize surface productivity in a Twin-Layer biofilm photobioreactor. *Algal Res* 2015;8:37–44.
- [193] Sun H, Kong Q, Geng Z, Duan L, Yang M, Guan B. Enhancement of cell biomass and activity of astaxanthin-rich *Haematococcus pluvialis*. *Bioresour Technol* 2015;186:67–73.
- [194] Sun H, Liu B, Lu X, Cheng KW, Chen F. Staged cultivation enhances biomass accumulation in the green growth phase of *Haematococcus pluvialis*. *Bioresour Technol* 2017;233(Complete):326–31.
- [195] Azizi M, Hejazi MA, Hashemi M. Supplementation with polyalcohols and sequential mixotrophy dilution photoinduction strategy boost the accumulation of astaxanthin by *Haematococcus pluvialis*. *Aquaculture* 2019;511(2):734225.
- [196] Li X, Wang X, Duan C, Yi S, Li J. Biotechnological production of astaxanthin from the microalga *Haematococcus pluvialis*. *Biotechnol Adv* 2020;43:107602.
- [197] An GH, Johnson EA. Influence of light on growth and pigmentation of the yeast *Phaffia rhodozyma*. *Antonie Leeuwenhoek* 1990 May;57(4):191–203.
- [198] de la Fuente JL, Rodríguez-Sáiz M, Schleissner C, Díez B, Peiro E, Barredo JL. High-titer production of astaxanthin by the semi-industrial fermentation of *Xanthophyllomyces dendrorhous*. *J Biotechnol* 2010;148(2–3):144–6.
- [199] Yamane Y, Higashida K, Nakashimada Y, Kakizono T, Nishio N. Influence of oxygen and glucose on primary metabolism and astaxanthin production by *phaffia rhodozyma* in batch and fed-batch cultures: kinetic and stoichiometric analysis. *Appl Environ Microbiol* 1997;63(11):4471–8.
- [200] Lu M, Lei J, Liu Y, Zhou P, Yu L. Kinetic model for optimal feeding strategy in astaxanthin production by *Xanthophyllomyces dendrorhous*. *Chin J Biotechnol* 2008;24(11):1937.
- [201] Batghare, Amit H, Singh Neha, Moholkar, et al. Investigations in ultrasound-induced enhancement of astaxanthin production by wild strain *Phaffia rhodozyma* MTCC 7536254. *Bioresource Technology Biomass Bioenergy Biowastes Conversion Technologies Biotransformations Production Technologies*; 2018. p. 166–73.
- [202] Zhang C, Zhang L, Liu J. Exogenous sodium acetate enhances astaxanthin accumulation and photoprotection in *Haematococcus pluvialis* at the non-motile stage. *J Appl Phycol* 2019;31(2):1001–8.
- [203] Kobayashi M, Kakizono T, Nagai S. Enhanced carotenoid biosynthesis by oxidative stress in acetate-induced cyst cells of a green unicellular alga, *Haematococcus pluvialis*. *Appl Environ Microbiol* 1993;59(3):867–73.
- [204] Yoshitomi T, Shimada N, Ki S, Hashizume M, Yoshimoto K. Polyethyleneimine-induced astaxanthin accumulation in the green alga *Haematococcus pluvialis* by increased oxidative stress. *J Biosci Bioeng* 2019;128(6):751–4.
- [205] Raman V, Ravi S. Effect of salicylic acid and methyl jasmonate on antioxidant systems of *Haematococcus pluvialis*. *Acta Physiol Plant* 2011;33(3):1043–9.
- [206] Shang M, Wei D, Zhao Y, Xu JW, Yu X. Enhanced astaxanthin production from *Haematococcus pluvialis* using butylated hydroxyanisole. *J Biotechnol* 2016;236:199–207.

- [207] Zhao Y, Yue C, Wei D, Tao L, Xu JW, Peng Z, et al. Butylated hydroxytoluene induces astaxanthin and lipid production in *Haematococcus pluvialis* under high-light and nitrogen-deficiency conditions. *Bioresour Technol* 2018;266:315–21.
- [208] Wang B, Pan X, Jia J, Xiong W, Manirafasha E, Ling X, et al. Strategy and regulatory mechanisms of glutamate feeding to enhance astaxanthin yield in *Xanthophyllomyces dendrorhous*. *Enzym. Microb Technol* 2019;125:45–52.
- [209] High-level production of astaxanthin by fed-batch culture of mutant strain *Phaffia rhodozyma* AJ-6-1. *J Microbiol Biotechnol* 2003;13(2):175–81.
- [210] Kim SK, Lee JH, Lee CH, Yoon YC. Increased carotenoid production in *Xanthophyllomyces dendrorhous* G276 using plant extracts. *J Microbiol* 2007;45(2):128.
- [211] Hara KY, Kageyama Y, Tanzawa N, Hirono-Hara Y, Wakabayashi K. Development of astaxanthin production from citrus peel extract using *Xanthophyllomyces dendrorhous*. *Environ Sci Pollut Control Ser* 2020:1–8.
- [212] Wang W, Yu L, Zhou P. Effects of different fungal elicitors on growth, total carotenoids and astaxanthin formation by *Xanthophyllomyces dendrorhous*. *Bioresour Technol* 2006;97(1):26–31.
- [213] Pang N, Chen S. Effects of C5 organic carbon and light on growth and cell activity of *Haematococcus pluvialis* under mixotrophic conditions. *Algal Res* 2017;21:227–35.
- [214] Hana N, Al-Shorgani N, Shukor H, AbdRahman N, Kalil M. Pre-optimization conditions for *Haematococcus pluvialis* growth. *Int J Adv Sci Eng Inf Technol* 2013;3:70–3.
- [215] Allewaert CC, Hiegle N, Strittmatter M, de Blok R, Guerra T, Gachon CMM, et al. Life history determinants of the susceptibility of the blood alga *Haematococcus* to infection by *Paraphysoderma sedebokerense* (Blastocladiomycota). *Algal Res* 2018;31:282–90.
- [216] Hwang S-W, Choi HI, Sim SJ. Acidic cultivation of *Haematococcus pluvialis* for improved astaxanthin production in the presence of a lethal fungus. *Bioresour Technol* 2019;278:138–44.
- [217] Johnson EA, Lewis MJ. Astaxanthin formation by the yeast *Phaffia rhodozyma*. *Jgenmicrobiology* 1979;115(1):173–83.
- [218] Meyer PS, Du Preez JC. Effect of culture conditions on astaxanthin production by a mutant of *Phaffia rhodozyma* in batch and chemostat culture. *Appl Microbiol Biotechnol* 1994;40(6):780–5.
- [219] Harith ZT, Lima M, Charalampopoulos D, Chatzifragkou A. Optimised production and extraction of astaxanthin from the yeast *Xanthophyllomyces dendrorhous*. *Microorganisms* 2020;8(3):430.
- [220] Kang CD, Lee JS, Park TH, Sim SJ. Complementary limiting factors of astaxanthin synthesis during photoautotrophic induction of *Haematococcus pluvialis*: C/N ratio and light intensity. *Appl Microbiol Biotechnol* 2007;74(5):987–94.
- [221] Evens TJ, Niedz RP, Kirkpatrick GJ. Temperature and irradiance impacts on the growth, pigmentation and photosystem II quantum yields of *Haematococcus pluvialis* (Chlorophyceae). *J Appl Phycol* 2008;20(4):411–22.
- [222] Giannelli L, Yamada H, Katsuda T, Yamaji H. Effects of temperature on the astaxanthin productivity and light harvesting characteristics of the green alga *Haematococcus pluvialis*. *J Biosci Bioeng* 2015;119(3):345–50.
- [223] Chai F, Wang Y, Mei X, Yao M, Chen Y, Liu H, et al. Heterologous biosynthesis and manipulation of crocetin in *Saccharomyces cerevisiae*. *Microb Cell Factories* 2017;16(1):54.
- [224] Shi F, Zhan W, Li Y, Wang X. Temperature influences β -carotene production in recombinant *Saccharomyces cerevisiae* expressing carotenogenic genes from *Phaffia rhodozyma*. *World J Microbiol Biotechnol* 2014;30(1):125–33.
- [225] Jiang G, Yang Z, Wang Y, Yao M, Yuan Y. Enhanced astaxanthin production in yeast via combined mutagenesis and evolution. *Biochem Eng J* 2020;156:107519.
- [226] Young PS, Binkley RM, Jun KW, Hee LM, Yup LS. Metabolic engineering of *Escherichia coli* for high-level astaxanthin production with high productivity. *Metab Eng* 2018;49:105–15.
- [227] Wang W, Yu L. Effects of oxygen supply on growth and carotenoids accumulation by *Xanthophyllomyces dendrorhous*. *Zeitschrift Fur Naturforschung C. J Biosci* 2009;64(11–12):853–8.
- [228] Liu YS, Wu JY. Use of n-hexadecane as an oxygen vector to improve *Phaffia rhodozyma* growth and carotenoid production in shake-flask cultures 2006;101:1033–8.
- [229] Li Y, Sommerfeld M, Chen F, Hu Q. Consumption of oxygen by astaxanthin biosynthesis: a protective mechanism against oxidative stress in *Haematococcus pluvialis* (Chlorophyceae). *J Plant Physiol* 2008;165(17):1783–97.
- [230] Christian D, Zhang J, Sawdon AJ, Peng CA. Enhanced astaxanthin accumulation in *Haematococcus pluvialis* using high carbon dioxide concentration and light illumination. *Bioresour Technol* 2018;256:548–51.
- [231] Wang J, Jiang W, Liang C, Zhu L, Li Y, Mo Q, et al. Overproduction of α -farnesene in *Saccharomyces cerevisiae* by farnesene synthase screening and metabolic engineering. *J Agric Food Chem* 2021;69(10):3103–13.
- [232] Clomburg JM, Qian S, Tan Z, Cheong S, Gonzalez R. The isoprenoid alcohol pathway, a synthetic route for isoprenoid biosynthesis 2019;116:12810–5.
- [233] Ferraz C, Leferink N, Kosov I, Scrutton N. Isopentenol utilization pathway for the production of linalool in *Escherichia coli* using an improved bacterial linalool/nerolidol synthase. *ChemBiochem* 2021;22(13):2325–34.
- [234] Luo Z, Liu N, Lazar Z, Chatzivasileiou A, Ward V, Chen J, et al. Enhancing isoprenoid synthesis in *Yarrowia lipolytica* by expressing the isopentenol utilization pathway and modulating intracellular hydrophobicity. *Metab Eng* 2020;61:344–51.
- [235] Hashimoto M, Hossain S. Fatty acids: from membrane ingredients to signaling molecules. 2018.
- [236] Nasution O, Lee YM, Kim E, Lee Y, Kim W, Choi W. Overexpression of OLE1 enhances stress tolerance and constitutively activates the MAPK HOG pathway in *Saccharomyces cerevisiae*. *Biotechnol Bioeng* 2017;114(3):620–31.
- [237] Zhang J-L, Bai Q-Y, Peng Y-Z, Fan J, Jin C-C, Cao Y-X, et al. High production of triterpenoids in *Yarrowia lipolytica* through manipulation of lipid components. *Biotechnol Biofuels* 2020;13(1):133.
- [238] Bu X, Lin J-Y, Cheng J, Yang D, Duan C-Q, Koffas M, et al. Engineering endogenous ABC transporter with improving ATP supply and membrane flexibility enhances the secretion of β -carotene in *Saccharomyces cerevisiae*. *Biotechnol Biofuels* 2020;13(1):168.
- [239] Cheng J, Li K, Yang Z, Lu H, Zhou J, Cen K. Gradient domestication of *Haematococcus pluvialis* mutant with 15% CO₂ to promote biomass growth and astaxanthin yield. *Bioresour Technol* 2016;216:340–4.
- [240] Butler T, McDougall G, Campbell R, Stanley M, Day J. Media screening for obtaining *Haematococcus pluvialis* red motile macrozooids rich in astaxanthin and fatty acids. *Biology* 2017;7(1):2.
- [241] Orosa M, Valero JF, Herrero C, Abalde J. Comparison of the accumulation of astaxanthin in *Haematococcus pluvialis* and other green microalgae under N-starvation and high light conditions. *Biotechnol Lett* 2001;23(13):1079–85.
- [242] Sun Z, Zhang Y, Sun L, Liu J. Light elicits astaxanthin biosynthesis and accumulation in the fermented ultrahigh-density *Chlorella zofingensis*. *J Agric Food Chem* 2019;67(19):5579–86.
- [243] Liu J, Sun Z, Gerken H, Liu Z, Jiang Y, Chen F. *Chlorella zofingensis* as an alternative microalgal producer of astaxanthin: biology and industrial potential. *Mar Drugs* 2014;12(6):3487–515.
- [244] Zhou P, Ye L, Xie W, Lv X, Yu H. Highly efficient biosynthesis of astaxanthin in *Saccharomyces cerevisiae* by integration and tuning of algal crtZ and bkt. *Appl Microbiol Biotechnol* 2015;99(20):8419–28.
- [245] Gao S, Tong Y, Wen Z, Zhu L, Ge M, Chen D, et al. Multiplex gene editing of the *Yarrowia lipolytica* genome using the CRISPR-Cas9 system. *J Ind Microbiol Biotechnol* 2016;43(8):1085–93.
- [246] Zhuang Y, Jiang G-L, Zhu M-J. Atmospheric and room temperature plasma mutagenesis and astaxanthin production from sugarcane bagasse hydrolysate by *Phaffia rhodozyma* mutant Y1. *Process Biochem* 2020;91:330–8.
- [247] Liu YS, Wu JY. Hydrogen peroxide-induced astaxanthin biosynthesis and catalase activity in *Xanthophyllomyces dendrorhous*. *Appl Microbiol Biotechnol* 2006;73(3):663–8.
- [248] Zhang J, Li Q, Liu J, Lu Y, Wang Y, Wang Y. Astaxanthin overproduction and proteomic analysis of *Phaffia rhodozyma* under the oxidative stress induced by TiO₂. *Bioresour Technol* 2020;311:123525.