

Heat Shock Protein 90 Inhibitors Repress Latent Membrane Protein 1 (LMP1) Expression and Proliferation of Epstein-Barr Virus-Positive Natural Killer Cell Lymphoma

Takayuki Murata^{1,9}, Seiko Iwata^{2,9}, Mohammed Nure Alam Siddiquey², Tetsuhiro Kanazawa², Fumi Goshima², Daisuke Kawashima¹, Hiroshi Kimura^{2*}, Tatsuya Tsurumi^{1*}

1 Division of Virology, Aichi Cancer Center Research Institute, Nagoya, Aichi, Japan, **2** Department of Virology, Nagoya University Graduate School of Medicine, Nagoya, Aichi, Japan

Abstract

Epstein-Barr virus (EBV) LMP1 is a major oncoprotein expressed in latent infection. It functions as a TNFR family member and constitutively activates cellular signals, such as NF κ B, MAPK, JAK/STAT and AKT. We here screened small molecule inhibitors and isolated HSP90 inhibitors, Radicicol and 17-AAG, as candidates that suppress LMP1 expression and cell proliferation not only in EBV-positive SNK6 Natural Killer (NK) cell lymphoma cells, but also in B and T cells. Tumor formation in immunodeficient NOD/Shi-scid/IL-2R γ ^{null} (NOG) mice was also retarded. These results suggest that HSP90 inhibitors can be alternative treatments for patients with EBV-positive malignancies.

Citation: Murata T, Iwata S, Siddiquey MNA, Kanazawa T, Goshima F, et al. (2013) Heat Shock Protein 90 Inhibitors Repress Latent Membrane Protein 1 (LMP1) Expression and Proliferation of Epstein-Barr Virus-Positive Natural Killer Cell Lymphoma. PLoS ONE 8(5): e63566. doi:10.1371/journal.pone.0063566

Editor: Joseph S. Pagano, The University of North Carolina at Chapel Hill, United States of America

Received: November 7, 2012; **Accepted:** April 3, 2013; **Published:** May 3, 2013

Copyright: © 2013 Murata et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: This work was supported by grants-in-aid for Scientific Research from the Ministry of Education, Science, Sports, Culture and Technology (no. 20390137, 21022055 to T. T.), the Ministry of Health, Labour, and Welfare (to T. T. & H. K.) and partly by the Uehara Memorial Research Fund (to T. T. & T. M.), the Takeda Science Foundation (to T. T. & T. M.), the Yasuda Medical Foundation, the Mochida Memorial Foundation for Medical and Pharmaceutical Research, the Senshin Medical Research Foundation, and the Kanae Foundation for Promotion of Medical Science (to T. M.). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

* E-mail: ttsurumi@aichi-cc.jp (TT); hkimura@med.nagoya-u.ac.jp (HK)

⁹ These authors contributed equally to this work.

Introduction

The Epstein-Barr virus (EBV) is a human gamma-herpesvirus that mainly infects and establishes latent infection in B lymphocytes, but can also infect other types of cells, including NK, T and epithelial cells. EBV infection has been implicated as a causal factor in a variety of malignancies, and the expression pattern of viral latent genes varies depending on the tissue of origin and the state of the tumors. Neoplasms such as Burkitt's lymphomas or gastric carcinomas express only EBV and EBNA1 (type I latency), whereas some Hodgkin lymphomas, nasopharyngeal carcinomas (NPC) and NK/T lymphomas express EBV, EBNA1, LMP1 and LMP2 genes (type II latency). In addition to the type II genes, EBNA2, EBNA3 and EBNA-LP are also expressed in immunosuppression-related lymphomas or lymphoblastoid cell lines (LCLs) (type III latency).

EBV is associated with various types of T or NK cell lymphoproliferative diseases (T/NK LPDs). A severe form of chronic active EBV disease (CAEBV), mainly found in East Asia including Japan, is caused by clonal expansion of EBV-infected T or NK cells [1–3]. Others include extranodal NK/T lymphoma, nasal type (ENKL), and aggressive NK cell leukemia (ANKL). Although such EBV-positive T/NK LPDs are relatively rare, therapeutic treatment for those disorders is challenging, and the

prognosis of those patients often can be dismal [4,5]. Therefore, development of effective and specific drugs is an important goal.

The EBV latent infection integral membrane protein 1 (LMP1) is frequently expressed in latent EBV infections, including NK/T lymphomas. Since it functions as a constitutive TNFR family member by aggregation in the plasma membrane, resulting in constitutive activation of cellular signaling through NF κ B, MAPK, JAK/STAT and AKT, LMP1 is assumed to be a major oncogene encoded by EBV [6–15].

Heat-shock protein 90 (HSP90) is an ATP-dependent molecular chaperone that is important for stability, quality control, protein interaction and functional maturation of cellular or viral client proteins. Because HSP90 is occasionally overexpressed and present in an activated form in cancer cells, and thereby supports proliferation of activated oncoproteins, including many cancer-associated kinases and transcription factors, it is regarded as an essential factor for oncogenic transformation [16,17]. Radicicol and 17-AAG are HSP90 inhibitors which interact directly within its ATP-binding pocket, preventing ATP binding and interaction with client proteins [18]. These inhibitors might thus have potential as anti-cancer drugs for malignancies that depend on particular driver oncogene products that are sensitive HSP90 clients [16,17,19]. For example, HSP90 inhibitors have shown




Figure 1. Representative example of screening for small molecule inhibitors that repress LMP1 expression. The EBV-positive NK lymphoma cell line, SNK6, was seeded and small molecule inhibitors were added to the media at concentrations of 10 and 3 μ M. After 72 h, cell RNA was collected and subjected to Real-Time RT-PCR using specific primers for LMP1 and GAPDH mRNAs. Relative LMP1 mRNA levels are shown after normalization to GAPDH mRNA levels. doi:10.1371/journal.pone.0063566.g001

promise as anti-myeloma agents in pre-clinical settings and are currently being evaluated in clinical trials [20].

In the present study, we screened small molecule inhibitors and isolated HSP90 inhibitors as candidates that suppress LMP1 expression and cell proliferation in EBV-positive SNK6 NK cell lymphoma cells. The inhibitors not only retarded tumor proliferation at the culture level but also tumor formation in immunodeficient NOD/Shi-scid/IL-2R γ ^{null} (NOG) mice. HSP90 inhibitors therefore may offer alternative treatments for EBV-positive malignancies.

Materials and Methods

Cell Culture and Reagents

An EBV-positive NK cell lymphoma line, SNK6, and an EBV-positive T cell line, SNT13, were maintained in RPMI1640 medium supplemented with 10% human serum (MP Biomedicals), 2 mM of Glutamax (GIBCO), 0.88 mM Oxalicacetic acid (SIGMA), 1 mM Sodium Pyruvate (GIBCO) and 700 U/ml of IL-2 (Primmune Inc.). SNK6 was originally established from a patient with ENKL and characterized by Nagata and others [21]. SNT13 is a $\gamma\delta$ T-cell clone established from a patient with CAEBV [22]. These cell lines of low passage numbers were kindly provided by N. Shimizu in December 2009. B95-8 cells [23] were maintained in RPMI1640 medium supplemented with 10% fetal bovine serum. The Screening Committee of Anticancer Drugs (SCADS), Japan, kindly provided a library of small molecule inhibitors.

Real-time PCR

For real-time RT-PCR, total cell RNA was purified using TriPure Isolation Reagent (Roche) and subjected to Real-Time RTPCR using a One Step SYBR PrimeScript RT-PCR Kit II (TaKaRa) with the Real-Time PCR System 7300 according to the manufacturer's instructions. PCR was performed as described earlier [24]. Primers used for the RT-PCR were as follows [25]: for GAPDH mRNA, 5'-TGCACCACCAACTGCTAGC-3' and 5'-GGCATGGACTGTGGTCATGAG-3'; for LMP1 mRNA,

5'-CTATTCCTTTGCTCTCATGC-3' and 5'-TGAGCAGGAGGTGATCATC-3'; and for EBNA1 mRNA, 5'-AGGTA-CAGGACCTGGAAATG-3' and 5'-CCTCGTCCATGGT-TATCACC-3'. Real-Time PCR with GAPDH primers was also performed to serve as an internal control for input RNA.

Antibodies and Immunoblotting

Anti-tubulin antibody was from Cell Signaling, and anti-LMP1 monoclonal antibody was reported previously [25]. For immunoblotting, cell proteins lysed in sample buffer were subjected to SDS-PAGE, followed by immunoblotting with the indicated antibodies as described previously [25].

Transplantation of SNK6 into NOD/Shi-scid/IL-2R γ ^{null} (NOG) mice

Female NOG mice were purchased from the Central Institute for Experimental Animals, Japan. Twenty four NOG mice at the age of 10 weeks were subcutaneously implanted with 5×10^6 SNK6 cells per mouse on day 0. From day 14, DMSO (vehicle) was injected into 12 mice, and the rest were treated with 17-AAG (6 times in two weeks, 50 mg/kg in total) intra-peritoneally. Peripheral blood was collected from the tail veins once in two weeks, and levels of EBV genomic DNA in whole blood were measured by Real-Time PCR as described previously [26]. Subcutaneous tumor masses were also measured with an external caliper and tumor volume was calculated using the formula: $\pi \times \text{short axis} \times \text{long axis} \times \text{height}/6$. Animal experiments were approved by the University Committee in accordance with the Guidelines for Animal Experimentation at Nagoya University.

Statistical analysis

Data shown are means \pm standard errors and were analyzed using SPSS for Windows version 18.0 (IBM Corporation, Chicago, IL, USA). The therapeutic results were analyzed using the Mann-Whitney *U* test between groups.




Figure 2. HSP90 inhibitors decrease LMP1 expression. (A, B) SNK6 cells were treated with Radicolol (R) or 17-AAG (AAG) at the concentrations of 3, 1, 0.3 or 0.1 μ M. After 72 h, cell RNA was collected and subjected to Real-Time RT-PCR. Relative LMP1 levels (A) and EBNA1 levels (B) are shown after normalization to GAPDH mRNA levels. (C) SNK6 cells were treated with Radicolol (R) or 17-AAG (AAG) at the concentrations of 3, 1, 0.3 or 0.1 μ M. After 72 h, cell proteins were collected and subjected to immunoblotting using anti-LMP1 and -Tubulin antibodies. (D, E) As in Fig. 2A, B95-8 (D) or SNT13 (E) cells were treated and subjected to Real-Time RT-PCR. Relative levels of LMP1 mRNA are shown after normalization to GAPDH mRNA levels. Each bar represents the mean and SD of three independent transfections. doi:10.1371/journal.pone.0063566.g002

Results

Identification of HSP90 inhibitors on screening for small molecule inhibitors that repress LMP1 transcription

CAEBV and ENKL committed to type II latency express the major EBV oncogene, LMP1. We therefore conducted a search for small molecular inhibitors that repress expression of LMP1. As an initial screen, the EBV-positive NK cell lymphoma, SNK6, was treated with chemicals or the vehicle DMSO at concentrations of 3 or 10 μ M for 3 days. Cellular RNAs were harvested and subjected to Real-Time RT-PCR. Among 300 small molecule

substances with identified targets, we found Radicolol and 17-AAG, inhibitors of HSP90, to decrease LMP1 transcripts (Fig. 1). In order to further evaluate the effect, SNK6 cells were administered 3, 1, 0.3, 0.1 μ M of Radicolol or 17-AAG (Fig. 2A–C). LMP1 levels decreased to 41% and 48% of control (DMSO) with 3 μ M of Radicolol and 17-AAG, respectively (Fig. 2A), with EBNA1 levels appearing relatively unaffected (Fig. 2B). We then examined protein levels of LMP1 in Fig. 2C. After 72 h with 3 or 1 μ M of Radicolol or 17-AAG, LMP1 protein levels were decreased in SNK6 cells (Fig. 2C), in a similar fashion with the transcript levels. LMP1 expression in B95-8 or EBV-positive T cell




Figure 3. HSP90 inhibitors suppress cell proliferation of the EBV-positive SNK6 NK lymphoma line. SNK6 cells were cultured with Radicicol (A) or 17-AAG (B) at concentrations of 3, 1, 0.3 or 0.1 μ M and cell numbers were counted on days 0, 2 and 4. Data are shown as the means of three independent replicates. doi:10.1371/journal.pone.0063566.g003

line (SNT13) was also reduced by these HSP90 inhibitors (Fig. 2D, E). For unknown reasons, 17-AAG markedly suppressed LMP1 expression from SNT13 cells even at lower concentrations (Fig. 2E).

Suppression of cell proliferation *in vitro* and *in vivo*

After confirming that HSP90 inhibitors repress LMP1 expression in EBV-positive lymphoma cells, we then examined whether the inhibitors might actually suppress cancerous growth. We first tested at the cell culture level (Fig. 3). Three or 1 μ M of Radicicol (Fig. 3A) or 17-AAG (Fig. 3B) could fully block proliferation of SNK6 cells depending on the concentration. Partial but appreciable inhibition was also exhibited at lower concentrations.

It has been already reported that HSP90 inhibitors block proliferation of EBV-positive malignancies, including NK/T lymphomas [27,28], although the earlier studies did not test LMP1 levels. For examination of effects *in vivo*, we here adopted a novel mouse xenograft model using severely immuno-deficient mice of the NOG strain [29], allowing assessment of whether the HSP90 inhibitors might repress EBV-positive malignant cells without seriously affecting other normal part of tissues and organs. The NOG mice were injected subcutaneously with 5×10^6 of SNK6 cells at day 0, and low doses of 17-AAG (50 mg/kg in total) or DMSO were administered into the abdominal cavity for 6 times between days 14 and 25. EBV genomic DNA titers in whole blood (Fig. 4A) and tumor sizes (Fig. 4B) were measured. As shown in Fig. 4, 17-AAG markedly reduced the viral titer in blood and growth of the NK lymphoma cells in NOG mice. Because engraftment of NK lymphomas in mice is not very efficient [29], only 50% of the mice developed subcutaneous tumor masses and others failed to nurture the lymphoma cells even in the DMSO




Figure 4. Effects of HSP90 inhibitors *in vivo*. HSP90 inhibitors repress EBV copy numbers in the whole blood (A) and tumor size (B) in mice implanted with EBV-positive NK cell lymphoma. Twenty-four NOG mice were subcutaneously implanted with 5×10^6 of SNK6 cells per mouse on day 0. From day 14, DMSO (vehicle) or 17-AAG (6 times in two weeks, 50 mg/kg in total) was injected to twelve mice each, intraperitoneally. Peripheral blood was collected from the tail veins once in two weeks, and levels of EBV genomic DNA in whole blood were measured by Real-Time PCR (A). Subcutaneous tumor masses (B) and LMP1 mRNA levels in the tumors (C) were also measured. * $p < 0.05$ (Mann-Whitney *U* test). doi:10.1371/journal.pone.0063566.g004

treatment group. Due to this low efficiency of transplantation, statistical significance could be exhibited only once for EBV DNA load (Fig. 4A, 14 weeks), but we gained the clear impression that 17-AAG worked more efficiently. This is an important difference when compared to LCLs, with which 100% of injected mice develop tumors, and thereby the efficacy of 17-AAG was exhibited more significantly [28]. We then compared the LMP1 expression




Figure 5. Suppression of LMP1 expression in EBV-positive NK cells by HSP90 inhibitors is likely dependent on IL-2. SNK6 cells, routinely cultured with IL-2, were washed extensively with PBS, and then cultured in the presence (black bars) or absence (gray bars) of IL-2 with Radicol (R) or 17-AAG (A) at concentrations of 3, 1, 0.3 or 0.1 μ M. After 72 h, cell RNAs were collected and subjected to Real-Time RT-PCR. Relative levels of LMP1 mRNA are shown after normalization to GAPDH mRNA levels. Each bar represents the mean and SD of three independent transfections.
doi:10.1371/journal.pone.0063566.g005

levels in the tumor developed in the absence or presence of 17-AAG (Fig. 4C). LMP1 transcript was lower in the tumor treated with 17-AAG, indicating that the inhibitor reduced LMP1 transcription *in vivo*, too. To summarize, our results imply high potential of HSP90 inhibitors for treating EBV-positive cancers, including EBV-positive NK lymphomas.

Discussion

ENKL is an aggressive type of cancer often associated with resistance to chemotherapy, and accordingly a poor prognosis. Therefore, treatments are needed that specifically target its molecular determinants. We here focused on expression of the major oncogene of EBV, LMP1, and found two HSP90 inhibitors, Radicol and 17-AAG, to both decrease LMP1 expression and cancerous growth of the EBV-positive SNK6 NK cell lymphoma line, *in vitro* and *in vivo*. Although the reduction in LMP1 levels by the inhibitors correlated with cell growth inhibition, we still cannot tell if low levels of LMP1 is responsible for the growth inhibition.

Jeon et al. previously reported that HSP90 inhibitors induce apoptotic cell death in EBV-positive NK/T lymphoma cells *in vitro* [27], but did not examine effects *in vivo*. Elsewhere, Sun et al. reported that HSP90 inhibitors block proliferation *in vitro* and *in vivo* mostly using EBV-positive LCLs [28], but NK/T lymphomas were not tested. Importantly, unlike ours, both of the previous papers did not include data on LMP1 levels.

LMP1 gene transcription differs between type II and type III latency infection. In the latter, LMP1 transcription is turned on by EBNA2 [30–32], whereas LMP1 expression is independent of EBNA2 in type II. In the previous work of Jeon and others induction of apoptosis by HSP90 inhibitors was considered to be through AKT signaling inhibition [27]. The decreased expression of LMP1 gene we observed here (Fig. 1, 2) might have been brought about through suppression of the AKT pathway. In latency II, including the ENKL case, it has been frequently reported that cytokines, such as IL-4, IL-6, IL-10, IL-13 and IL-

21, activate the JAK/STAT pathway, thereby inducing LMP1 gene expression through STAT [33–38]. Because JAK/STAT signaling can also be blocked by HSP90 inhibitors [39], it may also be involved. Actually, since our EBV-positive NK/T cells are cultured routinely with IL-2 and it is involved in LMP1 expression [40], the cytokine and its downstream signaling may also be implicated in the reduction by HSP90 inhibitors. To examine this, we cultured SNK6 cells with or without IL-2 and HSP90 inhibitors, and measured the levels of LMP1 mRNA (Fig. 5). As expected from the previous report [40], depletion of IL-2 down-regulated LMP1 levels to 5.5% of control, in DMSO-treated cells. Treatment with Radicol (R) or 17-AAG (A) at 1 μ M or higher caused reduced reduction of LMP1 levels in the presence of IL-2 (black bars), but it did not significantly decrease the levels without IL-2 (gray bars). This result suggests that HSP90 inhibitors suppress LMP1 expression, which is activated by IL-2, and that the cell signalings elicited by IL-2, such as JAK/STAT, are likely be responsible for the LMP1 reduction by HSP90 inhibitors. Besides JAK/STAT pathways, NF κ B signaling must also be notified, because it is known that IL-2 elicits NF κ B signaling [41], and HSP90 inhibitors can repress NF κ B [42,43]. In addition, C/EBP contributes to LMP1 expression in type II [25], and may also be regulated by HSP90 inhibitors.

In type III latency, although EBNA2 does not feature DNA binding activity, it enhances LMP1 promoter activity by acting as a cofactor. It associates with cellular transcriptional factors, including the Recombination signal Binding Protein J κ (RBP-J κ) [32,44] and PU-box 1 (PU.1) (also known as Spleen Forming Virus (SFV) Proviral Integration 1 (SPI1)) [30,31,45,46], which are then recruited onto the LMP1 promoter for transactivation. Because we found LMP1 expression was decreased by HSP90 inhibitors even in type III B95-8 cells (Fig. 2B), RBP-J κ or PU.1 may also be under the control of HSP90. In fact, PU.1 is known to be inhibited by HSP90 inhibitors [47,48].

In summary, we here observed reduced expression of LMP1 and simultaneous growth suppressive effects of Radicol and 17-AAG in EBV-positive lymphomas, especially NK lymphoma. Although the molecular mechanisms of how the HSP90 inhibitors block cell proliferation are elusive because HSP90 has a number of client proteins, our observation suggests that they could be potent therapeutic drugs for EBV-positive lymphomas. Rapid progress in the field of Hsp90 biology has brought about development of more potent and less toxic inhibitors. It is to be hoped that these may become useful as antiviral drugs against EBV-associated disorders.

Acknowledgments

We thank Dr. N. Shimizu for SNK6 and SNT13 cells. Panels of small molecule inhibitors were kindly provided by the Screening Committee of Anticancer Drugs (SCADS), which is supported by a Grant-in-Aid for Scientific Research on Innovative Areas, Scientific Support Programs for Cancer Research, from the Ministry of Education, Culture, Sports, Science and Technology of Japan. We would further like to express our appreciation to T. Kanda & T. Gamano for technical assistance.

Author Contributions

Conceived and designed the experiments: TM SI HK TT. Performed the experiments: TM SI MNAS TK FG DK. Analyzed the data: TM SI HK TT. Contributed reagents/materials/analysis tools: TM SI DK HK TT. Wrote the paper: TM HK TT.

References

- Kimura H, Hoshino Y, Kanegane H, Tsuge I, Okamura T, et al. (2001) Clinical and virologic characteristics of chronic active Epstein-Barr virus infection. *Blood* 98: 280–286.
- Kawa K, Okamura T, Yagi K, Takeuchi M, Nakayama M, et al. (2001) Mosquito allergy and Epstein-Barr virus-associated T/natural killer-cell lymphoproliferative disease. *Blood* 98: 3173–3174.
- Kimura H (2006) Pathogenesis of chronic active Epstein-Barr virus infection: is this an infectious disease, lymphoproliferative disorder, or immunodeficiency? *Rev Med Virol* 16: 251–261.
- Fox CP, Shannon-Lowe C, Rowe M (2011) Deciphering the role of Epstein-Barr virus in the pathogenesis of T and NK cell lymphoproliferations. *Herpesviridae* 2: 8.
- Kwong YL (2005) Natural killer-cell malignancies: diagnosis and treatment. *Leukemia* 19: 2186–2194.
- Soni V, Cahir-McFarland E, Kieff E (2007) LMP1 TRAFicking activates growth and survival pathways. *Adv Exp Med Biol* 597: 173–187.
- Lam N, Sugden B (2003) CD40 and its viral mimic, LMP1: similar means to different ends. *Cell Signal* 15: 9–16.
- Shair KH, Bendt KM, Edwards RH, Bedford EC, Nielsen JN, et al. (2007) EBV latent membrane protein 1 activates Akt, NFkappaB, and Stat3 in B cell lymphomas. *PLoS Pathog* 3: e166.
- Kulwichit W, Edwards RH, Davenport EM, Baskar JF, Godfrey V, et al. (1998) Expression of the Epstein-Barr virus latent membrane protein 1 induces B cell lymphoma in transgenic mice. *Proc Natl Acad Sci U S A* 95: 11963–11968.
- Lam N, Sugden B (2003) LMP1, a viral relative of the TNF receptor family, signals principally from intracellular compartments. *EMBO J* 22: 3027–3038.
- Dirmeier U, Hoffmann R, Kilger E, Schultheiss U, Brisenko C, et al. (2005) Latent membrane protein 1 of Epstein-Barr virus coordinately regulates proliferation with control of apoptosis. *Oncogene* 24: 1711–1717.
- Kilger E, Kieser A, Baumann M, Hammerschmidt W (1998) Epstein-Barr virus-mediated B-cell proliferation is dependent upon latent membrane protein 1, which simulates an activated CD40 receptor. *EMBO J* 17: 1700–1709.
- Kieser A, Kaiser C, Hammerschmidt W (1999) LMP1 signal transduction differs substantially from TNF receptor 1 signaling in the molecular functions of TRADD and TRAF2. *EMBO J* 18: 2511–2521.
- Paine E, Scheinman RI, Baldwin AS Jr, Raab-Traub N (1995) Expression of LMP1 in epithelial cells leads to the activation of a select subset of NF-kappa B/Rel family proteins. *J Virol* 69: 4572–4576.
- Uchida J, Yasui T, Takaoka-Shichijo Y, Muraoka M, Kulwichit W, et al. (1999) Mimicry of CD40 signals by Epstein-Barr virus LMP1 in B lymphocyte responses. *Science* 286: 300–303.
- Workman P, Burrows F, Neckers L, Rosen N (2007) Drugging the cancer chaperone HSP90: combinatorial therapeutic exploitation of oncogene addiction and tumor stress. *Ann N Y Acad Sci* 1113: 202–216.
- Neckers L, Workman P (2012) Hsp90 molecular chaperone inhibitors: are we there yet? *Clin Cancer Res* 18: 64–76.
- Whitesell L, Lindquist SL (2005) HSP90 and the chaperoning of cancer. *Nat Rev Cancer* 5: 761–772.
- Jhaveri K, Taldone T, Modi S, Chiosis G (2012) Advances in the clinical development of heat shock protein 90 (Hsp90) inhibitors in cancers. *Biochim Biophys Acta* 1823: 742–755.
- Usmani SZ, Chiosis G (2011) HSP90 inhibitors as therapy for multiple myeloma. *Clin Lymphoma Myeloma Leuk* 11 Suppl 1: S77–81.
- Nagata H, Konno A, Kimura N, Zhang Y, Kimura M, et al. (2001) Characterization of novel natural killer (NK)-cell and gammadelta T-cell lines established from primary lesions of nasal T/NK-cell lymphomas associated with the Epstein-Barr virus. *Blood* 97: 708–713.
- Zhang Y, Nagata H, Ikeuchi T, Mukai H, Oyoshi MK, et al. (2003) Common cytological and cytogenetic features of Epstein-Barr virus (EBV)-positive natural killer (NK) cells and cell lines derived from patients with nasal T/NK-cell lymphomas, chronic active EBV infection and hydroa vacciniforme-like eruptions. *Br J Haematol* 121: 805–814.
- Miller G, Lipman M (1973) Release of infectious Epstein-Barr virus by transformed marmoset leukocytes. *Proc Natl Acad Sci U S A* 70: 190–194.
- Murata T, Kondo Y, Sugimoto A, Kawashima D, Saito S, et al. (2012) Epigenetic Histone Modification of Epstein-Barr Virus BZLF1 Promoter during Latency and Reactivation in Raji Cells. *J Virol* 86: 4752–4761.
- Noda C, Murata T, Kanda T, Yoshiyama H, Sugimoto A, et al. (2011) Identification and characterization of CCAAT enhancer-binding protein (C/EBP) as a transcriptional activator for Epstein-Barr virus oncogene latent membrane protein 1. *J Biol Chem* 286: 42524–42533.
- Kimura M (2008) IRF2-binding protein-1 is a JDP2 ubiquitin ligase and an inhibitor of ATF2-dependent transcription. *FEBS Lett* 582: 2833–2837.
- Jeon YK, Park CH, Kim KY, Li YC, Kim J, et al. (2007) The heat-shock protein 90 inhibitor, geldanamycin, induces apoptotic cell death in Epstein-Barr virus-positive NK/T-cell lymphoma by Akt down-regulation. *J Pathol* 213: 170–179.
- Sun X, Barlow EA, Ma S, Hagemeyer SR, Duellman SJ, et al. (2010) Hsp90 inhibitors block outgrowth of EBV-infected malignant cells in vitro and in vivo through an EBNA1-dependent mechanism. *Proc Natl Acad Sci U S A* 107: 3146–3151.
- Imadome K, Yajima M, Arai A, Nakazawa A, Kawano F, et al. (2011) Novel mouse xenograft models reveal a critical role of CD4+ T cells in the proliferation of EBV-infected T and NK cells. *PLoS Pathog* 7: e1002326.
- Johannsen E, Koh E, Mosialos G, Tong X, Kieff E, et al. (1995) Epstein-Barr virus nuclear protein 2 transactivation of the latent membrane protein 1 promoter is mediated by J kappa and PU.1. *J Virol* 69: 253–262.
- Laux G, Adam B, Strobl LJ, Moreau-Gachelin F (1994) The Spi-1/PU.1 and Spi-B ets family transcription factors and the recombination signal binding protein RBP-J kappa interact with an Epstein-Barr virus nuclear antigen 2 responsive cis-element. *EMBO J* 13: 5624–5632.
- Grossman SR, Johannsen E, Tong X, Yalamanchili R, Kieff E (1994) The Epstein-Barr virus nuclear antigen 2 transactivator is directed to response elements by the J kappa recombination signal binding protein. *Proc Natl Acad Sci U S A* 91: 7568–7572.
- Kis LL, Gerasimcik N, Salamon D, Persson EK, Nagy N, et al. (2011) STAT6 signaling pathway activated by the cytokines IL-4 and IL-13 induces expression of the Epstein-Barr virus-encoded protein LMP-1 in absence of EBNA-2: implications for the type II EBV latent gene expression in Hodgkin lymphoma. *Blood* 117: 165–174.
- Kis LL, Salamon D, Persson EK, Nagy N, Scheeren FA, et al. (2010) IL-21 imposes a type II EBV gene expression on type III and type I B cells by the repression of C- and activation of LMP-1-promoter. *Proc Natl Acad Sci U S A* 107: 872–877.
- Kis LL, Takahara M, Nagy N, Klein G, Klein E (2006) IL-10 can induce the expression of EBV-encoded latent membrane protein-1 (LMP-1) in the absence of EBNA-2 in B lymphocytes and in Burkitt lymphoma- and NK lymphoma-derived cell lines. *Blood* 107: 2928–2935.
- Konforte D, Simard N, Paige CJ (2008) Interleukin-21 regulates expression of key Epstein-Barr virus oncoproteins, EBNA2 and LMP1, in infected human B cells. *Virology* 374: 100–113.
- Chen H, Lee JM, Zong Y, Borowitz M, Ng MH, et al. (2001) Linkage between STAT regulation and Epstein-Barr virus gene expression in tumors. *J Virol* 75: 2929–2937.
- Chen H, Hutt-Fletcher L, Cao L, Hayward SD (2003) A positive autoregulatory loop of LMP1 expression and STAT activation in epithelial cells latently infected with Epstein-Barr virus. *J Virol* 77: 4139–4148.
- Schoof N, von Bonin F, Trumper L, Kube D (2009) HSP90 is essential for Jak-STAT signaling in classical Hodgkin lymphoma cells. *Cell Commun Signal* 7: 17.
- Takahara M, Kis LL, Nagy N, Liu A, Harabuchi Y, et al. (2006) Concomitant increase of LMP1 and CD25 (IL-2-receptor alpha) expression induced by IL-10 in the EBV-positive NK lines SNK6 and KAI3. *Int J Cancer* 119: 2775–2783.
- Zhou J, Zhang J, Lichtenheld MG, Meadows GG (2002) A role for NF-kappa B activation in perforin expression of NK cells upon IL-2 receptor signaling. *J Immunol* 169: 1319–1325.
- Rakićina TV, Vasilevska IA, O'Dwyer PJ (2003) Additive interaction of oxaliplatin and 17-allylamino-17-demethoxygeldanamycin in colon cancer cell lines results from inhibition of nuclear factor kappaB signaling. *Cancer Res* 63: 8600–8605.
- Wang X, Ju W, Renouard J, Aden J, Belinsky SA, et al. (2006) 17-allylamino-17-demethoxygeldanamycin synergistically potentiates tumor necrosis factor-induced lung cancer cell death by blocking the nuclear factor-kappaB pathway. *Cancer Res* 66: 1089–1095.
- Waltzer L, Logeat F, Brou C, Israel A, Sergeant A, et al. (1994) The human J kappa recombination signal sequence binding protein (RBP-J kappa) targets the Epstein-Barr virus EBNA2 protein to its DNA responsive elements. *EMBO J* 13: 5633–5638.
- Sjoblom A, Jansson A, Yang W, Lain S, Nilsson T, et al. (1995) PU box-binding transcription factors and a POU domain protein cooperate in the Epstein-Barr virus (EBV) nuclear antigen 2-induced transactivation of the EBV latent membrane protein 1 promoter. *J Gen Virol* 76 (Pt 11): 2679–2692.
- Sjoblom A, Nerstedt A, Jansson A, Rymo L (1995) Domains of the Epstein-Barr virus nuclear antigen 2 (EBNA2) involved in the transactivation of the latent membrane protein 1 and the EBNA Cp promoters. *J Gen Virol* 76 (Pt 11): 2669–2678.
- Okawa Y, Hideshima T, Steed P, Vallet S, Hall S, et al. (2009) SNX-2112, a selective Hsp90 inhibitor, potently inhibits tumor cell growth, angiogenesis, and osteoclastogenesis in multiple myeloma and other hematologic tumors by abrogating signaling via Akt and ERK. *Blood* 113: 846–855.
- Moreau F, Buck I, Dicato M, Diederich M (2008) Radicol-mediated inhibition of Bcr-Abl in K562 cells induced p38-MAPK dependent erythroid differentiation and PU.1 down-regulation. *Biofactors* 34: 313–329.