

POSTER PRESENTATION

Open Access

Modelling dissociation constants of organic acids by local molecular parameters

Haiying Yu*, R Kühne, R-U Ebert, G Schüürmann

From 5th German Conference on Cheminformatics: 23. CIC-Workshop
Goslar, Germany. 8-10 November 2009

The dissociation constant pK_a defines the ionization degree of organic compounds. It is an important parameter that affects their toxicity and environmental behavior and fate. Among the present approaches to predict pK_a values of organic chemicals, increment methods show a good performance but are limited by missing values of special groups. The purpose of this study is to develop models for predicting the pK_a of organic acids directly from their molecular structures.

A quantum chemical method was introduced by employing local molecular parameters. Experimental pK_a values of more than 1000 organic acids, including phenols, aromatic carboxylic acids, aliphatic carboxylic acids and alcohols were selected, and all molecules were optimized using the semi-empirical AM1 Hamiltonian. Simple models with several descriptors for different subsets were calibrated by multilinear regression (MLR). Substituent positions on aromatic rings were also taken into account.

The obtained models yield good predictive squared correlation coefficients q^2 and superior performance compared with other quantum chemical models. The prediction capability is further evaluated using cross validation. The models unravel that the potential of oxygen atom conjoint to ionizable hydrogen atom to accept extra electronic charge is the single most important contribution to the dissociation of hydrogen atom.

Non-linear statistical analysis methods were also applied because of the non-linear character of the employed descriptors.

The study was supported by the China Scholarship Council and by the EU project OSIRIS (IP, contact No. 037017), which is gratefully acknowledged.

Published: 4 May 2010

doi:10.1186/1758-2946-2-S1-P10

Cite this article as: Yu *et al.*: Modelling dissociation constants of organic acids by local molecular parameters. *Journal of Cheminformatics* 2010 2(Suppl 1):P10.

Publish with **ChemistryCentral** and every scientist can read your work free of charge

“Open access provides opportunities to our colleagues in other parts of the globe, by allowing anyone to view the content free of charge.”

W. Jeffery Hurst, The Hershey Company.

- available free of charge to the entire scientific community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
<http://www.chemistrycentral.com/manuscript/>



ChemistryCentral

Department of Ecological Chemistry, Helmholtz Centre for Environmental Research – UFZ, Permoserstr. 15, 04318, Leipzig, Germany

