

RESEARCH

Open Access

Chabertia erschowi (Nematoda) is a distinct species based on nuclear ribosomal DNA sequences and mitochondrial DNA sequences

Guo-Hua Liu^{1,3†}, Lei Zhao^{1,2†}, Hui-Qun Song¹, Guang-Hui Zhao⁴, Jin-Zhong Cai⁵, Quan Zhao² and Xing-Quan Zhu^{1,2,3*}

Abstract

Background: Gastrointestinal nematodes of livestock have major socio-economic importance worldwide. In small ruminants, *Chabertia* spp. are responsible for economic losses to the livestock industries globally. Although much attention has given us insights into epidemiology, diagnosis, treatment and control of this parasite, over the years, only one species (*C. ovina*) has been accepted to infect small ruminants, and it is not clear whether *C. erschowi* is valid as a separate species.

Methods: The first and second internal transcribed spacers (ITS-1 and ITS-2) regions of nuclear ribosomal DNA (rDNA) and the complete mitochondrial (mt) genomes of *C. ovina* and *C. erschowi* were amplified and then sequenced. Phylogenetic re-construction of 15 Strongylida species (including *C. erschowi*) was carried out using Bayesian inference (BI) based on concatenated amino acid sequence datasets.

Results: The ITS rDNA sequences of *C. ovina* China isolates and *C. erschowi* samples were 852–854 bp and 862–866 bp in length, respectively. The mt genome sequence of *C. erschowi* was 13,705 bp in length, which is 12 bp shorter than that of *C. ovina* China isolate. The sequence difference between the entire mt genome of *C. ovina* China isolate and that of *C. erschowi* was 15.33%. In addition, sequence comparison of the most conserved mt small subunit ribosomal (*rms*) and the least conserved *nad2* genes among multiple individual nematodes revealed substantial nucleotide differences between these two species but limited sequence variation within each species.

Conclusions: The mtDNA and rDNA datasets provide robust genetic evidence that *C. erschowi* is a valid strongylid nematode species. The mtDNA and rDNA datasets presented in the present study provide useful novel markers for further studies of the taxonomy and systematics of the *Chabertia* species from different hosts and geographical regions.

Keywords: *Chabertia* spp, Nuclear ribosomal DNA, Internal transcribed spacer (ITS), Mitochondrial DNA, Phylogenetic analysis

Background

The phylum Nematoda includes many parasites that threaten the health of plants, animals and humans on a global scale. The soil-transmitted helminthes (including

roundworms, whipworms and hookworms) are estimated to infect almost one sixth of all humans, and more than a billion people are infected with at least one species [1]. *Chabertia* spp. are common gastrointestinal nematodes, causing significant economic losses to the livestock industries worldwide, due to poor productivity, failure to thrive and control costs [2-6]. In spite of the high prevalence of *Chabertia* reported in small ruminants [7], it is not clear whether the small ruminants harbour one or more than one species. Based on morphological features (e.g., cervical groove and cephalic vesicle) of adult worms, various *Chabertia* species have

* Correspondence: xingquanzhu1@hotmail.com

†Equal contributors

¹State Key Laboratory of Veterinary Etiological Biology, Key Laboratory of Veterinary Parasitology of Gansu Province, Lanzhou Veterinary Research Institute, Chinese Academy of Agricultural Sciences, Lanzhou, Gansu Province 730046, PR China

²College of Animal Science and Technology, Jilin Agricultural University, Changchun, Jilin Province 130118, PR China

Full list of author information is available at the end of the article

been described in sheep and goats in China, including *C. ovina*, *C. rishati*, *C. bovis*, *C. erschowi*, *C. gaohanensis* sp. nov and *C. shaanxiensis* sp. nov [8-10]. However, to date, only *Chabertia ovina* is well recognized as taxonomically valid [11,12]. Obviously, the identification and distinction of *Chabertia* to species using morphological criteria alone is not reliable. Therefore, there is an urgent need for suitable molecular approaches to accurately identify and distinguish closely-related *Chabertia* species from different hosts and regions.

Molecular tools, using genetic markers in mitochondrial (mt) genomes and the internal transcribed spacer (ITS) regions of nuclear ribosomal DNA (rDNA), have been used effectively to identify and differentiate parasites of different groups [13-16]. For nematodes, recent studies showed that mt genomes are useful genetic markers for the identification and differentiation of closely-related species [17,18]. In addition, employing ITS rDNA sequences, recent studies also demonstrated that *Haemonchus placei* and *H. contortus* are distinct species [19]; *Trichuris suis* and *T. trichiura* are different nematode species [20,21].

Using a long-range PCR-coupled sequencing approach [22], the objectives of the present study were (i) to characterize the ITS rDNA and mt genomes of *C. ovina* and *C. erschowi* from goat and yak in China, (ii) to compare these ITS sequences and mt genome sequences, and (iii) to test the hypothesis that *C. erschowi* is a valid species in phylogenetic analyses of these sequence data.

Methods

Parasites and isolation of total genomic DNA

Adult specimens of *C. ovina* (n = 6, coded CHO1-CHO6) and *C. erschowi* (n = 9, coded CHE1-CHE9) were collected, *post-mortem*, from the large intestine of a goat and a yak in Shaanxi and Qinghai Provinces, China, respectively, and were washed in physiological saline, identified morphologically [8,10], fixed in 70% (v/v) ethanol and stored at -20°C until use. Total genomic DNA was isolated separately from 15 individual worms using an established method [23].

Long-range PCR-based sequencing of mt genome

To obtain some mt sequence data for primer design, we PCR-amplified regions of *C. erschowi* of *cox1* gene by using a (relatively) conserved primer pair JB3-JB4.5 [24], *rrnL* gene was amplified using the designed primers *rrnLF* (forward; 5'-GAGCCTGTATTGGGTTCCAGTATGA-3') and *rrnLR* (reverse; 5'-AACTTTTTTTGATTTTCCTTCGTA-3'), *nad1* gene was amplified using the designed primers *nad1F* (forward; 5'-GAGCGTCATTTGTTGGGAAG-3') and *nad1R* (reverse; 5'-CCCCTCAGCAAAA TCAAAC-3'), *cytb* gene was amplified using the designed primers *cytbF* (forward; 5'-GGTACCTTTTTGGCTTT

TTATTATA-3') and *cytbR* (reverse; 5'-ATATGAACAGGGCTTATTATAGGAT-3') based on sequences conserved between *Oesophagostomum dentatum* and *C. ovina* Australia isolate. The amplicons were sequenced in both directions using BigDye terminator v.3.1, ABI PRISM 3730. We then designed primers (Table 1) to regions within *cox1*, *rrnL*, *nad1* and *cytb* and amplified from *C. ovina* (coded CHO1) in four overlapping fragments: *cox1-rrnL*, *rrnL-nad1*, *nad1-cytb* and *cytb-cox1*. Then we designed primers (Table 1) to regions within *cox1*, *rrnL*, *nad5*, *nad1*, *nad2* and *cytb* and amplified from *C. erschowi*

Table 1 Sequences of primers used to amplify mitochondrial DNA regions from *Chabertia erschowi* and *Chabertia ovina* from China

Primer	Sequence (5' to 3')
For <i>rrnS</i>	
CHOF	TCGTTTAGTGGGTATGTGTGGTTCT (for <i>C. ovina</i>)
CHOR	GCCTACTCCCTAACAAATGACGCTC (for <i>C. ovina</i>)
CHEF	GTGGTTTTAGGTTAGGGTTGAGTG (for <i>C. erschowi</i>)
CHER	ACGCTCATACAAAGTAATAACGCA (for <i>C. erschowi</i>)
For <i>nad2</i>	
CHOF	TTTGTGG(C/T)TAAGAGTGTT(G/A)GCTATT (for <i>C. ovina</i>)
CHOR	GAGCCGTAATCAAACATAGTAAATC (for <i>C. ovina</i>)
CHEF	TTTGTGG(C/T)TAAGAGTGTT(G/A)GCTATT (for <i>C. erschowi</i>)
CHER	ACCGTAATCAAACATAGTAAATCT (for <i>C. erschowi</i>)
For <i>C. ovina</i>	
COF	TGGTTGTGGTTTGGGCTCAT
rrnLR	ATGTCCTCACGCTAAGACTGCC
rrnLF	AGTTTGCTTCTGCCAGTGA
ND5R	ACCGTAACCTCGCCATCCTG
ND5F	ACGGCGTTAGTGGAGGAGGA
ND1R	CCACTAACCAACTCCCTTCACCC
ND1F	ATTGGTGCTTTGCGGGCCAGT
ND2R	CCATAAACCTTTAAACCTCCC
ND2F	TTGTTGGTTGGGAGACTATG
CYR	AAAGGGTCCTCAACCAAACA
CYF	CCTGTTGGGGACCTTCTATTG
COR	CCGCAGTAAATAAGCACGAGA
For <i>C. erschowi</i>	
COF	ACCGACGGCTTATGGAAT
rrnLR	AGTGCAACCAACATTATACCTT
rrnLF	TAAAGTTGCTTCTGCCAGTGA
ND1R	ATAATAGCCAACAAAAGCACCGACA
ND1F	CTTGTCGGTGCTTTGCG
CYR	CCGCCTCAATAAACATCTC
CYF	TGGTCCAGATTATTGAAGG
COR	TTACCCGTCAAATACAAAGT

(coded CHE1) in six overlapping fragments: *cox1-rrnL*, *rrnL-nad5*, *nad5-nad1*, *nad1-nad2*, *nad2-cytb* and *cytb-cox1*. The cycling conditions used were 92°C for 2 min (initial denaturation), then 92°C/10 s (denaturation), 50 -58°C (*C. erschowi*) or 56 -65°C (*C. ovina*)/30 s (annealing), and 60°C/10 min (extension) for 10 cycles, followed by 92°C for 2 min, then 92°C/10 s, 50 -58°C (*C. erschowi*) or 56 -65°C (*C. ovina*)/30 s, and 60°C/10 min for 20 cycles, with a cycle elongation of 10 s for each cycle and a final extension at 60°C/10 min. Each amplicon, which represented a single band in a 1.0% (w/v) agarose gel, following electrophoresis and ethidium-bromide staining, was column-purified and then sequenced using a primer walking strategy [22].

Sequencing of ITS rDNA and mt *rrnS* and *nad2*

The full ITS rDNA region including primer flanking 18S and 28S rDNA sequences was PCR-amplified from individual DNA samples using universal primers NC5 (forward; 5'-GTAGGTGAACCTGCGGAAGGATCATT-3') and NC2 (reverse; 5'-TTAGTTTCTTTTCCCTCCGCT-3') described previously [25]. The primers *rrnSF* and *rrnSR* (Table 1) designed to conserved mt genome sequences within the *rrnS* gene were employed for PCR amplification and subsequent sequencing of this complete gene (~ 700 bp) from multiple individuals of *Chabertia* spp. The primers *nad2F* and *nad2R* (Table 1) designed to conserved mt genome sequences within the *nad2* gene were employed for PCR amplification and subsequent sequencing of this complete gene (~ 900 bp) from multiple individuals of *Chabertia* spp..

Sequence analyses

Sequences were assembled manually and aligned against the complete mt genome sequences of *C. ovina* Australia isolate [26] using the computer program Clustal X 1.83 [27] to infer gene boundaries. Translation initiation and termination codons were identified based on comparison with that of *C. ovina* Australia isolate [26]. The secondary structures of 22 tRNA genes were predicted using tRNAscan-SE [28] and/or manual adjustment [29], and rRNA genes were identified by comparison with that of *C. ovina* Australia isolate [26].

Phylogenetic analyses

Amino acid sequences inferred from the 12 protein-coding genes of the two *Chabertia* spp. worms were concatenated into a single alignment, and then aligned with those of 14 other Strongylida nematodes (*Angiostrongylus cantonensis*, GenBank accession number NC_013065 [30]; *Angiostrongylus costaricensis*, NC_013067 [30]; *Angiostrongylus vasorum*, JX268542 [31]; *Aelurostrongylus abstrusus*, NC_019571 [32]; *Chabertia ovina* Australia isolate, NC_013831 [26]; *Cylicocyclus insignis*, NC_013808 [26]; *Metastrongylus pudendotectus*, NC_013813 [26]; *Metastrongylus salmi*, NC_013815

[26]; *Oesophagostomum dentatum*, FM161882 [17]; *Oesophagostomum quadrispinulatum*, NC_014181 [17]; *Oesophagostomum asperum*, KC715826 [33]; *Oesophagostomum columbianum*, KC715827 [33]; *Strongylus vulgaris*, NC_013818 [26]; *Syngamus trachea*, NC_013821 [26], using the Ancylostomatoidea nematode, *Necator americanus*, NC_003416 as the outgroup [29]. Any regions of ambiguous alignment were excluded using Gblocks (http://molevol.cmima.csic.es/castresana/Gblocks_server.html) [34] with the default parameters (Gblocks removed 1.6% of the amino acid alignments) and then subjected to phylogenetic analysis using Bayesian Inference (BI) as described previously [35,36]. Phylograms were drawn using the program Tree View v.1.65 [37].

Results

Nuclear ribosomal DNA regions of the two

Chabertia species

The rDNA region including ITS-1, 5.8S rDNA and ITS-2 were amplified and sequenced from *C. ovina* China isolates, and they were 852-854 bp (GenBank accession nos. KF913466-KF913471) in length, which contained 367-369 bp (ITS-1), 153 bp (5.8S rDNA) and 231-239 bp (ITS-2). These sequences were 862-866 bp in length for *C. erschowi* samples (GenBank accession nos. KF913448-KF913456), containing 375-378 bp (ITS-1), 153 bp (5.8S rDNA) and 239-245 bp (ITS-2).

Features of the mt genomes of the two *Chabertia* species

The complete mt genome sequence of *C. ovina* China isolate and *C. erschowi* were 13,717 bp and 13,705 bp in length, respectively (GenBank accession nos. KF660604 and KF660603, respectively). The two mt genomes contain 12 protein-coding genes (*cox1-3*, *nad1-6*, *nad4L*, *cytb*, *atp6*), 22 transfer RNA genes and two ribosomal RNA genes (*rrnS* and *rrnL*) (Table 2), but the *atp8* gene is missing (Figure 1). The protein-coding genes are transcribed in the same directions, as reported for *Oesophagostomum* spp. [17,33]. Twenty-two tRNA genes were predicted from the mt genomes, which varied from 55 to 63 bp in size. The two ribosomal RNA genes (*rrnL* and *rrnS*) were inferred; *rrnL* is located between tRNA-His and *nad3*, and *rrnS* is located between tRNA-Glu and tRNA-Ser^(UCN). Three AT-rich non-coding regions (NCRs) were inferred in the mt genomes (Table 2). For these genomes, the longest NCR (designated NC2; 250 bp for *C. ovina* China isolate and 240 bp for *C. erschowi* in length) is located between the tRNA-Ala and tRNA-Pro (Figure 1), have an A + T content of 83.75% and 84%, respectively.

Comparative analyses between *C. ovina* and *C. erschowi*


The mt genome sequence of *C. erschowi* was 13,705 bp in length, 12 bp shorter than that of *C. ovina* China

Table 2 Mitochondrial genome organization of *Chabertia erschowi* (CE) and *Chabertia ovina* China isolate (COC) and Australia isolate (COA)

Gene and region	Positions and nt sequence lengths (bp)			Initiation/termination codons		
	CE	COC	COA	CE	COC	COA
<i>cox1</i>	2-1579 (1578)	2-1579 (1578)	2-1579 (1578)	ATT/TAA	ATT/TAA	ATT/TAA
tRN A-Cys (C)	1583-1637 (55)	1583-1639 (57)	1583-1639 (57)			
tRNA-Met (M)	1639-1697 (59)	1640-1699 (60)	1640-1699 (60)			
tRNA-Asp (D)	1699-1758 (60)	1700-1758 (59)	1699-1759 (61)			
tRNA-Gly (G)	1760-1816 (57)	1759-1814 (56)	1757-1814 (58)			
<i>cox2</i>	1817-2512 (696)	1815-2510 (696)	1814-2509 (696)	ATT/TAA	ATA/TAA	ATA/TAA
tRNA-His (H)	2512-2566 (55)	2512-2568 (57)	2511-2567 (57)			
<i>rrnL</i>	2573-3542 (970)	2572-3533 (962)	2570-3531 (962)			
<i>nad3</i>	3543-3881 (339)	3534-3869 (327)	3532-3867 (336)	ATT/TAA	ATT/TAA	ATT/TAA
Non-coding region (NC1)	3882-3965 (84)	3870-3949 (80)	3868-3947 (80)			
<i>nad5</i>	3967-5548 (1582)	3950-5531 (1582)	3948-5529 (1582)	ATT/ TAA	ATT/T	ATT/TAT
tRNA-Ala (A)	5549-5603 (55)	5532-5588 (57)	5530-5586 (57)			
Non-coding region (NC2)	5604-5853 (250)	5589-5828 (240)	5587-5825 (239)			
tRNA-Pro (P)	5854-5909 (56)	5829-5882 (54)	5826-5880 (55)			
tRNA-Val (V)	5927-5982 (56)	5930-5984 (55)	5914-5970 (57)			
<i>nad6</i>	5983-6417 (435)	5985-6416 (432)	5970-6401 (432)	ATA/ TAA	TTG/TAA	TTG/TAA
<i>nad4L</i>	6420-6653 (234)	6418-6651 (234)	6402-6635 (234)	ATT/ TAA	ATT/TAG	ATT/TAG
tRNA-Trp (W)	6681-6736 (56)	6655-6712 (58)	6639-6697 (59)			
tRNA-Glu (E)	6739-6794 (56)	6740-6797 (58)	6725-6784 (60)			
<i>rrnS</i>	6797-7492 (696)	6798-7493 (696)	6780-7479 (700)			
tRNA-Ser UCN (S2)	7493-7547 (55)	7494-7548 (55)	7480-7536 (57)			
tRNA-Asn (N)	7547-7603 (57)	7548-7605 (58)	7535-7593 (59)			
tRNA-Tyr (Y)	7610-7666 (57)	7608-7664 (57)	7595-7652 (58)			
<i>nad1</i>	7667-8539 (873)	7665-8537 (873)	7652-8524 (873)	ATT/TAA	ATT/TAA	ATT/TAA
<i>atp6</i>	8539-9138 (600)	8538-9137 (600)	8525-9121 (597)	ATT/TAA	ATT/TAA	ATT/TAG
tRNA-Lys (K)	9150-9211 (62)	9144-9206 (63)	9128-9191 (64)			
tRNA-LeuUUR (L ₂)	9222-9276 (55)	9215-9269 (55)	9197-9252 (56)			
tRNA-Ser AGN (S1)	9277-9335 (59)	9270-9327 (58)	9252-9304 (53)			
<i>nad2</i>	9336-10175 (840)	9328-10167 (840)	9308-10147 (840)	GTT/TAA	ATT/TAA	ATA/TAA
tRNA-Ile (I)	10176-10234 (59)	10175-10235 (61)	10151-10211 (61)			
tRNA-Arg (R)	10235-10289 (55)	10240-10294 (55)	10215-10270 (56)			
tRNA-Gln (Q)	10290-10345 (56)	10299-10353 (55)	10271-10326 (56)			
tRNA-Phe (F)	10346-10403 (58)	10354-10412 (59)	10326-10385 (60)			
<i>cytb</i>	10404- 11516 (1113)	10413-11525 (1113)	10385-11497 (1113)	ATT/TAG	ATT/TAA	ATT/TAA
tRNA-Leu CUN (L1)	11517-11572 (56)	11529-11584 (56)	11501-11562 (62)			
<i>cox3</i>	11573-12338 (766)	11585-12350 (766)	11557-12327 (771)	ATT/T	ATT/T	ATT/TAA
tRNA-Thr (T)	12339-12397 (59)	12351-12404 (54)	12323-12377 (55)			
<i>nad4</i>	12398-13627 (1230)	12405-13634 (1230)	12376-13608 (1233)	TTG/TAA	TTG/TAA	ATT/TAA
Non-coding region (NC3)	13628 – 1 (75)	13635-1 (84)	13609-1 (75)			

isolate, and 23 bp longer than that of *C. ovina* Australia isolate. The arrangement of the mt genes (i.e., 13 protein genes, 2 *rrn* genes and 22 tRNA genes) and NCRs were

the same. A comparison of the nucleotide sequences of each mt gene as well as the amino acid sequences conceptually translated from individual protein-coding



genes of the two *Chabertia* are given in Table 3. The greatest nucleotide variation between the *C. ovina* China isolate and *C. erschowi* was in the *nad2* gene (19.4% and 17.92%), whereas least differences (7.33%) were detected in the *rrnS* gene, respectively (Table 3). The nucleotide


sequence difference between the entire mt genome of *C. ovina* China isolate and that of *C. erschowi* was 15.33%. Sequence difference between the entire mt genome of *C. ovina* Australia isolate and that of *C. erschowi* was 15.48%. Sequence difference between the entire mt genome of *C. ovina* China isolate and that of *C. ovina* Australia isolate was 4.28%.

The difference in the concatenated amino acid sequences of the 12 protein-coding genes of the *C. ovina* China isolate and those of *C. erschowi* was 9.36%, 10% between those of the *C. ovina* Australia isolate and those of *C. erschowi*, and 2.37% between those of the *C. ovina* China isolate and those of *C. ovina* Australia isolate. The amino acid sequence differences between each of the 12 protein-coding genes of the *C. ovina* Australia isolate and the corresponding homologues of *C. erschowi* ranged from 0.57-17.92%, with COX1 being the most conserved and NAD2 the least conserved proteins (Table 3). Phylogenetic analyses of concatenated amino acid sequence data sets, using *N. americanus* as the outgroup, revealed that the *Chabertia* and *Oesophagostomum* were clustered together, with absolute support (posterior probability (pp) = 1.00) support (Figure 2).

Sequence variation in complete *nad2* gene was assessed among 15 individuals of *Chabertia* from goats and yaks. Sequences of the six *C. ovina* China isolate individuals were the same in length (840 bp) (GenBank accession nos. KF913472-KF913477). Nucleotide variation among the six *C. ovina* China isolate individuals was detected at 18 sites (18/840; 2.1%). Sequences of the nine *C. erschowi* individuals were the same in length (840 bp) (GenBank accession nos. KF913484-KF913492). Nucleotide variation also

Table 3 Nucleotide and/or predicted amino acid (aa) sequence differences for mt protein-coding and ribosomal RNA genes among *Chabertia erschowi* (CE) and *Chabertia ovina* China isolate (COC) and Australia isolate (COA)

Gene	Nucleotide length (bp)			Nucleotide difference (%)			Number of aa			aa difference (%)		
	CE	COC	COA	CE/COC	CE/COA	COC/COA	CE	COC	COA	CE/COC	CE/COA	COC/COA
<i>atp6</i>	600	600	597	14.33	14.83	5.00	199	199	198	11.06	14.57	7.54
<i>nad1</i>	873	873	873	13.63	13.97	4.35	290	290	290	5.17	8.62	3.79
<i>nad2</i>	840	840	840	19.40	20.12	3.93	279	279	279	17.92	17.20	2.15
<i>nad3</i>	339	336	336	17.40	17.70	6.55	112	112	112	16.96	16.07	7.14
<i>nad4</i>	1230	1230	1233	17.64	18.82	5.43	409	409	410	14.67	17.80	4.88
<i>nad4L</i>	234	234	234	11.97	12.39	4.70	77	77	77	7.79	7.79	0
<i>nad5</i>	1582	1582	1582	17.51	17.32	4.87	527	527	527	14.42	13.66	2.47
<i>nad6</i>	435	432	432	19.08	19.31	5.56	144	143	143	15.97	17.36	1.40
<i>cox1</i>	1578	1578	1578	11.98	12.86	4.06	525	525	525	0.57	0.57	0
<i>cox2</i>	696	696	696	13.36	13.65	4.89	231	231	231	0.87	0.87	0
<i>cox3</i>	766	766	771	14.75	14.01	5.06	255	255	256	3.14	2.35	1.17
<i>cytb</i>	1113	1113	1113	16.80	16.89	4.67	370	370	370	9.73	9.73	1.08
<i>rrnS</i>	696	696	700	7.33	7.71	1.86	-	-	-	-	-	-
<i>rrnL</i>	970	962	962	13.61	13.92	3.20	-	-	-	-	-	-




occurred at 23 sites (23/840; 2.7%). All 15 alignments of the *nad2* sequences revealed that all individuals of *Chabertia* differed at 182 nucleotide positions (182/840; 21.7%). Phylogenetic analysis of the *nad2* sequence data revealed strong support for the separation of *C. ovina* and *C. erschowi* individuals into two distinct clades (Figure 3A).

Sequence variation in complete *rrnS* gene was assessed among 15 individuals of *Chabertia* from goat and yak. Sequences of the *rrnS* gene from the six *C. ovina* China isolate individuals were the same in length (696 bp) (GenBank accession nos. KF913478-KF913483). Nucleotide variation among the six *C. ovina* China isolate individuals was detected at seven sites (7/696; 1.0%).

Sequences of the *rrnS* gene from the nine *C. erschowi* individuals were the same in length (696 bp) (GenBank accession nos. KF913457-KF913465). Nucleotide variation also occurred at 6 sites (6/696; 0.9%). All 15 alignments of the *rrnS* sequences revealed that all individuals of *Chabertia* differed at 56 nucleotide positions (56/696; 8.05%). Phylogenetic analysis of the *rrnS* sequence data revealed strong support for the separation of *C. ovina* and *C. erschowi* individuals into two distinct clades (Figure 3B).

The ITS-1 and ITS-2 sequences from 10 individual adults of *C. ovina* China isolate were compared with that of 6 individual adults of *C. erschowi*. Sequence variations



were 0–2.9% (ITS-1) and 0–2.7% (ITS-2) within the two *Chabertia* species, respectively. However, the sequence differences were 6.3–8.2% (ITS-1) and 10.4–13.6% (ITS-2) between the *C. ovina* China isolate and *C. erschowi*.

Discussion

Chabertia spp. is responsible for economic losses to the livestock industries globally. Although several *Chabertia* species have been described from various hosts based on the microscopic features of the adult worms (e.g. cervical groove and cephalic vesicle), it is not clear whether *C. erschowi* is valid as a separate species due to unreliable morphological criteria. For this reason, we employed a molecular approach, so that comparative genetic analyses could be conducted.

In the present study, substantial levels of nucleotide differences (15.33%) were detected in the complete mt genome between *C. ovina* China isolate and *C. erschowi*, and 15.48% between *C. ovina* Australia isolate and *C. erschowi*. These mtDNA data provide strong support that *C. erschowi* represents a single species because a previous comparative study has clearly indicated that variation in mtDNA sequences between closely-related species were typically 10%–20% [13].

The difference in amino acid sequences of the concatenated 12 proteins encoded by the complete mt genome between *C. ovina* China isolate and *C. erschowi* is 9.36%, and 10% between the *C. ovina* Australia isolate and *C. erschowi*. This level of amino acid variation is higher than those of other nematodes. Previous studies of other congener nematodes have detected low level differences in 12 protein sequences. For example, differences in amino acid sequences between *A. duodenale* and *A. caninum* is 4.1% [29,38], and between *Toxocara malaysiensis* and *Toxocara cati* is 5.6% [39], and between *O. dentatum* and *O. quadrispinulatum* is 3.22% [17]. In addition, substantial levels of nucleotide differences (6.3%–8.2% in ITS-1 and 10.4–13.6% in ITS-2) were also detected between *C. ovina* China isolate and *C. erschowi*. These results also indicate that *C. erschowi* is a separate species from *C. ovina*. This proposal was further supported by phylogenetic analysis based on mtDNA sequences (Figure 3), although, to date, only small numbers of adult worms have been studied molecularly. Clearly, larger population genetic and molecular epidemiological studies should be conducted using the mt and nuclear markers defined in this study to further test this proposal/hypothesis.

Conclusion

The findings of this study provide robust genetic evidence that *C. erschowi* is a separate and valid species from *C. ovina*. The mtDNA and rDNA datasets reported

in the present study should provide useful novel markers for further studies of the taxonomy and systematics of *Chabertia* spp. from different hosts and geographical regions.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

XQZ and GHL conceived and designed the study, and critically revised the manuscript. GHL, LZ and HQS performed the experiments, analyzed the data and drafted the manuscript. GHZ, JZC and QZ helped in study design, study implementation and manuscript revision. All authors read and approved the final manuscript.

Acknowledgements

This work was supported in part by the International Science & Technology Cooperation Program of China (Grant No. 2013DFA31840), the "Special Fund for Agro-scientific Research in the Public Interest" (Grant No. 201303037) and the Science Fund for Creative Research Groups of Gansu Province (Grant No. 1210RJIA006).

Author details

¹State Key Laboratory of Veterinary Etiological Biology, Key Laboratory of Veterinary Parasitology of Gansu Province, Lanzhou Veterinary Research Institute, Chinese Academy of Agricultural Sciences, Lanzhou, Gansu Province 730046, PR China. ²College of Animal Science and Technology, Jilin Agricultural University, Changchun, Jilin Province 130118, PR China. ³College of Veterinary Medicine, Hunan Agricultural University, Changsha, Hunan Province 410128, PR China. ⁴College of Veterinary Medicine, Northwest A&F University, Yangling, Shannxi Province 712100, PR China. ⁵Qinghai Academy of Animal Science and Veterinary Medicine, Xining, Qinghai Province, PR China.

Received: 14 December 2013 Accepted: 18 January 2014

Published: 22 January 2014

References

1. Hotez PJ, Fenwick A, Savioli L, Molyneux DH: **Rescuing the bottom billion through control of neglected tropical diseases.** *Lancet* 2009, **373**:1570–1575.
2. Broughan JM, Wall R: **Faecal soiling and gastrointestinal helminth infection in lambs.** *Int J Parasitol* 2007, **37**:1255–1268.
3. Sweeny JP, Robertson ID, Ryan UM, Jacobson C, Woodgate RG: **Impacts of naturally acquired protozoa and strongylid nematode infections on growth and faecal attributes in lambs.** *Vet Parasitol* 2012, **184**:298–308.
4. Sweeny JP, Ryan UM, Robertson ID, Jacobsen C: **Molecular identification of naturally acquired strongylid infections in lambs—An investigation into how lamb age influences diagnostic sensitivity.** *Vet Parasitol* 2012, **187**:227–236.
5. Sweeny JP, Ryan UM, Robertson ID, Niemeyer D, Hunt PW: **Development of a modified molecular diagnostic procedure for the identification and quantification of naturally occurring strongylid larvae on pastures.** *Vet Parasitol* 2012, **190**:467–481.
6. Roeber F, Jex AR, Campbell AJ, Nielsen R, Anderson GA, Stanley KK, Gasser RB: **Establishment of a robotic, high-throughput platform for the specific diagnosis of gastrointestinal nematode infections in sheep.** *Int J Parasitol* 2012, **42**:1151–1158.
7. Tariq KA, Chishti MZ, Ahmad F, Shawl AS: **Epidemiology of gastrointestinal nematodes of sheep managed under traditional husbandry system in Kashmir valley.** *Vet Parasitol* 2008, **158**:138–143.
8. Xiong DS, Kong FY: ***Chabertia erschowi*.n.sp.—A new parasitic nematode of sheep and goat in China.** *J Beijing Agr Univ* 1956, **2**:115–122 (in Chinese).
9. Zhang JL: ***Chabertia shaanxiensis* sp. nov (Nematode: Strongylidae) from the cattle.** *J Anim Vet Adv* 1985, **16**:137–141.
10. Zhang LP, Li JB, Liu SG, An RY: **Comparison of the head structure of two *Chabertia* species by scanning electron microscope.** *J Chin Electron Micro Soc* 1998, **17**:345–346 (in Chinese).

11. Khalafalla RE, Elseify MA, Elbahy NM: Seasonal prevalence of gastrointestinal nematode parasites of sheep in Northern region of Nile Delta, Egypt. *Parasitol Res* 2011, **108**:337–340.
12. Sissay MM, Uggla A, Waller PJ: Prevalence and seasonal incidence of nematode parasites and fluke infections of sheep and goats in eastern Ethiopia. *Trop Anim Health Prod* 2007, **39**:521–531.
13. Blouin MS: Molecular prospecting for cryptic species of nematodes: mitochondrial DNA versus internal transcribed spacer. *Int J Parasitol* 2002, **32**:527–531.
14. Liu GH, Chen F, Chen YZ, Song HQ, Lin RQ, Zhou DH, Zhu XQ: Complete mitochondrial genome sequence data provides genetic evidence that the brown dog tick *Rhipicephalus sanguineus* (Acari: Ixodidae) represents a species complex. *Int J Biol Sci* 2013, **9**:361–369.
15. Jabbar A, Mohandas N, Jex AR, Gasser RB: The mitochondrial genome of *Protostrongylus rufescens* – implications for population and systematic studies. *Parasit Vectors* 2013, **6**:263.
16. Liu GH, Li C, Li JY, Zhou DH, Xiong RC, Lin RQ, Zou FC, Zhu XQ: Characterization of the complete mitochondrial genome sequence of *Spirometra erinaceieuropaei* (Cestoda: Diphylobothriidae) from China. *Int J Biol Sci* 2012, **8**:640–649.
17. Lin RQ, Liu GH, Hu M, Song HQ, Wu XY, Li MW, Zhang Y, Zou FC, Zhu XQ: *Oesophagostomum dentatum* and *Oesophagostomum quadrispinulatum*: characterization of the complete mitochondrial genome sequences of the two pig nodule worms. *Exp Parasitol* 2012, **131**:1–7.
18. Liu GH, Gasser RB, Su A, Nejsum P, Peng L, Lin RQ, Li MW, Xu MJ, Zhu XQ: Clear genetic distinctiveness between human- and pig-derived *Trichuris* based on analyses of mitochondrial datasets. *PLoS Negl Trop Dis* 2012, **6**:e1539.
19. Stevenson LA, Chilton NB, Gasser RB: Differentiation of *Haemonchus placei* from *H. contortus* (Nematoda: Trichostrongylidae) by the ribosomal DNA second internal transcribed spacer. *Int J Parasitol* 1995, **25**:483–488.
20. Cutillas C, Callejón R, de Rojas M, Tewes B, Ubeda JM, Ariza C, Guevara DC: *Trichuris suis* and *Trichuris trichiura* are different nematode species. *Acta Trop* 2009, **111**:299–307.
21. Liu GH, Zhou W, Nisbet AJ, Xu MJ, Zhou DH, Zhao GH, Wang SK, Song HQ, Lin RQ, Zhu XQ: Characterization of *Trichuris trichiura* from humans and *T. suis* from pigs in China using internal transcribed spacers of nuclear ribosomal DNA. *J Helminthol* 2014, **88**:64–68.
22. Hu M, Jex AR, Campbell BE, Gasser RB: Long PCR amplification of the entire mitochondrial genome from individual helminths for direct sequencing. *Nat Protoc* 2007, **2**:2339–2344.
23. Gasser RB, Hu M, Chilton NB, Campbell BE, Jex AJ, Otranto D, Cafarchia C, Beveridge I, Zhu X: Single-strand conformation polymorphism (SSCP) for the analysis of genetic variation. *Nat Protoc* 2006, **1**:3121–3128.
24. Bowles J, Blair D, Mcmanus DP: Genetic variants within the genus *Echinococcus* identified by mitochondrial DNA sequencing. *Mol Biochem Parasitol* 1992, **54**:165–174.
25. Zhu X, Chilton NB, Jacobs DE, Boes J, Gasser RB: Characterisation of *Ascaris* from human and pig hosts by nuclear ribosomal DNA sequences. *Int J Parasitol* 1999, **29**:469–478.
26. Jex AR, Hall RS, Littlewood DT, Gasser RB: An integrated pipeline for next generation sequencing and annotation of mitochondrial genomes. *Nucleic Acids Res* 2010, **38**:522–533.
27. Thompson JD, Gibson TJ, Plewniak F, Jeanmougin F, Higgins DG: The Clustal X windows interface: flexible strategies for multiple sequence alignment aided by quality analysis tools. *Nucleic Acids Res* 1997, **24**:4876–4882.
28. Lowe TM, Eddy SR, tRNAscan-SE: A program for improved detection of transfer RNA genes in genomic sequence. *Nucleic Acids Res* 1997, **25**:955–964.
29. Hu M, Chilton NB, Gasser RB: The mitochondrial genomes of the human hookworms, *Ancylostoma duodenale* and *Necator americanus* (Nematoda: Secernentea). *Int J Parasitol* 2002, **32**:145–158.
30. Lv S, Zhang Y, Zhang L, Liu Q, Liu HX, Hu L, Wei FR, Graeff-Teixeira C, Zhou XN, Utzinger J: The complete mitochondrial genome of the rodent intra-arterial nematodes *Angiostrongylus cantonensis* and *Angiostrongylus costaricensis*. *Parasitol Res* 2012, **111**:115–123.
31. Gasser RB, Jabbar A, Mohandas N, Schnyder M, Deplazes P, Littlewood DT, Jex AR: Mitochondrial genome of *Angiostrongylus vasorum*: comparison with congeners and implications for studying the population genetics and epidemiology of this parasite. *Infect Genet Evol* 2012, **12**:1884–1891.
32. Jabbar A, Jex AR, Mohandas N, Hall RS, Littlewood DT, Gasser RB: The mitochondrial genome of *Aelurostrongylus abstrusus*-diagnostic, epidemiological and systematic implications. *Gene* 2013, **516**:294–300.
33. Zhao GH, Hu B, Cheng WY, Jia YQ, Li HM, Yu SK, Liu GH: The complete mitochondrial genomes of *Oesophagostomum asperum* and *Oesophagostomum columbianum* in small ruminants. *Infect Genet Evol* 2013, **19**:205–211.
34. Talavera G, Castresana J: Improvement of phylogenies after removing divergent and ambiguously aligned blocks from protein sequence alignments. *Syst Biol* 2007, **56**:564–577.
35. Liu GH, Wang Y, Xu MJ, Zhou DH, Ye YG, Li JY, Song HQ, Lin RQ, Zhu XQ: Characterization of the complete mitochondrial genomes of two whipworms *Trichuris ovis* and *Trichuris discolor* (Nematoda: Trichuridae). *Infect Genet Evol* 2012, **12**:1635–1641.
36. Ronquist F, Huelsenbeck JP: MrBayes 3: Bayesian phylogenetic inference under mixed models. *Bioinformatics* 2003, **19**:1572–1574.
37. Page RD: TREEVIEW: an application to display phylogenetic trees on personal computers. *Comput Appl Biosci* 1996, **12**:357–358.
38. Jex AR, Waeschenbach A, Hu M, van Wyk JA, Beveridge I, Littlewood DT, Gasser RB: The mitochondrial genomes of *Ancylostoma caninum* and *Bunostomum phlebotomum*—two hookworms of animal health and zoonotic importance. *BMC Genomics* 2009, **10**:79.
39. Li MW, Lin RQ, Song HQ, Wu XY, Zhu XQ: The complete mitochondrial genomes for three *Toxocara* species of human and animal health significance. *BMC Genomics* 2008, **9**:224.

doi:10.1186/1756-3305-7-44

Cite this article as: Liu et al.: *Chabertia erschowi* (Nematoda) is a distinct species based on nuclear ribosomal DNA sequences and mitochondrial DNA sequences. *Parasites & Vectors* 2014 **7**:44.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

