



CASE REPORT

Open Access

An uncommon cause of seizures in children living in developed countries: neurocysticercosis -a case report

Irene Raffaldi^{1*}, Carlo Scolfaro¹, Federica Mignone¹, Sonia Aguzzi¹, Federica Denegri², Pier-Angelo Tovo¹

Abstract

Neurocysticercosis represents an important cause of seizures in children in endemic countries, such as Latin America, Asia and sub-Saharan Africa, while in Europe, especially in Italy, the cases of neurocysticercosis are anecdotal. We report the case of a 6 year old boy, born and lived for four years in Cameroon, who presented a right emiconvulsion. The diagnosis was neurocysticercosis. This case accentuates the need to consider neurocysticercosis in a child presenting with non febrile seizures, mainly if he emigrated from an area of high prevalence or if he had long-term stay in endemic regions.

Background

Neurocysticercosis represents an important cause of seizures in children in endemic countries. It is due to the brain involvement by the larval stage of the cestode *Taenia solium* (cysticerci). This parasite is commonly found in developing countries of Latin America, Asia and sub-Saharan Africa [1-5]. The prevalence of neurocysticercosis in some of these countries exceeds 10% [6]; conversely in Europe the cases of neurocysticercosis are anecdotal, especially caused by migratory flows from endemic zones or international travels. The case reported below describes a rare cause of seizures in a child who lives in a developed country.

Case presentation

Case report

A 6 year old boy who was born and lived in Cameroon for four years, in a rural area, was admitted to the Emergency Department with seizure lasting more than thirty minutes, not responsive to Diazepam 0.5 mg/Kg e.r. The patient had immigrated to Italy two years previously. No familiarity for seizures or headache. At the age of 1 year the child had experienced an acute and isolated febrile seizure with oculogyration. He had no recent history of traumatic or infective episodes, neither ingestion of drugs

or weight loss. When he woke up that morning, he had speaking difficulties, right deviation of the mouth, followed by right head and gaze deviation, right emiconvulsion, and loss of consciousness. At admission the child was afebrile, unconscious, with accelerated heart rate and respiratory difficulties. His head was deviated to the right, there was a generalized hypertonia with hyperextension of right arm and flexion of the left arm. The electroencephalogram (EEG) detected a slow mono-polymorphic activity on the left and central electroencephalographic leads. This activity was absent on the right leads. It suggested a post-critical focal cerebral suffering.

He was directly admitted to the Intensive Care Unit and he was successfully treated with Phenobarbital (5 mg/Kg i.v.). The magnetic resonance imaging (MRI) of the brain showed two cystic round lesions located one in the right lentiform nucleus and one in the left frontal-parietal lobe (Figure 1). In the following 36 hours the EEG displayed an improvement of child's cerebral conditions with the disappearance of the asymmetric slow activity reported previously. After 48 hours the child was transferred to our Department. Blood exams revealed high eosinophil cell count (720 cells/ μ L) and Ig E levels (217 UI/ml). Western blot assay detected specific antibodies against cysticercus (LDBIO DIAGNOSTICS, Lyon, France). Therefore the diagnosis of neurocysticercosis was made and the appropriated therapy was started: the child received, orally, albendazole

* Correspondence: ire_raffaldi@yahoo.it

¹Department of Pediatrics, Division of Infectious Diseases, University of Turin, Regina Margherita Children's Hospital, Piazza Polonia 94, 10126, Turin, Italy
Full list of author information is available at the end of the article




Figure 1 Brain MRI before specific treatment. (a) axial T2-weighted MRI shows an alteration in left frontal-parietal cortex attributing to vasogenic edema (b) coronal T1-weighted post contrast MRI shows a little ring-enhancing lesion in left frontal cortex.

(15 mg/Kg/day) in two divided doses and dexamethasone (2 mg/day) for eight days.

Six months later, the eosinophil and IgE levels got back to normal. At brain MRI calcified areas, corresponding to the previous lesions, were indicated, without any new active lesions (Figure 2).

In the following two years the patient remained seizure-free on oral antiepileptic therapy. Then the patient has been lost to follow up.

Discussion

Neurocysticercosis is the most common parasitic infection of central nervous system [7]. *Taenia solium*, responsible of the disease, can causes, in individuals, intestinal infection and cysticercosis. In the classic life cycle humans develop taeniasis after eating not well cooked pork meat, containing cysticerci. Larvae develop into adult tapeworm inside human's small intestine. The tape-worm releases daily eggs disseminated to the environment through feces. In developing countries, where sanitary regulations are often neglected, roaming pigs may ingest food polluted by eggs, becoming the intermediate host. Once in the pig's intestinal tract, eggs embrionate into oncospheres, which cross the intestinal wall and, through bloodstream, migrate to the target tissues (brain, muscle, subcutaneous surface). The cycle can be interrupted if individuals accidentally ingest eggs through food or water contaminated by feces. In this case, the parasite perceives the human as an intermediate host and disseminates to a variety of organs, including the central nervous system, where cysticerci develops [8].

The clinical manifestations of neurocysticercosis are variable, depending on the number, stage, location as well as the host response [9,10]. A large part of individuals holding the parasite in the central nervous system are asymptomatic [8]. In some patients symptoms may develop many years after the brain infection, as it has happened to our patient. The most frequent manifestations, due to parenchymal location, are seizures, learning disability, difficulty with balance, behavior changes. Extraparenchymal location of the parasite causes increased intracranial pressure with headache, vomiting and papilloedema [7,8,11].

Parenchymal forms of neurocysticercosis have a better prognosis than extraparenchymal forms, in term of remission of clinical signs [7]. Carpio and Hauser noticed that the risk of seizure recurrence is high after a first acute symptomatic fit but the risk seems related to the persistence of active brain lesions [12].

Histological demonstration of parasites from a brain biopsy should be the safe "gold standard" for the diagnosis of neurocysticercosis. Clearly this procedure is limited because of its invasiveness. Therefore neuroimaging remains the main instrument for the diagnosis. In the identification of extraparenchymal cysts MRI is more sensitive than Computed Tomography (CT) scan [10].

Immunological tests have low sensitivity and specificity, especially for single lesion. The enzyme-linked immunosorbent assay has 65% specificity and 50% sensitivity and it can give false negative results in case of parenchymal neurocysticercosis, inactive lesion or helminthic infections. The enzyme-linked immunoelectrotransfer blot




Figure 2 Post treatment brain MRI. Axial T2-weighted MRI shows edema's disappearance and calcified lesions, one in right lentiform nucleus and one in left frontal-parietal cortex.

assay, using purified glycoprotein antigens from *Taenia solium* cysticerci, has been reported to be highly specific (100%) and nearly 98% sensitive for patients with either multiple active parenchymal cysts or extraparenchymal neurocysticercosis [10,13].

The differential diagnoses include echinococcosis, pyogenic brain abscess, fungal abscess, tuberculoma, toxoplasmosis, a primary or metastatic tumor and infectious vasculitis [14].

Antiparasitic drugs are the mainstay of treatment; in particular, albendazole may be favourable for the treatment of parenchymal cysts because of its power to pass into cerebral spinal fluid [8]. Previous studies recommended the administration of albendazole at a dosage of 15 mg/Kg/die for 1 month, but later studies prove that a 1 week course is equally effective [14]. Corticosteroids are useful for reducing local edema and inflammation around dying parenchymal cysts; so they are often

administered together with antiparasitic drugs. Kalra displayed that the association of albendazole and dexamethasone increases complete or partial resolution of lesions and reduces the risk of recurrence of seizure among children presenting one or two ring-enhancing lesion on CT [15].

This case is interesting mainly for epidemiological reasons. In fact neurocysticercosis is endemic in Latin America, Africa and some Asiatic countries. In Europe, many cases have been reported in Portugal, Spain, Poland and Romania. In Italy it is a rare disease. In recent years no cases have been described, but with high rate of immigration from endemic areas (Africa and East Europe) this parasitosis will be found in our country too [16].

Conclusions

This case emphasizes the need to consider a parasitic infection of the central nervous system, particularly

neurocysticercosis, in a child with an onset of epilepsy especially if he had long-term stay in endemic regions, or he emigrated from an endemic area, although not recently. In case of relevant suspicion of neurocysticercosis it needs testing serum cysticercal antibody and making neuroimaging examinations.

Consent

Verbal informed consent was obtained from the patient's parents for publication of this case report and any accompanying images, at time of diagnosis; then the patient (with his family) has been lost to follow up.

Acknowledgements

We would thank Paola Sciortino for her precious help in the interpretation of the medical imaging.

Author details

¹Department of Pediatrics, Division of Infectious Diseases, University of Turin, Regina Margherita Children's Hospital, Piazza Polonia 94, 10126, Turin, Italy.

²Department of Neuroradiology, Orthopaedics and Traumatology Hospital, Via Zuretti 29, 10126, Turin, Italy.

Authors' contributions

IR, CS, FM, SA have made useful contribution in drafting the manuscript and in the revision of the literature.

PAT has been involved in revising it critically for important intellectual content. FD has participated in the diagnostic pathways. All authors read and approved the final manuscript.

Competing interests

The authors declare that they have no competing interests.

Received: 26 October 2010 Accepted: 25 January 2011

Published: 25 January 2011

References

1. Roman G, Sotelo J, Del Brutto O, Flisser A, Dumas M, Wadia N, Botero D, Cruz M, Garcia H, De Bittencourt PRM, Treilles I, Arriagada C, Lorenzana P, Nash TE, Spina-Franca A: **A proposal to declare neurocysticercosis an international reportable disease.** *Bull World Health Organ* 2000, **78**(3):399-406.
2. Grill J, Rakotomalala W, Andriantsimahavandy A, Boisier P, Guyon P, Roux J, Esterre P: **High prevalence of serological markers of cysticercosis among epileptic Malagasy children.** *Ann Trop Paediatr* 1996, **16**:185-191.
3. Singh S, Singh P: **Clinical profile and etiology of partial seizures in North Indian infants and children.** *J Epilepsy* 1997, **10**:32-36.
4. Gaffo AL, Guillén-Pinto D, Campos-Olazábal P, Burneo JG: **Cysticercosis as the main cause of partial seizures in children in Peru.** *Rev Neurol* 2004, **39**(10):924-926.
5. García HH, Gonzalez AE, Evans CA, Gilman RH: **Taenia solium cysticercosis.** In *Lancet*. Volume 362. Cysticercosis Working Group in Peru; 2003;(9383):547-556.
6. Ong S, Talan DA, Moran GJ, Mower W, Newdow M, Tsang VC, Pinner RW, EMERGENCY ID NET Study Group: **Neurocysticercosis in radiographically imaged seizure patients in U.S. Emergency Departments.** *Emerg Infect Dis* 2002, **8**(6):608-613.
7. Carpio A: **Neurocysticercosis: an update.** *Lancet Infect Dis* 2002, **2**(12):751-762.
8. Mody R, Nield LS, Stauffer W, Kamat D: **Seizure in a 20-month-old native of Minnesota: a case of neurocysticercosis.** *Pediatr Emerg Care* 2005, **21**(12):860-862.
9. Garcia HH, Del Brutto OH, Cysticercosis Working Group in Peru: **Neurocysticercosis: updated concepts about an old disease.** *Lancet Neurol* 2005, **4**(10):653-661.
10. Singh P, Singh S: **Neurocysticercosis in children.** *J Child Neurol* 2004, **19**(7):482-492.
11. San-Juan Orta D: **Clinical manifestations of neurocysticercosis.** *Neurologia* 2009, **24**(5):331-335.
12. Carpio A, Hauser WA: **Prognosis for seizure recurrence in patients with newly diagnosed neurocysticercosis.** *Neurology* 2002, **59**(11):1730-1734.
13. Ramos-Kuri M, Montoya RM, Padilla A, Govezensky T, Diaz ML, Sciueto E, Sotelo J, Larrade C: **Immunodiagnosis of neurocysticercosis. Disappointing performance of serology (enzyme-linked immunosorbent assay) in an unbiased sample of neurological patients.** *Arch Neurol* 1992, **49**(6):633-636.
14. Sinha S, Sharma BS: **Neurocysticercosis: a review of current status and management.** *J Clin Neurosci* 2009, **16**(7):867-876.
15. Kalra V, Dua T, Kumar V: **Efficacy of albendazole and short-course dexamethasone treatment in children with 1 or 2 ring-enhancing lesions of neurocysticercosis: a randomized controlled trial.** *J Pediatr* 2003, **143**(1):111-114.
16. Carangelo B, Erra S, Del Basso De Caro ML, Bucciero A, Vizioli L, Panagiotopoulos K, Cerillo A: **Neurocysticercosis. Case report.** *J Neurosurg Sci* 2001, **45**(1):43-46.

doi:10.1186/1824-7288-37-9

Cite this article as: Raffaldi et al.: An uncommon cause of seizures in children living in developed countries: neurocysticercosis -a case report. *Italian Journal of Pediatrics* 2011 **37**:9.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

