



Corrigendum

Corrigendum to ‘Roles of alternative RNA splicing of the Bif-1 gene by SRRM4 during the development of treatment-induced neuroendocrine prostate cancer’. [EBioMedicine 31 (2018) 267–275]



Yu Gan^{a,b,c}, Yanan Li^a, Zhi Long^{a,c}, Ahn R. Lee^a, Ning Xie^a, Jessica M. Lovnicki^a, Yuxin Tang^c, Xiang Chen^b, Jiaoti Huang^d, Xuesen Dong^{a,*}

^a Vancouver Prostate Centre, Department of Urologic Sciences, University of British Columbia, 2660 Oak Street, Vancouver, British Columbia V6H 3Z6, Canada


^b Department of Urology, Xiangya Hospital, Central South University, Changsha, China

^c Department of Urology, Third Xiangya Hospital, Central South University, Changsha, China

^d Department of Pathology, Duke University School of Medicine, Durham, NC, USA

The authors wish to point out that:

- i) There is a typo in one of the labels of Fig. 2a. In Fig. 2a (top right), “IHC Positive (H score < 0.3)” should be replaced with “IHC Negative (H score < 0.3)”;
- ii) There is an error in Fig. 4a, in which the bottom right micrograph (Bif-1b/c RISH=2) is inadvertently duplicated from bottom left micrograph (SRRM4 RISH=2). A corrected version of Fig. 4a is included below. The authors apologize for the errors.



These errors do not change the scientific conclusions of the article in any way.

DOI of original article: <http://dx.doi.org/10.1016/j.ebiom.2018.05.002>.

* Corresponding author.

E-mail addresses: yli@prostatecentre.com (Y. Li), zlong@prostatecentre.com (Z. Long), alee@prostatecentre.com (A.R. Lee), nxie@prostatecentre.com (N. Xie), jlovnicki@prostatecentre.com (J.M. Lovnicki), jiaoti.huang@duke.edu (J. Huang), xdong@prostatecentre.com (X. Dong).

<https://doi.org/10.1016/j.ebiom.2020.102741>

2352-3964/© 2020 The Author(s). Published by Elsevier B.V. This is an open access article under the CC BY-NC-ND license. (<http://creativecommons.org/licenses/by-nc-nd/4.0/>)