

POSTER PRESENTATION

Open Access

Short-term depression of inhibitory Purkinje cell synapses enhances gain modulation in the cerebellar nuclei


Dimitris Bampasakis^{1*}, Reinoud Maex², Neil Davey¹, Volker Steuber¹

From Twenty Second Annual Computational Neuroscience Meeting: CNS*2013
Paris, France. 13-18 July 2013

Information in neurons can be encoded by their action potential rate, thus making the transformation of input to output rate, the input-output (I-O) relationship, a core computational function. Introduction of a second input, often called modulatory input, can modify this I-O relationship in ways that correspond to different

arithmetic operations [1]. Here, we examine the modulation of the slope of the I-O relationship, also referred to as gain modulation.

Gain modulation can be based on a wide variety of biophysical mechanisms, with short-term depression (STD) of excitatory synapses being one of them [2].



* Correspondence: d.bampasakis@herts.ac.uk

¹Science and Technology Research Institute, University of Hertfordshire, Hatfield AL10 9AB, UK

Commonly, gain modulation is studied by examining the effect of tonic or synaptic inhibition on the excitatory I-O relationship. However, some projection neurons, like cerebellar Purkinje cells (PCs), are inhibitory. Therefore, the opposite scenario, in which the effect of inhibition on output rate is being modulated by an excitatory input, may occur as well. As a previous study found that inhibitory synaptic input variability can change the output rate of neurons in the cerebellar nuclei (CN) [3], the question arises how excitatory input can modulate this relationship.

Considering the excitatory input from mossy fibres (MF) onto CN neurons as modulatory, we investigated the effects on gain control exerted by STD of the inhibitory synapses that PCs make on a model CN neuron [3]. We found that STD at the inhibitory PC-CN synapse enhanced gain modulation (Figure 1). Thus, like STD at excitatory synapses, STD at inhibitory synapses can enable neurons to perform multiplicative operations on their inputs.

Author details

¹Science and Technology Research Institute, University of Hertfordshire, Hatfield AL10 9AB, UK. ²Department of Cognitive Sciences, École Normale Supérieure, Paris 75005, France.

Published: 8 July 2013

References

1. Silver RA: **Neuronal arithmetic.** *Nature Reviews Neuroscience* 2010, **11**:474-489.
2. Rothman JS, Cathala L, Steuber V, Silver RA: **Synaptic depression enables neuronal gain control.** *Nature* 2009, **457**:1015-1018.
3. Luthman J, Hoebeek FE, Maex R, Davey N, Adams R, De Zeeuw CI, Steuber V: **STD-dependent and independent encoding of input irregularity as spike rate in a computational model of a cerebellar nucleus neuron.** *Cerebellum* 2011, **10**:667-682.

doi:10.1186/1471-2202-14-S1-P374

Cite this article as: Bampasakis *et al.*: Short-term depression of inhibitory Purkinje cell synapses enhances gain modulation in the cerebellar nuclei. *BMC Neuroscience* 2013 **14**(Suppl 1):P374.

**Submit your next manuscript to BioMed Central
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

