



POSTER PRESENTATION

Open Access

# Diagnosis and prevalence of Congenital Adrenal Hyperplasia (CAH) in Austrian children screened or not screened for CAH

Katharina Luxenberger<sup>1</sup>, David Kasper<sup>2</sup>, Elke Fröhlich-Reiterer<sup>1</sup>, Elisabeth Suppan<sup>1</sup>, Gudrun Weinhandl<sup>1</sup>, Martin Borkenstein<sup>1\*</sup>

From 7th APPEs Biennial Scientific Meeting  
Nusa Dua, Bali. 14-17 November 2012

Prevalence of Congenital Adrenal Hyperplasia (CAH) is not exactly known in the Austria; a number of patients with CAH might not be diagnosed, especially males. CAH is in about 95 % of the cases due to a defect in the 21-hydroxylation ('classical CAH'). Newborn screening for CAH, based on the measurement of 17 $\alpha$ -hydroxyprogesterone (17-OHP) was shown to be efficient for diagnosis, and is part of the newborn screening programme in Austria since April 2001.

In our study we compared 2 groups of children:

Group A, children born in Styria (a province of Austria), 1992 – 2001, n = 119.001, m 61.256, and f 57.745;

Group B, children born in Styria 2002 – 2011, n = 103.228, m 52.722, and f 50.506

In group A (patients not screened), CAH was diagnosed in 8 children (m 4; f 4); 4 of them with simple virilising (SV) 21-OH deficiency (m 3; f 1) and 4 with salt wasting (SW) 21-OH deficiency (m 1; f 3).

In group B, 98,7 % of all newborns born in Styria could be screened by measuring 17-OHP in a dried blood spot on filter paper. Recall rate was 0,578 %. CAH was diagnosed in 10 children (m 3; f 7). 8 of them with SW (m 2; f 6), and 2 with 11 $\beta$ -hydroxylase deficiency (m 1; f 1).

Whereas group A displayed the expected Mendelian sex ratio, group B showed a strong female predominance (m 3; f 7)

Prevalence of CAH was 1:14.875 newborns in group A (not screened). In group B (newborns screened) prevalence was 1:10.132.

If one exclude the 2 patients with 11 $\beta$ -hydroxylase deficiency from group B, prevalence of 'classical CAH' was 1:14.875 in group A, compared to 1:12.903 in group B.

These data show that newborn screening for CAH seems to increase the rate of detection of CAH.

#### Authors' details

<sup>1</sup>Departments of Pediatrics and Adolescence Medicine, Medical Universities Graz, Graz, Austria. <sup>2</sup>Departments of Pediatrics and Adolescence Medicine, Medical Universities Vienna, Vienna, Austria.

Published: 3 October 2013

doi:10.1186/1687-9856-2013-S1-P111

Cite this article as: Luxenberger *et al.*: Diagnosis and prevalence of Congenital Adrenal Hyperplasia (CAH) in Austrian children screened or not screened for CAH. *International Journal of Pediatric Endocrinology* 2013 2013(Suppl 1):P111.

#### Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at  
[www.biomedcentral.com/submit](http://www.biomedcentral.com/submit)



<sup>1</sup>Departments of Pediatrics and Adolescence Medicine, Medical Universities Graz, Graz, Austria  
Full list of author information is available at the end of the article