


Israel's first national remote cardiac rehabilitation program: A retrospective analysis

DIGITAL HEALTH
Volume 9: 1–8
© The Author(s) 2023
Article reuse guidelines:
sagepub.com/journals-permissions
DOI: 10.1177/20552076231180762
journals.sagepub.com/home/dhj



Irene Nabutovsky^{1,2}, Daniel Breitner¹, Alexis Heller¹, Merav Moreno^{1,2},
Yoav Levine¹ , Yarin Klempfner^{1,2}, Mickey Scheinowitz^{1,3,4}
and Robert Klempfner^{1,1}

Abstract

Aims: Cardiac rehabilitation is an essential component of secondary prevention consistently unexploited by most eligible patients. Accordingly, the remote cardiac rehabilitation program (RCRP) was developed to create optimal conditions for remote instruction and supervision for patients to enable successful completion of the program.

Methods: This study comprised 306 patients with established coronary heart disease who underwent a 6-month RCRP. RCRP involves regular exercise, monitored by a smartwatch that relays data to the operations center and a mobile application on the patient's smartphone. A stress test was performed immediately before the RCRP and repeated after 3 months. The aims were to determine the effectiveness of the RCRP in improving aerobic capacity, and correlating the program goals and first-month activity, with attaining program goals during the last month.

Results: Participants were mostly male (81.5%), aged 58 ± 11 , enrolled in the main after a myocardial infarction or coronary interventions. Patients exercised aerobically for 183 min each week, 101 min (55% of total exercise) at the target heart rate. There was a significant improvement in exercise capacity, assessed by stress tests, metabolic equivalents which increased from 9.5 ± 3 to 11.4 ± 7 ($p < 0.001$). Independent predictors of RCRP goals were older age and more minutes of aerobic exercise during the first program month ($p < 0.05$).

Conclusion: Participants succeeded in performing guideline recommendations, resulting in a significant improvement in exercise capacity. Older age and increased volume of first month of exercise were significant factors associated with a greater likelihood to attain program goals.

Keywords

Remote cardiac rehabilitation, mobile application, adherence, telemedicine

Submission date: 19 December 2022; Acceptance date: 22 May 2023

Introduction

Cardiac rehabilitation (CR) is considered an essential component in the prevention of future heart episodes, re-hospitalizations, and death from all cardiac-related causes.^{1,2} Despite the documented evidence of the benefits of CR, less than one-third of patients actually begin rehabilitation.³ Of these, approximately 24% to 50% of patients terminate the program ahead of time and are therefore unable to achieve the desired outcome.⁴ The reasons for nonparticipation in rehabilitation programs are diverse. These include objective barriers (such as the

¹Faculty of Medicine, Tel Aviv University, Tel Aviv, Israel

²Cardiac Prevention and Rehabilitation Institute, Leviev Heart Center, Sheba Medical Center, Tel HaShomer, Israel

³Department of Biomedical Engineering, Faculty of Engineering, Sylvan Adams Sports Institute, School for Public Health, Sackler Faculty of Medicine, Tel Aviv University, Tel Aviv, Israel

⁴Neufeld Cardiac Research Institute, Sheba Medical Center, Tel HaShomer, Israel

Corresponding author:

Irene Nabutovsky, Cardiac Prevention and Rehabilitation Institute, Leviev Heart Center, Sheba Medical Center, Tel HaShomer, Israel 5265601.
Email: irene.n@sheba.health.gov.il



remoteness of the medical center, lack of accessible transportation, travel cost, unsuitable work hours, and socioeconomic factors), as well as behavioral and psychosocial reasons.^{5,6} Remote cardiac rehabilitation programs (RCRP) are typically performed in a patient's environment, using modern telecommunications technology tailored to the characteristics and needs of the individual patient. Thus, RCRP not only overcomes the many aforementioned barriers of traditional CR, but also achieves all the clinical goals set by traditional rehabilitation while simultaneously improving exercise adherence, physical activity levels, as well as reducing the relative cost of treatment.^{7,8}

In previous studies, the successful use of features provided by the RCRP platform in the early stages of the program resulted in better compliance and adherence.^{9,10} However, we found no mention of how patients' patterns of workouts and behavior in the first weeks of remote CR affected their subsequent adherence. The purpose of this study was to describe the achievements of the first RCR group within the Israeli Health system and to shed light on how potentially influential the first weeks of participation in the completion of the 3-month program.

Methods

RCRP

The first Israeli RCRP was developed at the Sheba Medical Center. It was approved by the Israeli Ministry of Health as a 3-month rehabilitation program for low-risk patients and is fully covered by the basic health insurance at no cost to patients. Low-risk definition according to the national guidelines is defined as lack of >5% ischemia per stress or pharmacological single-photon emission computed tomography (SPECT), left ventricular ejection fraction (LVEF) 50% or more, lack of sustained ventricular tachycardia, symptomatic atrial fibrillation or flutter or heart rates (HRs) > 120 beats per minute at rest. Low-risk definition excludes patients after cardiac arrest, heart failure with preserved systolic function and low functional capacity (metabolic equivalents (METs) < 6). A detailed description of our RCRP was published earlier.¹¹ Briefly, the program combined the use of advanced telecare technologies (platform by Datos Health) and smart wearable devices (mostly Garmin and Apple smartwatches) (Supplemental Appendix A- C) adapted to the characteristics and needs of each particular patient.

The program was geared to motivate the patient to perform workouts at home or in any convenient community facility, to lead a healthy lifestyle, as well as manage their clinical and psychological health. Necessary conditions for RCRP success included both motivational and technological aspects. For example, in addition to receiving automatic encouraging messages and reminders, the patient received a personal weekly call from a CR trainer to

discuss the training program and any problems that may have arisen. The patient received necessary technical and clinical support from a multiprofessional care team (physician, psychologist, physiologist, nurse).

Throughout the 18 months of the program, we collected data on behavior patterns, training, and goal achievements from the first low-risk group of 306 patients, 81.5% men, aged 57.59 (± 10.62) rehabilitating under the RCR at Sheba Medical Center in Israel. Basic characteristics, including a complete medical history, risk factors, and laboratory tests were collected. Training patterns were obtained via the smartwatch and then analyzed prospectively by the platform for 24 weeks. Objective improvement of an individual's functional capacity was assessed as the change between the first (2–3 weeks before starting rehabilitation) and the second (post 3 months of rehabilitation) exercise stress test (ergometry) using the same protocol and environment. Satisfaction with the program and care received was recorded and assessed using a questionnaire.

The main objective was to gauge the effectiveness of the RCRP protocol for any changes in exercise capacity assessed by treadmill exercise stress test using Bruce or modified Bruce protocol based on patient baseline functional capacity (history and impression) as assessed by the exercise physiologist. The same protocol was used for the second stress test after the 3-month program. The relative assessment of patients' adherence and achievement of the main "exercise goals" of the rehabilitation program, was also recorded. The "exercise goals" of the program were defined as the: (a) training performance of at least 150 aerobic minutes per week, of which; (b) 120 min in the target HR, set by the exercise physiologist; (c) two strength training sessions per week (after the first month); and (d) at least 8000 steps per day. And 100% achievement of these goals was considered satisfactory adherence. The individual target HR was determined in accordance with the recommendations of the national cardiac rehabilitation guidelines, based on the stress test results of each patient.

The study's second objective was to describe the differences in the baseline characteristics of patients that attained program goals versus the ones who did not. We conducted a retrospective analysis and comparison dividing all patients into two groups: those who achieved the main "exercise goals" of RCR and those who did not. The "exercise goals" for this comparison were specified as: (a) completion of the full 3-month program; (b) achievement of 600 aerobic minutes in the third month of the program (150 min weekly times 4 weeks); (c) achievement of 400 min in the target HR (120 min weekly times 4 weeks) in the third month.

The third objective was to identify potential independent predictors to better achieve the aforesaid exercise goals. We aimed to assess whether variables such as age, gender, baseline exercise capacity, indication for rehabilitation, use of the program's mobile app, as well as attaining goals (150

aerobic minutes per week and 120 min in the target HR) in the first month of the program, could serve as predictors for better future adherence during the third and final month of the program.

Evaluation of the patient's exercise performance was evaluated using the following exercise parameters:

1. The total number of weekly minutes of aerobic exercise (aerobic minutes).
2. The number of weekly aerobic minutes in the target HR.
3. The weekly percentage of training sessions at the target HR (preferably at least 70% of the total training time).
4. The weekly workout intensity (preferably at least 70% of the patient's individual maximum aerobic capacity, based on the stress test).
5. The assessment of perceived exertion on the Borg scale (patients were instructed to train within 12–14 moderate to vigorous on the scale).
6. Number of daily steps.
7. The number of weekly entries to the RCRP mobile phone application (patients were instructed to use the mobile app at least once every 1–2 days).

Statistical methods

Baseline characteristics are presented as median, mean \pm standard deviation (SD), or percentages as appropriate and comparisons were performed according to data type and its distribution. Characteristics of patients according to goals reached or not reached were compared using chi-square test for categorical data, and two-sided student t-test or Mann-Whitney tests for continuous data, as appropriate.

A paired sample t-test or Wilcoxon signed-rank test were used, according to the normality verification, in order to assess the differences between numeric values pre-RCRP and post-RCRP.

The effect size of the Delta between the METs pre-RCR and post-RCR were calculated as the difference of the METs means divided by the square root of the mean of the METs SD square.

A logistic regression model was constructed using the best subset method in order to determine independent predictors for the completion of the two program exercise goals during the third month of the program. The first goal was to reach 600 or more total aerobic minutes (any HR) per month, whereas the second goal was to reach 400 or more activity minutes in the individually designated target HR zone per month. The following covariables were introduced in the multivariate model using the best subset method: age, gender, pre-rehabilitation METs, and indication for CR. Predictions were based on patients' performance in the first month and some demographic data. Age and gender were taken as base features of all the models. For the predictive models, all the patients that dropped out of the program during the first month were excluded.

A p value < 0.05 was considered statistically significant and all tests were two sided. Statistical analyses were performed using the statistical software R foundation version 4.1.2.

Results

The study included data from the first 306 consecutive patients to complete RCRP. The patients joined the RCRP within 1 to 2 months after discharge from the hospital. Characteristics of the study group are summarized in Table 1.

Another comparison, shown in Supplemental Appendix D, presents the characteristics of the study group, 306 patients that participated in the RCRP, with the characteristics of a group of patients that completed a traditional center-based CR program (CCRP).

A total of 26% of this group were over 65 years of age. The main indications for CR were myocardial infarction (49%), percutaneous coronary interventions (48%) and coronary bypass surgery (12.5%). Participants had a preserved or normal systolic function and no high-risk criteria such as significant ischemia, effort angina, clinically significant ventricular arrhythmia, heart failure, or any evidence of clinical instability (mandatory national criteria to the RCRP entry). Participant adherence in the program, from the RCRP group, was 3.6 months on average, compared to CCRP participants that reached an average of 3.7 months in the rehabilitation program (Supplemental Appendix E).

The changes in exercise capacity

Following the RCR intervention, there was a significant and clinically meaningful improvement in exercise capacity, expressed by change in METs, assessed by pre-exercise and post-exercise stress tests using the same exercise protocol (Figure 1). Stress tests were performed by the same team of physiologists using the same treadmills, under the same conditions. The effect size of the exercise capacity improvement was 0.35.

RCRP exercise goals and patient performance

Overall, the total weekly minutes of aerobic training for 24 weeks was 183 min (median) for the entire population. Upon further analysis, the elderly group (age ≥ 65) achieved 208 min whereas the younger group achieved 186 min ($p < 0.01$). The median number of minutes of total aerobic minutes for 6 months was 183 min per week, which was 122% of the goal set by the physiologists. Additionally, time spent at the target HR was 101 min per week, which was only 84% of the 120 min exercise goal, usually divided into 4 exercise sessions. Figure 2 presents the median weekly aerobic activity of RCRP participants.

Exercise at the target HR was 55% of the total exercise time, which was less than our predefined goal of 70%, representing an intensity of 75% of the patient's

Table 1. Baseline characteristics of study population.

Variables	Mean (SE) or percentage (n)
Total population, n	306
Mean age (year) (SD, range)	57.7 (\pm 9.82)
Male	81.5% (n = 229)
Mean age group < 65 (years)	53.01 (\pm 8.48)
Mean age group \geq 65 (years)	70.16 (\pm 3.38)
Co-morbidities	
Metabolic	
Dyslipidemia	30.2% (n = 85)
Hypertension	25.6% (n = 72)
Diabetes mellitus	6% (n = 17)
Cardiovascular	
Myocardial infarction	49.1% (n = 138)
Atrial fibrillation/flutter	10.3% (n = 29)
Status postcoronary artery bypass graft	12.5% (n = 35)
Status postpercutaneous coronary intervention	48.8% (n = 137)
Status postvalve surgery	13.5% (n = 35)
Physical and functional status	
	Mean
BMI (kg/m ²)	27.76 (\pm 11.77)
Rest HR (bpm)	73 (\pm 12.3)
Maximum HR (baseline stress test)	140 (\pm 19.66)
Systolic blood pressure (mmHg)	131.18 (\pm 20.13)
Diastolic blood pressure (mmHg)	75.71 (\pm 12.12)
Maximum pre-METs (kcal/kg/min) baseline	9.49 (\pm 2.88)
Maximum pre-METs age group \geq 65	8.11 (\pm 2.8)
Maximum pre-METs age group < 65	9.98 (\pm 2.76)
Pre mean HR	121.45 (\pm 14.10)

BMI: body mass index; HR: heart rate; MET: maximal metabolic equivalent; SD: standard deviation; SE: standard error.




Figure 1. Changes in metabolic equivalents (METs) pre-RCRP and post-RCRP intervention compared to traditional CCRP. CCRP: center-based cardiac rehabilitation program; METs: metabolic equivalents; RCRP: Remote Cardiac Rehabilitation Program.

personalized maximum aerobic capacity (all median values). The subjective perception of effort (Borg scale) was 11.7 and the daily step count was 7700. Participants used the program's mobile application 3.8 times per week.

Patients' baseline characteristics and exercise goals attained

We evaluated the characteristics of those patients achieving the exercise goals set by the RCRP compared to those who did. Patients who achieved the exercise goals in the third month, were more likely to be older and have greater exercise activity during the first program month. Comparisons are summarized in Table 2.

Predicting exercise goals achievement during the third month of the program

Two main exercise goals were assessed using multivariate analysis: (1) would the patient achieve 600 min of monthly physical aerobic exercise and (2) would the patient achieve 400 min of monthly aerobic exercise at the target HR. Both outcomes were assessed during the third month of RCRP.

For every 100 min of training there was a 10% higher likelihood of achieving the 600 min aerobic goal during the third month. Similarly, for every 1000 steps the likelihood increased by 4% for the same outcome. Table 3 presents the results of the regression model.




Figure 2. The weekly aerobic exercise activity at target heart rate and below the target zone.

Among all the variables included in the multivariate model, only age and aerobic performance in the first weeks of RCRP were significant predictors, while variables such as gender, pre-rehabilitation METs, mobile phone application usage or workouts' average HR were not significant.

Discussion

The main findings of our study show that patients participating in RCRP can reach the programs' multiple exercise targets and improve their aerobic capacity. We also found that the patient's age and aerobic exercise volume during the program's first month are independently correlated with the program's goal achievement rate.

CR has been proved to be an underutilized service. It reduces cardiovascular mortality, hospital readmissions, and associated healthcare costs, while simultaneously improving health-related quality of life.¹² RCRP has also been demonstrated to be an effective and cost-efficient alternative to the traditional CR delivery model. It facilitates individualized support in almost any location.⁸

Overall, our program achieved its goals in terms of exercise adherence and was shown to significantly improve the functional capacity of its participants. These results are comparable with previous reports.¹³

In our study, we aimed to explore the differences in baseline characteristics between patients who managed to achieve the main goals of the RCRP and those who did not. We found that individuals who were deemed to be more successful by the end of the third month of the program tended to achieve greater aerobic performance in

the first month. The main predictors of achieving the RCRP goals during the third month include the age, and the number of aerobic minutes in the first month as well as steps. Accordingly, we can cautiously conclude that increased commitment to the program goals at the beginning of the rehabilitation is associated with better program goal completion. Importantly, this permits early identification of patients less likely to attain goals.

Despite ongoing research in this field, many studies have not documented a similar association between early aerobic minutes and RCRP goal completion. Initial active engagement in RCRP could be a significant predictor for future goal completion as well as attaining desired level of cardiovascular health. Several studies have shown that the timely start of rehabilitation after a cardiovascular event leads to better health outcomes for patients.¹⁴ Furthermore, other studies have even gone as far to say that delays in starting CR can cause avoidable harm in the long-term health of patients.¹⁵ This shows the already well documented importance of the timely start of CR programs.

However, just informing patients to join CR early after their cardiovascular event is insufficient on the part of the physician. Our results indicate that patients who demonstrated early active participation in the program (those who achieved greater aerobic minutes) better achieved the program's adherence goals. These results indicate that physicians should motivate patients early on in their remote-CR progress, as we found that the first few weeks are a crucial period for determining their future success in completing program goals. Younger patients could require special consideration since apparently it appears to be more difficult for them than for the elderly to fulfill all the requirements of the

Table 2. Observed differences between the baseline characteristics according to the RCRP “exercise goals.”

RCRP ‘exercise goals’	Baseline characteristics	Comparison groups (mean ± SD, %)		p Value
		Not reached	Reached	
Completed the full 3 months of RCRP		N = 108	N = 198	
	Age	55.77 ± 11.64	58.56 ± 9.92	0.011
	Weight	85.15 ± 17.45	81.50 ± 15.01	0.057
	METs	8.93 ± 2.84	9.79 ± 2.86	0.030
	Reached the aerobic goal (150 aerobic minutes per week) in the first month of the RCRP (%)	55 (n = 59)	75 (n = 148)	0.002
400 or more minutes of aerobic exercise attained in the third month of RCRP		N = 152	N = 46	
	Age	56.92 ± 10.25	63.75 ± 6.60	0.001
	Systolic BP	129.19 ± 17.87	136.89 ± 17.11	0.052
	Glucose	116.60 ± 24.81	134.13 ± 17.11	0.023
	Urea	36.37 ± 8.97	44.60 ± 24.56	0.031
	Hemoglobin	13.62 ± 1.74	12.31 ± 2.61	0.041
	HR at maximum effort	142.05 ± 19.16	135.81 ± 15.75	0.057
	Reached the aerobic goal (150 aerobic minutes per week) in the first month of the RCRP (%)	69 (n = 104)	95 (n = 44)	0.001
600 or more minutes of aerobic exercise attained in the third month of RCRP		N = 57	N = 141	
	Age	55.08 ± 11.0	59.95 ± 9.14	0.001
	Systolic BP	125.39 ± 19.22	133.82 ± 16.83	0.021
	Aerobic in target HR goal reached (120 min per week) in the first month of RCRP (%)	7 (n = 4)	25 (n = 35)	0.004

BP: blood pressure; HR: heart rate; MET: maximal metabolic equivalent; RCRP: Remote Cardiac Rehabilitation Program; SD: standard deviation.

RCRP.^{16,17} This can be explained both by the lack of time and by a less serious perception of their illness. We speculate that motivation was an important factor for early active participation in the program. Patients who achieved greater total aerobic minutes early on in the program could have had certain motivation that resulted in their better completing program goals. Another relevant CR study demonstrated that motivational programs appear helpful in increasing participation in early outpatient cardiac rehabilitation.¹⁸ Whether patients in our study who achieved greater total aerobic

minutes early on in the program were or were not motivated intrinsically or extrinsically is debatable, but further studies need to be performed in the future to determine which plays a bigger role in CR program completion.

However, it should be noted that, despite a good indicator of the total number of aerobic minutes, their aerobic exercise intensity was lower than desired: the percentage of time that patients spent in the target HR during the training was significantly lower than we expected. Also, the subjectivity when assessing training intensity was questionable.

Table 3. Predictive models for achieving “exercise goals” during the third month of RCRP.

Logistic regression model for $400 \geq \text{min of exercise in target HR during the third month of the program}$			
Variables	OR	CI	p Value
Age (per year)	1.09	1.03–1.15	0.001
Mean workout intensity (first month, per 10% of the HR reserve)	1.06	1.01–1.11	0.032
Total aerobic minutes (first month, per 100 min)	1.10	1.03–1.18	0.002
Logistic regression model for $\geq 600 \text{ min of exercise during the third month of the program}$			
Variables	OR	CI	p Value
Age (per year)	1.07	1.03–1.13	0.001
Total steps (first month, per 1000 steps)	1.04	1.00–1.08	0.034

CI: confidence interval; HR: heart rate; OR: odds ratio; RCRP: Remote Cardiac Rehabilitation Program.

In other words, patients who trained a lot, did not always do so effectively. Although the overall increase in functional capacity (METs) was good, it seems that patients were capable of more.

In addition, we found that older age was also associated with better achievement in all RCRP goals and was a predictor of adherence to these goals refuting the generally accepted opinion that digital programs are less suitable for the elderly.^{19,20}

The applicability of the research findings and limitations

The results of a study can have significant applicability in the healthcare industry. RCRP utilizes telemedicine and digital health technologies to provide patients with cardiovascular disease rehabilitation services in their homes, without the need for in-person visits to healthcare facilities.

The study results can be used to inform the development of new remote cardio-rehabilitation programs, as well as the refinement of existing ones. Healthcare providers and policymakers can use the study’s findings to design and implement effective remote cardio-rehabilitation programs to improve patient outcomes and reduce healthcare costs. In particular, we propose to focus efforts on better engagement with younger patients, as well as on better compliance with the program in the first weeks.

Additionally, the results can help clinicians to optimize their use of digital health technologies and telemedicine platforms to provide high-quality care to patients with cardiovascular disease and manage expectations regarding activity levels and change in functional capacity. Overall, the study results can contribute to the development of more patient-centered, accessible, and cost-effective models of care delivery that can improve health outcomes of individuals with cardiovascular disease.

Our study is not without limitations. This research is based on a single center experience and thus variations are to be expected. The study population enrolled was defined as low risk per national guidelines thus results cannot be extrapolated to all patients with cardiovascular conditions. Additionally, patients in our study did not work out as intensely as the program indicated (the average Borg was less than 12–14). Patients generally coped well with goals, however, we would have liked to see more intense training, such as more time spent in the target HR and a higher Borg scale. Furthermore, patients did not achieve goals that were related to their strength exercises (resistance training). This is partly due to a lack of time, patient preferences, and the instructions of medical staff who emphasized the value of aerobic exercise for cardiovascular health. This aspect has been identified in other similar RCRPs. An additional limitation worth mentioning is that patients in the study were mostly male (81.5%), making it difficult to extrapolate our results. In addition, the patients who participated in this study represent a low-risk group, which means that at baseline they already had good functional capacity and risk factors of cardiovascular disease were mostly within guideline recommended ranges. Because this is an observational study we can report associations and not address cautions. It is also important to note that subtle differences in the execution of the stress tests, and other less definable variables may have impacted the observed change (e.g., higher confidence of the patient and the care team itself in the second stress test).

Conclusion

To the best of our knowledge, there have been no previous attempts to understand how the patient’s behavior at the beginning of the RCRP could serve as a predictor of future program success. Understanding this information is instrumental in building more accurate, patient-oriented intervention programs in the future. Additionally, availability of this information at the outset will help us recognize certain characteristics which could contribute to or hinder their success in the program. The RCRP care team should pay special attention to the adherence and performance of physical activity in the first weeks of RCRP, because it is correlated with subsequent successful completion of the program.

Acknowledgements: The authors would like to thank the ARC Innovation Center and Medical Center beyond for their support.

Author contributions: IN led, designed, and conducted the study. DB, AH, and YL drafted and reviewed the manuscript. MM and YK collected and prepared the data alongside its statistical analysis. MS and IN interpreted the study results. RK oversaw the study and was involved in the study design, drafting, and reviewing. All authors approved the final version of the manuscript.

Declaration of conflicting interests: The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

Ethics approval: The present retrospective study was conducted according to the IRB approval instructions and GCP principles. Sheba Medical Center, approval ID: 1553-14-SMC.

Funding: The author(s) received no financial support for the research, authorship, and/or publication of this article.

Guarantor: IN.

Patient consent: Our study was conducted on de-identified data files extracted from the Sheba Medical Center data lake by hospitals approved personnel, according to the IRB approval granted (IRB approval number SMC 15-2699).

ORCID ID: Yoav Levine  <https://orcid.org/0000-0003-1198-6322>

Supplemental material: Supplemental material for this article is available online.

References

- Mampuya WM. Cardiac rehabilitation past, present and future: an overview. *Cardiovasc Diagn Ther* 2012 Mar; 2: 38.
- Dunlay SM, Pack QR, Thomas RJ, et al. Participation in cardiac rehabilitation, readmissions, and death after acute myocardial infarction. *Am J Med* 2014 Jun 1; 127: 538–546.
- Peters AE and Keeley EC. Trends and predictors of participation in cardiac rehabilitation following acute myocardial infarction: data from the behavioral risk factor surveillance system. *J Am Heart Assoc* 2017 Dec 29; 7: e007664.
- Turk-Adawi KI, Oldridge NB, Tarima SS, et al. Cardiac rehabilitation patient and organizational factors: what keeps patients in programs? *J Am Heart Assoc* 2013 Oct 21; 2: e000418.
- Bakhshayeh S, Sarbaz M, Kimiafar K, et al. Barriers to participation in center-based cardiac rehabilitation programs and patients' attitude toward home-based cardiac rehabilitation programs. *Physiother Theory Pract* 2019 Jun 3; 37: 158–168.
- Menezes AR, Lavie CJ, Milani RV, et al. Cardiac rehabilitation in the United States. *Prog Cardiovasc Dis* 2014 Mar 1; 56: 522–529.
- Rawstorn JC, Gant N, Direito A, et al. Telehealth exercise-based cardiac rehabilitation: a systematic review and meta-analysis. *Heart* 2016 Aug 1; 102: 1183–1192.
- Maddison R, Rawstorn JC, Stewart RA, et al. Effects and costs of real-time cardiac telerehabilitation: randomised controlled non-inferiority trial. *Heart* 2019 Jan 1; 105: 122–129.
- Varnfield M, Karunanithi M, Lee CK, et al. Smartphone-based home care model improved use of cardiac rehabilitation in post-myocardial infarction patients: results from a randomised controlled trial. *Heart* 2014 Nov 15; 100: 1770–1779.
- Whittaker F and Wade V. The costs and benefits of technology-enabled, home-based cardiac rehabilitation measured in a randomised controlled trial. *J Telemed Telecare* 2014 Oct; 20: 419–422.
- Nabutovsky I, Ashri S, Nachshon A, et al. Feasibility, safety, and effectiveness of a mobile application in cardiac rehabilitation. *IMAJ* 2020 Jun 1; 22: 357–363.
- Krishnamurthi N, Schopfer DW, Ahi T, et al. Predictors of patient participation and completion of home-based cardiac rehabilitation in the veterans health administration for patients with coronary heart disease. *Am J Cardiol* 2019 Jan 1; 123: 19–24.
- Regan EW, Handlery R, Stewart JC, et al. Integrating survivors of stroke into exercise-based cardiac rehabilitation improves endurance and functional strength. *J Am Heart Assoc* 2021; 10.
- Anderson L, Nguyen TT, Dall CH, et al. Exercise-based cardiac rehabilitation in heart transplant recipients. *Cochrane Database Syst Rev* 2017; 4.
- Hinde S, Harrison A, Bojke L, et al. Quantifying the impact of delayed delivery of cardiac rehabilitation on patients' health. *Eur J Prev Cardiol* 2020 Nov 1; 27: 1775–1781.
- Zhang L, Sobolev M, Piña IL, et al. Predictors of cardiac rehabilitation initiation and adherence in a multiracial urban population. *J Cardiopulm Rehabil Prev* 2017 Jan; 37: 30–38.
- Gaalema DE, Savage PD, Rengo JL, et al. Patient characteristics predictive of cardiac rehabilitation adherence. *J Cardiopulm Rehabil Prev* 2017 Mar; 37: 103–110.
- Pack QR, Johnson LL, Barr LM, et al. Improving cardiac rehabilitation attendance and completion through quality improvement activities and a motivational program. *J Cardiopulm Rehabil Prev* 2013 May; 33: 153.
- Magdalena M, Bujnowska-Fedak UG and Grata-Borkowska U. Use of telemedicine-based care for the aging and elderly: promises and pitfalls. *Smart Homecare Technol Telehealth* 2015 May; 3: 91–105.
- Smith A. *Older adults and technology use*. Pew Research Center, USA: Internet, Science & Tech, Pew Research Center, www.pewresearch.org/internet/2014/04/03/older-adults-and-technology-use/ (2013, accessed 14 March 2022).