

Meeting abstract

Open Access

Glucocorticoids regulate the human occludin gene through a single imperfect palindromic glucocorticoid response element

M Burek^{*1}, N Harke², J Leers³, S Kietz⁴, D Drenckhahn² and C Förster¹

Address: ¹University of Wuerzburg, Department of Anaesthesiology, Zentrum of Operative Medicine, Germany, ²University of Wuerzburg, Institute of Anatomy and Cell Biology, Würzburg, Germany, ³University of Giessen, Institute of Genetics, Giessen, Germany and ⁴University of Goettingen, Institute of Pediatrics, Göttingen, Germany

* Corresponding author

from 12th Joint Meeting of the Signal Transduction Society (STS). Signal Transduction: Receptors, Mediators and Genes Weimar, Germany. 29–31 October 2008

Published: 26 February 2009

Cell Communication and Signaling 2009, **7**(Suppl 1):A96 doi:10.1186/1478-811X-7-S1-A96

This abstract is available from: <http://www.biosignaling.com/content/7/S1/A96>

© 2009 Burek et al; licensee BioMed Central Ltd.

The 65 kDa protein occludin is an essential element of the blood-brain barrier. This integral membrane protein represents an important part of the tight junctions, which seal and protect the blood brain barrier against paracellular diffusion of solutes to the brain parenchyme and are therefore responsible for the high resistance and low permeability between cerebral capillary endothelial cells. However, the molecular basis for the regulation of occludin gene expression is only incompletely understood. In former projects we showed that treatment of a brain microvascular cell line, cEND, with glucocorticoids resulted in increased occludin expression in cell-cell-contacts. Induction of occludin expression by glucocorticoids was shown to be dependent on the glucocorticoid receptor. This study aims to identify the underlying molecular mechanism of gene expression and to identify potential glucocorticoid receptor binding sites within the occludin promoter, the glucocorticoid response elements (GRE). We identified one candidate GRE within the distal part of the occludin promoter that differs from the consensus GRE by the presence of a 4-basepair instead of a 3-basepair spacer between two highly degenerate halvesites (5'-ACATGTGTTTACAAAT-3'). Chromatin immunoprecipitation assay and site-directed mutagenesis confirmed binding of the glucocorticoid receptor to this site. We synthesized plasmid containing four copies of this GRE in the vector pGL-3 basic. After stimulation of the cells with hydrocortisone we could observe up to 8-fold increased activity in luciferase reporter assay. The need for glucocorticoid receptor dimerization to induce gene expression

was further confirmed by transfection studies using wild type and glucocorticoid receptor dimerization-deficient expression vectors, indicating that transactivation of occludin occur through the GRE.