

Short paper

Open Access

Feeding oxidized fat during pregnancy up-regulates expression of PPAR α -responsive genes in the liver of rat fetuses

Robert Ringseis, Anke Gutgesell, Corinna Dathe, Corinna Brandsch and Klaus Eder*

Address: Institut für Agrar- und Ernährungswissenschaften, Martin-Luther-Universität Halle-Wittenberg, Emil-Abderhalden-Straße 26, D-06108 Halle/Saale, Germany

Email: Robert Ringseis - robert.ringseis@landw.uni-halle.de; Anke Gutgesell - anke.gutgesell@landw.uni-halle.de; Corinna Dathe - corinna.dathe@landw.uni-halle.de; Corinna Brandsch - corinna.brandsch@landw.uni-halle.de; Klaus Eder* - klaus.eder@landw.uni-halle.de

* Corresponding author

Published: 12 March 2007

Received: 19 December 2006

Lipids in Health and Disease 2007, **6**:6 doi:10.1186/1476-511X-6-6

Accepted: 12 March 2007

This article is available from: <http://www.lipidworld.com/content/6/1/6>

© 2007 Ringseis et al; licensee BioMed Central Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

Background: Feeding oxidized fats causes activation of peroxisome proliferator-activated receptor α (PPAR α) in the liver of rats. However, whether feeding oxidized fat during pregnancy also results in activation of PPAR α in fetal liver is unknown. Thus, this study aimed to explore whether feeding oxidized fat during pregnancy causes a PPAR α response in fetal liver. Two experiments with pregnant rats which were administered three different diets (control; oxidized fat; clofibrate as positive control) in a controlled feeding regimen during either late pregnancy (first experiment) or whole pregnancy (second experiment) were performed.

Results: In both experiments pregnant rats treated with oxidized fat or clofibrate had higher relative mRNA concentrations of the PPAR α -responsive genes acyl-CoA oxidase (ACO), cytochrome P₄₅₀ 4A1 (CYP4A1), L-type carnitin-palmitoyl transferase I (L-CPT I), medium-chain acyl-CoA dehydrogenase (MCAD), and long-chain acyl-CoA dehydrogenase (LCAD) in the liver than control rats ($P < 0.05$). In addition, in both experiments fetuses of the oxidized fat group and the clofibrate group also had markedly higher relative mRNA concentrations of ACO, CYP4A1, CPT I, MCAD, and LCAD in the liver than those of the control group ($P < 0.05$), whereas the relative mRNA concentrations of PPAR α , SREBP-1c, and FAS did not differ between treatment groups. In the second experiment treatment with oxidized fat also reduced triacylglycerol concentrations in the livers of pregnant rats and fetuses ($P < 0.05$).

Conclusion: The present study demonstrates for the first time that components of oxidized fat with PPAR α activating potential are able to induce a PPAR α response in the liver of fetuses. Moreover, the present study shows that feeding oxidized fat during whole pregnancy, but not during late pregnancy, lowers triacylglycerol concentrations in fetal livers.

Background

In recent years the contribution of dietary oxidized fats to total energy intake has markedly increased in industrial-

ized countries, mainly, due to the increasing consumption of fast food which contain significant amounts of these heated and processed fats. We and others have shown that

oxidized fats are strong activators of peroxisome proliferator-activated receptor α (PPAR α) [1-3], a ligand-activated transcription factor that belongs to the family of nuclear receptors. PPAR α is centrally involved in the regulation of lipid homeostasis in the liver and is essential for normal liver function [4]. Upon activation of PPAR α by a ligand, which includes the fibrate class of hypolipidemic drugs such as clofibrate or fenofibrate, fatty acids, and eicosanoids, the transcription of genes containing a PPAR response element (PPRE) in its promoter region is induced. Typical PPAR α -responsive genes in the liver, which are also up-regulated by oxidized fats and its components such as oxidized fatty acids (e.g. hydroperoxy-fatty acids, cyclic fatty acids) [1-3,5,6], include a wide array of genes that are involved in peroxisomal and mitochondrial fatty acid β -oxidation such as L-type carnitine-palmitoyl transferase I (L-CPT I), acyl-CoA oxidase (ACO), cytochrome P₄₅₀ 4A1 (CYP4A1), medium-chain acyl-CoA dehydrogenase (MCAD), and long-chain acyl-CoA dehydrogenase (LCAD) [7,8]. PPAR α activation by oxidized fat was also shown to cause liver enlargement due to peroxisome proliferation and reduce triacylglycerol concentrations in liver, plasma, and very low-density lipoproteins in rats [3,9,10]. In part, the triacylglycerol-lowering effect of oxidized fat in the rat is presumably also due to the observed reduction in mRNA concentrations and activities of lipogenic enzymes such as fatty acid synthase (FAS) [9].

During pregnancy treatment of rats with the pharmacological PPAR α activator clofibrate has been shown to cause proliferation of peroxisomes [11,12], induction of peroxisomal enzyme activities [13], and induction of CYP4A1 in fetal liver [14] indicating that clofibrate is capable of activating PPAR α transplacentally. In addition, pathological changes in newborn rats born to mothers treated with clofibrate during pregnancy [15], and an impaired fetal growth of fenofibrate-treated pregnant rats have been reported [16]. Moreover, in rat and mouse liver epithelial cells treatment with the PPAR α agonist WY-14,643 caused up-regulation of proto-oncogenes [17-19], which has been attributed to the hepatocarcinogenic effect of peroxisome proliferators in rodents [18]. However, whether feeding oxidized fat to pregnant rats also results in activation of PPAR α and up-regulation of PPAR α target genes in fetal livers is unknown from the literature. Moreover, it is unknown whether components of oxidized fats such as oxidized fatty acids are able to substantially pass the placenta and enter the fetus, because it has been shown that the transplacental transport of fatty acids from the maternal diet is highly selective for individual fatty acids, e.g. long-chain polyunsaturated fatty acids such as docosahexaenoic acid and arachidonic acid are preferentially transported through the placenta at the expense of other less important fatty acids [20-22].

Therefore, the present study aimed to explore whether feeding oxidized fat during pregnancy causes a PPAR α response in fetal liver as estimated by the up-regulation of typical PPAR α -responsive genes such as ACO, L-CPT I, CYP4A1, MCAD, and LCAD and whether the induction of fatty acid catabolism might also affect fetal hepatic triacylglycerol concentrations. We also analyzed the mRNA abundance of the lipogenic transcription factor sterol regulatory-element binding protein (SREBP)-1c and its target gene FAS, because in previous studies administration of oxidized fats has also been demonstrated to reduce mRNA expression of lipogenic enzymes [9]. To address possible adverse effects of treatment with oxidized fat, we also determined the mRNA abundance of the proto-oncogenes c-myc, c-jun, and c-fos.

Since the duration of administration of oxidized fat during pregnancy might also influence the effect on lipid metabolism in the fetal liver, we performed two experiments which varied in the duration of administration of the oxidized fat. In the first experiment we investigated the effect of short-term administration (last 5 d of pregnancy) of oxidized fat on the PPAR α response in maternal and fetal liver, whereas in the second experiment the effect of long-term administration (whole pregnancy) was studied.

Results

Fatty acid composition and concentrations of lipid peroxidation products in the experimental fats

The oxidized fat in the short-term experiment had lower proportions of polyunsaturated fatty acids (C18:2) but higher proportions of saturated (C16:0, C18:0) and monounsaturated (C18:1) fatty acids compared to the control fat due to the heat treatment of sunflower oil during preparation of the oxidized fat (Table 1). In the long-term experiment the proportions of C18:2 and C18:1 in the oxidized fat were similar to those in the control fat due to adjustment of fatty acid composition of the control fat, whereas the proportions of saturated fatty acids were lower in the oxidized fat compared to the control fat (Table 2). In both experiments the oxidized fat had higher concentrations of peroxides, TBARS, and conjugated dienes, and a higher acid value than the fat of the control group. The concentrations of lipid peroxidation products (peroxide value and conjugated dienes) in the control fat were higher in the long-term experiment than in the short-term experiment. This is probably due to the fact that fats in the long-term experiment were included into a semi-synthetic diet which contained pro-oxidant minerals such as iron or copper whereas fats in the short-term experiment were administered the rats directly by gastric tube.

Table 1: Characteristics of the experimental fats of the short-term experiment

Treatment group Fat	Control	Clofibrate SFO ²	Oxidized fat ¹ oxidized SFO ²
Major fatty acids, g/100 g total fatty acids			
C16:0		6.1	9.1
C18:0		3.4	5.3
C18:1 (n-9)		32.6	38.3
C18:2 (n-6)		56.6	44.7
Peroxidation products			
Peroxide value, mEq O ₂ /kg		3.0	379
Acid value, g KOH/kg		0.4	5.8
Conjugated dienes, mmol/kg		< 0.1	274
TBARS ² , mmol/kg		1.1	13.1

¹Prepared by heating at a temperature of 60°C for 25 d.

²Abbreviations: SFO, sunflower oil; TBARS, thiobarbituric acid-reactive substances.

Final body weights and relative liver weights of pregnant rats

Body weight development was not affected by dietary treatment in both experiments due to the controlled feeding regimen applied. In the short-term experiment, final body weights of pregnant rats at d 21 of pregnancy did not differ between the three treatment groups (control, 300 ± 28 g; oxidized fat, 299 ± 24 g; clofibrate, 307 ± 17 g, mean ± SD). Pregnant rats treated with clofibrate or oxidized fat had higher relative liver weights than control rats ($P < 0.05$; control, 3.6 ± 0.2 g/100 g body weight; oxidized fat, 4.4 ± 0.3 g/100 g body weight; clofibrate, 4.9 ± 0.2 g/100 g body weight).

In the long-term experiment, final body weights of pregnant rats at d 21 of pregnancy were also not different between treatment groups (control, 342 ± 35 g; oxidized fat, 331 ± 40 g; clofibrate, 328 ± 18 g). Relative liver weights of pregnant rats treated with clofibrate or oxidized fat were also higher than those of control rats ($P < 0.05$; control, 3.4 ± 0.2 g/100 g body weight; oxidized fat, 4.4 ±

0.5 g/100 g body weight; clofibrate, 4.5 ± 0.3 g/100 g body weight).

Relative mRNA concentrations of PPAR α and PPAR α -responsive genes in livers of pregnant rats and fetuses

In the short-term experiment, relative mRNA concentrations of PPAR α in the liver of pregnant rats and fetuses were not different between treatment groups (pregnant rats: 1.00 ± 0.49, control group; 1.02 ± 0.48, oxidized fat group; 1.26 ± 0.22, clofibrate group; $P = 0.439$; fetuses: 1.00 ± 0.29, control group; 1.29 ± 0.50, oxidized fat group; 1.15 ± 0.74, clofibrate group; $P = 0.653$). Pregnant rats treated with oxidized fat had 2.4-, 3.0-, 2.5- and 2.1-fold higher relative mRNA concentrations of ACO, CYP4A1, MCAD, and LCAD, respectively, in the liver than control rats ($P < 0.05$; Fig. 1). The relative mRNA concentration of L-CPT I did not differ between both groups of rats. Treatment of pregnant rats with clofibrate resulted in 4.8-, 11-, 1.6-, 2.5- and 2.4-fold higher relative mRNA concentrations of ACO, CYP4A1, L-CPT I, MCAD, and LCAD, respectively, in the liver compared to control treatment ($P < 0.05$). Fetuses of the oxidized fat group had 6.3-

Table 2: Characteristics of the experimental fats of the long-term experiment

Treatment group Fat	Control	Clofibrate SFO ² : lard (54:46)	Oxidized fat oxidized SFO ²
Major fatty acids, g/100 g total fatty acids			
C16:0		14.9	10.2
C18:0		9.5	6.2
C18:1 (n-9)		28.9	34.4
C18:2 (n-6)		42.0	44.2
Peroxidation products			
Peroxide value, mEq O ₂ /kg		6.6	230
Conjugated dienes, mmol/kg		12.3	139
TBARS ² , mmol/kg		1.0	19

¹Prepared by heating at a temperature of 60°C for 25 d.

²Abbreviations: SFO, sunflower oil; TBARS, thiobarbituric acid-reactive substances.

, 9.0-, 6.4-, 1.5- and 2.1-fold higher relative mRNA concentrations of ACO, CYP4A1, L-CPT I, MCAD, and LCAD respectively, in the liver than those of the control group ($P < 0.05$), whereas fetuses of the clofibrate group had 20-, 51-, 12-, 2.8- and 3.0-fold higher relative mRNA concentration of ACO, CYP4A1, L-CPT I, MCAD, and LCAD, respectively in the liver than those of the control group ($P < 0.05$).

In the long-term experiment, relative mRNA concentrations of PPAR α in the liver of pregnant rats and fetuses also did not differ between treatment groups (pregnant rats: 1.00 ± 0.49 , control group; 1.56 ± 0.99 , oxidized fat group; 0.63 ± 0.29 , clofibrate group; $P = 0.053$; fetuses: 1.00 ± 0.70 , control group; 0.72 ± 0.20 , oxidized fat group; 0.75 ± 0.31 , clofibrate group; $P = 0.201$). Treatment of pregnant rats with oxidized fat resulted in 4.0-, 4.4-, 2.5-, 2.8- and 2.0-fold higher relative mRNA concentrations of ACO, CYP4A1, L-CPT I, MCAD, and LCAD, respectively, in the liver compared to control treatment ($P < 0.05$; Fig. 1). Pregnant rats treated with clofibrate had 4.8-, 8.3-, 2.0- and 1.9-fold higher relative mRNA concentrations of ACO, CYP4A1, MCAD, and LCAD, respectively, in the liver than control rats ($P < 0.05$), whereas the relative mRNA concentration of L-CPT I did not differ between both treatment groups. Fetuses of the oxidized fat group had 2.2-, 6.0-, 3.2- and 2.0-fold higher relative mRNA concentrations of ACO, CYP4A1, MCAD, and LCAD, respectively, in the liver than those of the control group ($P < 0.05$), whereas the relative mRNA concentration of L-CPT I did not differ between both treatment groups. Fetuses of the clofibrate group had 3.5-, 28-, 18- and 3.9-fold higher relative mRNA concentration of ACO, CYP4A1, L-CPT I, and LCAD, respectively, in the liver than those of the control group ($P < 0.05$), whereas the relative mRNA concentration of MCAD did not differ between both treatment groups.

Relative mRNA concentrations of SREBP-1c and FAS in livers of pregnant rats and fetuses

In the short-term experiment, treatment of pregnant rats with oxidized fat resulted in lower relative mRNA concentrations of FAS in the liver relative to control treatment ($P < 0.05$), whereas the relative mRNA concentration of SREBP-1c did not differ between these two groups (Fig. 2). Relative mRNA concentrations of SREBP-1c and FAS in the liver of pregnant rats treated with clofibrate did not differ from those of control rats. However, pregnant rats treated with clofibrate had higher relative mRNA concentrations of SREBP-1c in the liver than those fed oxidized fat ($P < 0.05$). In fetal livers relative mRNA concentrations of SREBP-1c and FAS did not differ between treatment groups.

In the long-term experiment, relative mRNA concentrations of SREBP-1c and FAS in the liver of pregnant rats and fetuses did not differ between the three treatment groups (Fig. 2).

Concentrations of triacylglycerols in livers of pregnant rats and fetuses

In the short-term experiment, concentrations of triacylglycerols in the liver of pregnant rats did not differ between the three treatment groups (Fig. 3). Fetuses of the clofibrate group had lower concentrations of triacylglycerols in the liver than those of the control and the oxidized fat group ($P < 0.05$); oxidized fat had no effect on the triacylglycerol concentration in fetal livers compared to control.

In the long-term experiment, treatment of pregnant rats with oxidized fat or clofibrate resulted in lower concentrations of triacylglycerols in the liver compared to control treatment ($P < 0.05$). Fetuses of the oxidized fat group and the clofibrate group also had lower concentrations of triacylglycerols in the liver than those of the control group ($P < 0.05$).


Relative mRNA concentrations of c-myc, c-jun, and c-fos in livers of pregnant rats and fetuses

In the short-term experiment, relative mRNA concentrations of c-myc, c-jun, and c-fos in the liver of pregnant rats and fetuses did not differ between the three treatment groups (Fig. 4).


In the long-term experiment, treatment of pregnant rats with oxidized fat resulted in higher relative mRNA concentrations of c-myc, c-jun, and c-fos relative to control treatment ($P < 0.05$). In fetal livers relative mRNA concentrations of c-myc, c-jun, and c-fos did not differ between treatment groups.

Discussion

The present study demonstrates for the first time that components of oxidized fat with PPAR α activating potential are able to induce a PPAR α response in the liver of fetuses. Moreover, the present study shows that feeding oxidized fat during whole pregnancy, but not during late pregnancy, lowers triacylglycerol concentrations in fetal livers. Hydroxy- and hydroperoxy-fatty acids such as hydroxyoctadecadienoic acid (HODE) and hydroperoxyoctadecadienoic acid (HPODE) occurring in oxidized fats are very potent PPAR α agonists [6,23,24]. These oxidized fatty acids are produced during the early stage of lipid peroxidation, and, due to their low thermodynamic stability, easily decompose at high temperatures. Thus, fats treated at low temperatures have markedly higher concentrations of these primary lipid peroxidation products than fats treated at high temperature [3]. Therefore, in order to provoke a significant PPAR α activating effect of the oxidized

**Figure 1**

Effects of treatment on relative mRNA concentrations of PPAR α -responsive genes in livers of pregnant rats and fetuses. Effect of short-term (d 16 – d 21 of pregnancy, left) and long-term (d 1 – d 21 of pregnancy, right) administration of three different diets (control, oxidized fat, clofibrate) during pregnancy on relative mRNA concentrations of ACO, CYP4A1, CPT I, MCAD, and LCAD in the liver of pregnant rats and fetuses at d 21 of pregnancy. Left, Bars represent mean \pm SD ($n = 9$ /group). Right, Bars represent mean \pm SD ($n = 12$ /group). Bars marked without a common superscript letter differ ($P < 0.05$). Results from one-way ANOVA (P -values): short-term experiment: ACO, $P = 0.0001$ (pregnant), $P = 0.001$ (fetus); CYP4A1, $P = 0.0001$ (pregnant), $P = 0.002$ (fetus); L-CPT I, $P = 0.013$ (pregnant), $P = 0.012$ (fetus); MCAD, $P = 0.042$ (pregnant), $P = 0.001$ (fetus); LCAD, $P = 0.001$ (pregnant), $P = 0.03$ (fetus); long-term experiment: ACO, $P = 0.0001$ (pregnant), $P = 0.015$ (fetus); CYP4A1, $P = 0.0001$ (pregnant), $P = 0.001$ (fetus); L-CPT I, $P = 0.016$ (pregnant), $P = 0.011$ (fetus); MCAD, $P = 0.001$ (pregnant), $P = 0.019$ (fetus); LCAD, $P = 0.001$ (pregnant), $P = 0.001$ (fetus).

**Figure 2**


Effects of treatment on relative mRNA concentrations of SREBP-1c and FAS in livers of pregnant rats and fetuses. Effect of short-term (d 16 – d 21 of pregnancy, left) and long-term (d 1 – d 21 of pregnancy, right) administration of three different diets (control, oxidized fat, clofibrate) during pregnancy on relative mRNA concentrations of SREBP-1c and FAS in the liver of pregnant rats and fetuses at d 21 of pregnancy. Left, Bars represent mean \pm SD ($n = 9$ /group). Right, Bars represent mean \pm SD ($n = 12$ /group). Bars marked without a common superscript letter differ ($P < 0.05$). Results from one-way ANOVA (P -values): short-term experiment: SREBP-1c, $P = 0.041$ (pregnant), $P = 0.058$ (fetus); FAS, $P = 0.026$ (pregnant), $P = 0.425$ (fetus); long-term experiment: SREBP-1c, $P = 0.526$ (pregnant), $P = 0.189$ (fetus); FAS, $P = 0.716$ (pregnant), $P = 0.334$ (fetus).

fat, we decided to use a fat treated at a relatively low temperature for a long period. Although we did not determine the concentrations of oxidized fatty acids such as 13-HODE or 13-HPODE in the oxidized fat, the high peroxide value and the high concentration of conjugated dienes indicate that the oxidized fats used in both experiments presumably had high concentrations of hydroxy- and hydroperoxy-fatty acids which may be particularly responsible for the PPAR α activating effect of oxidized fats. Nonetheless, other components of heated fats such as cyclic fatty acid monomers which also show a strong PPAR α response [5] might be also causative for PPAR α activation.

The lower concentrations of primary lipid peroxidation products (peroxides, conjugated dienes) in the oxidized

fat used in the long-term experiment compared to that used in the short-term experiment are probably explained by the fact that in the long-term experiment the lipid peroxidation products in the oxidized fat were determined after inclusion into the diet. Primary lipid peroxidation products easily decompose and are partially degraded in the presence of other diet components such as metal ions (e.g. iron, copper) acting as catalysts.

Since unspecific effects might have been caused by a different fatty acid composition of the experimental fats (the heating process caused a loss of polyunsaturated fatty acids), we aimed at equalizing in particular the concentration of the polyunsaturated fatty acid C18:2 (n-6) in the fresh fat and the oxidized fat in the second experiment which lasted during the whole pregnancy. Although the

**Figure 3**

Effects of treatment on triacylglycerol concentrations in livers of pregnant rats and fetuses. Effect of short-term (d 16 – d 21 of pregnancy, left) and long-term (d 1 – d 21 of pregnancy, right) administration of three different diets (control, oxidized fat, clofibrate) during pregnancy on triacylglycerol concentrations in the liver of pregnant rats and fetuses at d 21 of pregnancy. Left, Bars represent mean \pm SD (n = 9/group). Right, Bars represent mean \pm SD (n = 12/group). Bars marked without a common superscript letter differ ($P < 0.05$). Results from one-way ANOVA (P -values): short-term experiment: $P = 0.239$ (pregnant), $P = 0.025$ (fetus); long-term experiment: $P = 0.0001$ (pregnant), $P = 0.045$ (fetus).

concentrations of the long-chain saturated fatty acids C16:0 and C18:0 were consequently decreased in the oxidized fat compared to the fresh fat, we think that these differences are not responsible for the differences in the PPAR α response observed between the experimental groups, because saturated long chain-fatty acids were shown to bind and activate PPAR α only very weakly compared to polyunsaturated fatty acids such as C18:2 (n-6) [25]. Therefore, we assume that the PPAR α response to an oxidized fat might depend on the balance between unoxidized fatty acids with low PPAR α transactivation activity and oxidized fatty acids with high PPAR α transactivation activity because both types of fatty acids compete for the PPAR α -ligand binding site.

A further consequence of the use of lard in the control fat was that the cholesterol content of the control and the clofibrate diet slightly differed from that of the oxidized fat diet. However, based on an average cholesterol concentration of about 80 mg per 100 g lard the control and the clofibrate diet contained less than 0.004% cholesterol. Therefore, we assume that the slight difference in the cholesterol concentration between the experimental diets is negligible, especially since no relation between PPAR α activation and dietary cholesterol is known from the literature.

The present study clearly shows that feeding oxidized fat during pregnancy, similar to clofibrate, which was used as a positive control, causes not only a PPAR α response in

the liver of pregnant animals as shown by liver enlargement and up-regulation of PPAR α -responsive genes but also in the liver of the fetuses. Although the induction of the PPAR α response by oxidized fat in the fetal liver was not as pronounced as observed with clofibrate, the oxidized fat also caused a strong up-regulation of PPAR α -target genes of up to 9-fold in the livers of fetuses which was even more pronounced than the effect of oxidized fat in the livers of pregnant rats. Therefore, these findings suggest that not only pharmacological PPAR α activators but also components of oxidized fat are able to sufficiently pass the placenta and activate PPAR α in the fetal liver. This finding is novel since the placental transfer of these components of oxidized fat with PPAR α activating potential from the maternal diet to the fetus is unknown. Indeed, the transplacental transport of fatty acids is highly selective for individual fatty acids [20-22], but no data are available from the literature with respect to the placental passage of oxidized fatty acids which are presumably decisive for the PPAR α activating effect of oxidized fat. Thus, the present results indicate that these critical components of oxidized fats are also sufficiently transported across the placenta. Moreover, the observation from the short-term experiment that up-regulation of mRNA expression of ACO, CYP4A1 and L-CPT I by oxidized fat in the fetal liver was even more pronounced than in the liver of pregnant rats indicates that components of oxidized fat responsible for PPAR α activation are presumably transported through the placenta with high preference.




Figure 4
Effects of treatment on relative mRNA concentrations of proto-oncogenes in livers of pregnant rats and fetuses. Effect of short-term (d 16 – d 21 of pregnancy, left) and long-term (d 1 – d 21 of pregnancy, right) administration of three different diets (control, oxidized fat, clofibrate) during pregnancy on relative mRNA concentrations of c-myc, c-jun, and c-fos in the liver of pregnant rats and fetuses at d 21 of pregnancy. Left, Bars represent mean ± SD (n = 9/group). Right, Bars represent mean ± SD (n = 12/group). Bars marked without a common superscript letter differ ($P < 0.05$). Results from one-way ANOVA (P -values): short-term experiment: c-myc, $P = 0.880$ (pregnant), $P = 0.807$ (fetus); c-jun, $P = 0.362$ (pregnant), $P = 0.552$ (fetus); c-fos, $P = 0.669$ (pregnant), $P = 0.966$ (fetus); long-term experiment: c-myc, $P = 0.001$ (pregnant), $P = 0.420$ (fetus); c-jun, $P = 0.0001$ (pregnant), $P = 0.086$ (fetus); c-fos, $P = 0.048$ (pregnant), $P = 0.323$ (fetus).

Since activation of hepatic PPAR α by clofibrate or oxidized fat has been demonstrated to enhance the fatty acid oxidation capacity in the liver and to lower hepatic triacylglycerol concentrations in non-pregnant rats [1,3,9], we also studied the effect of oxidized fat on the concentrations of triacylglycerols in the fetal liver. Indeed, we could demonstrate that feeding the oxidized fat during pregnancy also reduced the concentrations of triacylglycerols in the fetal liver indicating that the transplacental induction of PPAR α responsive genes enhanced the peroxisomal and mitochondrial fatty acid oxidation capacity of the fetal liver. However, this effect has only been observed when the oxidized fat was fed during the whole pregnancy, but not when the oxidized fat was fed for the last 5 d of pregnancy only, although the PPAR α responsive genes were markedly up-regulated in fetal livers of pregnant rats treated during either whole or late pregnancy. This suggests that short-term administration of oxidized fat to pregnant rats causes significant alterations on the gene expression level, which have no implications on the phenotypic level, e.g. triacylglycerol concentrations. In contrast, short-term administration of clofibrate during pregnancy even revealed alterations on the phenotypic level as evidenced by reduced triacylglycerol concentrations in fetal livers. The latter might be attributed to the fact that clofibrate caused a more pronounced activation of PPAR α in fetal livers due to a higher affinity for PPAR α compared to oxidized fat leading to a marked induction of fatty acid oxidation and a significant lowering of triacylglycerol concentrations in fetal livers even after short-term exposure to clofibrate.

Interestingly, no effect of oxidized fat and even clofibrate on hepatic triacylglycerol concentrations following short-term treatment could be observed in pregnant rats, although clofibrate reduced triacylglycerol concentrations in fetal livers following short-term treatment. The failure of clofibrate or oxidized fat to reduce hepatic triacylglycerol concentrations in pregnant rats in the short-term experiment might be attributed to the fact that treatment was only performed during late pregnancy when significant alterations in lipid metabolism [e.g. hypertriacylglycerolemia as a consequence of enhanced adipose tissue lipolytic activity, enhanced liver production of VLDL particles, and decreased extrahepatic lipoprotein lipase activity [26-29]] occur in the pregnant animal. The enhanced arrival of free fatty acids, which also serve as ligands for PPAR α , in the liver during late pregnancy could decrease the availability for the stronger PPAR α activators clofibrate or components of oxidized fat with PPAR α activating potential to their PPAR α -ligand binding site. This in turn could reduce the capability of fibrates or oxidized fat to activate PPAR α and consequently, its metabolic effects. This might be important since a substantial body of evidence suggests that not all alterations in gene transcrip-

tion induced by pharmacological PPAR α activators are also induced by fatty acids [30], e.g. some genes that contain a PPRE do not respond to fatty acids but to high-affinity PPAR α activators [31-33]. Thus, administration of oxidized fat or clofibrate during late pregnancy might provoke differential effects on lipid metabolism than in virgin or early pregnant rats. The observation that treatment of rats with the triacylglycerol-lowering PPAR α agonist fenofibrate during late pregnancy even increased triacylglycerol concentrations in plasma, whereas in virgin rats treatment with fenofibrate caused a reduction in plasma triacylglycerol concentrations [16], is probably supportive of this assumption.

Whether the effect observed with oxidized fat or clofibrate might have been also influenced by the metabolic state, e.g. fasting vs. non-fasting, cannot be answered with certainty. In the present study we decided to perform gene expression analysis of PPAR α -responsive genes in the liver in the non-fasting state, because lipolysis of stored triacylglycerols in adipose tissue is strongly activated during fasting, resulting in a marked increase in plasma free fatty acid levels. These free fatty acids act as endogenous PPAR α ligands and might have competed with the exogenous ligands (e.g. clofibrate, oxidized fatty acids or cyclic fatty acid monomers) for the PPAR α -ligand binding site. Therefore, an altered ratio between endogenous free fatty acids and exogenous oxidized fatty acids as a consequence of the fasting state might have provoked a different PPAR α response in the liver of pregnant rats and fetuses than observed in the non-fasting state. However, further studies are required to definitely resolve this question.

Only slight effects have been observed in the present study with respect to the lipogenic transcription factor SREBP-1c and its target gene FAS. Namely, oxidized fat caused a slight reduction in the mRNA abundance of FAS in the liver of pregnant rats, which is consistent with recent findings in non-pregnant rats [9]. However, no effect of oxidized fat has been observed on gene expression of lipogenic enzymes in fetal livers indicating that the reduced hepatic triacylglycerol concentrations in fetuses from pregnant rats treated with oxidized fat are probably largely due to an enhanced fatty acid oxidation capacity due to transplacental activation of PPAR α . In part, the failure of oxidized fat on FAS gene expression in fetal livers might be explained by the fact that lipogenesis in fetal livers is generally very low and only increased during late pregnancy [34], and, therefore, probably does not respond to variations in the maternal diet.

Since pathological changes in newborn rats born to mothers treated with clofibrate during pregnancy have been reported [15], treatment with fenofibrate has been shown to impair fetal growth [16], and a link between induced

CYP4A1 expression, peroxisome proliferation, and carcinogenesis in rat livers has been described [35], we also addressed possible adverse effects of treatment with oxidized fat on fetal livers. With respect to the hepatocarcinogenic effect of peroxisome proliferators in rodents it has been suggested that enhanced DNA synthesis as a consequence of up-regulation of proto-oncogenes including *c-fos*, *c-jun*, and *c-myc* might be mechanistically involved [17-19]. In the present study an up-regulation of proto-oncogenes in the livers of pregnant rats has been observed in the long-term experiment, but not in the short-term experiment, indicating that short-term administration of oxidized fat has no impact on mRNA expression of proto-oncogenes. However, in fetal livers mRNA expression of proto-oncogenes was not affected by oxidized fat regardless of the duration of oxidized fat administration suggesting that the oxidized fat is uncritical with respect to hepatocarcinogenesis. Unexpectedly, treatment with clofibrate had no effect on proto-oncogene expression in pregnant rats and fetuses either in the short-term and the long-term experiment, although it has been reported that the high-affinity PPAR α -ligand WY-14,643 strongly up-regulated various proto-oncogenes in rat and mouse liver epithelial cells [18,19]. This differential action of WY-14,643 and clofibrate on proto-oncogene mRNA expression cannot be explained at the moment and, therefore, requires further research activities.

Although oxidized fat had no effect on proto-oncogene expression in fetal livers in the present study, oxidized fats might be considered critically in view of inducing oxidative stress in different tissues as shown in recent studies [36,37]. In addition, specific components of oxidized fats such as cyclic fatty acid monomers, which are formed in substantial amounts during domestic frying of frozen foods in sunflower oil [38], are probably toxic, e.g. earlier studies reported that mice receiving cyclic fatty acid monomers as well as rat pups from mothers fed cyclic fatty acid monomers had a higher death rate [39-41].

Conclusion

In conclusion, the present study demonstrates for the first time that components of oxidized fat with PPAR α activating potential contained in the maternal diet are able to induce a PPAR α response in the liver of fetuses as evidenced by an up-regulation of PPAR α target genes. In addition, the present study shows that feeding oxidized fat during whole pregnancy, but not during late pregnancy, lowers triacylglycerol concentrations in fetal livers, probably as a consequence of an enhanced peroxisomal and mitochondrial β -oxidation capacity. Although administration of oxidized fat during pregnancy had not impact on fetal proto-oncogene mRNA expression either after short-term or long-term administration, the observed pronounced transplacental PPAR α activation by

oxidized fat might be considered critically because of other recently reported adverse effects of treatment with PPAR α activators during pregnancy [15,16]. Therefore, further research should be encouraged with respect to possible detrimental effects of oxidized fat on fetal development.

Methods

Animals

Two experiments were carried out with female Sprague-Dawley rats obtained from Charles River (Sulzfeld, Germany). At 11 wk of age, the rats were mated by housing one male rat with two female rats. D 1 of pregnancy was assigned upon observation of sperm in the vaginal smears, at which time rats were randomly assigned to the treatment groups. The short-term experiment was performed from d 16 to d 21 of pregnancy and included 27 pregnant rats with an initial body weight (d 16 of pregnancy) of 297 ± 26 (Mean \pm SD) g, which were allotted to three groups of nine rats each. The long-term experiment was performed from d 1 to d 21 of pregnancy and included 36 rats with an initial body weight (d 1 of pregnancy) of 238 ± 27 g, which were allotted to three groups of twelve rats each. Pregnant rats were kept individually in Macrolon cages in a room maintained with controlled temperature ($23 \pm 1^\circ\text{C}$), humidity (50–60%), and lighting (0600 to 1800 h). All experimental procedures described followed established guidelines for the care and handling of laboratory animals [42] and were approved by the council of Saxony-Anhalt.

Diets

Short-term experiment

In the short-term experiment, rats received 2 mL of different experimental fats by gavage daily at 0800 h, and, additionally, fed a commercial standard rodent diet (Altromin, Lage, Germany). To standardize food intake, the diets were fed daily in controlled amounts of 16 g per d. The first group (control group) received sunflower oil, the second group (oxidized fat group) oxidized fat (see "preparation of the oxidized fat"), and the third group (clofibrate group) sunflower oil containing 75 mg clofibrate (Fluka, Buchs, Switzerland) equivalent to 250 mg clofibrate per kg body weight. The experimental fats were given for 5 d from d 16 of pregnancy to d 21 of pregnancy. The standard diet was completely consumed by all the rats. Thus, all the rats within this experiment consumed identical amounts of the food.

Long-term experiment

In the long-term experiment, semipurified diets, composed according to the recommendations of ASNS for rats during reproduction [43], were used. The diet consisted of (g/kg diet): casein, 200; cornstarch, 390; saccharose, 198; cellulose, 50; fat, 100; mineral mixture, 40; vitamin mix-

ture, 20; DL-methionine, 2. The type of fat was varied according to a one-factorial design. The first group (control group) received a mixture of sunflower oil and lard (54:46, w/w) was used. This ratio was chosen to equalize the fatty acid composition of the fresh fat with that of the oxidized fat, since the heating process caused a loss of polyunsaturated fatty acids, therefore, excluding that the treatment effects were caused by a different fatty acid composition of the experimental fats. The second group (oxidized fat group) received oxidized fat (see "preparation of the oxidized fat"). The third group (clofibrate group) received the same fat as in the control group, and clofibrate was added to the diet at a concentration of 5 g/kg. The vitamin E concentration of the diets was 50 mg α -tocopherol equivalents per kg diet. To adjust the vitamin E concentration of the diets, the native concentrations of tocopherols of the fats were analyzed. Based on the native concentrations of the fats, diets were supplemented individually with all-rac- α -tocopheryl acetate (the biopotency of all-rac- α -tocopheryl acetate is considered to be 67% of that of α -tocopherol). Diets were prepared by mixing the dry components with the fat and water and subsequent freeze drying. The residual water content of the diet was below 5 g/100 g of diet. In preliminary experiments rats fed diets with clofibrate or oxidized fat *ad libitum* consumed their diets over a longer period than rats fed control diets, which consequently shortened the fasting period and which itself has a pronounced effect on PPAR α -response. Therefore, we decided to administer food daily at 0800 h in controlled amounts to standardize intake and to ensure that rats from all treatment groups had a comparable fasting period. The amount of food administered was 20% less than the amounts of identical diets with fresh fats consumed *ad libitum* by rats in preliminary studies. The amount of food offered daily was increased continuously during the experiment from 14 g to 17 g. In this feeding system, the food offered was completely consumed by all the rats. Thus, all the rats within this experiment consumed identical amounts of food. The experimental diets were fed from d 1 of pregnancy to d 21 of pregnancy.

In both experiments, water was available *ad libitum* from nipple drinkers during the whole experiment.

Preparation of the oxidized fat

The oxidized fat was prepared by heating sunflower oil at a temperature of 60°C for 25 d. Sunflower oil was filled into a glass beaker and placed into a drying oven set at the intended temperature. Throughout the heating process, air was continuously bubbled through the fat at a flow rate of 650 ml/min. This treatment caused a loss of polyunsaturated fatty acids, and a complete loss of tocopherols and raised the concentrations of lipid peroxidation products in the fats. The extent of lipid peroxidation in the

oxidized fat was estimated by assaying the peroxide value (POV) [44], acid value [44], concentration of thiobarbituric acid substances (TBARS) [45], and concentration of conjugated dienes [46]. To assess lipid peroxidation products in the oxidized fat after inclusion into the diet (long-term experiment), the fat was extracted from aliquots of the diets with a mixture of hexane and isopropanol (3:2, v/v) and analysed for peroxide value, concentration of conjugated dienes, and TBARS.

Sample collection

4 h after the final portion had been administered the rats were anesthetized with diethyl ether and killed by decapitation. The liver and fetuses were excised immediately, and frozen with liquid nitrogen. In addition, livers from three randomly taken fetuses per pregnant rat were excised, and frozen with liquid nitrogen. All samples were stored at -80°C pending analysis.

Lipid analysis

Lipids of maternal livers and pools of fetal livers were extracted with a mixture of hexane and isopropanol (3:2, v/v) [47]. Total cholesterol and triacylglycerol concentrations were determined using enzymatic reagent kits obtained from Merck Eurolab (Darmstadt, Germany). Prior to enzymatic measurement, lipids of the extract were dissolved in Triton X-100 as described by De Hoff et al. [48]. Fatty acid composition of experimental fats was determined by GC-FID analysis of fatty acid methyl esters (FAME) as described previously in detail [49].

RNA isolation and real-time RT-PCR

For the determination of mRNA expression levels of PPAR α , CYP4A1, ACO, L-CPT I, MCAD, LCAD, SREBP-1c, FAS, c-myc, c-jun, and c-fos total RNA were isolated from maternal liver and fetal liver pools using Trizol™ reagent (Invitrogen, Karlsruhe, Germany) according to the manufacturer's protocol. RNA concentration and purity were estimated from the optical density at 260 and 280 nm, respectively. cDNA synthesis and relative quantification of target gene mRNA compared to the housekeeping gene GAPDH mRNA was determined by real-time detection RT-PCR as described previously [50]. Sequences of gene-specific primers obtained from Operon (Köln, Germany) were as follows (NCBI GenBank; forward, reverse): GAPDH ([NM_017008](#); 5'-GCA TGG CCT TCC GTG TTC C-3', 5'-GGG TGG TCC AGG GTT TCT TAC TC-3'), PPAR α ([NM_013196](#); 5'-CCC TCT CTC CAG CTT CCA GCC C-3', 5'-CCA CAA GCG TCT TCT CAG CCA TG-3'), CYP4A1 (M14972; 5'-CAG AAT GGA GAA TGG GGA CAG C-3', 5'-TGA GAA GGG CAG GAA TGA GTG G-3'), ACO (J02752; 5'-CTT TCT TGC TTG CCT TCC TTC TCC-3', 5'-GCC GTT TCA CCG CCT CGT A-3'), L-CPT I ([NM_031559](#); 5'-GGA GAC AGA CAC CAT CCA ACA TA-3', 5'-AGG TGA TGG ACT TGT CAA ACC-3'), MCAD ([NM_016986](#); 5'-CAA

GAG AGC CTG GGA ACT TG-3', 5'-CCC CAA AGA ATT TGC TTC AA-3'), LCAD ([NM_012819](#); 5'-AAG GAT TTA AGG GCA AGA AGC-3', 5'-GGA AGC GGA GGC GGA GTC-3'), SREBP-1c ([XM_213329](#); 5'-GGA GCC ATG GAT TGC ACA TT-3', 5'-AGG AAG GCT TCC AGA GAG GA-3'), FAS ([NM_017332](#); 5'-AGG TGC TAG AGG CCC TGC TA-3', 5'-GTG CAC AGA CAC CTT CCC AT-3'), c-myc ([NM_012603](#); 5'-CTG GAG TGA GAA GGG CTT TG-3', 5'-CAG CAG CTC GAA TTT CTT CC-3'), c-jun ([NM_021835](#); 5'-ACC AAG AAT TCC GTG ACG AC-3', 5'-CAA GGT CAT GCT CTG CTT CA-3'), and c-fos ([NM_022197](#); 5'-CAT CGG CAG AAG GGG CAA AGT AGA G-3', 5'-TGC CGG AAA CAA GAA GTC ATC AAA G-3').

Statistical analysis

Treatment effects were analyzed using one-way ANOVA. For significant *F*-values, means were compared by Fisher's multiple range test. Differences with *P* < 0.05 were considered significant.

List of abbreviations used

ACO, acyl-CoA oxidase; CYP4A1, cytochrome P₄₅₀ 4A1; FAS, fatty acid synthase; LCAD, long-chain acyl-CoA dehydrogenase; L-CPT I, L-type carnitin-palmitoyl transferase I; MCAD, medium-chain acyl-CoA dehydrogenase; PPAR α , peroxisome proliferator-activated receptor α ; PPRE, PPAR response element; SREBP-1c, sterol regulatory-element binding protein-1c.

Competing interests

The author(s) declare that they have no competing interests.

Authors' contributions

RR participated in the design of the study and in the interpretation of the results and prepared the manuscript.

AG and CD carried out the feeding experiments, quantification of lipid concentrations, and mRNA expression analysis.

CB participated in the design and coordination of the study, and interpretation of the results.

KE conceived of the study and its design, coordinated work, participated in the interpretation of the results, and helped to draft the manuscript.

All authors read and approved the final manuscript.

Acknowledgements

Performance of gastric tube feeding in the short-term experiment by Alexander Koch and Dr. Bettina König is gratefully acknowledged.

References

- Chao PM, Chao CY, Lin FJ, Huang CJ: **Oxidized frying oil up-regulates hepatic acyl-CoA oxidase and cytochrome P450 4A1 genes in rats and activates PPAR α .** *J Nutr* 2001, **131**:3166-3174.
- Chao PM, Hsu SC, Lin FJ, Li YJ, Huang CJ: **The up-regulation of hepatic acyl-CoA oxidase and cytochrome P450 4A1 mRNA expression by dietary oxidized frying oil is comparable between male and female rats.** *Lipids* 2004, **39**:233-238.
- Sülzle A, Hirche F, Eder K: **Thermally oxidized dietary fat upregulates the expression of target genes of PPAR α in rat liver.** *J Nutr* 2004, **134**:1375-1383.
- Costet P, Legendre C, More J, Edgar A, Galtier P, Pineau T: **Peroxisome proliferators-activated receptor α -isoform deficiency leads to progressive dyslipidemia with sexually dimorphic obesity and steatosis.** *J Biol Chem* 1998, **273**:29577-29585.
- Martin JC, Joffre F, Siess MH, Vernevauf MF, Collenot P, Genty M, Sebedio JL: **Cyclic fatty acid monomers from heated oil modify the activities of lipid synthesizing and oxidizing enzymes in rat liver.** *J Nutr* 2000, **130**:1524-1530.
- König B, Eder K: **Differential action of 13-HPODE on PPAR α downstream genes in rat Fao and human HepG2 hepatoma cell lines.** *J Nutr Biochem* 2006, **7**:410-418.
- Issemann I, Green S: **Activation of a member of the steroid hormone receptor superfamily by peroxisome proliferators.** *Nature* 1990, **347**:645-650.
- Schoonjans K, Staels B, Auwerx J: **The PPARs and their effects on lipid metabolism and adipocyte differentiation.** *Biochim Biophys Acta* 1996, **1302**:93-109.
- Eder K, Sülzle A, Skufca P, Brandsch C, Hirche F: **Effects of dietary thermoxidized fats on expression and activities of hepatic lipogenic enzymes in rats.** *Lipids* 2003, **38**:31-38.
- Chao PM, Yang MF, Tseng YN, Chang KM, Lu KS, Huang CJ: **Peroxisome proliferation in liver of rats fed oxidized frying oil.** *J Nutr Sci Vitaminol (Tokyo)* 2005, **51**:361-368.
- Stefanini S, Mauriello A, Farrace MGC, Cibelli A, Ceru MP: **Proliferative response of foetal liver peroxisomes to clofibrate treatment of pregnant rats. A quantitative evaluation.** *Biol Cell* 1989, **67**:299-305.
- Wilson GN, King T, Argyle JC, Garcia RF: **Maternal clofibrate administration amplifies fetal peroxisomes.** *Pediatr Res* 1991, **29**:256-262.
- Cibelli A, Stefanini S, Ceru MP: **Peroxisomal β -oxidation and catalase activities in fetal rat liver: effect of maternal treatment with clofibrate.** *Cell Mol Biol* 1988, **34**:191-205.
- Simpson AE, Brammar WJ, Pratten MK, Cockcroft N, Elcombe CR: **Placental transfer of the hypolipidemic drug, clofibrate, induces CYP4A expression in 18.5-day fetal rats.** *Drug Metab Dispos* 1996, **24**:547-554.
- Amels S, Fazekas-Todea I, Sandor S: **The prenatal noxious effect of blood cholesterol level lowering compound.** *Rev Roum Morphol Embryol* 1974, **19**:37-43.
- Soria A, Bocos C, Herrera E: **Opposite metabolic response to fenofibrate treatment in pregnant and virgin rats.** *J Lipid Res* 2002, **43**:74-81.
- Ledwith BJ, Manam S, Troilo P, Joslyn DJ, Galloway SM, Nichols WW: **Activation of immediate-early gene expression by peroxisome proliferators in vitro.** *Mol Carcinog* 1993, **8**:20-27.
- Ledwith BJ, Johnson TE, Wagner LK, Pauley CJ, Manam S, Galloway SM, Nichols WW: **Growth regulation by peroxisome proliferators: opposing activities in early and late G1.** *Cancer Res* 1996, **56**:3257-3264.
- Miller RT, Glover SE, Stewart WS, Corton JC, Popp JA, Cattley RC: **Effect on the expression of c-met, c-myc and PPAR-alpha in liver and liver tumors from rats chronically exposed to the hepatocarcinogenic peroxisome proliferator WY-14,643.** *Carcinogenesis* 1996, **17**:1337-1341.
- Herrera E, Amusquivar E, Lopez-Soldado I, Ortega : **Maternal lipid metabolism and placental lipid transfer.** *Horm Res* 2006, **65**:59-64.
- Haggarty P, Ashton J, Joynson M, Abramovich DR, Page K: **Effect of maternal polyunsaturated fatty acid concentration on transport by human placenta.** *Biol Neonate* 1999, **75**:350-359.
- Haggarty P: **Placental regulation of fatty acid delivery and its effect on fetal growth – a review.** *Placenta* 2002, **23**(Suppl A):S28-38.

23. Mishra A, Chaudhary A, Sethi S: **Oxidized omega-3 fatty acids inhibit NF- κ B activation via a PPAR α -dependent pathway.** *Arterioscler Thromb Vasc Biol* 2004, **24**:1621-1627.
24. Delerive P, Furman C, Teissier E, Fruchart JC, Duriez P, Staels B: **Oxidized phospholipids activate PPAR α in a phospholipase A2-dependent manner.** *FEBS Lett* 2000, **471**:34-38.
25. Forman BM, Chen J, Evans RM: **Hypolipidemic drugs, polyunsaturated fatty acids, and eicosanoids are ligands for peroxisome proliferator-activated receptors α and δ .** *Proc Natl Acad Sci USA* 1997, **94**:4312-4317.
26. Herrera E: **Metabolic adaptations in pregnancy and their implications for the availability of substrates to the fetus.** *Eur J Clin Nutr* 2000, **54**(Suppl 1):S47-51.
27. Elliott JA: **The effect of pregnancy on the control of lipolysis in fat cells isolated from human adipose tissue.** *Eur J Clin Invest* 1975, **5**(2):159-163.
28. Chaves JM, Herrera E: **In vitro glycerol metabolism in adipose tissue from fasted pregnant rats.** *Biochem Biophys Res Commun* 1978, **85**:1299-1306.
29. Martin-Hidalgo A, Holm C, Belfrage P, Schotz MC, Herrera E: **Lipoprotein lipase and hormone-sensitive lipase activity and mRNA in rat adipose tissue during pregnancy.** *Am J Physiol* 1994, **266**:E930-935.
30. Duplus E, Glorian M, Forest C: **Fatty acid regulation of gene transcription.** *J Biol Chem* 2000, **275**:30749-30752.
31. Cho HP, Nakamura MT, Clarke SD: **Cloning, expression, and nutritional regulation of the mammalian delta-6 desaturase.** *J Biol Chem* 1999, **274**(1):417-477.
32. Cho HP, Nakamura M, Clarke SD: **Cloning, expression, and fatty acid regulation of the human delta-5 desaturase.** *J Biol Chem* 1999, **274**:37335-37339.
33. Ren B, Thelen A, Jump DB: **Peroxisome proliferator-activated receptor alpha inhibits hepatic S14 gene transcription. Evidence against the peroxisome proliferator-activated receptor alpha as the mediator of polyunsaturated fatty acid regulation of s14 gene transcription.** *J Biol Chem* 1996, **271**:17167-17173.
34. Herrera E, Amusquivar E: **Lipid metabolism in the fetus and the newborn.** *Diabetes Metab Res Rev* 2000, **16**:202-210.
35. Reddy JK, Rao MS: **Oxidative DNA damage caused by persistent peroxisome proliferation: its role in hepatocarcinogenesis.** *Mutat Res* 1989, **214**:63-68.
36. Garrido-Polonio C, Garcia-Linares MC, Garcia-Arias MT, Lopez-Varela S, Garcia-Fernandez MC, Terpstra AH, Sanchez-Muniz FJ: **Thermally oxidised sunflower-seed oil increases liver and serum peroxidation and modifies lipoprotein composition in rats.** *Br J Nutr* 2004, **92**:257-265.
37. Keller U, Brandsch C, Eder K: **Supplementation of vitamins C and E increases the vitamin E status but does not prevent the formation of oxysterols in the liver of guinea pigs fed an oxidised fat.** *Eur J Nutr* 2004, **43**:353-359.
38. Romero A, Bastida S, Sanchez-Muniz FJ: **Cyclic fatty acid monomer formation in domestic frying of frozen foods in sunflower oil and high oleic acid sunflower oil without oil replenishment.** *Food Chem Toxicol* 2006, **44**:1674-1681.
39. Saito M, Kaneda T: **Studies on the relationship between the nutritive value and the structure of polymerized oils.** *Jpn J Oil Chem Soc* 1976, **25**:13-20.
40. Sebedio JL, Grandgirard A: **Cyclic fatty acids: natural sources, formation during heat treatment, synthesis and biological properties.** *Prog Lipid Res* 1989, **28**:303-336.
41. Sébédio JL, Chardigny JM, Juaneda P, Giraud MC, Nour M, Christie WW, Dobson G: **Nutritional impact and selective incorporation of cyclic fatty acid monomers in rats during reproduction.** In *21st World Congress of the International Society for Fat Research Oils-Fats-Lipids Volume 2*. Edited by: Castenmiller WAM, Barnes PJ, The Hague The Netherlands; 1995:307-310.
42. National Research Council: **Guide for the Care and Use of Laboratory Animals.** In *Publication no. 85-23 (rev.)* National Institutes of Health, Washington, DC; 1985.
43. Reeves PG, Nielsen FH, Fahey GC Jr: **A1N-93 purified diets for laboratory rodents: final report of the American Institute of Nutrition ad hoc writing committee on the reformulation of the A1N-76A rodent diet.** *J Nutr* 1993, **123**:1939-1951.
44. Deutsche Gesellschaft für Fettwissenschaft: **Einheitsmethoden zur Untersuchung von Fetten, Fettprodukten, Tensiden und verwandten Stoffen.** Wissenschaftliche Verlagsgesellschaft Stuttgart Germany 1994.
45. Sidwell CG, Salwin H, Benca M, Mitchell JH Jr: **The use of thiobarbituric acid as a measure of fat oxidation.** *J Am Oil Chem Soc* 1954, **31**:603-606.
46. Recknagel RO, Glende EA Jr: **Spectrophotometric detection of lipid conjugated dienes.** *Methods Enzymol* 1984, **105**:331-337.
47. Hara A, Radin NS: **Lipid extraction of tissues with a low toxicity solvent.** *Anal Biochem* 1978, **90**:420-426.
48. De Hoff JL, Davidson JH, Kritchevsky V: **An enzymatic assay for determining free and total cholesterol in tissues.** *Clin Chem* 1978, **24**:433-435.
49. Brandsch C, Ringseis R, Eder K: **High dietary iron concentrations enhance the formation of cholesterol oxidation products in the liver of adult rats fed salmon oil with minimal effects on antioxidant status.** *J Nutr* 2002, **132**:2263-2269.
50. Ringseis R, Eder K: **Effects of dietary fat and oxidized cholesterol on gene expression in rat liver as assessed by cDNA expression array analysis.** *Eur J Nutr* 2005, **44**:231-241.

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
http://www.biomedcentral.com/info/publishing_adv.asp

