

## Research



**Cite this article:** Kocot KM, Todt C, Mikkelsen NT, Halanych KM. 2019 Phylogenomics of Aplacophora (Mollusca, Aculifera) and a solenogaster without a foot. *Proc. R. Soc. B* **286**: 20190115.  
<http://dx.doi.org/10.1098/rspb.2019.0115>

Received: 19 January 2019  
Accepted: 17 April 2019

**Subject Category:**  
Evolution

**Subject Areas:**  
taxonomy and systematics

**Keywords:**  
Neomeniomorpha, Solenogastres,  
Chaetodermomorpha, Caudofoveata,  
*Apodomenia*

**Author for correspondence:**  
Kevin M. Kocot  
e-mail: [kmkocot@ua.edu](mailto:kmkocot@ua.edu)

Electronic supplementary material is available online at <https://dx.doi.org/10.6084/m9.figshare.c.4483301>.

# Phylogenomics of Aplacophora (Mollusca, Aculifera) and a solenogaster without a foot

Kevin M. Kocot<sup>1</sup>, Christiane Todt<sup>2</sup>, Nina T. Mikkelsen<sup>3</sup>  
and Kenneth M. Halanych<sup>4</sup>

<sup>1</sup>The University of Alabama and the Alabama Museum of Natural History, 500 Hackberry Lane, Tuscaloosa, AL 35487, USA

<sup>2</sup>Rådgivende Biologer AS, Edvard Griegs vei 3, 5059 Bergen, Norway

<sup>3</sup>University Museum of Bergen, The Natural History Collections, University of Bergen, Allégaten 41, 5007 Bergen, Norway

<sup>4</sup>Department of Biological Sciences, Auburn University, Auburn, AL 36849, USA

KMK, 0000-0002-8673-2688

Recent molecular phylogenetic investigations strongly supported the placement of the shell-less, worm-shaped aplacophoran molluscs (Solenogastres and Caudofoveata) and chitons (Polyplacophora) in a clade called Aculifera, which is the sister taxon of all other molluscs. Thus, understanding the evolutionary history of aculiferan molluscs is important for understanding early molluscan evolution. In particular, fundamental questions about evolutionary relationships within Aplacophora have long been unanswered. Here, we supplemented the paucity of available data with transcriptomes from 25 aculiferans and conducted phylogenomic analyses on datasets with up to 525 genes and 75 914 amino acid positions. Our results indicate that aplacophoran taxonomy requires revision as several traditionally recognized groups are non-monophyletic. Most notably, Cavibelonia, the solenogaster taxon defined by hollow sclerites, is polyphyletic, suggesting parallel evolution of hollow sclerites in multiple lineages. Moreover, we describe *Apodomenia enigmatica* sp. nov., a bizarre new species that appears to be a morphological intermediate between Solenogastres and Caudofoveata. This animal is not a missing link, however; molecular and morphological studies show that it is a derived solenogaster that lacks a foot, mantle cavity and radula. Taken together, these results shed light on the evolutionary history of Aplacophora and reveal a surprising degree of morphological plasticity within the group.

## 1. Introduction

The two groups of worm-like aplacophoran molluscs, Solenogastres (= Neomeniomorpha) and Caudofoveata (= Chaetodermomorpha), have perplexed biologists since their discovery [1,2]. Aplacophorans are characterized by a narrow or completely reduced foot, a unique posterior dorsoterminal sensory organ, and a small mantle cavity restricted to the posterior-most part of the body. Solenogastres and caudofoveates both completely lack a shell, but instead are covered in a dense coat of spiny or scale-like calcareous sclerites [3–8].

Aplacophorans have generally been regarded as early-branching molluscs and therefore have been central to questions surrounding the origin and early evolution of the phylum. Whether Solenogastres and Caudofoveata constitute a monophyletic taxon, Aplacophora [4,9,10], or a 'basal', paraphyletic grade [5,8,11–14], has been debated [6,7,15]. Recent molecular studies [16–18] have strongly supported monophyly of Aplacophora and a sister group relationship of Aplacophora and Polyplacophora (chitons), consistent with the Aculifera hypothesis [4]. Analyses of fossils (e.g. [19]) and evolutionary developmental

approaches [20,21] have provided further evidence for this hypothesis. Support for Aculifera has had an important impact on understanding of plesiomorphic characteristics of Mollusca [16] as it suggests the last common ancestor of the phylum was a large-bodied, chiton-like animal, but many more questions remain unanswered. Although aplacophorans are not the sister taxon to all other molluscs as previously thought [8,13,14,22,23], resolving aplacophoran phylogeny is critical to understanding early molluscan evolution, as it could help reveal the evolutionary polarity of key morphological characters for Aplacophora, Aculifera and Mollusca as a whole.

Caudofoveate taxonomy is based primarily on characteristics of the sclerites and radula. Around 130 species have been described and three families are traditionally recognized [6,24]. Limifossoridae has been hypothesized to show the most plesiomorphic morphological characters among caudofoveates [25–27], mainly a solenogaster-like (distichous) radula with two teeth per row and a simple body shape. Within the more diverse Solenogastres, classification is based primarily on characters of the sclerites, cuticle, radula, ventrolateral foregut glands and reproductive anatomy [24,28,29]. Presently, around 280 species in 24 families and four orders are recognized, but the actual diversity within the group is estimated to be considerably higher [6]. According to the taxonomy established by Salvini-Plawen [28], the orders Pholidoskepia and Neomeniamorpha are grouped together in a higher taxon called Aplotegmentaria. The small-bodied, scale-bearing Pholidoskepia have been regarded as ‘primitive’ solenogasters [28,30,31]. The remaining two orders, Cavibelonia and Sterrofustia, are grouped together in a higher taxon called Pachytegumentaria.

Chitons have a fairly rich fossil record [32] and their phylogeny is at least generally understood [33–36]. However, no *bona fide* solenogaster or caudofoveate fossils are known [19,37–40], and cladistic morphological analyses examining solenogaster phylogeny [5,41,42] have generally failed to recover most higher-level taxa monophyletic, suggesting that the existing taxonomy does not reflect the evolutionary history of the group or that the morphological data analysed lack sufficient phylogenetic signal to reconstruct aplacophoran relationships. Recent molecular studies employing datasets dominated by nuclear ribosomal and mitochondrial genes [43–45] have greatly improved understanding of relationships within Caudofoveata. However, nuclear ribosomal genes are GC rich in Solenogastres [46,47] and universal primers for mitochondrial genes do not work well on some aplacophorans [45]. Here, we employed a phylogenomic approach to reconstruct a broad-scale phylogenetic framework for Aplacophora. In the light of the reconstructed phylogenetic framework, including a newly described and highly unusual lineage, we assessed the monophyly of traditionally recognized aplacophoran taxa and implications for understanding early molluscan evolution.

## 2. Material and methods

### (a) Taxon sampling and morphological work

We aimed to sample transcriptome data from as many recognized aplacophoran taxa and as broad a range of morphological disparity as possible (electronic supplementary material, tables S1 and S2). The identification of specimens and

data collection for description of *Apodomenia enigmatica* sp. nov. involved examination of sclerites, radulae (if present) and internal anatomy following standard approaches of [48,49]. Scanning electron microscopy was conducted on dried, uncoated specimens using a Phenom Pro with an accelerating voltage of 5 kV. When possible, voucher specimens of species sampled herein were deposited into the University Museum of Bergen or the Alabama Museum of Natural History (see below).

### (b) Molecular techniques


Because prey nucleic acid contamination in solenogasters has been problematic in previous molecular studies [46,47], specimens were starved in the laboratory prior to preservation whenever possible (electronic supplementary material, table S1). Specimens of all taxa were preserved in RNAlater and stored at  $-80^{\circ}\text{C}$  or frozen at  $-80^{\circ}\text{C}$ . Different RNA extraction approaches were employed depending on sample size (electronic supplementary material, table S1). Total RNA concentration and purity were estimated using a NanoDrop 2000 (Thermo Scientific) and RNA quality was evaluated on a 1% SB agarose gel. For most taxa, cDNA library preparation and sequencing was performed as described in [50]. For *Chaetoderma nitidulum*, *Falcidens sagittiferus*, *Stylomenia sulcodoryata* and *Tonicella lineata*, total RNA was sent to Macrogen (South Korea) for Illumina stranded library preparation and sequencing using 1/4 lane of an Illumina HiSeq 2500 with  $2 \times 100$  bp paired-end sequencing.

### (c) Dataset assembly

For most taxa, digital normalization and assembly were performed as described in [50]. For taxa sequenced at Macrogen and publicly available *Rhyssoplax* and *Pholidoskepia* sp. (misidentified as *Chaetoderma* sp. by Zapata *et al.* [51]; see [52]), read trimming, digital normalization and assembly were performed using the 3/2014 version of TRINITY. Contigs from all taxa were translated with TRANSDCODER and translated sequences shorter than 50 amino acids (AAs) were deleted.

For orthology inference, we employed HAMStR 13 [53], using a custom core orthologue set based on transcriptome data from *Alexandromenia crassa*, *A. enigmatica*, *Helluoherpia aegiri*, *Leptoichiton* sp., *Neomenia carinata*, *Prochaetoderma californicum*, *Simrothiella margaritacea* and the genome of *Lottia gigantea* following [50]. In cases where one of the first or last 20 characters of an AA sequence was an X, all characters between the X and that end of the sequence were deleted and treated as missing data. Each gene was then aligned with MAFFT [54] and alignments were trimmed with ALISCORE [55] and ALICUT [56] to remove ambiguously aligned regions. A consensus sequence was inferred for each alignment using infoalign [57] and the percentage of positions of a sequence that differed from the consensus of the alignment were calculated using the infoalign’s ‘change’ calculation. Any sequence with a value greater than 75 was deleted. Sequence regions containing lesser than or equal to 20 AAs in length surrounded by 10 or more gaps on either side were deleted. We deleted sequences that did not overlap with all other sequences in the alignment by greater than or equal to 20 AAs, starting with the shortest sequence.

In some cases, a taxon was represented in an alignment by two or more sequences. We built trees in FASTTREE 2 [58] using the ‘slow’ option and used PHYLOTREEPRUNER [59] to select the best sequence for each taxon. Only genes sampled for 20+ taxa after pruning with PhyloTreePruner were retained. To further screen for paralogy and contamination, we used TRESPEx [60] to search for gene trees where select, well-established monophyletic groups (Conchifera, Polyplacophora, Pholidoskepia, Amphimeniidae, Neomeniidae and Prochaetodermatidae) were recovered non-monophyletic with strong support (bootstrap support greater than 95) and excluded those 12 genes from further



**Figure 1.** Phylogeny of Aplacophora based on 200 best genes in terms of branch-length heterogeneity. BI topology shown with posterior probabilities/RAxML/IQ-TREE bootstrap support values less than 1.0/100/100 shown at each node (see electronic supplementary material, figures S2–S7 for ML topologies). Coloured bars show the proportion of genes sampled for each taxon. (Online version in colour.)

consideration. This yielded a complete data matrix with 525 genes that was 75 914 AAs long with 30.43% missing data (electronic supplementary material, figure S1A). We also measured branch-length heterogeneity (LB) score as calculated in TRESpEX to identify genes most likely to be susceptible to long-branch attraction and assembled a data matrix with the best 200 genes according to LB, which was 30 185 AAs long with 30.7% missing data (electronic supplementary material, figure S1B).

#### (d) Phylogenetic analyses

Phylogenetic analyses were conducted for all data matrices using maximum likelihood (ML) in RAxML 7.3.8 [61] with the '-f a' flag, which specifies a search for best-scoring ML tree and a rapid bootstrap analysis in one program run. Each matrix was partitioned by gene and analysed with the PROTGAMMALGF model. Nodal support was assessed with 100 rapid bootstraps (-N 100).

ML analyses were also performed on all matrices in IQ-TREE [62] using the site-heterogeneous PMSF model [63] (-m LG + C60 + G + F) with the RAxML bipartitions tree provided as the required guide tree (-ft). Nodal support was assessed with 1000 rapid bootstraps (-bb 1000).

Bayesian inference (BI) analysis was conducted in PHYLOBAYES MPI 1.2f [64] with the site-heterogeneous CAT-GTR-G4 model. Because of the computational intensity of BI, only the matrix of the 200 least branch-length heterogeneous genes was analysed using this method. Four parallel chains were run for approximately 8000 cycles each with the first 2000 trees discarded as burn-in. A 50% majority rule consensus tree was computed from the remaining trees from each chain. PHYLOBAYES bpcomp

maxdiff of 0.1565 and meandiff of 0.0046 indicated that all chains had converged.

#### (e) DNA barcoding

We sequenced cytochrome *c* oxidase subunit I (COI) from specimens of *A. enigmatica* sp. nov. spanning its known geographical range following the laboratory approaches of [45] or by transcriptome sequencing as described above. ML-corrected substitutions per site were calculated in MEGA 7 using the maximum composite likelihood parameter with a  $\gamma$  parameter of 1.0 [65].

## 3. Results

#### (a) Phylogenetic analyses

Because aplacophorans have exhibited relatively long branches in previous phylogenomic studies [16,17] and *Falcidens caudatus* was on an extremely long branch in the ML analysis of all 525 genes (electronic supplementary material, figures S2 and S3), we conducted analyses of all 525 genes excluding *F. caudatus* (electronic supplementary material, figures S4 and S5) and sorted genes by LB as calculated in TRESpEX [60] and assembled and analysed a reduced dataset of just the 200 genes with the lowest branch-length heterogeneity (figure 1; electronic supplementary material, figures S6 and S7 and tables S3 and S4). Because analyses of the complete dataset (electronic supplementary material, figures S2 and S3) yielded similar results as that of the

reduced dataset, we focus our discussion on analyses of the reduced dataset and highlight notable differences when applicable. Details on data matrices analysed are presented in electronic supplementary material, tables S3 and S4.

Analyses of the dataset with reduced LB strongly supported Polyplacophora (BI posterior probability [pp]/RAxML bootstrap support [bs]/IQ-TREE bs = 1.00/100/100), Aplacophora (1.00/86/100), Solenogastres (1.00/100/100) and Caudofoveata (0.99/78/100). Within Solenogastres, Cavibelonia is polyphyletic. Amphimieniidae was recovered as the sister taxon of all other sampled lineages of Solenogastres with maximal support. The remaining cavibelonians along with the one sampled representative of Sterrofustia (*Phyllomenia* sp.), Neomeniamorpha and *A. enigmatica* sp. nov. formed a maximally supported clade, which was recovered as the sister taxon of Pholidoskepia. Within this clade, *Phyllomenia* formed a clade with Pruvotinidae with maximal support. The clade of Neomeniamorpha, *Phyllomenia* and Pruvotinidae was recovered as the sister group to a well-supported clade consisting of the remaining 'cavibelonian' taxa: *Epimения*, *Entonomenia*, Proneomeniidae and Simrothiellidae; support for placement of this clade was also variable (0.77/66/94).

We recovered Pholidoskepia monophyletic with full support in all analyses. Dondersiidae was recovered with maximal support in all analyses. However, relationships among families differed among analyses. In the BI analysis, Macellomeniidae was recovered sister to Dondersiidae with relatively strong support (pp = 0.98). Macellomeniidae was recovered sister to Gymnomeniidae in the ML analyses, but with weak support (electronic supplementary material, figures S6 and S7). Meiomeniidae was recovered in a clade with Gymnomeniidae with moderate support in BI (pp = 0.96) but as the sister taxon of all other pholidoskepians in ML with moderate to weak support (electronic supplementary material, figures S6 and S7).

Within Caudofoveata, we sampled at least one member of each recognized family and recovered a well-supported Chaetodermatidae (*Falcidens* + *Chaetoderma*) with maximal support (1.00/100). *Chaetoderma* was nested within *Falcidens* with *C. nitidulum* and *Falcidens caudatus* forming a clade with maximal support.

### (b) *Apodomenia enigmatica* sp. nov.

Distinguishing between the two major lineages of Aplacophora is generally straightforward: caudofoveates lack a foot but have an anterior muscular structure called the oral shield, whereas solenogastres have a narrow, midventral foot and lack an oral shield. However, during two recent Antarctic research expeditions, specimens of an aplacophoran, which lacks both a foot and an oral shield, were found inside *Rossella* sp. sponges (electronic supplementary material, table S5). We sequenced COI from six specimens (GenBank MK404651–MK404656) spanning the known geographical range of the species. Only 15 of 625 nucleotide positions in the amplified region were variable (ML-corrected substitutions per site = 0.008; electronic supplementary material, figure S8), suggesting that all of the sampled specimens belong to the same species.

Apodomeniidae fam. nov.

Diagnosis: cuticle thick, sclerites acicular and in one layer; foot reduced; radula and ventrolateral foregut glands lacking; spawning duct with extraepithelial gland cells.

*Apodomenia* gen. nov.

Diagnosis: sclerites solid acicular spines; radula and ventrolateral foregut glands lacking; foot lacking, foot groove covered by the cuticle and sclerites; common vestibulo-buccal opening; secondary genital opening unpaired; mantle cavity reduced.

Etymology: '*Apodo*' from *apodus* (lat.) 'lacking a foot'; '*menia*' is a common suffix for solenogaster genus names that is derived from '*-mene*' (gr.) referring to the moon or crescent.

*Apodomenia enigmatica* sp. nov.

Type species for *Apodomenia* gen. nov., by monotypy.


Diagnosis: Body up to about 16 cm long, slender and very stiff. Ventral groove and foot lacking. Cuticle thick, with robust mantle sclerites arranged in a right angle to body surface. Sclerites are flattened solid spines. Vestibulum with a few simple sensory papillae. Mouth opening within vestibulum. Radula and ventrolateral foregut glands lacking. Midgut with paired anteriodorsal caecum; without regular constrictions. Spawning ducts partly fused, ciliated and surrounded by extraepithelial gland cells, the cell bodies of which lie distally to a thick muscular coat. One pair of branched seminal vesicles. Mantle cavity highly reduced, lacking respiratory folds.

Type material: Holotype (ZMBN 129503): two histological section series (anterior + posterior). Paratype 1 (ZMBN 129501): one histological section series (anterior). Paratype 2 (ZMBN 129505): large specimen incomplete at posterior end, fixed in 4% formalin and preserved in 70% ethanol. Paratype 3 (ZMBN 129502): posterior end broken, anterior end dissected, fixed in 4% formalin and preserved in 70% ethanol. Paratype 4 (ALMNH 21269): one complete specimen broken at midbody, fixed in 4% formalin, stained with phosphomolybdic acid and preserved in 70% ethanol. Paratype 5 (ALMNH 21270): stained with phosphomolybdic acid, and preserved in 95% ethanol. Sample data for all specimens collected are presented in electronic supplementary material, table S5. Holotype and paratypes 1–3 are deposited in the University Museum of Bergen (ZMBN) and paratypes 5–6 are deposited in the Alabama Museum of Natural History (ALMNH).

Type locality: Wright's Gulf, Antarctica (73°17.7997 S, 129°11.5466 W) at 506 m in association with *Rossella* sp. Collected 25 January 2013.

Etymology: '*enigmatica*', from lat. *enigmaticus*, *-a*, *-um*, meaning mysterious, refers to the highly unusual morphology and lifestyle of the species.

Description: animals uniformly cylindrical-elongate, tending to curl up spirally when disturbed and during fixation. Largest specimen found (paratype 4; figure 2a) 155 mm long, with a maximum diameter of 8 mm; tip of posterior end missing. Animals completely covered in thick cuticle pierced by evenly sized massive, flattened spines. Spines arranged at a right angle to the body surface, resulting in an overall velvety appearance. Cuticle translucent and thus, on a closer look, body surface appears rather spiny even though only the tips of the spines protrude from the cuticle (figure 2b,c). In living animals, yellowish organs (gonad and midgut) and red hemolymph visible through the integument. Ventral furrow lacking, but sometimes the ventral side close to the anterior end appears slightly flattened. Areas with thin cuticle and distinctly smaller spines surrounding mouth and area around the anus and genital opening (figure 2d,e).




**Figure 2.** *Apodomenia enigmatica* sp. nov. (a) Habitus with (broken) posterior end above (paratype 4). Scale bar, 8 mm. (b) Scanning electron micrograph of section through the midbody of ALMNH 21271. Scale bar, 200  $\mu\text{m}$ . (c–e) Holotype, light microscopic images. (c) Cross-section through the body wall in the midgut region. Scale bars, 600  $\mu\text{m}$ . sc, sclerites embedded in the cuticle; bw, body wall. (d) Anterior body in a ventral view with vestibulum (v). (e) Posterior body in a ventral view with sclerite-free area where the anus (a) and gonopore (go) are situated (reduced mantle cavity). (f–h) Histological sections of holotype. (f) Anterior body with the pharynx (ph) and anterior midgut caecum (mg). Scale bar, 500  $\mu\text{m}$ . (g) Posterior body in the region anterior to the pericardium, with branched seminal receptacles (rs). Scale bar, 200  $\mu\text{m}$ . (h) Hindgut region showing the rectum (re) and paired spawning ducts (spd) surrounded by glands (gl). Scale bar, 200  $\mu\text{m}$ . cu, cuticle; de, detritus; gon, gonad; gpd, gonopericardioduct; mg, midgut; pc, pericardium; pd, pericardioduct; sc, sclerite. (Online version in colour.)

The following descriptions are based on the holotype, an adult specimen with an approximate length of 55 mm and a maximum body diameter of 4 mm. Epidermis 50–60  $\mu\text{m}$  thick, lacking glandular cells or papillae. Spines secreted by single enlarged epidermal cells, which elongate into the cuticle and lift the bases of spines. Cuticle up to 250  $\mu\text{m}$  thick. Animal generally uniform in thickness, but close to anterior and posterior body ends, the ventral cuticle is thinner. Epidermal sclerites are solid, flattened spines up to 800  $\mu\text{m}$  long and up to  $70 \times 30 \mu\text{m}$  at the base. Epidermis underlain by thick layers of circular and longitudinal musculature (figure 2f).

Figure 3 shows reconstructions of the anterior and posterior body regions of the holotype based on histology. The mouth opening is located in a small vestibulum, which bears a few papillae (folds). Foregut epithelium high and glandular (figure 2f). No foregut glands were observed. A radula is lacking. Pharynx slightly longer than the maximum height of the anterior body. Pharynx with muscular sheet and posteriorly constricted by strong circular musculature; narrow opening between the pharynx and midgut. Midgut wide and uniform, lined by large digestive and glandular

cells (figure 2f). Long, paired, antiodorsal caecum and a short antioventral caecum present. Midgut filling most of the long tubular body and, near the posterior body end, it narrowing to a short ciliated rectum. Anus posterior to the genital opening and surrounded by an area covered in thin cuticle and short sclerites. Remarkably, no mantle cavity is present.

The dorsal paired gonad is well developed, holding both oocytes and spermatocytes. The gonad reaching to the anterior body end, dorsally to the midgut caecum; the median gonad walls fused (figure 2f). Pericardioducts (figure 2g) short and paired; distinctly ciliated. They fuse just anterior to the relatively narrow pericardium (figure 2h), which contains a large, muscular heart ventricle. Short pericardioducts connecting to voluminous spawning ducts that run posteriorly and fuse with each other ventrally to the rectum. Paired seminal receptacles consisting of long and slender ciliated ducts, which anteriorly branch into a number of chambers (figure 2g). Both paired and fused parts of the spawning duct lined with ciliated epithelium and surrounded by a thick coat of extraepithelial gland cells, the cell bodies of which come to lie outside a strong



**Figure 3.** Lateral reconstruction of the internal anatomy of the holotype of *A. enigmatica* sp. nov. (a) Anterior body. The anteriormost part of the nervous system (cer?) was ambiguous. (b) Posterior body. A depression in the cuticle dorsally to the anus could be interpreted as a dorsoterminal sense organ (dts?), but this is doubtful. a, anus; cu, cuticle; gpd, gonopercardioduct; go, gonopore; gon, gonade; ln, lateral nerve cord; mg, midgut; pc, pericardium; pd, pericardioduct; pgl, pedal gland; ph, pharynx; phw, pharynx wall; r, rectum; rs, seminal receptacle; spd, spawning duct; v, vestibulum; vbo, vestibular opening; ve, heart ventricle; vn, ventral nerve cord. (Online version in colour.)

circular muscle layer (figure 2h). Single gonopore situated just in front of anal opening and surrounded by an area covered in thin cuticle and small sclerites.

Four major nerve cords run through the entire body, a ventral and a lateral pair. At the posterior end, the ventral and lateral chords of each side are joined by connectives. Reconstruction of the anterior nervous system (including a cerebral ganglion) was not possible based on the two section series available.

## 4. Discussion

Here, we present a phylogenetic framework for Aplacophora that differs dramatically from the current taxonomy of the group and describe a remarkable new solenogaster lacking most of the characters traditionally used to diagnose

Mollusca, significantly expanding known morphological variation in Aplacophora. The unusual morphology of *Apodomenia* initially led us to hypothesize that it represents a 'missing link' between Solenogastres and Caudofoveata. However, all analyses firmly place this species well within Solenogastres. Thus, our results indicate that the foot was secondarily lost at least twice in aplacophoran evolution. Some Palaeozoic chiton-like animals lacking a foot [37–40] have been hypothesized to be stem-group caudofoveates [66,67]. As these animals had chiton-like shells, this hypothesis would suggest independent loss of shells in Caudofoveata and Solenogastres. Although we agree that the available evidence support a chiton-like ancestor for Aplacophora [20,68], independent reduction in the foot in Caudofoveata and *Apodomenia* raises the possibility that Palaeozoic chiton-like taxa without a foot represent additional independent losses. In addition to lacking a foot, *A. enigmatica* is without a mantle cavity or radula, making it among the most extreme deviations from the 'hypothetical archetypal mollusc' [69] known. *Apodomenia enigmatica* sp. nov. demonstrates the striking plasticity of the aplacophoran body plan despite the superficially uniform (worm-shaped) appearance of many members of the group.

Within Solenogastres, we show that several traditionally recognized higher-level taxa (e.g. Aplotegmentaria, Pachytegumentaria and Cavibelonia) are not monophyletic. Cavibelonia was originally defined by the presence of hollow, acicular sclerites [28]. However, some cavibelonians have a scleritome combining scales with hollow acicular sclerites (e.g. Acanthomeniidae) and other species have solid, flattened sclerites (e.g. *Helicoradomenia* spp. [70]). Other characters used in solenogaster taxonomy, such as the radula and ventrolateral foregut glands, are quite variable among taxa ascribed to Cavibelonia. Thus, recovering this clade as polyphyletic was not shocking. Notably, even Salvini-Plawen, who erected the group, expressed his doubts about its validity [8]. Our results are consistent with either multiple independent origins of hollow sclerites (in Amphimeniidae, Pruvotinidae and the last common ancestor of the Epimeniidae/Rhopalomeniidae/Pruvotinidae/Simrothielidae clade) as hypothesized by Salvini-Plawen [8] or multiple independent losses of hollow sclerites (in Neomeniamorpha, Pholidoskepia, Sterrofustia and *Apodomenia*). All solenogasters have solid scales (at least along the foot and around the dorsoterminal sensory organ, if present) and, at least in *Epimonia* and *Proneomenia*, solid scales cover the body of post-larval animals and are later replaced by hollow sclerites [48,71]. We hypothesize that hollow acicular sclerites were present in the last common ancestor of Solenogastres and were modified independently in pholidoskepians, whose scale-like sclerites were likely selected for as an adaptation to a meiofaunal lifestyle, neomeniids, whose harpoon-shaped sclerites appear to grow via a slight modification of the developmental program that produces hollow sclerites in cavibelonians.

Smith *et al.* [17] sequenced an unidentified species of Solenogastres from Greenland. We recollected this species from the same locality and identified it by histology as a pruvotiniid (ZMBN 129506–129508). Pruvotinidae was recovered as the sister taxon to the one sampled representative of Sterrofustia, *Phyllomenia*. Sterrofustia is distinguished from the cavibelonian family Pruvotinidae exclusively by the presence of solid sclerites. Pruvotinidae is otherwise a large,

diverse group with species that span a wide range of morphological variation (summarized by García-Alvarez & Salvini-Plawen [24]). Notably, the meiofaunal *Hypomenia sanjuanensis* exhibits a continuum of sclerites with internal cavities ranging in size from those with a cavity that fills around half the volume of the sclerite to those with no hollow cavity at all [49]. Thus, we view the status of *Sterrofustia* as an order within Solenogastres to be questionable.

Pholidoskepia has been viewed as the extant lineage of Solenogastres with the most plesiomorphic morphological characters [5]. This, combined with the hypothesis that Solenogastres is the sister group to all other Mollusca (e.g. [14]), which is now generally rejected, may have prompted the hypothesis that the last common ancestor of Mollusca was a small, pholidoskepien-like animal [23]. Our results placing large-bodied taxa throughout Solenogastres and Pholidoskepia on a relatively long branch are more consistent with recent work in suggesting the last common ancestor was a relatively large-bodied, chiton-like animal [16] and that the mostly small-bodied Pholidoskepia are relatively derived [72].

We sampled four of the six currently recognized families of Pholidoskepia, and recovered the group monophyletic with strong support. All relationships were strongly supported in BI but placement of Meiomeniidae and Macellomeniidae were weakly supported in ML. Gymnomeniidae has been thought to be closely related to Meiomeniidae as the two families are distinguished almost exclusively on the basis of body size and the number of different sclerite types present [24]. Characters shared by these two taxa include the pedal commissure sac (a unique statocyst-like, geotactic sense organ), an almost complete lack of a basal lamina in the epidermis and a very thin cuticle together resulting in a very fragile integument, ventrolateral foregut glands lacking ducts and the persistence of protonephridia in postlarval or even adult animals [73]. Interestingly, a pedal commissure sac has recently also been found in a meiofaunal dondersiid species [74]. Strong support for a clade of *Macellomenia* and Dondersiidae from BI makes sense in the light of morphology (e.g. same radula type in both families). Sampling of additional members of Pholidoskepia will hopefully help to resolve this issue in the future.

Our results may also shed light on earlier discussions on the plesiomorphic radula type of solenogasters, aplacophorans and molluscs in general. Eernisse & Kerth [75] and Scheltema *et al.* [76] suggested a bipartite (distichous) radula with a medially split radula membrane and two radula teeth or plates in each row to represent the ancestral state. This viewpoint was based on preliminary results on the fossil *Wiwaxia corrugata* and on ontogenetic data for selected chiton and gastropod species. Scheltema [77] later included new fossil findings into her updated interpretation and suggested that a unipartite radula (radula membrane not medially split) with an unpaired central rhachidian tooth and several teeth per row most probably represents the plesiomorphic state for Mollusca. The lack of a rhachidian tooth in aplacophorans is thus interpreted as a derived character. Most interestingly, several early-branching solenogaster clades in our trees do have a unipartite radula, where the single tooth could be homologous to a rhachidian tooth. This includes Amphimeniidae as well as Dondersiidae and Macellomeniidae within Pholidoskepia. Members of Proneomeniidae also have a monopartite radula, but with numerous teeth attached to the radular membrane (polystichous radula). This radula type appears most similar to the radula

of other molluscs with a rasping radula, but there seems to be some variation concerning the presence of an unpaired central tooth. Considering the placement of Proneomeniidae, the polystichous radula is unlikely to be a plesiomorphy for Solenogastres. Complete radula reduction can be found not only in *Apodomenia* sp. nov., but in various groups, including all Neomeniidae and many Dondersiidae.

Within Caudofoveata, Limifossoridae exhibits a putatively plesiomorphic distichous radula and a simple body shape [25–27]. However, our results place Prochaetodermatidae sister to Limifossoridae + Chaetodermatidae, consistent with recent studies [44,45]. Prochaetodermatids are small, mostly deep-sea aplacophorans that differ from other caudofoveates by having a paired oral shield, a pair of cuticular jaws, and a radula with two lateral teeth and an undivided radular membrane with a central plate. Interestingly, the long branches separating Prochaetodermatidae and Chaetodermatidae + Limifossoridae show substantial genetic divergence between the two clades. Our results also confirm earlier results indicating that *Chaetoderma* is nested within *Falcidens* [43,44].

## 5. Conclusion

Our results have significantly altered understanding of the evolutionary history and morphological diversity of Aplacophora. Molecular phylogenetics practically turns upside-down previous hypotheses of phylogenetic relationships in both Solenogastres (a large-bodied cavibelonian taxon as the sister group to all other solenogasters) and Caudofoveata (Prochaetodermatidae and not Limifossoridae as sister to all other caudofoveates). Especially in Solenogastres, our results are consistent with a shift from support for the Testaria hypothesis (small-sized pholidoskepien taxa display the most ancestral morphology within Mollusca [22]) to the Aculifera hypothesis (ancestral molluscs were relatively large-bodied, polyplacophoran-like animals [4]). Consequently, evolution of recent aplacophoran molluscs appears to have included several steps of reduction in morphological characters, including the shell(s), digestive gland, broad rasping radula and kidney. Even more extreme reduction is observed in the anomalous *Apodomenia*, which lacks all major characters usually used to define Mollusca.

In addition to advancing understanding of aplacophoran phylogeny, we have dramatically expanded on the previously limited amount of molecular sequence data from aculiferan molluscs by producing deeply sequenced, high-quality Illumina transcriptomes. Our hope is that these data will be of use to researchers addressing a wide variety of questions. We are optimistic that future studies with improved taxon sampling of key lineages not sampled herein (e.g. Acanthomeniidae) will continue to provide insight into the phylogeny and evolution of Aplacophora and Aculifera, thereby shedding more light on the early evolution of Mollusca as a whole.

**Competing interests.** We declare we have no competing interests.

**Funding.** This study was supported in part by awards from the US National Science Foundation to K.M.K. (DEB-1210518 and DBI-1306538) and K.M.H. (DEB-1036537, OCE-1155188, ANT-1043745) and an award from the Norwegian Taxonomy Initiative (project no. 70184222) to C.T.

**Data accessibility.** Data available as part of the electronic supplementary material.

**Acknowledgements.** We thank Akiko Okusu for sharing specimens of *Epimienia babai* and Trish Morse for assistance with collecting meiofaunal solenogasters from Friday Harbor and sharing specimens. We thank the crew and scientists of BioSkag II aboard R/V *Håkon Mosby*, the crew of R/V *Hans Brattström* (University of Bergen), the crew and scientists of the Icy Inverts cruises aboard R/V *Lawrence M. Gould* and *Nathaniel B. Palmer*, the crew and scientists of the IceAGE cruises aboard R/V *Meteor* and R/V *Poseidon*, the crew and scientists of the BoWls cruises aboard R/V *Oceanus*, the crew and volunteers of the University of Washington Friday Harbor Laboratories R/V *Centennial*, and the FHL administration and staff for

supporting specimen collection. We thank Elena Gerasimova who trained KMK in histology and helped with the histological sectioning of specimens. We thank Andrea Kohn and Pam Brannock for help with RNA extraction. We thank M. Carmen Cobo Llovo, Maddie McCutcheon, Rebecca Varney and Meghan Yap-Chiongco for helpful comments on an earlier version of this manuscript. Finally, we thank our departed colleagues Christoffer Schander, Amélie Scheltema and Luitfried von Salvini-Plawen for helpful discussions and inspiration that led to this work. This is Auburn University Marine Biology Program contribution no. 186 and Mollette Lab contribution no. 90.

## References

- Tullberg T. 1875 Neomenia: a new genus of invertebrate animals. *K. Svenska Vetens. Akad. Handlingar* **13**, 1–12.
- Lovén S. 1844 Chaetoderma, ett nytt maskslägte. *Öfversigt af Kongl Vetenskaps-akademiens förhandlingar* **1**, 116.
- Scheltema AH. 1978 Position of the class Aplacophora in the phylum Mollusca. *Malacologia* **17**, 99–109.
- Scheltema AH. 1993 Aplacophora as progenetic aculiferans and the coelomate origin of molluscs as the sister taxon of Sipuncula. *Biol. Bull.* **184**, 57–78. (doi:10.2307/1542380)
- Salvini-Plawen L. 2003 On the phylogenetic significance of the aplacophoran Mollusca. *Iberus* **21**, 67–97.
- Todt C. 2013 Aplacophoran mollusks—still obscure and difficult? *Am. Malacol. Bull.* **31**, 1–7. (doi:10.4003/006.031.0110)
- Todt C, Okusu A, Schander C, Schwabe E. 2008 Solenogastres, Caudofoveata, and Polyplacophora. In *Phylogeny and evolution of the Mollusca* (eds WF Ponder, DR Lindberg), pp. 71–96.
- Salvini-Plawen L. 1985 Early evolution and the primitive groups. In *The Mollusca* (eds E Trueman, M Clarke), pp. 59–150. Orlando, FL: Academic Press.
- Scheltema AH. 1996 Phylogenetic position of Sipuncula, Mollusca and the progenetic Aplacophora. In *Origin and evolutionary radiation of the mollusca*, pp. 53–58. Oxford, UK: Oxford University Press.
- Ivanov DL. 1996 Origin of Aculifera and problems of monophyly of higher taxa in molluscs. In *Origin and evolutionary radiation of the Mollusca* (ed. JD Taylor), pp. 59–65. New York, NY: Oxford University Press.
- Salvini-Plawen L. 1980 A reconsideration of systematics in the Mollusca (phylogeny and higher classification). *Malacologia* **19**, 249–278.
- Salvini-Plawen L. 1981 On the origin and evolution of the Mollusca. *Atti Conv. Lincei* **49**, 235–293.
- Salvini-Plawen L, Steiner G. 1996 Synapomorphies and plesiomorphies in higher classification of Mollusca. In *Origin and evolutionary radiation of the Mollusca* (ed. JD Taylor), pp. 29–51. Oxford, UK: Oxford University Press.
- Haszprunar G. 2000 Is the Aplacophora monophyletic? A cladistic point of view. *Am. Malacol. Bull.* **15**, 115–130.
- Kocot KM. 2013 Recent advances and unanswered questions in deep molluscan phylogenetics. *Am. Malacol. Bull.* **31**, 1–14. (doi:10.4003/006.031.0112)
- Kocot KM *et al.* 2011 Phylogenomics reveals deep molluscan relationships. *Nature* **477**, 452–456. (doi:10.1038/nature10382)
- Smith SA, Wilson NG, Goetz FE, Feehery C, Andrade SCS, Rouse GW, Giribet G, Dunn CW. 2011 Resolving the evolutionary relationships of molluscs with phylogenomic tools. *Nature* **480**, 364–367. (doi:10.1038/nature10526)
- Vinther J, Sperling EA, Briggs DEG, Peterson KJ. 2012 A molecular palaeobiological hypothesis for the origin of aplacophoran molluscs and their derivation from chiton-like ancestors. *Proc. R. Soc. B* **279**, 1259–1268. (doi:10.1098/rspb.2011.1773)
- Vinther J, Parry L, Briggs DEG, Van Roy P. 2017 Ancestral morphology of crown-group molluscs revealed by a new Ordovician stem aculiferan. *Nature* **542**, 471. (doi:10.1038/nature21055)
- Scherholz M, Redl E, Wollesen T, Todt C, Wanninger A. 2013 Aplacophoran mollusks evolved from ancestors with polyplacophoran-like features. *Curr. Biol.* **23**, 2130–2134. (doi:10.1016/j.cub.2013.08.056)
- Scherholz M, Redl E, Wollesen T, Todt C, Wanninger A. 2015 From complex to simple: myogenesis in an aplacophoran mollusk reveals key traits in aculiferan evolution. *BMC Evol. Biol.* **15**, 201. (doi:10.1186/s12862-015-0467-1)
- Salvini-Plawen LV, Steiner G. 2014 The Testaria concept (Polyplacophora + Conchifera) updated. *J. Nat. Hist.* **48**, 2751–2772. (doi:10.1080/00222933.2014.964787)
- Haszprunar G, Schander C, Halaných K. 2008 Relationships of higher molluscan taxa. In *Phylogeny and evolution of the Mollusca* (eds WF Ponder, DR Lindberg), pp. 19–32. Berkeley, CA: University of California Press.
- García-Álvarez O, Salvini-Plawen L. 2007 Species and diagnosis of the families and genera of Solenogastres (Mollusca). *Iberus* **25**, 73–143.
- Salvini-Plawen L. 1972 Zur Morphologie und Phylogenie der Mollusken: Die Beziehungen der Caudofoveata und der Solenogastres als Aculifera, als Mollusca und als Spiralia. *Z. Wiss. Zool.* **184**, 205–394. (doi:10.1007/bf00406755)
- Wolter K. 1992 Ultrastructure of the radula apparatus in some species of aplacophoran molluscs. *J. Mollusc. Stud.* **58**, 245–256. (doi:10.1093/mollusc/58.3.245)
- Ivanov D. 1986 New species of Caudofoveata from the Tchukto Sea (Mollusca, Aplacophora). *Arch. Zool. Mus. Moscow State Univ.* **24**, 83–95.
- Salvini-Plawen L. 1978 Antarktische und subantarktische Solenogastres. *Zoologica* **44**, 1–315.
- Handl CH, Todt C. 2005 Foregut glands of Solenogastres (Mollusca): anatomy and revised terminology. *J. Morphol.* **265**, 28–42. (doi:10.1002/jmor.10336)
- Salvini-Plawen L. 1985 New interstitial Solenogastres (Mollusca). *Stylogologia* **1**, 101–108.
- Haszprunar G. 1992 The first molluscs—small animals. *Ital. J. Zool.* **59**, 1–16.
- Puchalski SS, Eernisse DJ, Johnson CC. 2008 The effect of sampling bias on the fossil record of chitons (Mollusca, Polyplacophora). *Am. Malacol. Bull.* **25**, 87–95. (doi:10.4003/0740-2783-25.1.87)
- Okusu A, Schwabe E, Eernisse DJ, Giribet G. 2003 Towards a phylogeny of chitons (Mollusca, Polyplacophora) based on combined analysis of five molecular loci. *Org. Divers. Evol.* **3**, 281–302. (doi:10.1078/1439-6092-00085)
- Sigwart JD, Schwabe E, Saito H, Samadi S, Giribet G. 2011 Evolution in the deep sea: a combined analysis of the earliest diverging living chitons (Mollusca: Polyplacophora: Lepidopleurida). *Invertebr. Syst.* **24**, 560–572. (doi:10.1071/IS10028)
- Sigwart JD, Stoeger I, Knebelberger T, Schwabe E. 2013 Chiton phylogeny (Mollusca: Polyplacophora) and the placement of the enigmatic species *Chorioplax grayi* (H. Adams & Angus). *Invertebr. Syst.* **27**, 603–621. (doi:10.1071/IS13013)
- Irisarri I, Eernisse DJ, Zardoya R. 2014 Molecular phylogeny of Acanthochitonina (Mollusca: Polyplacophora: Chitonida): three new mitochondrial genomes, rearranged gene orders and systematics. *J. Nat. Hist.* **48**, 2825–2853. (doi:10.1080/00222933.2014.963721)
- Sutton MD. 2001 A three-dimensionally preserved fossil polychaete worm from the Silurian of Herefordshire, England. *Proc. R. Soc. Lond. B* **268**, 2355–2363. (doi:10.1098/rspb.2001.1788)
- Sutton MD, Briggs DEG, Siveter DJ, Siveter DJ. 2004 Computer reconstruction and analysis of the vermiform



- mollusc *Acaenoplax hayae* from the Herefordshire Lagerstätte (Silurian, England), and implications for molluscan phylogeny. *Palaeontology* **47**, 293–318. (doi:10.1111/j.0031-0239.2004.00374.x)
39. Sutton MD, Briggs DEG, Siveter DJ, Siveter DJ, Sigwart JD. 2012 A Silurian armoured aplacophoran and implications for molluscan phylogeny. *Nature* **490**, 94–97. (doi:10.1038/nature11328)
  40. Sutton MD, Sigwart JD. 2012 A chiton without a foot. *Palaeontology* **55**, 401–411. (doi:10.1111/j.1475-4983.2011.01126.x)
  41. Scheltema AH, Schander C. 2000 Discrimination and phylogeny of solenogaster species through the morphology of hard parts (Mollusca, Aplacophora, Neomeniomorpha). *Biol. Bull.* **198**, 121–151. (doi:10.2307/1542810)
  42. Scheltema AH, Schander C, Kocot KM. 2012 Hard and soft anatomy in two genera of Dondersiidae (Mollusca, Aplacophora, Solenogastres). *Biol. Bull.* **222**, 233–269. (doi:10.1086/BBLv222n3p233)
  43. Mikkelsen NT, Todt C. 2018 One or many? Molecular versus morphological diversity in the aplacophoran *Chaetoderma nitidulum* Lovén, 1844 (Mollusca: Caudofoveata). *J. Mollusc. Stud.* **84**, 113–131. (doi:10.1093/mollusc/eyy009)
  44. Mikkelsen NT, Kocot KM, Halanych KM. 2018 Mitogenomics reveals phylogenetic relationships of caudofoveate aplacophoran molluscs. *Mol. Phylogenet. Evol.* **127**, 429–436. (doi:10.1016/j.ympev.2018.04.031)
  45. Mikkelsen NT, Todt C, Kocot KM, Halanych KM, Willassen E. 2019 Molecular phylogeny of Caudofoveata (Mollusca) challenges traditional views. *Mol. Phylogenet. Evol.* **132**, 138–150. (doi:10.1016/j.ympev.2018.10.037)
  46. Okusu A, Giribet G. 2003 New 18S rRNA sequences from neomenioid aplacophorans and the possible origin of persistent exogenous contamination. *J. Mollusc. Stud.* **69**, 385. (doi:10.1093/mollusc/69.4.385)
  47. Meyer A, Todt C, Mikkelsen N, Lieb B. 2010 Fast evolving 18S rRNA sequences from Solenogastres (Mollusca) resist standard PCR amplification and give new insights into mollusk substitution rate heterogeneity. *BMC Evol. Biol.* **10**, 70. (doi:10.1186/1471-2148-10-70)
  48. Todt C, Kocot KM. 2014 New records for the solenogaster *Proneomenia sluiteri* (Mollusca) from Icelandic waters and description of *Proneomenia custodiens* sp. n. *Pol. Polar Res.* **35**, 291–310. (doi:10.2478/popore-2014-0012)
  49. Kocot KM, Todt C. 2014 Three new meiofaunal solenogaster species (Mollusca: Aplacophora) from the north-east Pacific. *J. Nat. Hist.* **48**, 3007–3031. (doi:10.1080/00222933.2014.961987)
  50. Kocot KM *et al.* 2017 Phylogenomics of Lophotrochozoa with consideration of systematic error. *Syst. Biol.* **66**, 256–282.
  51. Zapata F, Wilson NG, Howison M, Andrade S.C.S, Jörgen KM, Schrödl M, Goetz FE, Giribet G, Dunn CW. 2014 Phylogenomic analyses of deep gastropod relationships reject Orthogastropoda. *Proc. R. Soc. B* **281**, 20141739. (doi:10.1098/rspb.2014.1739)
  52. Zapata F, Wilson NG, Howison M, Andrade SC, Jörgen KM, Schrödl M, Goetz FE, Giribet G, Dunn CW. 2015 Correction to Phylogenomic analyses of deep gastropod relationships reject Orthogastropoda. *Proc. R. Soc. B* **282**, 20142941. (doi:10.1098/rspb.2014.2941)
  53. Ebersberger I, Strauss S, Von Haeseler A. 2009 HaMStR: profile hidden markov model based search for orthologs in ESTs. *BMC Evol. Biol.* **9**, 157. (doi:10.1186/1471-2148-9-157)
  54. Katoh K, Kuma K-I, Toh H, Miyata T. 2005 MAFFT version 5: improvement in accuracy of multiple sequence alignment. *Nucleic Acids Res.* **33**, 511–518. (doi:10.1093/nar/gki198)
  55. Misof B, Misof K. 2009 A Monte Carlo approach successfully identifies randomness in multiple sequence alignments: a more objective means of data exclusion. *Syst. Biol.* **58**, 21–34. (doi:10.1093/sysbio/syp006)
  56. Kück P. 2009 *ALICUT: a perlScript which cuts ALISCORE identified RSS*. Version 2. Bonn, Germany: Department of Bioinformatics, Zoologisches Forschungsmuseum A Koenig (ZFMK).
  57. Rice P, Longden I, Bleasby A. 2000 EMBOS: the European molecular biology open software suite. *Trends Genet.* **16**, 276–277. (doi:10.1016/S0168-9525(00)02024-2)
  58. Price MN, Dehal PS, Arkin AP. 2010 FastTree 2—approximately maximum-likelihood trees for large alignments. *PLoS ONE* **5**, e9490. (doi:10.1371/journal.pone.0009490)
  59. Kocot KM, Citarella MR, Moroz LL, Halanych KM. 2013 PhyloTreePruner: a phylogenetic tree-based approach for selection of orthologous sequences for phylogenomics. *Evol. Bioinform.* **9**, 429. (doi:10.4137/EBO.S12813)
  60. Struck TH. 2014 TreSpEx—detection of misleading signal in phylogenetic reconstructions based on tree information. *Evol. Bioinform.* **10**, S14239. (doi:10.4137/EBO.S14239)
  61. Stamatakis A. 2006 RAXML-VI-HPC: maximum likelihood-based phylogenetic analyses with thousands of taxa and mixed models. *Bioinformatics* **22**, 2688. (doi:10.1093/bioinformatics/bt1446)
  62. Nguyen L-T, Schmidt HA, Haeseler AV, Minh BQ. 2015 IQ-TREE: a fast and effective stochastic algorithm for estimating maximum-likelihood phylogenies. *Mol. Biol. Evol.* **32**, 268–274. (doi:10.1093/molbev/msu300)
  63. Wang H-C, Minh BQ, Susko E, Roger AJ. 2018 Modeling site heterogeneity with posterior mean site frequency profiles accelerates accurate phylogenomic estimation. *Syst. Biol.* **67**, 216–235. (doi:10.1093/sysbio/syx068)
  64. Lartillot N, Lepage T, Blanquart S. 2009 PhyloBayes 3: a Bayesian software package for phylogenetic reconstruction and molecular dating. *Bioinformatics* **25**, 2286. (doi:10.1093/bioinformatics/btp368)
  65. Kumar S, Nei M, Dudley J, Tamura K. 2008 MEGA: a biologist-centric software for evolutionary analysis of DNA and protein sequences. *Brief. Bioinform.* **9**, 299–306. (doi:10.1093/bib/bbn017)
  66. Vinther J. 2014 A molecular palaeobiological perspective on aculiferan evolution. *J. Nat. Hist.* **48**, 2805–2823. (doi:10.1080/00222933.2014.963185)
  67. Vinther J. 2015 The origins of molluscs. *Palaeontology* **58**, 19–34. (doi:10.1111/pala.12140)
  68. Scheltema AH, Ivanov DL. 2002 An aplacophoran postlarva with iterated dorsal groups of spicules and skeletal similarities to Paleozoic fossils. *Invertebr. Biol.* **121**, 1–10. (doi:10.1111/j.1744-7410.2002.tb00124.x)
  69. Lindberg DR, Ghiselin MT. 2003 Fact, theory and tradition in the study of molluscan origins. *Proc. Calif. Acad. Sci.* **54**, 663–686.
  70. Kingsley R, Froelich J, Marks C, Spicer L, Todt C. 2012 Formation and morphology of epidermal sclerites from a deep-sea hydrothermal vent solenogaster *Helicoradomenia* sp. (Solenogastres, Mollusca). *Zoomorphology* **132**, 1–9. (doi:10.1007/s00435-012-0168-x)
  71. Okusu A. 2002 Embryogenesis and development of *Epimenia babai* (Mollusca Neomeniomorpha). *Biol. Bull.* **203**, 87–103. (doi:10.2307/1543461)
  72. Bergmeier FS, Haszpruner G, Todt C, Jörgen KM. 2016 Lost in a taxonomic Bermuda Triangle: comparative 3D-microanatomy of cryptic mesopsammic Solenogastres (Mollusca). *Org. Divers. Evol.* **16**, 613–639. (doi:10.1007/s13127-016-0266-6)
  73. Todt C, Wanninger A. 2010 Of tests, trochs, shells, and spicules: development of the basal mollusk *Wirenia argentea* (Solenogastres) and its bearing on the evolution of trochozoan larval key features. *Front. Zool.* **7**, 6. (doi:10.1186/1742-9994-7-6)
  74. Klink SP, Bergmeier FS, Neusser TP, Jörgen KM. 2015 Stranded on a lonely island: description of *Dondersia(?)* Todtae sp. nov., the first shelf solenogaster (Mollusca, Aplacophora) from the Azores. *Açoreana* **10**, 603–618.
  75. Eernisse DJ, Kerth K. 1988 The initial stages of radular formation in chitons (Mollusca: Polyplacophora). *Malacologia* **28**, 95–103.
  76. Scheltema AH, Kerth K, Kuzirian AM. 2003 Original molluscan radula: comparisons among Aplacophora, Polyplacophora, Gastropoda, and the Cambrian fossil *Wiwaxia corrugata*. *J. Morphol.* **257**, 219–245. (doi:10.1002/jmor.10121)
  77. Scheltema AH. 2014 The original molluscan radula and progenesis in Aplacophora revisited. *J. Nat. Hist.* **48**, 2855–2869. (doi:10.1080/00222933.2014.959573)