

Seroprevalence of Antibodies against *Anisakis simplex* Larvae among Health-Examined Residents in Three Hospitals of Southern Parts of Korea

Jung Kim¹, Jin Ok Jo¹, Seon Hee Choi², Min Kyoung Cho², Hak Sun Yu², Hee Jae Cha¹ and Meesun Ock^{1,*}

¹Department of Parasitology and Genetics, Kosin University College of Medicine, Busan 602-702, Korea; ²Department of Parasitology, Pusan National University School of Medicine, Busan 602-739, Korea

Abstract: The present study was performed to estimate the seroprevalence of larval *Anisakis simplex* infection among the residents health-examined in 3 hospitals in southern parts of Korea. A total of 498 serum samples (1 serum per person) were collected in 3 hospitals in Busan Metropolitan city, Masan city, and Geoje city in Gyeongsangnam-do (Province) and were examined by IgE-ELISA and IgE-western blotting with larval *A. simplex* crude extract and excretory-secretory products (ESP). The prevalence of antibody positivity was 5.0% and 6.6% with ELISA against crude extracts and ESP, respectively. It was also revealed that infection occurred throughout all age groups and higher in females than in males. A specific protein band of 130 kDa was detected from 10 patients with western blot analysis against crude extract and ESP among those who showed positive results by ELISA. Our study showed for the first time the seroprevalence of anisakiasis in Korea. The allergen of 130 kDa can be a candidate for serologic diagnosis of anisakiasis.

Key words: *Anisakis simplex*, seroepidemiology, excretory-secretory product, crude extract, ELISA, western blotting

INTRODUCTION

The nematode *Anisakis simplex* is a representative parasite for marine mammals. Almost all marine fish and cephalopods can become infected with the third stage larva (L3) of *A. simplex*. Human infections occur upon the ingestion of marine fish or cephalopods. The ingested *A. simplex* larvae by humans penetrate the stomach wall causing acute abdominal pain, nausea, and vomiting within a few hours. When the larvae invade the gastric or intestinal mucosa, inflammatory reaction often results in ulcer or eosinophilic granuloma. Recently, it has become clear that anisakiasis is often associated with strong allergic reactions ranging from urticaria to anaphylactic shock [1-4].

Chai et al. [5] reported that the prevalence of anisakiasis has remarkably increased throughout the world in the last 30 years. One of the main reasons for this increase is attributed to the preference for raw and slightly cooked food. This trend will bring about the rise of infectious diseases caused by parasitic

infections, like *A. simplex* larvae, through marine fishery products. Although anisakiasis might occur in any country where the people eat raw or undercooked fish or squids, the majority of cases have been diagnosed in Korea, Japan, Spain, and some other countries because of their eating habits [6-9].

The prevalence of *A. simplex* infection has been reported to be different depending on the countries and areas. Several studies have reported that more than 10% of gastrointestinal anisakiasis showed allergic symptoms related to fish consumption. The anisakid infection rate in Spain was known to be around 0.43% in Galicia area and 12.4% in the population of Madrid [10,11]. An epidemiological study in Japan has shown that anisakiasis was more frequent in coastal populations and among 20- to 50-year-old males. In addition, these patients were reported to engage in fishing industry or inhabit coastal areas [12]. The high prevalence of *A. simplex* larval infection of fish and cephalopods has been reported in various fish species and squids [13-15]. Favored fishes of the Korean people, such as mackerels, cods, Alaska pollacks, scabbard fish, and squids were heavily infected with *A. simplex* L3 [16,17]. These reports suggested possible association between fish consumption and allergic responses in Korean people. Nevertheless, seroepidemiologic surveys among Koreans have not been accomplished.

To obtain data of seroprevalence against *A. simplex* allergens

•Received 10 December 2010, revised 3 April 2011, accepted 11 April 2011.

*Corresponding author (msoc@ns.kosinmed.or.kr)

© 2011, Korean Society for Parasitology

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

among Koreans, we analyzed blood samples from residents in the southern parts of Korea by ELISA and Western blot analysis using crude extract and excretory-secretory products (ESP) from *A. simplex* L3.

MATERIALS AND METHODS

Subjects of investigation

We prepared 498 sera from blood samples collected in 3 hospitals, each located in Busan Metropolitan city, Masan city, and Geoje city. The study population were selected from the patients admitted for health examinations. They had no history of allergic symptoms, including allergic rhinitis, urticaria, atopic dermatitis, and asthma (total IgE < 100 IU/ml). Serum samples were obtained by clotting and centrifugation of the blood at room temperature and were stored at -70°C. The collection of sera for use in these studies was approved by the Human Subjects Investigation Committee of Kosin University, Busan, Korea.

Preparation of *Anisakis* antigens

A. simplex L3 larvae were collected manually from the viscera, flesh, and body cavities of naturally infected mackerels (*Scomber japonicus*) and thoroughly washed with PBS. The crude extract of *A. simplex* L3 larvae was prepared from 300 larvae (about 1 g). Larvae were frozen in liquid nitrogen and smashed by mortar. To extract proper amounts of proteins, the protein extraction solution was added and stored on ice for 30 min according to the manufacturer's instructions (PRO_PREP™, iNtRON Biotechnology, Seoul, Korea). The supernatant was collected after centrifugation at maximum speed for 10 min at 4°C. The amount of protein was measured by the Bradford method.

ESP were prepared as described below. The collected larvae were incubated in PBS for 24 hr to remove salts and other foreign bodies. Larvae were incubated in DMEM (Dulbecco's Modified Eagle Medium) with gentamycin (150 mg/ml) and vancomycin (10 mg/ml) at 0.5 ml/larva, 37°C for 48 hr. The media were collected and centrifuged at 2,500 rpm for 20 min and concentrated with Amicon stirred cells, cut off molecular weight less than 10,000 (Millipore Corp, Massachusetts, USA).

IgE ELISA

The crude extract and ESP of *A. simplex* L3 larvae were used as antigens. Both of them were diluted for the ELISA Starter Kit

(Koma Biotech, Seoul, Korea) as 1 g/ml and 100 µl of aliquots were added each well of 96 well ELISA plate. Antigens were incubated overnight at 4°C and washed with Tris-Buffered Saline with tween-20 (TBST) 3 times. Blocking solution was added each well and incubated for 1 hour at room temperature. Samples were incubated with 1:5 diluted patients' sera and 1:500 diluted peroxidase conjugated goat anti-human IgE (Sigma Aldrich, St. Louis, Missouri, USA) as the secondary antibody. For color reactions, 100 µl of TMB substrate solution was added and stopped the reactions when color reaction was enough to measure. Absorbance was measured at 450 nm by ELISA reader (Emax, Molecular Devices, Downingtown, Pennsylvania, USA). A mean OD value ± 3SD of the negative control sera was set as a cut-off value. Negative control sera were collected from 10 healthy persons aged 20-25 years who had no allergic history and the total IgE level was less than 100 IU/ml.

Immunoblot

Antigens from crude extracts and ESP were run through protein gel electrophoresis. Proteins were diluted with dissociation buffer (3:1) and boiled at 100°C for 5 min. Boiled proteins were run through 12% SDS-polyacrylamide gel and transferred on a nitrocellulose membrane by Mini-Protean II (Bio-Rad, Richmond, California, USA) at 25 volt for 1.5 hr. Transferred proteins were checked by ponceau staining and blocked with PBS containing 5% skim milk at room temperature for 1 hr. After 3 times washing with PBST for 10 min, membranes were incubated with diluted ELISA positive patients' sera (1:10) at 4°C overnight. Blotted membranes were washed 3 times with PBST for 10 min and incubated with horseradish peroxidase-labeled goat anti-human IgE (1:500) (Sigma) at room temperature for 1 hr. After 3 times washing with PBST for 10 min, color reaction was conducted by adding 4-chloro-1-naphthol (Sigma) or DAB substrate (Pierce, Rockford, Illinois, USA) on the blotted membrane.

RESULTS

The study population consisted of 269 females and 229 males. The ages of patients were from teens to 98 years old. Among them, 107 were in their twenties, and was the largest group. The numbers of thirties, forties, fifties, sixties, seventies, and eighties and older were 81, 71, 55, 55, 45, and 50, respectively (Table 1). To analyze the pattern of antigens from crude extracts and ESP of *A. simplex* L3, proteins were run through SDS-

Table 1. Age group distribution of subjects and results of ELISA and western blot with *Anisakis simplex* L3 larva crude extract antigen and excretory-secretory antigen

Age	No. of subjects			ELISA positive		WB ^a positive	
	Total	Female	Male	CE ^b	ES ^c	CE	ES
10-19	34	23	11	4	3	0	0
20-29	107	67	40	6	5	0	1
30-39	81	54	27	5	6	0	4
40-49	71	26	45	3	4	0	1
50-59	55	17	38	2	1	1	1
60-69	55	15	40	1	3	0	1
70-79	45	29	16	4	7	1	0
80-	50	38	12	0	4	0	0
Total	498	269	229	25	33 (20 ^d)	2 ^e	8 ^e

^aWestern blot, ^bcrude extracts, ^cexcretory-secretory product, ^dthe number of positive females, ^e130 kDa band positive.

PAGE. Proteins were distributed from molecular weight of 200 kDa to 10 kDa (Fig. 1).

The optical density (OD) by ELISA of the 498 patients revealed 0.03 ± 0.084 (mean \pm SD), and 0.01 ± 0.108 for antigens from crude extract and ESP, respectively. The seropositive rate was 5.0% with crude extract antigen while that with ESP antigen was 6.6% (33 positives) by ELISA among the 498 patients (by cut-off OD value of 0.10). Twenty were females among 33 positive patients (Table 1). The OD value of the positive group was 0.28 ± 0.114 and 0.23 ± 0.117 , and that of negative group was 0.01 ± 0.042 and 0.01 ± 0.079 for antigens from crude extracts and ESP, respectively. The specific serum IgE level against *A. simplex* crude extracts and ESP showed various distributions between the study populations. The serum samples from Geoje city showed more extensive distribution and higher values than those from the other 2 cities (Fig. 2).

Western blotting carried out for ELISA positive sera revealed a specific band of about 130 kDa in 10 patients in both crude extract and ESP (Fig. 3). The mean OD of ELISA for 10 patients who showed the specific band in western blot was 0.28 ± 0.221 . The regional distribution of positive serum samples by Western blotting showed that 8 positive patients were in Geoje city, and 2 patients were in Masan city. Eight were females among all 10 positive sera.

DISCUSSION

In our study, we analyzed the serum samples obtained from residents of Busan Metropolitan city, Masan city, and Geoje city by ELISA and western blotting for the seroepidemiologic




Fig. 1. SDS-PAGE of excretory-secretory products (ES) from *Anisakis simplex* L3 larvae and somatic crude extract (CE). M: Molecular marker. Arrow: 130 kDa.

study of *A. simplex* L3 infection. The positive rates by ELISA showed 5.0% and 6.6% for antigens from crude extracts and ESP, respectively. Western blot analysis of ELISA positive sera revealed the specific band of about 130 kDa in 10 patients in both crude extract and ESP. This protein is considered similar to Ani s 7 (139 kDa) which was proposed by Rodriguez et al. [18], which was reported to be the typical antigen for *A. simplex* L3. Our work provided the first data on the seroprevalence of anti-*Anisakis* IgE sensitization in Korea.

In Spain, the prevalence of anisakiasis varied depending on location, such as 12.4% in Madrid and 0.43% in Galicia [10,11]. Valinas et al. [10] suggested that the low positive infection rate (0.43%) of anisakis allergy in Galicia, Spain, was due to the difference between live or dead larvae. They postulated that only live *A. simplex* L3 can cause anisakis allergy. However, other reports set forth a counterargument that it can be different according to the habit of fish consumption, genetic background, and diagnostic methods [19].

Many factors can affect the seroepidemiological prevalence. The types of antigens and diagnostic methods, however, are the most important factors influencing seroepidemiological inves-




Fig. 2. Distribution of specific serum IgE against *Anisakis simplex* somatic crude extracts (CE) and excretory-secretory products (ES). It was expressed as the OD value of ELISA in residents of Geoje city, Masan city, and Busan Metropolitan city. Each point represents a single serum. Cut-off OD value is 0.10.




Fig. 3. Western blot analysis of *Anisakis simplex* L3 excretory-secretory products (A, C) and crude extracts (B) with sera of patients. A,B: sera from Geoje city, C: sera from Masan city. The 130 kDa protein (arrow) was detected in 10 sera among ELISA positive samples. M: Molecular marker.

tigations. Although the crude extracts and ESP are frequently used for serologic diagnosis, ESP were reported as the more potent and clinically important allergens for diagnosis [20-23]. Our results also showed that ESP was more potent allergen than the crude extract. However, cross reactions with other allergens, such as intestinal and blood-tissue nematodes, should be considered in regard to seropositive prevalence [24,25].

Many excretory-secretory and somatic *Anisakis*-specific antigens have been reported, including Ani s 1 (21 kDa) [26], Ani s 2 (100 kDa, paramyosin) [27], Ani s 3 (41 kDa, tropomyosin) [28-29], Ani s 4 (9 kDa, cysteine protease inhibitor) [30-31], Ani s 5 (15 kDa protein homologous with the SXP/RAL-2 family proteins), and Ani s 6 (7 kDa, serine protease inhibitor) [32]. Recent studies also reported functionally unknown antigens, including Ani s 8 (15 kDa), Ani s 9 (14 kDa), and Ani s 10 (21 kDa protein with unknown function; Allergen Nomenclature Sub-Committee; <http://www.allergen.org>). We identified a 130 kDa protein with western blotting among ELISA positive patients. This protein showed a similar molecular weight with Ani s 7. Ani s 7 (139 kDa) was identified to have a 1096-amino acid fragment 7 (GenBank: EF158010) and 19 repeats of a novel CX17-25 CX 9-22 CX8 CX6 tandem repeat motif [33]. Ani s 7 was reported probably the most important major excretory-secretory allergens since they were recognized by 100% of infected patients. In addition, Anti-Ani s 7 IgE antibodies were reported that they were induced by live *Anisakis* larvae peaked at about day 30 post-infection and then decreased slowly over the course of 2 months [33]. Our results also suggested that Ani s 7 can be a potent serodiagnostic antigen in Korean patients.

In conclusion, we recognized that the seropositive rate of

anisakiasis was 5.0% and 6.6% with ELISA against crude extracts and ESP, respectively. The specific 130 kDa protein was confirmed by Western blot analysis among ELISA positive serum samples. This protein was similar with Ani s 7 in molecular weight and can be a candidate for diagnosis of anisakiasis among Koreans.

ACKNOWLEDGEMENTS

This work was supported by the Korea Research Foundation Grant funded by the Korean Government (MOEHRD, Basic Research Promotion Fund) (KRF-2006-E00037).

REFERENCES

- Kasuya S, Hamano H, Izumi S. Mackerel induced urticaria and *Anisakis*. *Lancet* 1990; 335: 665.
- Daschner A, Pascual CY. *Anisakis simplex*: sensitization and clinical allergy. *Curr Opin Allergy Clin Immunol* 2005; 5: 281-285.
- Moneo I, Caballero ML, Rodriguez-Perez R, Rodriguez-Mahillo AI, Gonzalez-Muñoz M. Sensitization to the fish parasite *Anisakis simplex*: clinical and laboratory aspects. *Parasitol Res* 2007; 101: 1051-1055.
- Audicana MT, Kennedy MW. *Anisakis simplex*: from obscure infectious worm to inducer of immune hypersensitivity. *Clin Microbiol Rev* 2008; 21: 360-379.
- Chai JY, Murrell KD, Lymbery AJ. Fish-borne parasitic zoonoses; status and issues. *Int J Parasitol* 2005; 35: 1233-1254.
- Im KI, Shin HJ, Kim BH, Moon SI. Gastric anisakiasis cases in Cheju-do, Korea. *Korean J Parasitol* 1995; 33: 179-186.
- López-Serrano MC, Gomez AA, Daschner A, Moreno-Ancillo A, de Parga JM, Caballero MT, Barranco P, Cabañas R. Gastroallergic anisakiasis: findings in 22 patients. *J Gastroenterol Hepatol* 2000; 15: 503-506.
- Nawa Y, Hatz C, Blum J. Sushi delights and parasites: the risk of fishborne and foodborne parasitic zoonoses in Asia. *Clin Infect Dis* 2005; 41: 1297-1303.
- Choi SJ, Lee JC, Kim MJ, Hur GY, Shin SY, Park HS. The clinical characteristics of *Anisakis* allergy in Korea. *Korean J Intern Med* 2009; 24: 160-163.
- Valinas B, Lorenzo S, Eiras A, Figueriras A, Sanmartín ML, Ubeira FM. Prevalence and risk factors for IgE sensitization to *Anisakis simplex* in a Spanish population. *Allergy* 2001; 56: 667-671.
- Puente P, Anadon AM, Rodero M, Romaria F, Ubeira FM, Cuellar C. *Anisakis simplex*; the high prevalence in Madrid (Spain) and its relation with fish consumption. *Exp Parasitol* 2008; 118: 271-274.
- Kimura S. Positive ratio of allergen specific IgE antibodies in serum, from a large scale study. *Rinsho Byori* 2001; 49: 376-380.
- Chun SK, Chung BK, Ryu BS. Studies on *Anisakis* sp. (1) on the infection state of anisakiasis like larvae isolated from various marine fishes. *Korean J Fish Aquat Sci* 1968; 1: 99-105.
- Abollo E, Gestal C, Pascual S. *Anisakis* infestation in marine fish and cephalopods from Galician waters: an updated perspective. *Parasitol Res* 2001; 87: 492-499.
- Shih HH, Ku CC, Wang CS. *Anisakis simplex* (Nematoda: Anisakidae) third-stage larval infections of marine cage cultured cobia, *Rachycentron canadum* L., in Taiwan. *Vet Parasitol* 2010; 171: 277-285.
- Ma HW, Jiang TJ, Quan FS, Chen XG, Wang HD, Zhang YS, Cui MS, Zhi WY, Jiang DC. The infection status of anisakid larvae in marine fish and cephalopods from the Bohai Sea, China and their taxonomical consideration. *Korean J Parasitol* 1997; 35: 19-24.
- Choi SH, Kim J, Jo JO, Cho MK, Yu HS, Cha HJ, Ock MS. *Anisakis simplex* larvae: infection status in marine fish and cephalopods purchased from the Cooperative Fish Market in Busan, Korea. *Korean J Parasitol*. 2011; 49: 39-44.
- Rodríguez E, Anadón AM, García-Bodas E, Romarís F, Iglesias R, Gárate T, Ubeira FM. Novel sequences and epitopes of diagnostic value derived from the *Anisakis simplex* Ani s 7 major allergen. *Allergy* 2008; 63: 219-225.
- Falcão H, Lunet N, Neves E, Barros H. Do only live larvae cause *Anisakis simplex* sensitization? *Allergy* 2002; 57: 44.
- Arlian LG, Morgan MS, Quirce S, Marañón F, Fernández-Caldas E. Characterization of allergens of *Anisakis simplex*. *Allergy* 2003; 58: 1299-1303.
- Baeza ML, Rodríguez A, Matheu V, Rubio M, Tornero P, de Barrio M, Herrero T, Santaolalla M, Zubeldia JM. Characterization of allergens secreted by *Anisakis simplex* parasite: clinical relevance in comparison with somatic allergens. *Clin Exp Allergy* 2004; 34: 296-302.
- Kim JS, Kim KH, Cho S, Park HY, Cho SW, Kim YT, Joo KH, Lee JS. Immunochemical and biological analysis of allergenicity with excretory-secretory products of *Anisakis simplex* third stage larva. *Int Arch Allergy Immunol*. 2005; 136: 320-328.
- Moneo I, Caballero ML, Rodriguez-Perez R, Rodriguez-Mahillo AI, Gonzalez-Muñoz M. Sensitization to the fish parasite *Anisakis simplex*; clinical and laboratory aspects. *Parasitol Res* 2007; 101: 1051-1055.
- Sakanari JA, Mckerrow JH. Anisakiasis. *Clin Microbiol Rev* 1989; 2: 278-284.
- Cho JK, Cho SW. Shared epitope for monoclonal IR162 between *Anisakis simplex* larvae and *Clonorchis sinensis* and cross-reactivity between antigens. *J Parasitol* 2000; 86: 1145-1149.
- Moneo I, Caballero ML, Gómez F, Ortega E, Alonso MJ. Isolation and characterization of a major allergen from the fish parasite *Anisakis simplex*. *J Allergy Clin Immunol* 2000; 106: 177-182.
- Shimakura K, Miura H, Ikeda K, Ishizaki S, Nagashima Y, Shirai T, Kasuya S, Shiomi K. Purification and molecular cloning of a major allergen from *Anisakis simplex*. *Mol Biochem Parasitol* 2004; 135: 69-75.
- Asturias JA, Eraso E, Martínez A. Cloning and high level expres-

- sion in *Escherichia coli* of an *Anisakis simplex* tropomyosin isoform. *Mol Biochem Parasitol* 2000; 108: 263-267.
29. Asturias JA, Eraso E, Moneo I, Martínez A. Is tropomyosin an allergen in *Anisakis*? *Allergy* 2000; 55: 898-899.
30. Moneo I, Caballero ML, González-Muñoz M, Rodríguez-Mahillo AI, Rodríguez-Perez R, Silva A. Isolation of a heat-resistant allergen from the fish parasite *Anisakis simplex*. *Parasitol Res* 2005; 96: 285-289.
31. Rodríguez-Mahillo AI, Gonzalez-Muñoz M, Gomez-Aguado E, Rodríguez-Perez R, Corcuera MT, Caballero ML, Moneo I. Cloning and characterization of the *Anisakis simplex* allergen Ani s 4 as a cysteine-protease inhibitor. *Int J Parasitol* 2007; 37: 907-917.
32. Kobayashi Y, Ishizaki S, Shimakura K, Nagashima Y, Shiomi K. Molecular cloning and expression of two new allergens from *Anisakis simplex*. *Parasitol Res* 2007; 100: 1233-1241.
33. Anadón AM, Romarís F, Escalante M, Rodríguez E, Gárate T, Cuéllar C, Ubeira FM. The *Anisakis simplex* Ani s 7 major allergen as an indicator of true *Anisakis* infections. *Clin Exp Immunol* 2009; 156: 471-478.