








# Non-Traumatic Subdural Hematoma and Cancer: A Cohort Study

Søren Hauge Okholm <sup>1</sup>, Dávid Nagy <sup>1</sup>, Dóra Körmendiné Farkas <sup>1</sup>, Cecilia Hvitfeldt Fuglsang <sup>1</sup>, Frederikke Schønfeldt Troelsen <sup>1</sup>, Victor W Henderson <sup>1,2</sup>, Henrik Toft Sørensen <sup>1</sup>

<sup>1</sup>Department of Clinical Epidemiology, Department of Clinical Medicine, Aarhus University, Aarhus, Denmark; <sup>2</sup>Departments of Epidemiology & Population Health and of Neurology & Neurological Sciences, Stanford University, Stanford, CA, USA

Correspondence: Henrik Toft Sørensen, Department of Clinical Epidemiology, Aarhus University Hospital and Aarhus University, Olof Palmes Allé 43-45, Aarhus N, 8200, Denmark, Tel +45 87 16 82 15, Email [hts@clin.au.dk](mailto:hts@clin.au.dk)

**Introduction:** Cancer may increase the risk of bleeding. However, whether subdural hematoma is a marker of occult cancer remains unknown. We examined the association between non-traumatic subdural hematoma and cancer risk in a cohort study.

**Materials and Methods:** Using Danish nationwide health registries, we identified 2713 patients with non-traumatic subdural hematoma and no previous cancer diagnosis, who were hospitalized between April 1, 1996 and December 31, 2019. We computed age-, sex-, and calendar year-standardized incidence ratios (SIRs) as the ratio of the observed to expected number of patients with cancer by using national incidence rates as reference as a measure of relative risk.

**Results:** We identified 77 cancer cases within the first year of follow-up and 272 cancer cases thereafter. The one-year risk of cancer was 2.8% (95% confidence interval: 2.2–3.5), and the one-year SIR was 1.7 (95% confidence interval: 1.3–2.1). During the subsequent years, the SIR was 1.0 (95% confidence interval: 0.9–1.1). The relative risk was elevated for some hematological and liver cancers.

**Conclusion:** The risk of a new cancer diagnosis was clearly increased in patients with non-traumatic subdural hematoma compared with the general population during the first year of follow-up. However, the absolute risk was low, thus limiting the clinical relevance of pursuing early cancer detection in these patients.

**Keywords:** cancer, cohort study, epidemiology, non-traumatic subdural hematoma, population-based

## Introduction

The association between cancer and coagulopathy is well established, because tumor cells may induce a hypercoagulable state that leads to clotting as well as vessel injury.<sup>1</sup> Thus, patients with cancer have an elevated risk of both thrombotic and hemorrhagic events.<sup>2–4</sup> Hence, bleeding and venous thromboembolism may be the first symptoms of an occult malignancy.<sup>5–9</sup> Trauma is the most common cause of subdural hematoma, while the etiologies of non-traumatic or spontaneous subdural hematoma include coagulopathy, vascular lesions, and arachnoid cysts.<sup>10,11</sup> Due to coagulopathy and intratumoral hemorrhage, patients with cancer may present with non-traumatic subdural hematoma more frequently than the general population.<sup>12,13</sup>

Existing research on non-traumatic subdural hematoma as a presenting symptom of occult cancer is scarce, and it is unclear whether, and to what extent, non-traumatic subdural hematoma may be the precursor of some types of recessive cancer. We therefore conducted a nationwide cohort study to examine the risk of a cancer diagnosis in patients with non-traumatic subdural hematoma.

## Methods

Denmark has a tax-supported health care system that grants free and equal access to hospitals for its approximately 5.8 million residents.<sup>14</sup> We used the Danish National Patient Registry (DNPR) and the Danish Cancer Registry to conduct this population-based cohort study in Denmark. Data for every individual are linked via the unique 10-digit civil

registration number given to each Danish resident by the Central Registry System.<sup>15</sup> The civil registration number is issued at birth or upon immigration and is linked to each individual until death or emigration.<sup>16</sup>

The DNPR covers all Danish hospitals since 1977.<sup>17</sup> The Danish Cancer Registry has recorded any incident malignant neoplasms in the Danish population since 1943.<sup>18</sup>

We used diagnosis codes according to the 8th revision of the International Classification of Diseases (ICD) until 1993 and the 10th revision of ICD from 1994 onwards,<sup>17</sup> except for cancer codes, all of which were converted into the 10th revision of ICD. We then categorized the different types of cancer by using the classification according to the Danish Health Data Authority.<sup>18</sup>

Through the DNPR, we identified all patients with a first-time hospital-based diagnosis of non-traumatic subdural hematoma from April 1, 1996, to December 31, 2019. Patients who had a cancer diagnosis before the non-traumatic subdural hematoma were excluded from the analysis ( $n=537$ ). We used the DNPR to acquire data on the date of diagnosis of non-traumatic subdural hematoma and comorbidity at the time of diagnosis.

Each patient was characterized by sex, age, year of non-traumatic subdural hematoma diagnosis, comorbidity, and whether they were treated in inpatient, outpatient hospital clinic, or emergency department settings.

We followed all patients with non-traumatic subdural hematoma from the diagnosis date until the first occurrence of any cancer, death, emigration, or the end of study on December 31, 2019. Because early detection might represent occult cancer, we divided the follow-up period into two subgroups: (1) The first year of follow-up after the non-traumatic subdural hematoma diagnosis and (2) the years thereafter. As a measure of absolute risk of cancer for patients with non-traumatic subdural hematoma, we calculated the cumulative incidence and 95% confidence interval (CI) for each subgroup of cancer, treating death as a competing risk. Based on the assumption that cancers diagnosed within the first year were present at the time of the non-traumatic subdural hematoma, by calculating the reciprocal excess risk, we computed the number of patients who would need to be examined for cancer at the time of diagnosis to identify one excess cancer case.

As a measure of relative risk, age-, sex-, and calendar year-standardized incidence ratios (SIRs) with 95% CI were computed comparing number of cancer cases among patients with non-traumatic subdural hematoma with the expected number of cases, according to national cancer incidence rates.<sup>18</sup> As with the cumulative incidences, we calculated the SIRs for the first year of follow-up and the subsequent years separately. In addition, we calculated SIRs for the first three months and the subsequent months in a sensitivity analysis to investigate the impact of the one-year cut-off on our results. We stratified our analyses by sex, age, and year of non-traumatic subdural hematoma diagnosis, cancer type, and the hospital department to which the patients were admitted.

## Results

We identified 2713 patients with an inpatient (84.3%), outpatient hospital clinic, or emergency department diagnosis of non-traumatic subdural hematoma. In total, 1260 (46.4%) of patients underwent surgical treatment within 30 days after the diagnosis of non-traumatic subdural hematoma. The majority were men (60.7%), and the median age was 71.2 years (interquartile range: 58.9–80.7 years) at the time of diagnosis. The median follow-up time was 3.6 years (interquartile range: 0.2–8.6 years).

Seventy-seven patients with non-traumatic subdural hematoma were diagnosed with cancer within the first year of follow-up, corresponding to an absolute risk of any cancer within the first year of 2.8% (95% CI 2.2–3.5) (Table 1). Thirty-one excess cancer cases were detected during the first year, thus indicating that 63 patients with non-traumatic subdural hematoma would need to be examined to detect one excess cancer case. The SIR within the first year was 1.7 (95% CI 1.3–2.1). The relative risk for male patients (SIR = 1.6 [95% CI 1.2–2.1]) was slightly lower than that for female patients (SIR = 1.9 [95% CI 1.2–2.8]). Patients with hematoma before the age of 60 had a higher relative risk than patients in older age groups. We found strong associations with lung cancer (SIR = 1.6 [95% CI 0.6–3.2]), meningeal cancer (SIR = 25.8 [95% CI 10.4–53.2]), non-Hodgkin lymphoma (SIR = 4.7 [95% CI 1.9–9.8]), and myeloid leukemia (SIR = 16.3 [95% CI 5.3–37.9]) among patients with non-traumatic subdural hematoma during the first year of follow-up. The SIR within the first three months was 3.3 (95% CI 2.3–4.4).

**Table 1** Absolute Risks and Standardized Incidence Ratios of Site-Specific Cancers During the First Year of Follow-Up Among Patients with Non-Traumatic Subdural Hematoma, Denmark, 1996–2019. Only Sites with Five or More Recorded Cancer Cases are Presented

	Observed (n)	Expected (n)	Risk % (95% CI)	SIR (95% CI)
Overall	77	45.3	2.8 (2.2–3.5)	1.7 (1.3–2.1)
Lungs, bronchi, and trachea	7	4.5	0.3 (0.1–0.5)	1.6 (0.6–3.2)
Prostate	5	6.2	0.2 (0.1–0.4)	0.8 (0.3–1.9)
Membranes of the brain and spinal meninx	7	0.3	0.3 (0.1–0.5)	25.8 (10.4–53.2)
Non-Hodgkin malignant lymphoma	7	1.5	0.3 (0.1–0.5)	4.7 (1.9–9.8)
Myeloid leukemia	5	0.3	0.2 (0.1–0.4)	16.3 (5.3–37.9)
Basal cell skin carcinoma	11	9.7	0.4 (0.2–0.7)	1.1 (0.6–2.0)
Other skin cancers (excluding basal cell skin carcinoma)	5	2.6	0.2 (0.1–0.4)	1.9 (0.6–4.4)
All other cancers	30	20.3	1.1 (0.8–1.6)	1.5 (1.0–2.1)

**Abbreviations:** CI, confidence interval; SIR, standardized incidence ratio.

In the subsequent years of follow-up, 272 patients with hematoma were diagnosed with cancer, corresponding to an absolute risk of 27.8% (95% CI 19.1–37.2) (Table 2). The SIR for cancer in the following years was 1.0 (95% CI 0.9–1.1). The relative risk was similar for males (SIR = 0.9 [95% CI 0.8–1.1]) and females (SIR = 1.1 [95% CI 0.9–1.3]). We

**Table 2** Absolute Risks and Standardized Incidence Ratios of Site-Specific Cancers During Subsequent Years of Follow-Up Among Patients with Non-Traumatic Subdural Hematoma, Denmark, 1996–2019. Only Sites with Five or More Recorded Cancer Cases are Presented. The Median Follow-Up Time Was 3.6 Years (Interquartile Range: 0.2–8.6 Years)

	Observed (n)	Expected (n)	Risk % (95% CI)	SIR (95% CI)
Overall	272	283.9	27.8 (19.1–37.2)	1.0 (0.9–1.1)
Colorectal	29	30.1	2.3 (1.4–3.5)	1.0 (0.6–1.4)
Liver	6	2.6	0.5 (0.2–1.0)	2.3 (0.9–5.0)
Pancreas	7	5.9	0.5 (0.2–1.0)	1.2 (0.5–2.4)
Lungs, bronchi, and trachea	29	26.6	6.3 (1.1–18.5)	1.1 (0.7–1.6)
Malignant melanoma	8	7.6	0.6 (0.3–1.1)	1.0 (0.5–2.1)
Other skin cancers (excluding basal cell skin carcinoma)	16	18.3	1.8 (0.7–3.6)	0.9 (0.5–1.4)
Basal cell skin carcinoma	57	64.9	4.6 (2.9–6.8)	0.9 (0.7–1.1)
Breast	13	16.0	1.4 (0.6–2.6)	0.8 (0.4–1.4)
Prostate	32	36.9	2.4 (1.6–3.4)	0.9 (0.6–1.2)
Urinary bladder	11	15.3	0.8 (0.4–1.4)	0.7 (0.4–1.3)
Brain	7	3.6	0.5 (0.2–1.0)	1.9 (0.8–4.0)
Non-Hodgkin malignant lymphoma	12	9.6	1.1 (0.5–2.0)	1.3 (0.7–2.2)
Lymphoid leukemia	6	3.3	0.4 (0.2–0.8)	1.8 (0.7–4.0)
Metastases and non-specified cancer in lymph nodes	10	4.5	1.9 (0.5–5.1)	2.2 (1.1–4.1)
All other cancers	29	38.7	3.0 (1.8–4.6)	0.7 (0.5–1.1)

**Abbreviations:** CI, confidence interval; SIR, standardized incidence ratio.

found an elevated relative risk of liver cancer (SIR = 2.3 [95% CI 0.9–5.0]), brain cancer (SIR = 1.9 [95% CI 0.8–4.0]), non-Hodgkin malignant lymphoma (SIR = 1.3 [95% CI 0.7–2.2]), lymphoid leukemia (SIR = 1.8 [95% CI 0.7–4.0]), and non-specified cancer in lymph nodes (SIR = 2.2 [95% CI 1.1–4.1]) in the subsequent years of follow-up.

## Discussion

In this Danish cohort study, we found a dichotomous pattern of risk with time. The absolute risk of any cancer was 2.8% within the first year of follow-up after a non-traumatic subdural hematoma. This percentage corresponds to a 1.7-fold increased relative risk of any cancer during the same period, compared with that in the general population. We observed no overall association between non-traumatic subdural hematoma and cancer risk in the following years of follow-up, but we did observe an elevated risk of certain cancers, notably liver cancer.

Non-traumatic subdural hematoma might be associated with cancer diagnoses for several reasons. Heightened diagnostic efforts and the physiological effects of occult cancer probably explain the association during the first three months and one year of follow-up. The early periods of increased cancer diagnosis were not followed by a potential compensatory deficit pattern in cancer risk; consequently, a diagnostic bias is not likely. The elevated long-term risk of liver cancer after the first year of follow-up suggests that non-traumatic subdural hematoma and cancer might share risk factors such as alcohol use. The site-specific relative risks indicated associations among non-traumatic subdural hematoma, hematological cancers, and brain cancer, probably because coagulation disturbances and potential tumor growth in the bloodstream may occur with cancers in the brain and meninges.

Although a strength of our study is that the cohort was population-based and had virtually no losses to long-term follow-up, our data lack clinical details, particularly regarding shared risk factors for non-traumatic subdural hematoma and cancer, except for age, sex, and comorbidities. Moreover, our risk estimates are somewhat imprecise. Cancer diagnoses in the Danish Cancer Registry are of high quality.<sup>18</sup> The recorded non-traumatic subdural hematoma diagnoses might, in some cases, have been misclassified. If non-differential, such misclassification would tend to underestimate the strength of the associations.

In conclusion, our data indicated an increased risk of cancer within the first year after a diagnosis of non-traumatic subdural hematoma. However, the absolute risk during the first year of follow-up was low, which refrained us from recommending cancer detection guidelines for the clinical care of patients with non-traumatic subdural hematoma.

## Ethics Approval and Informed Consent

This study was conducted in accordance with the Declaration of Helsinki. The study was approved by the National Board of Health and by the Danish Data Protection Agency. According to Danish law, approval from the Danish Committee on Health Research Ethics was not necessary. Since this is a register-based study, written consent was not required.

## Author Contributions

All authors made a significant contribution to the work reported, whether that is in the conception, study design, execution, acquisition of data, analysis and interpretation, or in all these areas; took part in drafting, revising or critically reviewing the article; gave final approval of the version to be published; have agreed on the journal to which the article has been submitted; and agree to be accountable for all aspects of the work.

## Funding

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

## Disclosure

Professor Henrik Toft Sørensen reports The Department of Clinical Epidemiology, Aarhus University, receives funding for other studies from the European Medicines Agency and companies in the form of institutional research grants to (and administered by) Aarhus University. None of these studies have any relation to the present study. The authors state that they have no conflicts of interest. Results from this paper were presented at the International Conference on Thrombosis

and Hemostasis Issues in Cancer (ICTHIC) in 2021 as a poster presentation. The related abstract was published in *Thrombosis Research* in April 2021:10.1016/S0049-3848(21)00272-3.

## References

1. Caine GJ, Stonelake PS, Lip GY, Kehoe ST. The hypercoagulable state of malignancy: pathogenesis and current debate. *Neoplasia*. 2002;4(6):465–473. doi:10.1038/sj.neo.7900263
2. Falanga A, Marchetti M, Vignoli A. Coagulation and cancer: biological and clinical aspects. *J Thromb Haemost*. 2013;11(2):223–233. doi:10.1111/jth.12075
3. Reeves BN, Key NS. Acquired hemophilia in malignancy. *Thromb Res*. 2012;129(Suppl 1):S66–S68. doi:10.1016/S0049-3848(12)70019-1
4. Ay C, Pabinger I, Cohen AT. Cancer-associated venous thromboembolism: burden, mechanisms, and management. *Thromb Haemost*. 2017;117(2):219–230. doi:10.1160/TH16-08-0615
5. Astin M, Griffin T, Neal RD, Rose P, Hamilton W. The diagnostic value of symptoms for colorectal cancer in primary care: a systematic review. *Br J Gen Pract*. 2011;61(586):e231–e243.
6. Clarke MA, Long BJ, Del Mar Morillo A, Arbyn M, Bakkum-Gamez JN, Wentzensen N. Association of endometrial cancer risk with postmenopausal bleeding in women: a systematic review and meta-analysis. *JAMA Intern Med*. 2018;178(9):1210–1222. doi:10.1001/jamainternmed.2018.2820
7. Prandoni P, Falanga A, Piccioli A. Cancer and venous thromboembolism. *Lancet Oncol*. 2005;6(6):401–410. doi:10.1016/S1470-2045(05)70207-2
8. Ording AG, Veres K, Farkas DK, Adelborg K, Sørensen HT. Risk of cancer in patients with epistaxis and haemoptysis. *Br J Cancer*. 2018;118(6):913–919. doi:10.1038/bjc.2017.494
9. Nørgaard M, Veres K, Ording AG, Djurhuus JC, Jensen JB, Sørensen HT. Evaluation of hospital-based hematuria diagnosis and subsequent cancer risk among adults in Denmark. *JAMA Netw Open*. 2018;1(7):e184909. doi:10.1001/jamanetworkopen.2018.4909
10. Talalla A, McKissock W. Acute “spontaneous” subdural hemorrhage. An unusual form of cerebrovascular accident. *Neurology*. 1971;21(1):19–25. doi:10.1212/WNL.21.1.19
11. Coombs JB, Coombs BL, Chin EJ. Acute spontaneous subdural hematoma in a middle-aged adult: case report and review of the literature. *J Emerg Med*. 2014;47(3):e63–e68. doi:10.1016/j.jemermed.2014.04.026
12. Velander AJ, DeAngelis LM, Navi BB. Intracranial hemorrhage in patients with cancer. *Curr Atheroscler Rep*. 2012;14(4):373–381. doi:10.1007/s11883-012-0250-3
13. Reichman J, Singer S, Navi B, et al. Subdural hematoma in patients with cancer. *Neurosurgery*. 2012;71(1):74–79. doi:10.1227/NEU.0b013e3182517938
14. Schmidt M, Schmidt SAJ, Adelborg K, et al. The Danish health care system and epidemiological research: from health care contacts to database records. *Clin Epidemiol*. 2019;11:563–591. doi:10.2147/CLEP.S179083
15. Frank L. Epidemiology. When an entire country is a cohort. *Science*. 2000;287(5462):2398–2399. doi:10.1126/science.287.5462.2398
16. Schmidt M, Pedersen L, Sørensen HT. The Danish civil registration system as a tool in epidemiology. *Eur J Epidemiol*. 2014;29(8):541–549. doi:10.1007/s10654-014-9930-3
17. Schmidt M, Schmidt SA, Sandegaard JL, Ehrenstein V, Pedersen L, Sørensen HT. The Danish national patient registry: a review of content, data quality, and research potential. *Clin Epidemiol*. 2015;7:449–490. doi:10.2147/CLEP.S91125
18. Gjerstorff ML. The Danish cancer registry. *Scand J Public Health*. 2011;39(7\_suppl):42–45. doi:10.1177/1403494810393562

### Clinical Epidemiology

Dovepress

### Publish your work in this journal

Clinical Epidemiology is an international, peer-reviewed, open access, online journal focusing on disease and drug epidemiology, identification of risk factors and screening procedures to develop optimal preventative initiatives and programs. Specific topics include: diagnosis, prognosis, treatment, screening, prevention, risk factor modification, systematic reviews, risk & safety of medical interventions, epidemiology & biostatistical methods, and evaluation of guidelines, translational medicine, health policies & economic evaluations. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use.

Submit your manuscript here: <https://www.dovepress.com/clinical-epidemiology-journal>