

Eriophyoid mites from Qinghai Province, northwestern China with descriptions of nine new species (Acari, Eriophyoidea)

Hao-Sen Li^{1,†}, Xiao-Feng Xue^{1,‡}, Xiao-Yue Hong^{1,§}

¹ Department of Entomology, Nanjing Agricultural University, Nanjing, Jiangsu 210095, China

[†] [urn:lsid:zoobank.org:author:44F5179A-8C66-4CCF-AFAC-E4CD486ECAA8](https://doi.org/urn:lsid:zoobank.org:author:44F5179A-8C66-4CCF-AFAC-E4CD486ECAA8)

[‡] [urn:lsid:zoobank.org:author:C10A90B5-91B5-4267-BA89-4EF78F465573](https://doi.org/urn:lsid:zoobank.org:author:C10A90B5-91B5-4267-BA89-4EF78F465573)

[§] [urn:lsid:zoobank.org:author:6EA91087-981C-4749-BF02-1DFCFBA04F9A](https://doi.org/urn:lsid:zoobank.org:author:6EA91087-981C-4749-BF02-1DFCFBA04F9A)

Corresponding author: Xiao-Yue Hong (xyhong@njau.edu.cn)

Academic editor: A. Bochkov | Received 23 January 2012 | Accepted 10 May 2012 | Published 21 May 2012

[urn:lsid:zoobank.org:pub:3648605E-11DD-4E95-8D86-9E91E90055EA](https://doi.org/urn:lsid:zoobank.org:pub:3648605E-11DD-4E95-8D86-9E91E90055EA)

Citation: Li H-S, Xue X-F, Hong X-Y (2012) Eriophyoid mites from Qinghai Province, northwestern China with descriptions of nine new species (Acari, Eriophyoidea). ZooKeys 196: 47–107. doi: 10.3897/zookeys.196.2726

Abstract

Eriophyoid mites from Qinghai Province, northwestern China were studied herein. Up to now, only six species have been reported from Qinghai Province. In field surveys, 17 eriophyoid mite species were collected, among which nine species were found new to science. The new species and their host plants are listed as follows: *Acaphyllisa tuberculumae* **sp. n.** on *Populus* sp. (Salicaceae); *Proiectus xiningensis* **sp. n.** on *Pinus* sp. (Pinaceae); *Phyllocoptes beishaniensis* **sp. n.** on *Spiraea mongolica* Maxim. (Rosaceae); *Tetra pruniana* **sp. n.** on *Prunus tomentosa* Thunb. (Rosaceae) Rupr. (Berberidaceae); *Tetra pyriana* **sp. n.** on *Pyrus calleryana* Decne. (Rosaceae); *Tetra simonia* **sp. n.** on *Populus simonii* Carr. (Salicaceae); *Diptacus berberinus* **sp. n.** on *Berberis amurensis* Rupr. (Berberidaceae); *Diptacus mengdaensis* **sp. n.** on *Lonicera elisae* Franch. (Caprifoliaceae); *Rhyncaphytoptus spinus* **sp. n.** on *Lonicera rupicola* Hook. f. et Thoms. (Caprifoliaceae). *Aculops ulmi* Hong & Xue, 2005 was re-described.

Keywords

Eriophyoid mites, Qinghai, taxonomy, new species

Introduction

Qinghai Province (89°35'E–103°04'E, 13°39'N–39°19'N), located in the northwest of the People's Republic of China, is a part of the Qinghai-Tibet Plateau, with an average elevation of over 3000m (1650m–6860m). The average temperature ranges from 0.4°C to 7.4°C. (Qinghai Province Government Website).

Up to now, six species of eriophyoid mites from Qinghai have been reported. They are *Aceria paramacrodontis* Kuang, 1988 on *Lycium* sp. (Solanaceae), *Aceria qinghaiensis* Kuang, 1997 on *Salix babylonica* L. (Salicaceae), *Aculodes salicis* Kuang, 1997 on *Salix babylonica* L. (Salicaceae), *Aculops xiningensis* Kuang, 2000 on *Malus pumila* P. Mill. (Rosaceae), *Aculus huangzhongensis* Kuang, 2000 on *Syringa oblata* Lindl. (Oleaceae) and *Tetraspinus syringae* Lin & Kuang, 2001 on *Syringa oblata* Lindl. (Oleaceae). In July 2007, field surveys were conducted in Qinghai Province, northwestern China. Twenty-five eriophyoid mite samples were collected and 17 eriophyoid mite species were identified, among which nine species were found new to science. No species reported earlier from Qinghai were collected in this survey. In total, there are 23 species of the Eriophyoidea from Qinghai Province, belonging to two families and 12 genera. A list of eriophyoid mites from Qinghai Province is given (Table 1).

Table 1. List of eriophyoid mites and their hosts in Qinghai Province.

Family	Subfamily	Tribe	Species	Host	
Eriophyidae	Eriophyinae	Aceriini	<i>Aceria paramacrodontis</i> Kuang, 1988	<i>Lycium</i> sp. (Solanaceae)	
			<i>Aceria qinghaiensis</i> Kuang, 1997	<i>Salix babylonica</i> L. (Salicaceae)	
	Phyllocoptinae	Acaricalini	<i>Acaphyllisa tuberculumae</i> sp. n.	<i>Populus</i> sp. (Salicaceae)	
		Phyllocoptini	<i>Proiectus xiningensis</i> sp. n.	<i>Pinus</i> sp. (Pinaceae)	
			<i>Phyllocoptes beishaniensis</i> sp. n.	<i>Spiraea mongolica</i> Maxim. (Rosaceae)	
			<i>Phyllocoptes asperatae</i> Song, Xue & Hong, 2006	<i>Picea meyeri</i> Rehd. Et Wils. (Pinaceae)	
			<i>Phyllocoptes dangchangi</i> Song, Xue & Hong, 2006	<i>Picea</i> sp. (Pinaceae)	
			<i>Phyllocoptes gansunensis</i> Kuang & Luo, 1998	<i>Potentilla parvifolia</i> Fisch. ap. Lehm. (Rosaceae)	
			<i>Phyllocoptiruta platycladusa</i> Xue, Song, Amrine & Hong, 2007	<i>Juniperus chinensis</i> L. (Cupressaceae)	
			Anthocoptini	<i>Aculus changbais</i> Xue, Song & Hong, 2008	<i>Salix chaenomeloides</i> Kimura (Salicaceae)
				<i>Aculus huangzhongensis</i> Kuang, 2000	<i>Syringa oblata</i> Lindl. (Oleaceae)
		<i>Aculodes salicis</i> Kuang, 1997		<i>Salix babylonica</i> L. (Rosaceae)	
		<i>Aculops umli</i> Hong & Xue, 2005		<i>Ulmus</i> sp. (Ulmaceae)	

Family	Subfamily	Tribe	Species	Host
			<i>Aculops xiningensis</i> Kuang, 2000	<i>Malus pumila</i> P. Mill. (Rosaceae)
			<i>Tetraspinus syringae</i> Lin & Kuang, 2001	<i>Syringa oblata</i> Lindl. (Oleaceae)
			<i>Tetra pinnatifidae</i> Xue, Song & Hong, 2006	<i>Prunus armeniaca</i> Linn. (Rosaceae)
			<i>Tetra pruniana</i> sp. n.	<i>Prunus tomentosa</i> Thunb. (Rosaceae)
			<i>Tetra pyriana</i> sp. n.	<i>Pyrus calleryana</i> Decne. (Rosaceae)
				<i>Pyrus betulifolia</i> Bunge. (Rosaceae)
		<i>Tetra simonia</i> sp. n.	<i>Populus simonii</i> Carr. (Salicaceae)	
Diptilomiopidae	Diptilomiopinae		<i>Diptacus berberinus</i> sp. n.	<i>Berberis amurensis</i> Rupr. (Berberidaceae)
			<i>Diptacus mengdaensis</i> sp. n.	<i>Lonicera elisae</i> Franch. (Caprifoliaceae)
	Rhyncaphytopinae		<i>Rhyncaphytoptus spinus</i> sp. n.	<i>Lonicera rupicola</i> Hook. f. et Thoms. (Caprifoliaceae)
			<i>Rhyncaphytoptus ulmi</i> Xin & Dong, 1981	<i>Ulmus</i> sp. (Ulmaceae)

Materials and methods

In the field, eriophyoid mites were collected by the aid of a hand-lens (30X) from the lower surface of host plant leaves. Eriophyoid mites, together with host plants, were immersed in 75% alcohol and kept in vials. Each vial was marked with the collection data, such as specimen number, collection date, host plant, mite color, location, collector, and mite relationship to host plant. The collection data were also recorded in the collection notebook for further use. The host plants were kept in a plant specimen folder in a dry environment.

The morphological terminology used here follows that of Lindquist (1996) and the generic classification was made according to Amrine et al. (2003). Slides were mounted using Keifer's F-medium and modified Berlese medium (Amrine and Manson 1996). Specimens were measured based on the methods outlined by de Lillo et al. (2010). Specimens were examined with a Zeiss A2 (Germany) research microscope with phase contrast and semi-schematic drawings were made. Photos of slide mounted mites were taken with the same microscope (100× oil immersion objective with 10× eyepieces), connected to a computer using Axiovision image analysis software. For each species, the holotype female measurement precedes the corresponding range for paratypes (given in parentheses). All measurements are in micrometres (µm), and are lengths when not otherwise specified. All the type materials are deposited at Arthropod/Mite collection of, the Department of Entomology, Nanjing Agricultural University, Jiangsu Province, China.

Taxonomy

Family Eriophyidae Nalepa, 1898

Subfamily Eriophyinae Nalepa, 1898

Tribe Aceriini Amrine & Stasny, 1994

Genus *Aceria* Keifer, 1944

***Aceria paramacrodonis* Kuang, 1988**

http://species-id.net/wiki/Aceria_paramacrodonis

Aceria paramacrodonis Kuang 1988: 49–50, figures 1–6.

Aceria paramacrodonis; Amrine and Stasny 1994: 73.

Aceria paramacrodonis; Kuang 1995: 61, figure 45.

Aceria paramacrodonis; Amrine and Stasny 1996: 295–304.

Aceria paramacrodonis; Hong and Zhang 1996: 25, figure 43.

Aceria paramacrodonis; Song et al. 2008: 4.

Host. *Lycium* sp. (Solanaceae).

Relation to host. Leaf gall; mites produce pocket galls on the lower side of leaves.

Distribution. China (Gansu, Ningxia, Qinghai, Shandong).

***Aceria qinghaiensis* Kuang, 1997**

http://species-id.net/wiki/Aceria_qinghaiensis

Aceria qinghaiensis Kuang and Pang 1997: 231–232, figures 6–11.

Aceria qinghaiensis; Kuang et al. 2005: 31–32, figure 29.

Aceria qinghaiensis; Song et al. 2008: 14.

Host. *Salix babylonica* L. (Salicaceae).

Relation to host. The mites produce pockets on the lower surface of the leaves.

Distribution. China (Gansu, Qinghai).

Subfamily Phyllocoptinae Nalepa, 1892

Tribe Acaricalini Amrine & Stasny, 1994

Genus *Acaphyllisa* Keifer, 1978

***Acaphyllisa tuberculumae* sp. n.**

urn:lsid:zoobank.org:act:90E42ADF-5C3A-449D-A4C1-BA012B3D09F1

http://species-id.net/wiki/Acaphyllisa_tuberculumae

Figures 1–3

Description. Female. (n = 8) Body fusiform, light yellow, 171 (171–195), 72 (70–75) wide. **Gnathosoma** 21 (20–21), projecting obliquely down, suboral plate present, pe-




Figure 1. *Acaphyllisa tuberculumae* sp. n.: **D** dorsal view of female **CG** coxae and female genitalia **CMG** coxae and male genitalia.

dipalp coxal seta (*ep*) 4 (4–5), dorsal pedipalp genual seta (*d*) 7 (7–8), cheliceral stylets 16 (16–18). **Prodorsal shield** 49 (45–49), 53 (53–60) wide, subtriangular; frontal lobe 6 (5–8); median, admedian and submedian lines present, median line ending at




Figure 2. *Acaphyllisa tuberculumae* sp. n.: V ventral view of female em empodium IG female internal genitalia L1 leg I L2 leg II.




Figure 3. *Acaphyllisa tuberculumae* sp. n.: **A** dorsal view of female **B** ventral view of female **C** dorsal view of female posterior part **D** ventral view of female posterior part **E** prodorsal shield **F** coxae and female genitalia.

basal 1/2 of prodorsal shield, admedian lines connected at basal 1/2 and 2/3 of prodorsal shield, forming three cells on each side of the median line. Scapular tubercles ahead of rear shield margin, 2 (2–3), 18 (18–19) apart, scapular setae (*sc*) 8 (6–8), projecting centrad. **Coxigenital region** with 10 smooth annuli. Coxisternal plates with granules, anterolateral setae on coxisternum **I** (*lb*) 7 (7–8), 14 (14–15) apart, proximal setae on coxisternum **I** (*la*) 19 (19–21), 11 (10–11) apart, proximal setae on coxisternum **II** (*2a*) 51 (51–53), 26 (26–28) apart, tubercles *lb* and *la* 8 (8–9) apart, tubercles *la* and *2a* 8 (8–9) apart. Prosternal apodeme combined, 7 (6–7). **Leg I** 36 (35–36), femur 10 (9–10), basiventral femoral seta (*bv*) 8 (8–10); genu 5 (5–6), antaxial genual seta (*l'*) 29 (29–31); tibia 8 (7–8), paraxial tibial seta (*l'*) 7 (7–8), located at 1/4 from dorsal base; tarsus 8 (7–8), seta *ft'* 17 (17–19), seta *ft''* 22 (22–23), seta *u'* 5 (5–6); tarsal empodium (*em*) 6 (6–7), divided, 2-rayed on each side, tarsal solenidion (ω) 7 (6–7), knobbed. **Leg II** 28 (28–33), femur 10 (9–10), basiventral femoral seta (*bv*) 10 (9–10); genu 5 (4–5), antaxial genual seta (*l''*) 6 (6–8); tibia 6 (6–7); tarsus 7 (6–7), seta *ft'* 5 (5–7), seta *ft''* 21 (21–22), seta *u'* 5 (5–6); tarsal empodium (*em*) 6 (5–6), divided, 2-rayed on each side, tarsal solenidion (ω) 6 (6–7), knobbed. **Opisthosoma** dorsally with 55 (55–57) annuli, with round microtubercles, ventrally with 77 (74–77) annuli, with round microtubercles. Setae *c2* 31 (28–31) on ventral annulus 14 (14–16), 56 (54–56) apart; setae *d* 55 (55–60) on ventral annulus 34 (33–34), 38 (33–38) apart; setae *e* 17 (16–17) on ventral annulus 52 (52–53), 20 (18–20) apart; setae *f* 28 (25–28) on ventral annulus 71 (69–71), 26 (25–26) apart. Setae *h1* 2 (2–3), *h2* 70 (70–75). **Female genitalia** 18 (17–18), 24 (24–25) wide, coverflap smooth, setae *3a* 33 (33–35), 14 (14–15) apart.

Male. (n = 1) Body fusiform, light yellow, 150, 57 wide. **Gnathosoma** 17, projecting obliquely down, suboral plate present, pedipalp coxal seta (*ep*) 5, dorsal pedipalp genual seta (*d*) 5, cheliceral stylets 13. **Prodorsal shield** has the same design as female, 44, 48 wide, subtriangular; frontal lobe 5. Scapular tubercles ahead of rear shield margin, 3, 18 apart, scapular setae (*sc*) 6, projecting centrad. **Coxigenital region** with 10 smooth annuli. Coxisternal plates with granules, anterolateral setae on coxisternum **I** (*lb*) 7, 12 apart, proximal setae on coxisternum **I** (*la*) 18, 10 apart, proximal setae on coxisternum **II** (*2a*) 45, 23 apart, tubercles *lb* and *la* 7 apart, tubercles *la* and *2a* 7 apart. Prosternal apodeme combined, 5. **Legs** with usual series of setae. **Leg I** 29, femur 9, basiventral femoral seta (*bv*) 8; genu 5, antaxial genual seta (*l''*) 31; tibia 7, paraxial tibial seta (*l'*) 6, located at 1/4 from dorsal base; tarsus 6, seta *ft'* 17, seta *ft''* 18, seta *u'* 4; tarsal empodium (*em*) 6, divided, 2-rayed on each side, tarsal solenidion (ω) 6, knobbed. **Leg II** 28, femur 10, basiventral femoral seta (*bv*) 8; genu 4, antaxial genual seta (*l''*) 8; tibia 6; tarsus 6, seta *ft'* 5, seta *ft''* 18, seta *u'* 4; tarsal empodium (*em*) 5, divided, 2-rayed on each side, tarsal solenidion (ω) 6, knobbed. **Opisthosoma** dorsally with 51 annuli, with round microtubercles, ventrally with 66 annuli, with round microtubercles. Setae *c2* 21 on ventral annulus 12, 42 apart; setae *d* 42 on ventral annulus 27, 28 apart; setae *e* 16 on ventral annulus 43, 16 apart; setae *f* 25 on ventral annulus 61, 21 apart. Setae *h1* 2, *h2* 60. **Male genitalia** forming a “Y” like structure in the middle, 19 wide, setae *3a* 22, 15 apart.

Type material. Holotype, female (slide number NJAUEri789B, marked Holotype), from *Populus* sp. (Salicaceae), Xining City, Qinghai Province, P. R. China, 36° 38'18"N, 101°45'27"E, elevation 2241m, 21 July 2007, coll. Xiao-Feng Xue. **Paratypes**, 7 females and 1 male (slide number NJAUEri789B), with the same data as holotype.

Relation to host. Vagrant on leaf lower surface. No damage to the host was observed.

Etymology. The specific designation *tuberculumae* is from the character of dorsal opisthosomal microtubercles, "tuberculum" in Latin; masculine in gender.

Differential diagnosis. This species is similar to *Acaphyllisa populi* Xue & Hong, 2006, but can be differentiated from the latter by prodorsal shield with six cells in the middle (prodorsal shield without cells in *A. populi*), opisthosoma dorsally with round microtubercles (opisthosoma dorsally with elliptical microtubercles only on ridges in *A. populi*), female genitalia coverflap smooth (female genital coverflap with 10 longitudinal ridges in *A. populi*).

Tribe Phyllocoptini Nalepa, 1892

Genus *Proiectus* Huang, 2001

Proiectus xiningensis sp. n.

urn:lsid:zoobank.org:act:21DA68FE-18B9-4305-B088-20BF3A865A3C

http://species-id.net/wiki/Proiectus_xiningensis

Figures 4–7

Description. Female. (n = 5) Body fusiform, light yellow, 248 (223–308), 100 (100–110) wide, 90 (90–91) thick. **Gnathosoma** 33 (33–34), projecting obliquely down, suboral plate present, pedipalp coxal seta (*ep*) 3 (3–5), dorsal pedipalp genual seta (*d*) 15 (11–15), cheliceral stylets 33 (33–34). **Prodorsal shield** 73 (65–73), 100 (100–110) wide, subtriangular, with a large projection on each lateral margin 7 (6–7); frontal lobe broad 24 (22–24); median, admedian and submedian lines obscure, median and admedian lines connected at base. Scapular tubercles ahead of rear shield margin, 2 (2–3), 30 (27–30) apart, scapular setae (*sc*) 8 (6–8), projecting centrad. **Coxigenital region** with 17 (14–17) annuli, with round microtubercles, with deep seam under coxisternal plate II. Coxisternal plates with short lines, anterolateral setae on coxisternum I (*lb*) 10 (10–11), 18 (18–19) apart, proximal setae on coxisternum I (*la*) 22 (20–22), 13 (12–13) apart, proximal setae on coxisternum II (*2a*) 52 (50–52), 32 (32–33) apart, tubercles *lb* and *la* 11 (10–11) apart, tubercles *la* and *2a* 10 (10–11) apart. Prosternal apodeme separated, 4 (4–5). **Leg I** 43 (40–45), femur 9 (8–10), basiventral femoral seta (*bv*) 14 (12–14); genu 9 (8–10), antaxial genual seta (*l'*) 28 (28–34); tibia 15 (13–15), paraxial tibial seta (*l*) 5 (5–6), located at 1/2 from dorsal base; tarsus 10 (9–10), seta *ft'* 17 (17–18), seta *ft''* 30 (30–32), seta *u'* 5 (5–6); tarsal empodium (*em*) 8 (7–8), simple, 5-rayed, tarsal solenidion (ω) 10 (9–10), knobbed. **Leg II** 40 (40–41), femur 13 (13–16), basiventral femoral seta (*bv*) 13 (10–13); genu 6 (6–8), antaxial genual seta (*l'*) 8 (7–8); tibia 10 (10–12); tarsus 8 (8–9), seta *ft'* 9 (8–9),




Figure 4. *Projctus xiningensis* sp. n.: **D** dorsal view of female **CMG** coxae and male genitalia **CG** coxae and female genitalia.




Figure 5. *Proiectus xiningensis* sp. n.: **L** lateral view of female **LO** lateral microtubercles **IG** female internal genitalia **em** empodium **L1** leg I **L2** leg II.




Figure 6. *Proiectus xiningensis* sp. n.: **A** dorsal view of female **B** ventral view of female **C** lateral microtubercles **D** empodium **E** dorsal view of female posterior part **F** ventral view of female posterior part **G** leg I and leg II.




Figure 7. *Proiectus xiningensis* sp. n.: **H** lateral view of female **I** coxae and female genitalia **J** coxae and male genitalia **K** lateral view of female posterior part **L** prodorsal shield **M** female internal genitalia.

seta ft'' 27 (27–28), seta u' 5 (5–6); tarsal empodium (em) 8 (7–8), simple, 5-rayed, tarsal solenidion (ω) 10 (9–10), knobbed. **Opisthosoma** dorsally with 41 (39–41) annuli, with weak filamentous microtubercles, ventrally with 92 (92–100) annuli, with round microtubercles. Setae $c2$ 21 (21–22) on ventral annulus 15 (15–20), 70 (70–73) apart; setae d 100 (80–100) on ventral annulus 34 (34–39), 37 (37–46) apart; setae e 55 (55–65) on ventral annulus 56 (56–61), 21 (21–22) apart; setae f 33 (30–33) on ventral annulus 84 (84–90), 28 (28–29) apart. Setae $h1$ 7 (6–7), $h2$ 65 (55–65). **Female genitalia** 28 (28–30), 33 (33–34) wide, coverflap with 19 longitudinal ridges, setae $3a$ 10 (10–12), 23 (21–22) apart.

Male. (n = 1) Body fusiform, light yellow, 250, 90 wide. **Gnathosoma** 34, projecting obliquely down, suboral plate present, pedipalp coxal seta (ep) 3, dorsal pedipalp genual seta (d) 11, cheliceral stylets 50. **Prodorsal shield** has the same design as female, 70, 90 wide, subtriangular; with a large projection on each lateral margin, 6; frontal lobe broad, 21. Scapular tubercles ahead of rear shield margin, 2, 23 apart, scapular setae (sc) 6, projecting centrad. **Coxigenital region** with 16 annuli, with round microtubercles, with deep seam under coxisternal plate II. Coxisternal plates with short lines, anterolateral setae on coxisternum I ($1b$) 10, 15 apart, proximal setae on coxisternum I ($1a$) 15, 12 apart, proximal setae on coxisternum II ($2a$) 52, 29 apart, tubercles $1b$ and $1a$ 10 apart, tubercles $1a$ and $2a$ 11 apart. Prosternal apodeme separated, 5. **Leg I** 40, femur 10, basiventral femoral seta (bv) 13; genu 8, antaxial genual seta (l'') 31; tibia 13, paraxial tibial seta (l) 5, located at 1/2 from dorsal base; tarsus 8, seta ft' 17, seta ft'' 30, seta u' 5; tarsal empodium (em) 7, simple, 5-rayed, tarsal solenidion (ω) 9, knobbed. **Leg II** 37, femur 15, basiventral femoral seta (bv) 12; genu 6, antaxial genual seta (l'') 7; tibia 10; tarsus 8, seta ft' 7, seta ft'' 25, seta u' 5; tarsal empodium (em) 7, simple, 5-rayed, tarsal solenidion (ω) 9, knobbed. **Opisthosoma** dorsally with 41 annuli, with weak filamentous microtubercles, ventrally with 94 annuli, with round microtubercles. Setae $c2$ 21 on ventral annulus 18, 60 apart; setae d 80 on ventral annulus 36, 38 apart; setae e 60 on ventral annulus 58, 20 apart; setae f 27 on ventral annulus 88, 25 apart. Setae $h1$ 6, $h2$ 65. **Male genitalia** 25 wide, setae $3a$ 12, 22 apart.

Type material. Holotype, female (slide number NJAUEri790, marked Holotype), from *Pinus* sp. (Pinaceae), Xining City, Qinghai Province, P. R. China, 36°38'18"N, 101°45'27"E, elevation 2241m, 21 July 2007, coll. Xiao-Feng Xue. **Paratypes**, 4 females and 1 male (slide number NJAUEri790), with the same data as holotype.

Relation to host. Vagrant on terminal part of the needles. No damage to the host was observed.

Etymology. The specific designation *xiningensis* is from the place name Xining City, where this new species was collected; feminine in gender.

Differential diagnosis. This species is similar to *Proiectus thunbergis* Xue, Song, Amrine & Hong, 2007, but can be differentiated from the latter by median and admedian lines of prodorsal shield simple (median and admedian lines with granules in *P. thunbergis*), opisthosoma dorsally annuli with weak filamentous microtubercles (opisthosoma dorsally annuli with round microtubercles in *P. thunbergis*), tarsal empodium (em) 5-rayed (4-rayed in *P. thunbergis*).

Genus *Phyllocoptes* Nalepa, 1887***Phyllocoptes beishaniensis* sp. n.**

urn:lsid:zoobank.org:act:0ABE4028-A29D-4B4A-B7FE-133603A01C99

http://species-id.net/wiki/Phyllocoptes_beishaniensis

Figures 8–11

Description. Female. (n = 13) Body fusiform, light yellow, 168 (160–178), 63 (66–67) wide, 61 (61–68) thick. **Gnathosoma** 20 (20–25), projecting obliquely down, suboral plate present, pedipalp coxal seta (*ep*) 6 (5–6), dorsal pedipalp genual seta (*d*) 6 (6–7), cheliceral stylets 16 (16–20). **Prodorsal shield** 48 (46–48), 65 (60–65) wide, subtriangular; frontal lobe 10 (9–10); median, admedian and submedian lines present, median line ending at basal 1/2 of prodorsal shield and connected with admedian lines at basal 1/4. Scapular tubercles ahead of rear shield margin, 2 (2–3), 22 (21–22) apart, scapular setae (*sc*) 10 (8–11), projecting centrad. **Coxigenital region** with 5 (4–5) smooth annuli. Coxisternal plates smooth, anterolateral setae on coxisternum I (*lb*) 8 (7–8), 13 (13–14) apart, proximal setae on coxisternum I (*la*) 21 (17–21), 10 (10–11) apart, proximal setae on coxisternum II (*2a*) 42 (42–45), 30 (30–31) apart, tubercles *lb* and *la* 9 (9–10) apart, tubercles *la* and *2a* 10 (10–11) apart. Prosternal apodeme combined 6 (6–7). **Leg I** 33 (33–34), femur 11 (10–11), basiventral femoral seta (*bv*) 12 (12–13); genu 6 (5–6), antaxial genual seta (*l''*) 23 (22–23); tibia 8 (8–9), paraxial tibial seta (*l'*) 7 (6–7), located at 1/3 from dorsal base; tarsus 6 (6–7), seta *ft'* 18 (18–19), seta *ft''* 22 (22–23), seta *u'* 5 (5–6); tarsal empodium (*em*) 8 (8–9), simple, 6-rayed, tarsal solenidion (ω) 7 (6–7), knobbed. **Leg II** 31 (29–31), femur 10 (10–11), basiventral femoral seta (*bv*) 10 (9–10); genu 6 (5–6), antaxial genual seta (*l''*) 10 (8–10); tibia 6 (6–7); tarsus 6 (6–7), seta *ft'* 7 (6–7), seta *ft''* 21 (21–22), seta *u'* 5 (5–6); tarsal empodium (*em*) 8 (8–9), simple, 6-rayed, tarsal solenidion (ω) 8 (8–9), knobbed. **Opisthosoma** dorsally with 45 (45–53) annuli, smooth; ventrally with 52 (51–52) annuli, with round microtubercles. Setae *c2* 30 (29–30) on ventral annulus 10 (9–13), 55 (54–55) apart; setae *d* 60 (55–60) on ventral annulus 20 (19–20), 30 (30–31) apart; setae *e* 40 (40–45) on ventral annulus 31 (30–31), 15 (15–16) apart; setae *f* 25 (25–27) on ventral annulus 46 (45–46), 24 (24–25) apart. Setae *h1* 5 (4–5), *h2* 80 (80–85). **Female genitalia** 17 (17–18), 22 (22–23) wide, coverflap with 10 longitudinal ridges, setae *3a* 56 (55–56), 16 (16–17) apart.

Male. (n = 9) Body fusiform, light yellow, 169–195, 56–67 wide. **Gnathosoma** 19–22, projecting obliquely down, suboral plate present, pedipalp coxal seta (*ep*) 4–5, dorsal pedipalp genual seta (*d*) 6–7, cheliceral stylets 17–18. **Prodorsal shield** has the same design as female, 42–50, 49–56 wide, subtriangular; frontal lobe 8–9. Scapular tubercles ahead of rear shield margin, 2–3, 18–19 apart, scapular setae (*sc*) 8–9, projecting centrad. **Coxigenital region** with 5 smooth annuli. Coxisternal plates smooth, anterolateral setae on coxisternum I (*lb*) 5–6, 12–15 apart, proximal setae on coxisternum I (*la*) 13–14, 10–11 apart, proximal setae on coxister-




Figure 8. *Phyllocoptes beishaniensis* sp. n.: **D** dorsal view of female CMG coxae and male genitalia **em** empodium **L1** leg I **L2** leg II.




Figure 9. *Phyllocoptes beishaniensis* sp. n.: **L** lateral view of female **LO** lateral microtubercles **IG** female internal genitalia **CG** coxae and female genitalia




Figure 10. *Phyllocoptes beishaniensis* sp. n.: **A** dorsal view of female **B** ventral view of female **C** lateral microtubercles **D** empodium **E** dorsal view of female posterior part **F** ventral view of female posterior part **G** leg I and leg II.




Figure 11. *Phyllocoptes beishaniensis* sp. n.: **H** lateral view of female **I** lateral view of female posterior part **J** female internal genitalia **K** prodorsal shield **L** coxae and female genitalia **M** coxae and male genitalia.

num **II** (2a) 22–27, 27–31 apart, tubercles *1b* and *1a* 8–9 apart, tubercles *1a* and *2a* 9–10 apart. Prosternal apodeme combined, 5–7. **Leg I** 27–32, femur 10–11, basiventral femoral seta (*bv*) 11–12; genu 5–6, antaxial genual seta (*l'*) 21–22; tibia 6–7, paraxial tibial seta (*l*) 5–6, located at 1/3 from dorsal base; tarsus 6–7, seta *ft'* 18–19, seta *ft''* 21–22, seta *u'* 5–6; tarsal empodium (*em*) 7–9, simple, 6-rayed, tarsal solenidion (ω) 6–7, knobbed. **Leg II** 27–30, femur 9–10, basiventral femoral seta (*bv*) 9–10; genu 5–6, antaxial genual seta (*l'*) 7–9; tibia 5–6; tarsus 5–6, seta *ft'* 5–6, seta *ft''* 17–19, seta *u'* 5–6; tarsal empodium (*em*) 7–8, simple, 6-rayed, tarsal solenidion (ω) 7–8, knobbed. **Opisthosoma** dorsally with 52 annuli, smooth, ventrally with 58 annuli, with round microtubercles. Setae *c2* 21–23 on ventral annulus 11, 46–53 apart; setae *d* 29–30 on ventral annulus 21, 25–28 apart; setae *e* 25–28 on ventral annulus 35, 12–13 apart; setae *f* 20 (20–23) on ventral annulus 54, 20–21 apart. Setae *h1* 4–5, *h2* 42–45. **Male genitalia** 19–21 wide, setae *3a* 13–14, 17–19 apart.

Type material. **Holotype**, female (slide number NJAUEri815, marked Holotype), from *Spiraea mongolica* Maxim. (Rosaceae), Beishan National Forest Park, Huzhu County, Qinghai Province, P. R. China, 36°53'35"N, 102°25'56"E, elevation 2610m, 22 July 2007, coll. Xiao-Feng Xue. **Paratypes**, 7 females and 2 males (slide number NJAUEri815), with the same data as holotype.

Relation to host. Vagrant on leaf lower surface.

Etymology. The specific designation *beishaniensis* is from the place name Beishan National Forest Park, where the new species were collected; feminine in gender.

Differential diagnosis. This species is similar to *Phyllocoptes adalius* (Keifer, 1939) from *Rosa* sp., but can be differentiated from the latter by prodorsal shield front lobe stout (prodorsal shield front lobe pointed in *P. adalius*), dorsal opisthosoma annuli smooth (opisthosoma annuli entirely covered with spinuliferous microtubercles in *P. adalius*), Coxisternal plates smooth (coxisternal plates with short lines in *P. adalius*).

***Phyllocoptes asperatae* Song, Xue & Hong, 2006**

http://species-id.net/wiki/Phyllocoptes_asperatae

Phyllocoptes asperatae Song et al. 2006: 36–38, figure 2.

Phyllocoptes asperatae; Xue et al. 2009:134.

Phyllocoptes asperatae; Song et al. 2009c: 37.

Material examined. 4 females and 2 males (slide number NJAUEri810), from a new host, *Picea meyeri* Rehd. Et Wils. (Pinaceae), Beishan National Forest Park, Huzhu County, Qinghai Province, P. R. China, 36°53'35"N, 102°25'56"E, elevation 2610m, 22 July 2007, coll. Xiao-Feng Xue.

Host. *Picea asperata* Mast. (Pinaceae); *Picea meyeri* Rehd. Et Wils. (Pinaceae)

Relation to host. Vagrant on leaf surface. No damage to the host was observed.

Distribution. China (Shaanxi, Qinghai).

***Phyllocoptes dangchangi* Song, Xue & Hong, 2006**

http://species-id.net/wiki/Phyllocoptes_dangchangi

Phyllocoptes dangchangi Song et al. 2006: 38–40, figure 3.

Phyllocoptes dangchangi; Song et al. 2008: 29.

Phyllocoptes dangchangi; Xue et al. 2009:134.

Phyllocoptes dangchangi; Song et al. 2009c: 36.

Material examined. 11 females and 1 male (slide number NJAUeri811), from *Picea* sp. (Pinaceae), Beishan National Forest Park, Huzhu County, Qinghai Province, P. R. China, 36°53'35"N, 102°25'56"E, elevation 2610m, 22 July 2007, coll. Xiao-Feng Xue.

Host. *Picea asperata* Mast. (Pinaceae); *Picea* sp. (Pinaceae).

Relation to host. Vagrant on terminal part of the needles. No damage to the host was observed.

Distribution. China (Gansu, Qinghai).

***Phyllocoptes gansuensis* Kuang & Luo, 1998**

http://species-id.net/wiki/Phyllocoptes_gansuensis

Phyllocoptes gansuensis Kuang et al. 1998: 201–203, figures 25–29.

Phyllocoptes gansuensis; Huang 2001: 45.

Phyllocoptes gansuensis; Kuang et al. 2005: 66–67, figure 64.

Phyllocoptes gansuensis; Song et al. 2008: 30.

Phyllocoptes gansuensis; Xue et al. 2009:134.

Phyllocoptes gansuensis; Song et al. 2009c: 37.

Material examined. 8 females and 2 males (slide number NJAUeri824), from a new host, *Potentilla parvifolia* Fisch. ap. Lehm. (Rosaceae), Beishan National Forest Park, Huzhu County, Qinghai Province, P. R. China, 36°53'35"N, 102°25'56"E, elevation 2610m, 22 July 2007, coll. Xiao-Feng Xue.

Host. *Potentilla glabra* Lodd. (Rosaceae); *Potentilla parvifolia* Fisch. et. Lehm. (Rosaceae).

Relation to host. Vagrant on leaf lower surface. No damage to the host was observed.

Distribution. China (Gansu, Qinghai).

Genus *Phyllocoptruta* Keifer, 1938***Phyllocoptruta platyclada* Xue, Song, Amrine & Hong, 2007**

http://species-id.net/wiki/Phyllocoptruta_platyclada

Phyllocoptruta platyclada Xue et al. 2007: 340–342, figure 3.

Phyllocoptruta platyclada; Xue et al. 2010: 698.

Material examined. 8 females (slide number NJAUEri814), from a new host, *Juniperus chinensis* L. (Cupressaceae), Beishan National Forest Park, Huzhu County, Qinghai Province, P. R. China, 36°53'35"N, 102°25'56"E, elevation 2610m, 22 July 2007, coll. Xiao-Feng Xue.

Host. *Platycladus orientalis* (Linn.) Franco (Cupressaceae); *Juniperus chinensis* L. (Cupressaceae).

Relation to host. Vagrant on terminal part of the needles. No damage to the host was observed.

Distribution. China (Shaanxi, Qinghai).

Tribe Anthocoptini Amrine & Stasny, 1994**Genus *Aculus* Keifer, 1959*****Aculus changbais* Xue, Song & Hong, 2008**

http://species-id.net/wiki/Aculus_changbais

Aculus changbais Xue et al. 2008: 41–42, figure 3.

Aculus changbais; Song et al. 2009b: 3.

Material examined. 8 females and 1 male (slide number NJAUEri778), from a new host, *Salix chaenomeloides* Kimura (Salicaceae), Mengda Natural Reserve, Xunhua County, Qinghai Province, P. R. China, 35°47'38"N, 102°40'40"E, elevation 2523m, 19 July 2007, coll. Xiao-Feng Xue.

Host. *Salix gracilistyla* Miq. (Salicaceae); *Salix chaenomeloides* Kimura (Salicaceae).

Relation to host. Vagrant on leaf lower surface. No damage to the host was observed.

Distribution. China (Jilin, Qinghai).

***Aculus huangzhongensis* Kuang, 2000**

http://species-id.net/wiki/Aculus_huangzhongensis

Aculus huangzhongensis Kuang 2000: 392–393, figures 13–18.

Aculus huangzhongensis; Kuang et al. 2005: 91–92, figure 91.

Aculus huangzhongensis; Song et al. 2009b: 2.

Host. *Syringa oblata* Lindl. (Oleaceae).

Relation to host. Vagrant on leaf surface. No damage to the host was observed.

Distribution. China (Qinghai).

Genus *Aculodes* Keifer, 1966

Aculodes salicis Kuang, 1997

http://species-id.net/wiki/Aculodes_salicis

Aculodes salicis Kuang 1997: 232–233, figures 12–15.

Aculodes salicis; Skoracka 2003: 43.

Aculodes salicis; Kuang et al. 2005: 80–81, figure 79.

Host. *Salix babylonica* L. (Salicaceae).

Relation to host. Forming galls on the leaf of the host.

Distribution. China (Qinghai), Poland.

Genus *Aculops* Keifer, 1966

Aculops ulmi Hong & Xue, 2005

http://species-id.net/wiki/Aculops_ulmi

Figures 12–15

Aculops ulmi Hong and Xue 2005: 205, 209.

Redescription. Female. (n = 10) Body fusiform, light yellow, 192 (192–230), 70 (62–72) wide, 80 (80–81) thick. **Gnathosoma** 23 (23–25), projecting obliquely down, suboral plate present, pedipalp coxal seta (*ep*) 3 (2–4), dorsal pedipalp genual seta (*d*) 6 (5–6), cheliceral stylets 22 (22–25). **Prodorsal shield** 33 (33–34), 48 (48–51) wide, subtriangular; median, admedian and submedian lines present, median line ending at basal 1/4 of prodorsal shield, median and admedian lines connected at basal 1/4 of prodorsal shield, admedian and submedian lines connected at basal 2/3 of prodorsal shield, forming two cells on both sides of median line. Scapular tubercles on rear shield margin, 4 (4–5), 26 (26–28) apart, scapular setae (*sc*) 55 (55–60), projecting posteriorly, knobbed at the end. **Coxigenital region** with 7 (6–7) annuli, with triangular microtubercles. Coxisternal plates with short lines and granules, anterolateral setae on coxisternum **I** (*Ib*) 13 (10–13), 14 (14–15) apart, proximal setae on coxisternum **I** (*Ia*) 30 (27–30), 11 (11–13) apart, proximal setae on coxisternum **II** (*2a*) 54 (54–57), 26 (26–28) apart, tubercles *Ib* and *Ia* 7 (7–8) apart,




Figure 12. *Aculops ulmi* Hong & Xue: **D** dorsal view of female **L1** leg I **L2** leg II **em** empodium **IG** female internal genitalia.




Figure 13. *Aculops ulmi* Hong & Xue: L lateral view of female LO lateral microtubercles CG coxae and female genitalia.




Figure 14. *Aculops ulmi* Hong & Xue: **A** dorsal view of female **B** ventral view of female **C** lateral view of female **D** dorsal view of female posterior part **E** ventral view of female posterior part **F** lateral microtubercles **G** lateral view of female posterior part **H** female internal genitalia.




Figure 15. *Aculops ulmi* Hong & Xue: **I** female genitalia **J** female genitalia from holotype.

tubercles *1a* and *2a* 9 (9–10) apart. Prosternal apodeme combined, 7 (5–11). **Leg I** 31 (31–35), femur 10 (9–12), basiventral femoral seta (*bv*) 12 (11–14); genu 5 (5–6), antiaxial genual seta (*l'*) 22 (22–29); tibia 7 (7–9), paraxial tibial seta (*l*) 6 (6–8), located at 1/3 from dorsal base; tarsus 8 (8–11), seta *ft'* 16 (16–18), seta *ft''* 20 (20–25), seta *u'* 6 (5–6); tarsal empodium (*em*) 6 (6–7), simple, 2-rayed, tarsal solenidion (ω) 9 (8–10), slightly knobbed. **Leg II** 28 (28–31), femur 9 (9–11), basiventral femoral seta (*bv*) 12 (11–12); genu 5 (5–6), antiaxial genual seta (*l'*) 9 (9–13); tibia 5 (5–7); tarsus 9 (9–10), seta *ft'* 7 (6–7), seta *ft''* 18 (18–20), seta *u'* 5 (4–5); tarsal empodium (*em*) 6 (6–7), simple, 2-rayed, tarsal solenidion (ω) 8 (8–11), slightly knobbed. **Opisthosoma** dorsally with 35 (22–38) annuli, with triangular microtubercles, ventrally with 55 (55–56) annuli, with triangular microtubercles. Setae *c2* 16 (14–16) on ventral annulus 10 (10–11), 61 (61–69) apart; setae *d* 57 (55–65) on ventral annulus 21 (21–22), 50 (48–50) apart; setae *e* 14 (12–19) on ventral annulus 33 (32–34), 23 (23–24) apart; setae *f* 27 (26–30) on ventral annulus 51 (49–53), 21 (19–21) apart. Setae *h1* 3 (3–4), *h2* 90 (85–90). **Female genitalia** 11 (11–14), 23 (22–23) wide, coverflap with 8 longitudinal ridges, setae *3a* 17 (17–22), 17 (16–17) apart.

Male. Unknown.

Type material. Hong and Xue (2005) described types as follows: Holotype female (slide number 17.viii.2003), from *Ulmus* sp. (Ulmaceae), Xingtai city, Hebei Province, P. R. China, coll. Xiao-Feng Xue. Paratypes (slide number 17.2003). 9 females and 2 males. All types here were re-examined. Female genitalia coverflap with 10–12 longitudinal ridges.

Additional material. 6 females (slide number NJAUERi792) from *Ulmus* sp. (Ulmaceae), Xining, Qinghai Province, P. R. China, 36°38'18"N, 101°45'27"E, elevation 2241m, 21 July 2007, coll. Xiao-Feng Xue.

Relation to host. Vagrant on leaf lower surface. No damage to the host was observed.

Distribution. China (Hebei, Qinghai).

Notes. Instead of the original description: female genitalia coverflap smooth, female genitalia coverflap is with 8 longitudinal ridges in this redescription.

***Aculops xiningensis* Kuang, 2000**

http://species-id.net/wiki/Aculops_xiningensis

Aculops xiningensis Kuang 2000: 392, figures 7–12.

Aculops xiningensis; Kuang et al. 2005: 86–87, figure 86.

Host. *Malus pumila* P. Mill. (Rosaceae).

Relation to host. Vagrant on leaf lower surface. No damage to the host was observed.

Distribution. China (Qinghai).

Genus *Tetraspinus* Boczek, 1961***Tetraspinus syringae* Lin & Kuang, 2001**

http://species-id.net/wiki/Tetraspinus_syringae

Tetraspinus syringae Lin and Kuang 2001: 351–353, figures 11–16.

Tetraspinus syringae; Kuang et al. 2005: 125–126, figure 128.

Host. *Syringa oblata* Lindl. (Oleaceae).

Relation to host. Vagrant on leaf lower surface. No damage to the host was observed.

Distribution. China (Qinghai).

Genus *Tetra* Keifer, 1944***Tetra pinnatifidae* Xue, Song & Hong, 2006**

http://species-id.net/wiki/Tetra_pinnatifidae

Tetra pinnatifidae Xue et al. 2006a: 6–8, figure 2.

Material examined. 8 females and 2 males (slide number NJAUEr1793) from a new host, *Prunus armeniaca* Linn. (Rosaceae), Xining, Qinghai Province, P. R. China, 36°38'18"N, 101°45'27"E, elevation 2241m, 21 July 2007, coll. Xiao-Feng Xue.

Host. *Crataegus pinnatifida* Bunge (Rosaceae); *Prunus armeniaca* Linn. (Rosaceae).

Relation to host. Vagrant on leaf lower surface. No damage to the host was observed.

Distribution. China (Shaanxi, Qinghai).

***Tetra pruniana* sp. n.**

urn:lsid:zoobank.org:act:D0B42704-F828-4F47-9265-47B3EF8E603A

http://species-id.net/wiki/Tetra_pruniana

Figures 16–18

Description. Female. (n = 9) Body fusiform, light yellow, 215 (210–225), 77 (75–78) wide. **Gnathosoma** 21 (21–23), projecting obliquely down, suboral plate present,

pedipalp coxal seta (*ep*) 4 (4–5), dorsal pedipalp genual seta (*d*) 7 (7–8), cheliceral stylets 16 (16–18). **Prodorsal shield** 54 (54–55), 75 (75–78) wide, subtriangular; frontal lobe 12 (11–12); median and submedian lines absent, admedian lines connected by two weak transverse lines. Scapular tubercles near rear shield margin, 3 (3–4), 34 (32–34) apart, scapular setae (*sc*) 14 (13–15), projecting posteriorly. Rear shield with wave-like margin. **Coxigenital region** with 9 annuli. Coxisternal plates with short lines, anterolateral setae on coxisternum I (*lb*) 11 (10–13), 14 (13–14) apart, proximal setae on coxisternum I (*la*) 20 (20–25), 10 (10–11) apart, proximal setae on coxisternum II (*2a*) 50 (45–50), 28 (28–29) apart, tubercles *lb* and *la* 7 (7–8) apart, tubercles *la* and *2a* 10 (10–12) apart. Prosternal apodeme combined, 8 (6–8). **Leg I** 34 (32–35), femur 12 (12–13), basiventral femoral seta (*bv*) 12 (12–14); genu 7 (6–7), antaxial genual seta (*l'*) 20 (19–20); tibia 10 (9–10), paraxial tibial seta (*l*) 6 (5–6), located at 1/3 from dorsal base; tarsus 7 (7–8), seta *ft'* 19 (19–21), seta *ft''* 25 (24–25), seta *u'* 5 (4–5); tarsal empodium (*em*) 7 (6–7), simple, 4-rayed, tarsal solenidion (ω) 6 (6–7), knobbed. **Leg II** 27 (27–31), femur 10 (10–11), basiventral femoral seta (*bv*) 12 (12–13); genu 5 (5–6), antaxial genual seta (*l'*) 8 (7–8); tibia 6 (6–7); tarsus 6 (6–7), seta *ft'* 6 (6–7), seta *ft''* 25 (23–25), seta *u'* 5 (5–6); tarsal empodium (*em*) 6 (5–6), simple, 4-rayed, tarsal solenidion (ω) 6 (6–7), knobbed. **Opisthosoma** dorsally with 25 (24–25) annuli, with weak filamentous microtubercles, ventrally with 53 (53–59) annuli, with round microtubercles. Setae *c2* 25 (24–26) on ventral annulus 9 (9–11), 58 (58–59) apart; setae *d* 60 (57–65) on ventral annulus 22 (22–24), 33 (32–33) apart; setae *e* 18 (17–20) on ventral annulus 37 (37–40), 18 (17–18) apart; setae *f29* (29–33) on ventral annulus 50 (50–56), 25 (22–25) apart. Setae *h1* 3 (2–3), *h2* 85 (80–95). **Female genitalia** 15 (15–17), 25 (24–25) wide, coverflap with 10 longitudinal ridges, setae *3a* 18 (16–19), 17 (17–18) apart.

Male. (n = 6) Body fusiform, light yellow, 170–186, 73–74 wide. **Gnathosoma** 16–17, projecting obliquely downwards, suboral plate present, pedipalp coxal seta (*ep*) 4–5, dorsal pedipalp genual seta (*d*) 7–8, cheliceral stylets 14–16. **Prodorsal shield** has the same design as female, 47–50, 72–73 wide, subtriangular; frontal lobe 10–12. Scapular tubercles on rear shield margin, 3–4, 30–33 apart, scapular setae (*sc*) 7–9, projecting posteriorly. **Coxigenital region** with 9 smooth annuli. Coxisternal plates with short lines, anterolateral setae on coxisternum I (*lb*) 10–11, 13–14 apart, proximal setae on coxisternum I (*la*) 21–22, 10–11 apart, proximal setae on coxisternum II (*2a*) 45–48, 28–30 apart, tubercles *lb* and *la* 7–8 apart, tubercles *la* and *2a* 9–10 apart. Prosternal apodeme combined, 7–8. **Leg I** 34–35, femur 10–12, basiventral femoral seta (*bv*) 9–10; genu 6–7, antaxial genual seta (*l'*) 19–22; tibia 9–10, paraxial tibial seta (*l*) 6–7, located at 1/3 from dorsal base; tarsus 7–8, seta *ft'* 15–17, seta *ft''* 21–24, seta *u'* 4–5; tarsal empodium (*em*) 7–8, simple, 4-rayed, tarsal solenidion (ω) 6–7, knobbed. **Leg II** 28–29, femur 9–10, basiventral femoral seta (*bv*) 10–11; genu 5–6, antaxial genual seta (*l'*) 6–7; tibia 7–8; tarsus 6–7, seta *ft'* 6–7, seta *ft''* 22–25, seta *u'* 5–6; tarsal empodium (*em*) 6–7, simple, 4-rayed, tarsal solenidion (ω) 7–8, knobbed. **Opisthosoma** dorsally with 26–27 annuli, with weak filamentous microtubercles, ventrally with 53–54 annuli, with round microtubercles. Setae *c2* 18–20 on ventral annulus 10–11, 55–65 apart; setae *d* 32–35 on ventral annulus 20–22, 30–45




Figure 16. *Tetra pruniana* sp. n.: **D** dorsal view of female **em** empodium **L1** leg I **L2** leg II.

apart; setae *e* 18–20 on ventral annulus 33–35, 18–30 apart; setae *f* 25–28 on ventral annulus 50–51, 20–22 apart. Setae *h1* 2–3, *h2* 80–90. **Male genitalia** 13–14, 23–28 wide, setae *3a* 17–18, 20–21 apart.




Figure 17. *Tetra pruniana* sp. n.: V ventral view of female IG female internal genitalia CMG coxae and male genitalia.

Type material. **Holotype**, female (slide number NJAUEri808, marked Holo-type), from *Prunus tomentosa* Thunb. (Rosaceae), Xining City, Qinghai Province, P. R. China, 36°38'18"N, 101°45'27"E, elevation 2241m, 21 July 2007, coll. Xiao-Feng Xue. **Paratypes**, 2 females and 4 males (slide number NJAUEri808), with the same data as holotype.




Figure 18. *Tetra pruniana* sp. n.: **A** dorsal view of female **B** ventral view of female **C** dorsal view of female posterior part **D** ventral view of female posterior part **E** prodorsal shield **F** coxae and female genitalia **G** female internal genitalia **H** male genitalia **I** empodium.

Relation to host. Vagrant on leaf lower surface. No damage to the host was observed.

Etymology. The specific designation *pruniana* is from the generic name of host plant, *Prunus*; feminine in gender.

Differential diagnosis. This species is similar to *Tetra pinnatifidae* Xue, Song & Hong, 2006a, but can be differentiated from the latter by median and submedian lines absent (median and submedian lines present in *T. pinnatifidae*), admedian lines connected by two weak transverse lines (admedian lines separated in *T. pinnatifidae*).

***Tetra pyriana* sp. n.**

urn:lsid:zoobank.org:act:99B1B2A9-DB3F-466C-8900-951BB3B35485

http://species-id.net/wiki/Tetra_pyriana

Figures 19–22

Description. Female. (n = 11) Body fusiform, light yellow, 180 (169–185), 73 (72–76) wide, 65 (64–65) thick. **Gnathosoma** 20 (20–23), projecting obliquely down, pedipalp coxal seta (*ep*) 4 (3–4), dorsal pedipalp genual seta (*d*) 6 (5–7), cheliceral stylets 20 (20–21). **Prodorsal shield** 47 (47–49), 68 (68–72) wide, subtriangular; frontal lobe 12 (11–13); median, admedian and submedian lines present, median line obscure and ending at basal 1/3 of prodorsal shield. Scapular tubercles on rear shield margin, 4 (3–4), 36 (33–36) apart, scapular setae (*sc*) 12 (12–13), projecting posteriorly. **Coxigenital region** with 13 (12–13) smooth annuli. Coxisternal plates with short lines, anterolateral setae on coxisternum I (*lb*) 7 (7–9), 12 (11–12) apart, proximal setae on coxisternum I (*la*) 21 (20–21), 10 (9–10) apart, proximal setae on coxisternum II (*2a*) 40 (40–45), 27 (25–27) apart, tubercles *lb* and *la* 6 (6–7) apart, tubercles *la* and *2a* 10 (9–10) apart. Prosternal apodeme combined, 7 (6–7). **Leg I** 32 (30–32), femur 12 (11–12), basiventral femoral seta (*bv*) 11 (10–11); genu 6 (5–6), antaxial genual seta (*l''*) 20 (20–21); tibia 7 (7–8), paraxial tibial seta (*l'*) 5 (5–6), located at 1/3 from dorsal base; tarsus 8 (7–8), seta *ft'* 19 (17–19), seta *ft''* 25 (21–25), seta *u'5* (4–5); tarsal empodium (*em*) 6 (6–7), simple, 4-rayed, tarsal solenidion (ω) 6 (6–7), knobbed. **Leg II** 28 (27–28), femur 11 (10–11), basiventral femoral seta (*bv*) 8 (8–11); genu 5 (4–5), antaxial genual seta (*l''*) 8 (7–8); tibia 7 (6–7); tarsus 7 (6–7), seta *ft'* 5 (5–6), seta *ft''* 20 (19–20), seta *u'4* (4–5); tarsal empodium (*em*) 6 (6–7), simple, 4-rayed, tarsal solenidion (ω) 5 (5–6), knobbed. **Opisthosoma** dorsally with 32 (30–32) annuli, with filamentous microtubercles, ventrally with 58 (58–60) annuli, with round microtubercles. Setae *c2* 25 (24–30) on ventral annulus 11 (11–13), 57 (54–57) apart; setae *d* 45 (45–50) on ventral annulus 22 (22–24), 32 (32–33) apart; setae *e* 18 (17–18) on ventral annulus 37 (37–41), 17 (16–17) apart; setae *f29* (28–30) on ventral annulus 53 (53–54), 25 (24–25) apart. Setae *h1* 3 (2–3), *h2* 85 (80–90). **Female genitalia** 15 (14–16), 22 (22–23) wide, coverflap with 11 (10–12) longitudinal ridges, setae *3a* 22 (19–24), 14 (14–15) apart.




Figure 19. *Tetra pyriana* sp. n.: **D** dorsal view of female **LO** lateral microtubercles **em** empodium **L1** leg I **L2** leg II.




Figure 20. *Tetra pyriana* sp. n.: L lateral view of female IG female internal genitalia CG coxae and female genitalia CMG coxae and male genitalia.



Figure 21. *Tetra pyriana* sp. n.: **A** dorsal view of female **B** ventral view of female **C** lateral microtubercles **D** empodium **E** dorsal view of female posterior part **F** ventral view of female posterior part **G** leg I and leg II.




Figure 22. *Tetra pyriana* sp. n.: **H** lateral view of female **I** lateral view of female posterior part **J** female internal genitalia **K** prodorsal shield **L** coxae and female genitalia **M** coxae and male genitalia.

Male. (n = 2) Body fusiform, light yellow, 150–176 63–65 wide. **Gnathosoma** 21–22, projecting obliquely down, pedipalp coxal seta (*ep*) 2–3, dorsal pedipalpal genual seta (*d*) 6–7, cheliceral stylets 19–20. **Prodorsal shield** has the same design as female, 42–47, 63–65 wide, subtriangular; frontal lobe 10–11. Scapular tubercles on rear shield margin, 3–4, 30–31 apart, scapular setae (*sc*) 11–13, projecting posteriorly. **Coxigenital region** with 12–13 smooth annuli. Coxisternal plates with short lines, anterolateral setae on coxisternum I (*lb*) 7–8, 12–13 apart, proximal setae on coxisternum I (*la*) 18–20, 9–10 apart, proximal setae on coxisternum II (*2a*) 45–46, 24–26 apart, tubercles *lb* and *la* 6–7 apart, tubercles *la* and *2a* 7–9 apart. Prosternal apodeme combined, 5–6. **Leg I** 29–32, femur 7–9, basiventral femoral seta (*bv*) 9–12; genu 5–6, antaxial genual seta (*l''*) 20–23; tibia 6–8, paraxial tibial seta (*l'*) 5–6, located at 1/3 from dorsal base; tarsus 7–8, seta *ft'* 17–18, seta *ft''* 21–23, seta *u'* 3–4; tarsal empodium (*em*) 6–7, simple, 4-rayed, tarsal solenidion (*ω*) 5–6, knobbed. **Leg II** 25–27, femur 7–9, basiventral femoral seta (*bv*) 9–10; genu 4–5, antaxial genual seta (*l''*) 6–8; tibia 7–8; tarsus 6–7, seta *ft'* 2–3, seta *ft''* 20–21, seta *u'* 2–3; tarsal empodium (*em*) 6–7, simple, 4-rayed, tarsal solenidion (*ω*) 6–7, knobbed. **Opisthosoma** dorsally with 29–31 annuli, with filamentous microtubercles, ventrally with 60–61 annuli, with round microtubercles. Setae *c2* 24–25 on ventral annulus 13–14, 50–51 apart; setae *d* 36–37 on ventral annulus 24–25, 29–30 apart; setae *e* 15–16 on ventral annulus 41–42, 16–17 apart; setae *f* 28–29 on ventral annulus 55–56, 20–21 apart. Setae *h1* 2–3, *h2* 85–90. **Male genitalia** 13–14, 19–20 wide, setae *3a* 17–18, 16–17 apart.

Type material. **Holotype**, female (slide number NJAUEri796, marked Holotype), from *Pyrus calleryana* Decne. (Rosaceae), Xining, Qinghai Province, P. R. China, 36°38'18"N, 101°45'27"E, elevation 2241m, 21 July 2007, coll. Xiao-Feng Xue. **Paratypes**, 10 females and 2 males (slide number NJAUEri796), with the same data as holotype.

Additional material. 13 females (slide number NJAUEri807), from *Pyrus betulifolia* Bunge. (Rosaceae), Xining, Qinghai Province, P. R. China, 36°38'18"N, 101°45'27"E, elevation 2241m, 21 July 2007, coll. Xiao-Feng Xue; 14 females and 1 male (slide number NJAUEri806), from *Pyrus calleryana* Decne. (Rosaceae), Xining, Qinghai Province, P. R. China, 36°38'18"N, 101°45'27"E, elevation 2241m, 21 July 2007, coll. Xiao-Feng Xue.

Relation to host. Vagrant on leaf lower surface. No damage to the host was observed.

Etymology. The specific designation *pyriana* is from the generic name of host plant, *Pyrus*; feminine in gender.

Differential diagnosis. This species is similar to *Tetra pinnatifidae* Xue, Song & Hong, 2006a, but can be differentiated from the latter by median line weak and ending at basal 1/3 of prodorsal shield (median normal and does not disappear at basal 1/3 of prodorsal shield in *T. pinnatifidae*), coxigenital region with smooth annuli (coxigenital region annuli with round microtubercles in *T. pinnatifidae*).

***Tetra simonia* sp. n.**

urn:lsid:zoobank.org:act:11D2D601-440B-4A7D-AD50-18F7B91B3A33

http://species-id.net/wiki/Tetra_simonia

Figures 23–26

Description. Female. (n = 14) Body fusiform, light yellow, 196 (196–224), 72 (72–77) wide, 78 (78) thick. **Gnathosoma** 22 (20–22), projecting obliquely down, pedipalp coxal seta (*ep*) 3 (3–4), dorsal pedipalp genual seta (*d*) 12 (10–12), cheliceral stylets 19 (19–20). **Prodorsal shield** 46 (42–46), 72 (72–77) wide, subtriangular; frontal lobe 11 (11–12); median, admedian and submedian lines robust and connected, forming an “M” shape in the middle and two “H” shapes anterior to the “M”. Scapular tubercles on rear shield margin, 3 (3–4), 46 (44–46) apart, scapular setae (*sc*) 14 (14–15), projecting posteriorly. **Coxigenital region** with 14 (14–15) smooth annuli. Coxisternal plates smooth, anterolateral setae on coxisternum I (*lb*) 11 (11–12), 14 (14–19) apart, proximal setae on coxisternum I (*la*) 17 (17–23), 12 (12–13) apart, proximal setae on coxisternum II (*2a*) 53 (53–58), 28 (27–31) apart, tubercles *lb* and *la* 8 (8–9) apart, tubercles *la* and *2a* 9 (9–10) apart. Prosternal apodeme combined, 6 (6–8). **Leg I** 34 (34–37), femur 12 (11–12), basiventral femoral seta (*bv*) 12 (12–13); genu 8 (8–9), antaxial genual seta (*l''*) 24 (22–24); tibia 13 (11–13), paraxial tibial seta (*l'*) 7 (6–7), located at 1/3 from dorsal base; tarsus 7 (7–8), seta *ft'* 22 (21–22), seta *ft''* 23 (22–23), seta *u'* 6 (6–7); tarsal empodium (*em*) 6 (6–7), simple, 4-rayed, tarsal solenidion (ω) 6 (6–7), knobbed. **Leg II** 30 (30–34), femur 11 (11–12), basiventral femoral seta (*bv*) 13 (12–13); genu 6 (6–7), antaxial genual seta (*l''*) 11 (11–13); tibia 9 (7–9); tarsus 7 (6–7), seta *ft'* 5 (5–6), seta *ft''* 22 (22–24), seta *u'* 6 (6–7); tarsal empodium (*em*) 6 (6–7), simple, 4-rayed, tarsal solenidion (ω) 6 (6–7), knobbed. **Opisthosoma** dorsally with 31 (30–31) annuli, with dark shading on rear annular margins, ventrally with 58 (58–65) annuli, with round microtubercles. Setae *c2* 36 (36–42) on ventral annulus 11 (11–13), 52 (52–56) apart; setae *d* 77 (71–77) on ventral annulus 21 (21–23), 33 (33–38) apart; setae *e* 23 (20–23) on ventral annulus 37 (37–39), 18 (18–19) apart; setae *f* 40 (36–40) on ventral annulus 51 (51–54), 30 (30–32) apart. Setae *h1* 5 (4–5), *h2* 150 (135–150). **Female genitalia** 20 (18–22), 25 (24–25) wide, coverflap with 10 (10–13) longitudinal ridges, setae *3a* 17 (17–18), 18 (17–18) apart.

Male. (n = 1) Body fusiform, light yellow, 188, 74 wide. **Gnathosoma** 21, projecting obliquely downwards, pedipalp coxal seta (*ep*) 3, dorsal pedipalp genual seta (*d*) 11, cheliceral stylets 19. **Prodorsal shield** has the same design as female, 42, 74 wide, subtriangular; frontal lobe 11. Scapular tubercles on rear shield margin, 44 apart, scapular setae (*sc*) 13, projecting posteriorly. **Coxigenital region** with 14 smooth annuli. Coxisternal plates smooth, anterolateral setae on coxisternum I (*lb*) 13, 14 apart, proximal setae on coxisternum I (*la*) 23, 12 apart, proximal setae on coxisternum II (*2a*) 49, 30 apart, tubercles *lb* and *la* apart 8, tubercles *la* and *2a* 10 apart. Prosternal apodeme combined, 6. **Leg I** 35, femur 11, basiventral femoral seta (*bv*) 12; genu 8, antaxial genual seta (*l''*)




Figure 23. *Tetra simonia* sp. n.: D dorsal view of female LO lateral microtubercles em empodium L1 leg I L2 leg II.




Figure 24. *Tetra simonia* sp. n.: L lateral view of female IG female internal genitalia CG coxae and female genitalia CMG coxae and male genitalia.




Figure 25. *Tetra simonia* sp. n.: **A** dorsal view of female **B** ventral view of female **C** lateral microtubercles **D** empodium **E** dorsal view of female posterior part **F** ventral view of female posterior part **G** leg I and leg II.




Figure 26. *Tetra simonia* sp. n.: **H** lateral view of female **I** lateral view of female posterior part **J** female internal genitalia **K** prodorsal shield **L** coxae and female genitalia **M** coxae and male genitalia.

26; tibia 12, paraxial tibial seta (*l*) 6, located at 1/3 from dorsal base; tarsus 7, seta *ft*' 21, seta *ft*" 24, seta *u*' 6; tarsal empodium (*em*) 6, simple, 4-rayed, tarsal solenidion (ω) 6, knobbed. **Leg II** 32, femur 10, basiventral femoral seta (*bv*) 14; genu 6, antaxial genual seta (*l*') 11; tibia 8; tarsus 8, seta *ft*' 5, seta *ft*" 18, seta *u*' 6; tarsal empodium (*em*) 6, simple, 4-rayed, tarsal solenidion (ω) 6, knobbed. **Opisthosoma** dorsally with 31 annuli, with dark shading on rear annular margins, ventrally with 67 annuli, with round microtubercles. Setae *c* 2 42 on ventral annulus 13, 58 apart; setae *d* 70 on ventral annulus 27, 40 apart; setae *e* 25 on ventral annulus 47, 20 apart; setae *f* 40 on ventral annulus 63, 30 apart. Setae *h* 1 5, *h* 2 130. **Male genitalia** 15, 25 wide, setae *3a* 18, 19 apart.

Type material. **Holotype**, female (slide number NJAUEri799, marked Holotype), from *Populus simonii* Carr. (Salicaceae), Xining, Qinghai Province, P. R. China, 36°38'18"N, 101°45'27"E, elevation 2241m, 21 July 2007, coll. Xiao-Feng Xue. **Paratypes**, 13 females and 1 male (slide number NJAUEri799), with the same data as holotype.

Additional material. 4 females (slide number NJAUEri789A), from *Populus simonii* Carr. (Salicaceae), Xining, Qinghai Province, P. R. China, 36°38'18"N, 101°45'27"E, elevation 2241m, 21 July 2007, coll. Xiao-Feng Xue. 4 females and 2 males (slide number NJAUEri823), from *Populus simonii* Carr. (Salicaceae), Beishan National Forest Park, Huzhu County, Qinghai Province, P. R. China, 36°53'35"N, 102°25'56"E, elevation 2610m, 22 July 2007, coll. Xiao-Feng Xue.

Relation to host. Vagrant on leaf lower surface. No damage to the host was observed.

Etymology. The specific designation *simonia* is from the species name of host plant, *simonii*; feminine in gender.

Differential diagnosis. This species is similar to *Tetra smilaxis* Xue, Song & Hong, 2006a, but can be differentiated from the latter by opisthosoma with dark shading on rear annular margins (dark shading absent in *T. smilaxis*), prodorsal shield as wide as opisthosoma (opisthosoma wider than prodorsal shield in *T. smilaxis*), admedian lines connected at basal 1/3 but separate at basal 2/3 of prodorsal shield (admedian lines connected at basal 1/3 and basal 2/3 of prodorsal shield in *T. smilaxis*).

Family Diptilomiopidae Keifer, 1944

Subfamily Diptilomiopinae Keifer, 1944

Genus *Diptacus* Keifer, 1951

Diptacus berberinus sp. n.

urn:lsid:zoobank.org:act:3EF374D2-74CD-46A0-833C-A80FC1EAE1E2

http://species-id.net/wiki/Diptacus_berberinus

Figures 27–30

Description. Female. (n = 9) Body fusiform, light yellow, 283 (280–360), 110 (102–110) wide, 115 (114–115) thick. **Gnathosoma** 26 (25–27), projecting downwards, pedipalp coxal seta (*ep*) 6 (5–6), dorsal pedipalp genual seta (*d*) 12 (11–12), cheliceral stylets 65 (65–66). **Prodorsal shield** 40 (40–46), 80 (77–80) wide, with wide and broad

frontal lobe, 7 (7–8); median, admedian and submedian lines present, admedian lines connected at the basal 1/3 and 2/3 of prodorsal shield, forming 3 cells on each side, submedian lines connected with the median and admedian at the basal 2/3 of prodorsal shield, forming the cell-like pattern at anterior shield margin. Scapular tubercles ahead of rear shield margin, 3 (2–3), 30 (27–30) apart, scapular setae (*sc*) 5 (4–5), projecting centrad to forward. **Coxigenital region** with 13 (13–15) annuli, with triangular microtubercles. Coxisternal plate I with granules, coxisternal plate II smooth, anterolateral setae on coxisternum I (*Ib*) 20 (18–20), 17 (17–18) apart, proximal setae on coxisternum I (*Ia*) 43 (43–45), 19 (17–19) apart, proximal setae on coxisternum II (*2a*) 70 (70–80), 49 (43–56) apart, tubercles *Ib* and *Ia* 12 (12–13) apart, tubercles *Ia* and *2a* 17 (14–17) apart. Prosternal apodeme separated, 5 (5–6). **Leg I** 65 (60–65), femur 20 (20–22), basiventral femoral seta (*bv*) absent; genu 8 (8–9), antaxial genual seta (*l''*) 48 (48–52); tibia 18 (17–19), paraxial tibial seta (*l*) 11 (10–11), located at 1/2 from dorsal base; tarsus 11 (11–12), seta *ft'* 30 (29–30), seta *ft''* 40 (37–40), seta *u'* 7 (6–7); tarsal empodium (*em*) 11 (10–11), divided, 7-rayed on each side, tarsal solenidion (ω) 10 (10–14), knobbed. **Leg II** 55 (54–55), femur 20 (19–20), basiventral femoral seta (*bv*) absent; genu 8 (7–8), antaxial genual seta (*l''*) 18 (15–18); tibia 17 (16–17); tarsus 11 (10–11), seta *ft'* 11 (10–11), seta *ft''* 44 (44–50), seta *u'* 8 (7–8); tarsal empodium (*em*) 11 (10–11), divided, 7-rayed on each side, tarsal solenidion (ω) 10 (10–11), knobbed. **Opisthosoma** dorsally with 59 (54–62) annuli, smooth, ventrally with 106 (101–106) annuli, with triangular microtubercles. Setae *c2* 115 (110–115) on ventral annulus 18 (18–20), 74 (74–75) apart; setae *d* 100 (100–120) on ventral annulus 40 (37–40), 56 (49–56) apart; setae *e* 60 (55–60) on ventral annulus 65 (60–65), 31 (29–35) apart; setae *f65* (60–70) on ventral annulus 93 (87–93), 35 (35–37) apart. Setae *h1* 2 (1–2), *h2* 103 (95–165). **Female genitalia** 35 (31–40), 35 (34–42) wide, coverflap with short lines on base, and 4 longitudinal ridges in 2 ranks, 1 ridge near the base and 3 ridges at distal margin, setae *3a* 12 (11–15), 24 (24–25) apart.

Male. (n = 1) Body fusiform, light yellow, 269, 86 wide. **Gnathosoma** 60, projecting downwards, pedipalp coxal seta (*ep*) 5, dorsal pedipalp genual seta (*d*) 11, cheliceral stylets 65. **Prodorsal shield** has the same design as female, 38, 71 wide, with wide and broad frontal lobe, 7. Scapular tubercles ahead of rear shield margin, 3, 27 apart, scapular setae (*sc*) 4, projecting centrad to forward. **Coxigenital region** with 14 annuli, with triangular microtubercles. Coxisternal plate I with granules, coxisternal plate II smooth, anterolateral setae on coxisternum I (*Ib*) 22, 16 apart, proximal setae on coxisternum I (*Ia*) 36, 16 apart, proximal setae on coxisternum II (*2a*) 60, 41 apart, tubercles *Ib* and *Ia* 11 apart, tubercles *Ia* and *2a* 12 apart. Prosternal apodeme separated, 6. **Leg I** 46, femur 17, basiventral femoral seta (*bv*) absent; genu 7, antaxial genual seta (*l''*) 46; tibia 13, paraxial tibial seta (*l*) 10, located at 1/2 from dorsal base; tarsus 8, seta *ft'* 30, seta *ft''* 37, seta *u'* 6; tarsal empodium (*em*) 10, divided, 7-rayed on each side, tarsal solenidion (ω) 10, knobbed. **Leg II** 38, femur 17, basiventral femoral seta (*bv*) absent; genu 7, antaxial genual seta (*l''*) 15; tibia 13; tarsus 6, seta *ft'* 10, seta *ft''* 38, seta *u'* 7; tarsal empodium (*em*) 10, divided, 7-rayed on each side, tarsal solenidion (ω) 11, knobbed. **Opisthosoma** dorsally with 56 annuli, smooth, ventrally with 83 annuli, with triangular microtuber-




Figure 27. *Diptacus berberinus* sp. n.: **D** dorsal view of female **IG** female internal genitalia **LO** lateral microtubercles **L1** leg I **L2** leg II **em** empodium.




Figure 28. *Diptacus berberinus* sp. n.: L lateral view of female CMG coxae and male genitalia CG coxae and female genitalia.




Figure 29. *Diptacus berberinus* sp. n.: **A** dorsal view of female **B** ventral view of female **C** lateral microtubercles **D** empodium **E** dorsal view of female posterior part **F** ventral view of female posterior part **G** leg I and leg II.




Figure 30. *Diptacus berberinus* sp. n.: **H** lateral view of female **I** coxae and female genitalia **J** lateral view of female posterior part **K** female internal genitalia **L** coxae and male genitalia **M** prodorsal shield.

cles. Setae *c*2 95 on ventral annulus 15, 63 apart; setae *d* 100 on ventral annulus 30, 49 apart; setae *e* 55 on ventral annulus 46, 27 apart; setae *f* 60 on ventral annulus 71, 25 apart. Setae *h*1 2, *h*2 120. **Male genitalia** 23, 30 wide, setae *3a* 10, 23 apart.

Type material. **Holotype**, female (slide number 783, marked Holotype), from *Berberis amurensis* Rupr. (Berberidaceae), Mengda Natural Reserve, Xunhua County, Qinghai Province, P. R. China, 35° 47' 38" N, 102° 40' 40" E, elevation 2523m, 19 July 2007, coll. Xiao-Feng Xue. **Paratypes**, 8 females and 1 male (slide number 783), with the same data as holotype.

Relation to host. Vagrant on leaf lower surface. No damage to the host was observed.

Etymology. The specific designation *berberinus* is from the generic name of host plant, *Berberis*; masculine in gender.

Differential diagnosis. This species is similar to *Diptacus maddenis* Song, Xue & Hong, 2007a, but can be differentiated from the latter by opisthosomal dorsal annuli smooth (opisthosomal dorsal annuli with elongated microtubercles in *D. maddenis*), female genital coverflap with short lines at the base (genital coverflap with granules in *D. maddenis*), tarsal empodium 7-rayed (4-rayed in *D. maddenis*).

***Diptacus mengdaensis* sp. n.**

urn:lsid:zoobank.org:act:12904959-7EE9-4780-BCC1-EF886FA54F98

http://species-id.net/wiki/Diptacus_mengdaensis

Figures 31–33

Description. Female. (n = 13) Body fusiform, light yellow, 215 (210–232), 104 (104–105) wide, 71 (71–74) thick. **Gnathosoma** 25 (24–25), projecting downwards, pedipalp coxal seta (*ep*) 4 (4–5), dorsal pedipalp genual seta (*d*) 14 (14–15), cheliceral stylets 62 (61–62). **Prodorsal shield** 50 (50–51), 76 (71–76) wide, with wide and broad frontal lobe, 7 (7–8); median, admedian and submedian lines present, admedian lines connected at the base of prodorsal shield, ending at basal 1/3 of prodorsal shield. Scapular tubercles ahead of rear shield margin, 3 (3–4), 34 (34–35) apart, scapular setae (*sc*) 6 (6–7), projecting centrad. **Coxigenital region** with 16 (15–16) annuli, with microtubercles. Coxisternal plates with short lines, anterolateral setae on coxisternum I (*lb*) 16 (16–18), 18 (18–20) apart, proximal setae on coxisternum I (*la*) 25 (25–26), 17 (17–18) apart, proximal setae on coxisternum II (*2a*) 60 (60–70), 44 (44–45) apart, tubercles *lb* and *la* 12 (12–13) apart, tubercles *la* and *2a* 16 (16–17) apart. Prosternal apodeme separated, 10 (9–10). **Leg I** 60 (58–60), femur 16 (16–17), basiventral femoral seta (*bv*) absent; genu 10 (8–10), antaxial genual seta (*l''*) 52 (52–53); tibia 17 (16–17), paraxial tibial seta (*l*) 10 (9–10), located at 1/3 from dorsal base; tarsus 11 (10–11), seta *ft'* 30 (25–30), seta *ft''* 40 (32–40), seta *u'* 6 (5–6); tarsal empodium (*em*) 8 (8–9), divided, 5-rayed at each side, tarsal solenidion (*ω*) 9 (9–10), knobbed. **Leg II** 54 (50–54), femur 18 (17–18), basiventral femoral seta (*bv*) absent; genu 7 (7–8), antaxial genual seta (*l''*) 16 (15–16); tibia 11 (9–11); tarsus 8 (8–10), seta *ft'* 12 (12–13), seta *ft''* 34 (34–34), seta *u'* 5 (5–6); tarsal empodium (*em*) 9 (8–9),




Figure 31. *Diptacus mengdaensis* sp. n.: **D** dorsal view of female **LO** lateral microtubercles **em** empodium **L1** leg I **L2** leg II.




Figure 32. *Diptacus mengdaensis* sp. n.: L lateral view of female IG female internal genitalia CG coxae and female genitalia.




Figure 33. *Diptacus mengdaensis* sp. n.: **A** dorsal view of female **B** ventral view of female **C** dorsal view of female posterior part **D** ventral view of female posterior part **E** prodorsal shield **F** lateral microtubercles **G** empodium **H** coxae and female genitalia.

divided, 5-rayed at each side, tarsal solenidion (ω) 9 (9–10), knobbed. **Opisthosoma** dorsally with 44 (44–48) annuli, smooth, ventrally with 112 (112–115) annuli, with round microtubercles. Setae *c*2 55 (54–55) on ventral annulus 16 (16–18), 76 (70–76) apart; setae *d* 90 (90–93) on ventral annulus 39 (39–42), 51 (51–53) apart; setae *e* 70 (65–70) on ventral annulus 67 (67–69), 29 (29–31) apart; setae *f* 50 (50–55) on ventral annulus 98 (98–101), 37 (36–37) apart. Setae *h*1 2 (2–3), *h*2 152 (150–152). **Female genitalia** 24 (24–25), 37 (37–39) wide, coverflap with 4 longitudinal ridges in 2 ranks, 1 near the base and 3 at distal margin, setae *3a* 12 (10–12), 20 (20–21) apart.

Male. Unknown.

Type material. **Holotype**, female (slide number NJAUEri777, marked Holotype), from *Lonicera elisae* Franch. (Caprifoliaceae), Mengda Natural Reserve, Xunhua County, Qinghai Province, P. R. China, 35° 47' 38" N, 102° 40' 40" E, elevation 2523m, 19 July 2007, coll. Xiao-Feng Xue. **Paratypes**, 12 females (slide number NJAUEri777), with the same data as holotype.

Relation to host. Vagrant on leaf lower surface. No damage to the host was observed.

Etymology. The specific designation *mengdaensis* is from the place name Mengda Natural Reserve, where this new species was collected; feminine in gender.

Differential diagnosis. This species is similar to *Diptacus lonicerae* Kuang, 2001, but can be differentiated from the latter by prodorsal shield with admedian lines and submedian lines separated (admedian lines and submedian lines connected in *D. lonicerae*), prodorsal shield frontal lobe wide and broad (frontal lobe small in *D. lonicerae*), female genital coverflap with 4 longitudinal ridges in 2 ranks, 1 near the base and 3 far from the base (female genital coverflap with 6–8 longitudinal ridges in *D. lonicerae*).

Subfamily Rhyncaphytoptinae Roivainen, 1953

Genus *Rhyncaphytoptus* Keifer, 1939

Rhyncaphytoptus ulmi Xin & Dong, 1981

http://species-id.net/wiki/Rhyncaphytoptus_ulmi

Rhyncaphytoptus ulmi Xin and Dong 1981: 216–217, figures 2–3.

Rhyncaphytoptus ulmi; Amrine and Stasny 1994: 277.

Rhyncaphytoptus ulmi; Amrine and Stasny 1996: 300.

Rhyncaphytoptus ulmi; Hong and Zhang 1996: 79, figures 188–1–188–2.

Rhyncaphytoptus ulmi; Kuang et al. 2005: 155–156, figure 158.

Rhyncaphytoptus ulmi; Xue et al. 2006b: 3.

Rhyncaphytoptus ulmi; Song et al. 2007b: 59.

Rhyncaphytoptus ulmi; Xue et al. 2009: 3.

Material examined. 16 females (slide number NJAUEri792B and NJAUEri795), from *Ulmus* sp. (Ulmaceae), Xining, Qinghai Province, P. R. China, 36°38'18"N, 101°45'27"E, elevation 2241m, 21 July 2007, coll. Xiao-Feng Xue.

Host. *Ulmus* sp. (Ulmaceae).

Relation to host. Vagrant on leaf lower surface. No damage to the host was observed.

Distribution. China (Jiangsu, Gansu, Jilin, Liaoning, Shaanxi, Shandong, Xinjiang, Qinghai).

***Rhyncaphytoptus spinus* sp. n.**

urn:lsid:zoobank.org:act:F3227702-A360-40D4-9367-E395D5F361DE

http://species-id.net/wiki/Rhyncaphytoptus_spinus

Figures 34–36

Description. Female. (n = 6) Body fusiform, light yellow, 270 (232–323), 82 (82–91) wide, 105 (104–105) thick. **Gnathosoma** 58 (56–59), projecting downwards, suboral plate present, pedipalp coxal seta (*ep*) 4 (4–5), dorsal pedipalp genual seta (*d*) 10 (9–11), palp tarsus ventral seta (*v*) 4 (3–4), cheliceral stylets 71 (71–72). **Prodorsal shield** 38 (38–40), 60 (57–64) wide, with broad frontal lobe, 7 (6–7); median, admedian and submedian lines present, median, admedian lines connected at basal 1/3 and 2/3 of prodorsal shield, forming three cells each both side. Scapular tubercles ahead of rear shield margin, 6 (5–6), 34 (34–40) apart, scapular setae (*sc*) 21 (21–25), projecting forward, knobbed at the end. **Coxigenital region** with 15 (14–15) annuli, with microtubercles. Coxisternal plates smooth, anterolateral setae on coxisternum **I** (*Ib*) 28 (26–28), 16 (16–18) apart, proximal setae on coxisternum **I** (*Ia*) 46 (44–46), 10 (10–13) apart, proximal setae on coxisternum **II** (*2a*) 80 (71–80), 30 (30–36) apart, tubercles *Ib* and *Ia* 6 (5–7) apart, tubercles *Ia* and *2a* 9 (9–13) apart. Prosternal apodeme combined, 6 (6–7). **Leg I** 44 (43–49), femur 16 (16–17), basiventral femoral seta (*bv*) 18 (15–20); genu 8 (8–9), antaxial genual seta (*l''*) 30 (30–32); tibia 12 (12–13), paraxial tibial seta (*l'*) 14 (14–17), located at 1/3 from dorsal base; tarsus 9 (9–10), seta *ft'* 26 (24–28), seta *ft''* 35 (34–35), seta *u'8* 6 (6–8); tarsal empodium (*em*) 12 (12–13), simple, 8-rayed, tarsal solenidion (ω) 11 (10–11), tapered. **Leg II** 41 (41–46), femur 14 (14–16), basiventral femoral seta (*bv*) 18 (18–19); genu 7 (7–8), antaxial genual seta (*l''*) 10 (10–11); tibia 11 (10–11); tarsus 8 (8–10), seta *ft'* 12 (12–13), seta *ft''* 34 (34–34), seta *u'7* 6 (6–7); tarsal empodium (*em*) 12 (12–13), simple, 8-rayed, tarsal solenidion (ω) 11 (10–11), tapered. **Opisthosoma** dorsally with 38 (38–42) annuli, with long spiny microtubercles, ventrally with 98 (97–98) annuli, with triangle microtubercles. Setae *c2* 33 (31–33) on ventral annulus 17 (17–19), 76 (71–76) apart; setae *d* 97 (95–97) on ventral annulus 38 (38–40), 53 (53–69) apart; setae *e* 41 (41–45) on ventral annulus 60 (59–61), 31 (31–39) apart; setae *f* 36 (36–38) on ventral annulus 92 (92–93), 28 (28–30) apart. Setae *h1* 5 (5–6), *h2* 110 (110–115). **Female genitalia** 21 (18–21), 32 (32–35) wide, coverflap smooth, setae *3a* 75 (70–75), 20 (20–23) apart.

Male. Unknown.

Type material. **Holotype**, female (slide number NJAUeri820, marked Holotype), from *Lonicera rupicola* Hook. f. et Thoms. (Caprifoliaceae), Beishan National Forest




Figure 34. *Rhyncaphytoptus spinus* sp. n.: **D** dorsal view of female **L1** leg I **L2** leg II **IG** female internal genitalia **em** empodium.




Figure 35. *Rhyncaphytoptus spinus* sp. n.: V ventral view of female CG coxae and female genitalia LO lateral microtubercles.




Figure 36. *Rhyncaphytoptus spinus* sp. n.: **A** dorsal view of female **B** ventral view of female **C** prodorsal shield **D** coxae and female genitalia **E** dorsal view of female posterior part **F** ventral view of female posterior part **G** empodium **H** leg I and leg II **I** female internal genitalia.

Park, Huzhu County, Qinghai Province, P. R. China, 36°53'35"N, 102°25'56"E, elevation 2610m, 22 July 2007, coll. Xiao-Feng Xue. **Paratypes**, 5 females (slide number NJAUEri820), with the same data as holotype.

Relation to host. Vagrant on leaf lower surface. No damage to the host was observed.

Etymology. The specific designation *spinus* is from the character of the dorsal opisthosomal microtubercles, spiny, “spina, spinus” in Latin; masculine in gender.

Differential diagnosis. This species is similar to *Rhyncaphytoptus guanegounis* Song, Xue & Hong, 2007b, but can be differentiated from the latter by median, ad-median and submedian lines present on prodorsal shield (prodorsal shield smooth in *R. guanegounis*), prodorsal shield with wide and broad frontal lobe (prodorsal shield with long and broad frontal lobe in *R. guanegounis*), opisthosomal dorsal annuli with spiny microtubercles (opisthosomal dorsal annuli smooth in *R. guanegounis*), tarsal empodium (*em*) 8-rayed (6-rayed in *R. guanegounis*).

Discussion

Although Qinghai has climate, vegetation and biological diversity similar to Tibet, we did not find any eriophyoid mite species already reported in Tibet (Song et al. 2011). On the contrary, 6 of 23 species were reported from Gansu Province or Shaanxi Province, east neighboring provinces. As of 2010, 932 eriophyoid mite species have been described from China (Hong et al. 2010), and the number is still increasing. Qinghai has about 8% of the land of China, but only about 2% eriophyoid mites were reported to date. Furthermore, only two investigations have been conducted. More systematic collections of eriophyoid mites from Qinghai Province are needed in the near future.

Acknowledgements

This research was funded by the National Natural Science Foundation of China (31172132). We thank Xiao Han of NJAU for early review of the manuscript. We also thank Professor Xin-Hua Li of the Dept of Botany at NJAU for identifying host plants. We are very grateful to Professor James W. Amrine Jr. for his valuable suggestions on the manuscript.

References

Amrine JW Jr, Stasny TA (1994) Catalog of the Eriophyoidea (Acarina: Prostigmata) of the world. Indira Publishing House, West Bloomfield, Michigan, U.S.A., 804 pp.

- Amrine JW Jr, Manson DCM (1996) Preparation, mounting and descriptive study of eriophyoid mites. In: Lindquist EE, Sabelis MW, Bruin J (Eds) Eriophyoid Mites: Their Biology, Natural Enemies and Control. Elsevier, World Crop Pests, 6, 383–396.
- Amrine JW Jr, Stasny TA (1996) Corrections to the catalog of the Eriophyoidea (Acarina: Prostigmata) of the world. *International Journal of Acarology* 22(4): 295–304. doi: 10.1080/01647959608684108
- Amrine JW Jr, Stasny TA, Flechtmann CHW (2003) Revised keys to world genera of Eriophyoidea (Acari: Prostigmata). Indira Publishing House, West Bloomfield, MI, USA., 244 pp.
- de Lillo E, Craemer C, Amrine JW Jr, Nuzzaci E G (2010) Recommended procedures and techniques for morphological studies of Eriophyoidea (Acari: Prostigmata). *Experimental and Applied Acarology* 51: 283–307. doi: 10.1007/s10493-009-9311-x
- Hong XY, Zhang ZQ (1996) The eriophyoid mites of China: an illustrated catalog and identification keys (Acari: Prostigmata: Eriophyoidea). Associated Publishers, Florida, U.S.A., 318 pp.
- Hong XY, Xue XF (2005) Four new species of *Aculops* Keifer (Acari: Eriophyoidea: Eriophyidae) from China. *Oriental Insects* 39: 203–211. doi: 10.1080/00305316.2005.10417433
- Hong XY, Xue XF, Song ZW (2010) Eriophyoidea of China: a review of progress, with a checklist. *Zoosymposia* 4: 57–93.
- Huang KW (2001) The eriophyid mites of Taiwan: description of eighty-six species from the Tengchih Area. *Bulletin of National Museum of Natural Science* 14: 1–84.
- Keifer HH (1939) Eriophyid Studies VII. *Bulletin of the California Department of Agriculture*, 28: 484–505.
- Kuang HY (1995) Economic insect fauna of China. Fasc. 44. (Acari: Eriophyoidea) (I). Science Press, Beijing, China, 198pp+pls7.
- Kuang HY, Pang HL (1997) Four new species of the Eriophyinae (Acari: Eriophyidae) from China. *Entomotaxonomia* 19(3): 230–234.
- Kuang HY, Gong GJ, Luo GH (1998) A new genus and five new species of Phyllocoptinae from China (Acari: Eriophyoidea). *Acta Entomologica Sinica* 41(2): 197–203.
- Kuang HY (1988) Two new species of the genus *Aceria* from China (Acariformes: Eriophyidae). *Acta Zootaxonomica Sinica* 13(1): 49–51.
- Kuang HY (2000) Four new species of Phyllocoptinae from China (Acari: Eriophyoidea). *Acta Zootaxonomica Sinica* 25(4): 391–394.
- Kuang HY (2001) Two new species of *Diptacus* from China. *Entomotaxonomia* 23(2): 153–156.
- Kuang HY, Luo GH, Wang AW (2005) Fauna of Eriophyid mites from China (II) (Acari: Eriophyoidea). China Forestry Publishing House, Beijing, China, 176 pp.
- Lin FB, Kuang HY (2001) Four new species of Phyllocoptinae from China (Acari: Eriophyoidea). *Acta Entomologica Sinica* 44(3): 350–355.
- Lindquist EE (1996) External anatomy and notation of structures In: Lindquist EE, Sabelis MW, Bruin J (Eds) Eriophyoid mites: their biology, natural enemies and control. Elsevier, World Crop Pests 6: 3–31.
- Qinghai Provincial Government Website: <http://www.qh.gov.cn/> [accessed 8 March 2012]
- Skoracka A (2003) New species of *Aculodes* (Acari: Eriophyoidea) from grasses in Poland. *Acta Zoologica Academiae Scientiarum Hungaricae* 49(1): 43–60.

- Song ZW, Xue XF, Hong XY (2006) A review of Chinese *Phyllocoptes* Nalepa (Acari: Eriophyoidea: Eriophyidae: Phyllocoptinae) with descriptions of five new species. *Zootaxa* 1167: 31–45.
- Song ZW, Xue XF, Hong XY (2007a) Description of seven new species in the genus *Diptacus* Keifer (Acari: Eriophyoidea: Diptilomiopidae: Diptilomiopinae) from northwestern China. *Zootaxa* 1429: 39–53.
- Song ZW, Xue XF, Hong XY (2007b) Four new species and a new synonym in the genus *Rhyncaphytoptus* Keifer (Acari: Eriophyoidea: Diptilomiopidae) from China. *Zootaxa* 1520: 57–68.
- Song ZW, Xue XF, Hong XY (2008) Eriophyoid mite fauna (Acari: Eriophyoidea) of Gansu Province, northwestern China with descriptions of twelve new species. *Zootaxa* 1756: 1–48.
- Song ZW, Xue XF, Hong XY (2009a) Seven new species of Phyllocoptini from Tibet Autonomous Region, China (Acari: Eriophyidae: Phyllocoptinae). *Zootaxa* 2291: 1–23.
- Song ZW, Xue XF, Hong XY (2009b) Six new species of *Aculus* Keifer from Tibet Autonomous Region, China (Acari: Eriophyoidea: Eriophyidae: Phyllocoptinae). *Zootaxa* 2305: 1–23.
- Song ZW, Xue XF, Hong XY (2009c) Seven new species of *Phyllocoptes* Nalepa from Tibet Autonomous Region, China (Acari: Eriophyoidea: Phyllocoptinae). *Zootaxa* 2313: 35–60.
- Song ZW, Xue XF, Hong XY (2011) Eriophyoid mite fauna (Acari: Eriophyoidea) of the Tibet Autonomous Region, southwestern China with descriptions of six new species. *International Journal of Acarology* 37(5): 367–390. doi: 10.1080/01647954.2010.519723
- Xin JL, Dong HQ (1981) Two new species of the genus *Rhyncaphytoptus* from China. *Fudan Journal (National Science)* 20(2): 216–219.
- Xue XF, Song ZW, Hong XY (2006a) A taxonomic study of the genus *Tetra* Keifer (Acari: Eriophyidae: Phyllocoptinae: Anthocoptini) from Shaanxi Province, China with descriptions of nine new species. *Zootaxa* 1249: 1–22.
- Xue XF, Song ZW, Hong XY (2006b) Six new species of Rhyncaphytoptinae from northwestern China (Acari: Eriophyoidea: Diptilomiopidae). *Zootaxa* 1225: 1–19.
- Xue XF, Hong XY (2006) Eriophyoid mite fauna from Henan Province, central China (Acari: Eriophyoidea) with descriptions of five new species. *Zootaxa* 1204: 1–30.
- Xue XF, Song ZW, Amrine JW Jr, Hong XY (2007) Eriophyoid mites on coniferous plants in China with descriptions of a new genus and five new species (Acari: Eriophyoidea). *International Journal of Acarology* 33(4): 342–355. doi: 10.1080/01647950708683695
- Xue XF, Song ZW, Hong XY (2008) Eight new species of the genus *Aculus* Keifer (Acari: Eriophyidae) from China. *Zootaxa* 1721: 35–52.
- Xue XF, Song ZW, Hong XY (2009) Five new species of *Phyllocoptes* from China (Acari: Eriophyidae). *International Journal of Acarology* 35(2): 133–148. doi: 10.1080/01647950902934234
- Xue XF, Song ZW, Hong XY (2010) Review of *Phyllocoptruta*, with descriptions of two new species (Acari: Eriophyoidea). *Annals of the Entomological Society of America* 103(5): 697–705. doi: 10.1603/AN10019