



The Impact of COVID-19 on Surgical Training: the Past, the Present and the Future

Marina Yiasemidou^{1,2}

Received: 15 May 2021 / Accepted: 19 May 2021 / Published online: 12 June 2021
© The Author(s) 2021

Abstract

The COVID-19 pandemic and infection control measures had an unavoidable impact on surgical services. During the first wave of the pandemic, elective surgery, endoscopy, and ‘face-to-face’ clinics were discontinued after recommendations from professional bodies. In addition, training courses, examinations, conferences, and training rotations were postponed or cancelled. Inadvertently, infection control and prevention measures, both within and outside hospitals, have caused a significant negative impact on training. At the same time, they have given space to new technologies, like telemedicine and platforms for webinars, to blossom. While the recovery phase is well underway in some parts of the world, most surgical services are not operating at full capacity. Unfortunately, some countries are still battling a second or third wave of the pandemic with severely negative consequences on surgical services. Several studies have looked into the impact of COVID-19 on surgical training. Here, an objective overview of studies from different parts of the world is presented. Also, evidence-based solutions are suggested for future surgical training interventions.

Keywords COVID-19 · Training · Simulation · Virtual reality

Within 6 months of the declaration of a coronavirus SARS-CoV-2 pandemic, nearly six million cases have been confirmed worldwide and around 400,000 deaths were recorded [1]. Countries responded with unprecedented infection control and prevention measures, in an effort to contain the spread of the virus [2]. These included restrictions in unnecessary movement, widespread use of protective equipment, and sealing off national or regional borders [3].

The pandemic and infection control measures had an unavoidable impact on surgical services [4]. During the first wave of the pandemic, elective surgery, endoscopy, and ‘face-to-face’ clinics were discontinued after recommendations from professional bodies [5–10]. In addition, training courses [11, 12], examinations, conferences, and training rotations were postponed or cancelled [13]. Due to the reports of worryingly high postoperative complications, during the early days of the pandemic, several emergency

pathologies were treated conservatively (e.g. appendicitis and diverticulitis) [14–17]. A further reduction of training opportunities was caused by the attempts to minimise staff members in operating theatres [18]. Many hospitals applied a ‘consultant-only’ operating policy, in order to reduce operating times and hence staff exposure [18].

Inadvertently, infection control and prevention measures, both within and outside hospitals, have caused a significant negative impact on training [19]. At the same time, they have given space to new technologies, like telemedicine and platforms for webinars, to blossom [19, 20]. Several studies have looked into the impact of COVID-19 on surgical training [19, 21–29]. Here, the author will attempt a comprehensive summary of their findings.

An Overview

A great number of studies have reported a detrimental effect on surgical training. The majority of them described it ‘severe’ or ‘significant’ and some even ‘catastrophic’ [19, 21, 22, 28, 30–43]. These reports come both from developed [13, 21, 22, 24, 26, 31–34, 44–52] and developing countries [28, 30, 35, 53–55] with very different healthcare systems.

✉ Marina Yiasemidou
marinayiasemidou@gmail.com

¹ NIHR Academic Clinical Lecturer in General Surgery, Hull York Medical School, Hull, UK

² ST7 Colorectal Surgery, Bradford Teaching Hospitals, Bradford, UK

Moreover, the results were uniform between a variety of surgical specialties; general surgery [13, 19, 21, 30, 32, 34, 37, 41, 42, 44, 47, 56, 57], orthopaedics [19, 24, 58], ENT (ear, nose, throat) surgery [19, 31, 50], vascular surgery [19, 22, 26], urology [22, 26, 53], oral and maxillofacial surgery [19, 33], and neurosurgery [19, 54, 59].

Most of the evidence provided are extracted from surveys and editorials [19, 21–29]. While most surveys have a decent response rate and a great number of responders [19, 21–29], surveys are considered to provide low-level evidence [60]. There are several reasons for this. Establishing population size estimates are important for assessing the generalizability of the results of any research attempt [60]. However, there is no ideal method for population size estimation. Different methods are subject to different biases; even employing multiple approaches in an attempt to minimise these resulted in a wide variance of estimates [61]. As such, several assumptions have to be made in order to establish the ideal response rate, inevitably introducing biases [62].

In surveys, the principle of random recruitment may be violated, if authors select responders with specific characteristics from their networks [62]. Surveys are also liable to recall bias and sampling bias [62]. For instance, in the current scenario of assessing the impact of COVID-19 on surgical training, responders may misremember the number of operations performed. Moreover, in order to assess the impact of COVID-19, responders may have compared their current situation to 1 year ago, which enhances the probability of recall bias.

A degree of sampling data is unavoidable. For example, surgeons that have no access to an internet network lack access to surveys that are distributed electronically [19, 55]. Specifically, for international surveys that are distributed in the English language, sampling bias is introduced as the non-English speakers will be unable to complete the survey.

Despite the potential introduction of bias, the ‘unanimous’ results from all the surveys strengthen the findings [19, 21–29]. Also, in agreement is the demonstrated reduction of performed cases on the electronic logbook of trainees [63], which provides an ‘independent’ confirmation of the effect demonstrated in the surveys.

Areas of Training Affected

Theatre Exposure

Operating theatre exposure was significantly reduced [13, 19, 30, 31, 44, 53], with some authors reporting a reduction of case numbers as great as 50% [41]. Several factors contributed to this. Elective surgery came to a standstill during the first wave of the pandemic [64–66]. Emergency admissions and emergency surgery were also reduced [67].

The reduction of theatre staff numbers and ‘consultant-only operating’ policies have further reduced trainee exposure to surgery [18].

Laparoscopic surgery was identified as an aerosol-generating procedure and was hence considered high risk [68, 69]. This caused anxiety amongst theatre staff and surgeons and prompted guidelines for avoidance of laparoscopic surgery [68, 69]. While there were some exceptions [70, 71], a large number of hospitals discontinued or refrained from minimally invasive surgery [69]. As such, trainees did not get the opportunity to enhance their minimal invasive surgical skills.

Immediately after the first wave of the pandemic, the surgical community engaged in a recovery phase for surgery [72–74]. This entailed the formation of COVID-19-free pathways (‘green’ pathways) for elective patients and vigorous testing of patients and staff [74]. These processes, although necessary, are time consuming and often lead to surgical departments operating at less than 100% capacity [75]. Even to this day, delays are noticed in patient care due to testing and difficulty in self-isolating prior to elective surgery [75]. All these contribute to reduced theatre cases and hence less training opportunities for surgical learners.

The rapid development of vaccines provided a slither of hope [72] for an accelerated pathway to theatre; however, there are large discrepancies in their distribution globally [76]. Also, there are concerns that they do not prevent asymptomatic transmission [77]. Therefore, vaccination can only be used as an adjunct to self-isolation rather than a replacement.

In addition to delays outside the theatre complex, several authors report reduced workflow within it as well [78]. Several strategies were employed to increase theatre workflow during the pandemic and recovery phase [78]. Whether these are effective remains to be seen [78].

Undoubtedly, the extra steps to get the patients to theatre reduced workflow and the establishment of COVID-19-free surgical pathways [74] and have reduced the number of cases performed on an operating list. As a result, training opportunities are reduced, not only during the pandemic but the recovery phase as well.

Outpatient Clinics

During the COVID-19 pandemic, outpatient clinics were either cancelled or converted from ‘face-to-face’ to virtual consultations [21, 22, 30, 34, 46, 55, 58, 66, 79–82]. This, initially, decreased the exposure of trainees to supervised outpatient clinics, further limiting their training opportunities [19]. On the other hand, once the initial transition period had passed, many found several advantages in virtual clinics [58, 81], accrediting them for the continuation of outpatient clinic exposure for trainees [58]. Some authors suggest that

with some modifications in existing curricula, alternative interventions such as virtual clinics can form part of the routine training experience [83].

The notion of virtual outpatient clinics is at least 10 years old [84]. Their implementation saw a reduction in patients seen in ‘face-to-face’ clinic [85], improving patient experience [86]. Patient satisfaction was also high, reported in some cases as 97% [87]. A recent systematic review and meta-analysis by Chaudhry et al. [86] included 12 studies, 8 of which were randomised controlled trials comparing surgeon and patient satisfaction with ‘face-to-face’ and virtual clinics. They found no statistical difference in surgeon satisfaction (pooled OR 0.38 [95% CI 0.07 to 2.19]; $p = 0.28$) or in patient-reported clinical outcomes [86]. Patients reported time savings, both in respect to travel time (17 min shorter [95% CI 2 to 32]; $p = 0.03$) and other waiting times (180 min shorter [95% CI 78 to 281]; $p < 0.001$) [86]. Moreover, a randomised controlled trial by Llorens et al. [88] comparing an in-person and telemedicine clinic demonstrated significant cost savings with the virtual intervention.

Besides the expert surgeons who report at least equal satisfaction with virtual as with in-person clinics, trainees would like to see telemedicine and virtual clinics remaining as part of their practice even after the pandemic [19]. Phone or other virtual consultations can be as easily supervised as ‘face-to-face’ consultations, therefore not impeding the potential of a learning experience.

Conferences, Training Courses, and Teaching Sessions

As a result of infection control measures aiming to reduce large gatherings, conferences, training sessions, and teaching were postponed or took place on a virtual platform [19, 56]. In a global survey conducted by our team, the trainees express a relative dissatisfaction with the virtual platforms [19]. They cited technical challenges (lack of hardware and access to a high-quality network), lack of engagement and/or interaction and inappropriate timing, as the reasons for their dissatisfaction [19]. Despite that, in the same survey, trainees did recognise the potential of these educational processes and state that they would like to see virtual conferences and courses remain as part of training after the pandemic [19]. The survey was conducted from the 23 April to the 15 May 2020, which was rather early on in the pandemic. At that time, the surgical world had to quickly adjust to the new teaching methods and perhaps was not as adequately prepared to provide high-quality virtual resources.

Surveys conducted subsequently showed extremely good trainee satisfaction with virtual conferences and teaching sessions [89–95]. Specifically, Ottesen et al. [89]

reported that the virtual platform exceeded expectations of 85.7% of attendees and 100% would participate in future virtual events. There were also reports of virtual events believed to be superior to traditional conferences [90, 93].

One notable exception is the 2021 paper by Woodruff et al. [96]. While they accept that the results of phase 2/3 clinical trials are adequately reported in virtual conferences, they report fewer overall submissions [96]. They are particularly concerned that this may lead to fewer presentations of observational and post hoc analyses of clinical trials, often presented by residents, fellows, and trainees [96]. Conference presentations are essential for career progression and form an integral part of job applications [96]. The authors of this study are worried that virtual conferences hinder presentation and public speaking skills for trainees [96]. They also point out the missed opportunity for ad hoc spontaneous networking which often result in collaboration and mentorship [96]. These are all valid points which need to be addressed. The authors see hybrid conferences as a potential solution for the future [96].

Endoscopy

Endoscopy sessions were also discontinued during the first wave of the pandemic [19, 23, 25]. This was due to concerns about viral contamination between both patients and providers of endoscopy [97]. Studies have shown that there is indeed a substantial risk of exposure and infection with respiratory diseases that can be spread via an airborne route [98]. Endoscopists are often exposed to infectious biologic samples during procedures [99]. This is particularly true due to the short physical distance between patient and endoscopist during procedures. This distance is shorter than 6 feet; the distance that SARS droplets from infectious patients can reach [100].

While endoscopy sessions were reinstated during the recovery phase of the pandemic, the numbers of procedures are reduced, again causing a negative impact on training [25, 27, 101]. Pawlak et al. [27] conducted an international survey assessing the impact of COVID-19 on endoscopy training. 93.8% out of 770 respondents reported a reduction in endoscopy case volume, with a median percentage reduction of 99% (interquartile range, 85–100%) [27]. The reduction was greatest for colonoscopy procedures [27]. The restrictions concerned not only case volume but also trainee activity (i.e. procedures were performed by experts only) [27]. A survey conducted amongst UK trainees showed similar paucity in endoscopy training [101]. The reasons cited for this were changes to institutional policy that excluded trainees from procedures (75.8%), low case volume (56.8%), and redeployment to another clinical area (47.7%) [101].

Recovery of Training

Our group have worked on a framework for training recovery based on the results of a global survey that we conducted [19]. It emerged that trainees had concerns about the lack of guidance from training stakeholders and would like to see their mentors and trainers prioritising training at every opportunity possible [19]. Based on this and the opinion of experts, we proposed a four-stage recovery plan. This consists of:

Guidance from national/international training stakeholders.

Involvement of trainees, trainers, regional training programme representatives, the hospital managerial team, and the digital support team in order to discuss local implementation of guidelines and necessary adjustments that may be required locally.

Formation of implementation team who will carry out the plan set up by above teams.

Auditing and adjusting the plan by engaging in a ‘trial and error’ process [19].

Alternative Methods of Training

The severe reduction of case volume is apparent in studies evaluating the impact of COVID-19 on surgical training [4, 19, 29, 39, 63, 82, 83, 101]. Therefore, methods outside the operating theatre must be sought as an adjunct to conventional training, to enhance surgical skills. Surgical simulation has been utilised for years in the surgical and other fields and was shown to be effective in enhancing surgical skills, particularly for novices [102–106]. In addition to simulation, methods like mental practice and ‘warm up’ before surgery may enhance the learning experience in the operating theatre [107–110]. Concerns about poor fidelity have now degree been resolved due to modern additive technologies such as 3D reconstruction from CT or MR images or 3D printing [107–110]. Moreover, new embalming methods made cadaveric simulation more accessible, by reducing the storage requirements and making cadavers ‘reusable’ [105, 111].

Immersive technologies are also useful for training during the pandemic [112]. These refer to virtual reality (VR), augmented reality (AR), and mixed reality (MR) [112]. Perhaps their biggest advantage is that they provide Omni-Learning; the ability to learn anywhere, anytime, with anyone [113]. AR uses holograms projected into the real-world environment [112]. This could include three-dimensional (3D) object transmission which can be viewed by a remote headset user [114]. The author in no way is suggesting that these can replace operating theatre experience, but can

exponentially increase the didactic impact of every theatre session. Technologies such as these can allow for real-time streaming of operations during which the trainee can have the same optic output as their trainer (see what they see) [112], something which is of great importance in identifying efficiently and promptly the appropriate planes of dissection. Knowledge that can be put in good use the next time they are in theatre.

Conclusion

There is little doubt that COVID-19 has significantly decreased training opportunities for surgeons [4, 19, 29, 39, 63, 82, 83, 101]. This was partially counteracted by the introduction of alternative teaching methods such as virtual teaching platforms [19, 43, 95, 96]. However, there is a long way to go to ensure that surgical training is not heavily impacted long term. This effort needs to be coordinated by training authorities nationally, with the involvement of trainees in decision-making. Alternative teaching methods should be used, not to replace, but to enhance the scarce training opportunities in existence.

Declarations

Competing Interests The author declares no competing interests.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article’s Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article’s Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

1. WHO (2020) Coronavirus disease (COVID-19) outbreak. <https://www.who.int/westernpacific/emergencies/covid-19>. Accessed 20 July 2020
2. Xiao Y, Torok ME (2020) Taking the right measures to control COVID-19. *Lancet Infect Dis* 20(5):523–524. [https://doi.org/10.1016/S1473-3099\(20\)30152-3](https://doi.org/10.1016/S1473-3099(20)30152-3)
3. Ruyi L (2020) Ministry of Communications: it is strictly forbidden to seal highways, highways, and quarantine stations. *Beijing Daily Client*. <https://baijiahao.baidu.com/s?id=1658210115200722778&wfr=spider&for=pc>. Accessed 17 Feb 2020

4. Teoh JY, Ong WLK, Gonzalez-Padilla D, Castellani D, Dubin JM, Esperto F, Campi R, Gudar K, Talwar R, Okhunov Z, Ng CF, Jain N, Gauhar V, Wong MC, Wroclawski ML, Tanidir Y, Rivas JG, Tiong HY, Loeb S, UroSoMe Working G (2020) A global survey on the impact of COVID-19 on urological services. *Eur Urol* 78(2):265–275. <https://doi.org/10.1016/j.eururo.2020.05.025>
5. Ali MJ, Hegde R, Nair AG, Bajaj MS, Betharia SM, Bhattacharjee K, Chhabra AK, Das JK, Dudeja G, Grover AK, Honavar SG, Kim U, Mahesh L, Mukherjee B, Sethi A, Sharma M, Singh U (2020) All India Ophthalmological Society - Oculoplastics Association of India consensus statement on preferred practices in oculoplasty and lacrimal surgery during the COVID-19 pandemic. *Indian J Ophthalmol* 68(6):974–980. https://doi.org/10.4103/ijoo.IJO_1415_20
6. Brat GA, Hersey S, Chhabra K, Gupta A, Scott J (2020) Protecting surgical teams during the COVID-19 outbreak: a narrative review and clinical considerations. *Ann Surg*. <https://doi.org/10.1097/SLA.0000000000003926>
7. Pignatti M, Pinto V, Miralles MEL, Giorgini FA, Cannamela G, Cipriani R (2020) How the COVID-19 pandemic changed the plastic surgery activity in a regional referral center in Northern Italy. *J Plast Reconstr Aesthet Surg* 73(7):1348–1356. <https://doi.org/10.1016/j.bjps.2020.05.002>
8. Poulouse BK, Phieffer LS, Mayerson J, Like D, Forrest LA, Rahmanian A, Bellamy B, Guertin M, Pawlik TM (2021) Responsible return to essential and non-essential surgery during the COVID-19 pandemic. *J Gastrointest Surg* 25(5):1105–1107. <https://doi.org/10.1007/s11605-020-04673-9>
9. Thaler M, Khosravi I, Hirschmann MT, Kort NP, Zagra L, Epinette JA, Liebensteiner MC (2020) Disruption of joint arthroplasty services in Europe during the COVID-19 pandemic: an online survey within the European Hip Society (EHS) and the European Knee Associates (EKA). *Knee Surg Sports Traumatol Arthrosc* 28(6):1712–1719. <https://doi.org/10.1007/s00167-020-06033-1>
10. Zheng MH, Boni L, Fingerhut A (2020) Minimally invasive surgery and the novel coronavirus outbreak: lessons learned in China and Italy. *Ann Surg* 272(1):e5–e6. <https://doi.org/10.1097/SLA.0000000000003924>
11. Elangovan S, Mahrous A, Marchini L (2020) Disruptions during a pandemic: gaps identified and lessons learned. *J Dent Educ* 84(11):1270–1274. <https://doi.org/10.1002/jdd.12236>
12. Hevia V, Lorca J, Hevia M, Dominguez A, Lopez-Plaza J, Artiles A, Alvarez S, Sanchez A, Fraile A, Lopez-Fando L, Sanz E, Ruiz M, Alcaraz E, Burgos FJ (2020) COVID-19 pandemic: impact and rapid reaction of urology. *Actas Urol Esp* 44(7):450–457. <https://doi.org/10.1016/j.acuro.2020.04.006>
13. Al-Jabir A, Kerwan A, Nicola M, Alsafi Z, Khan M, Sohrabi C, O'Neill N, Iosifidis C, Griffin M, Mathew G, Agha R (2020) Impact of the coronavirus (COVID-19) pandemic on surgical practice - part 1. *Int J Surg* 79:168–179. <https://doi.org/10.1016/j.ijso.2020.05.022>
14. Brothers L, Shaw GL, Wright EL (1993) Durations of extended mental rehearsals are remarkably reproducible in higher level human performances. *Neurol Res* 15(6):413–416. <https://doi.org/10.1080/01616412.1993.11740175>
15. Chiofalo B, Baiocco E, Mancini E, Vocaturo G, Cuttillo G, Vincenzoni C, Bruni S, Bruno V, Mancari R, Vizza E (2020) Practical recommendations for gynecologic surgery during the COVID-19 pandemic. *Int J Gynaecol Obstet* 150(2):146–150. <https://doi.org/10.1002/ijgo.13248>
16. Iyengar K, Vaish A, Vaishya R (2020) Revisiting conservative orthopaedic management of fractures during COVID-19 pandemic. *J Clin Orthop Trauma* 11(4):718–720. <https://doi.org/10.1016/j.jcot.2020.05.010>
17. Suwanwongse K, Shabarek N (2020) Successful conservative management of acute appendicitis in a coronavirus disease 2019 (COVID-19) patient. *Cureus* 12(4):e7834. <https://doi.org/10.7759/cureus.7834>
18. Royal College of Surgeons of England. COVID-19: good practice for surgeons and surgical teams. <https://www.rcseng.ac.uk/standards-and-research/standards-and-guidance/good-practice-guides/coronavirus/covid-19-good-practice-for-surgeons-and-surgical-teams/> Accessed 25th May 2020
19. Research Education Innovation in Surgery (REINS) initiative COVID-19 group, Yiasemidou M, Tomlinson J, Chetter I, Shenkar BC (2021) Impact of the SARS-CoV-2 (COVID-19) crisis on surgical training: global survey and a proposed framework for recovery. *BJS Open* 5(2):zraa051. <https://doi.org/10.1093/bjsopen/zraa051>
20. Laloo R, Giorga A, Williams A, Biyani CS, Yiasemidou M (2020) Virtual surgical education for core surgical trainees in the Yorkshire deanery during the COVID-19 pandemic. *Scott Med J* 65(4):138–143. <https://doi.org/10.1177/0036933020951927>
21. Hope C, Reilly JJ, Griffiths G, Lund J, Humes D (2021) The impact of COVID-19 on surgical training: a systematic review. *Tech Coloproctol* 25(5):505–520. <https://doi.org/10.1007/s10151-020-02404-5>
22. Johnson AP, Wohlaer MV, Mouawad NJ, Malgor RD, Coogan SM, Sheahan MG 3rd, Singh N, Cuff RF, Woo K, Coleman DM, Shalhub S (2021) The impact of the COVID-19 pandemic on vascular surgery trainees in the United States. *Ann Vasc Surg* 72:182–190. <https://doi.org/10.1016/j.avsg.2020.09.045>
23. Kumar S, Prenner S, Kochman ML (2020) The impact of COVID-19 on endoscopic training. *Am J Gastroenterol* 115(7):1142–1143. <https://doi.org/10.14309/ajg.0000000000000716>
24. Megaloikonomos PD, Thaler M, Igoumenou VG, Bonanzinga T, Ostojic M, Couto AF, Diallo J, Khosravi I (2020) Impact of the COVID-19 pandemic on orthopaedic and trauma surgery training in Europe. *Int Orthop* 44(9):1611–1619. <https://doi.org/10.1007/s00264-020-04742-3>
25. Mejia Perez LK, Sharma N (2020) Endoscopy training during COVID-19. *Gastrointest Endosc* 92(4):988. <https://doi.org/10.1016/j.gie.2020.06.006>
26. Mousa AY, Broce M (2020) The impact of COVID-19 on vascular training. *J Vasc Surg* 72(1):380–381. <https://doi.org/10.1016/j.jvs.2020.04.469>
27. Pawlak KM, Kral J, Khan R, Amin S, Bilal M, Lui RN, Sandhu DS, Hashim A, Bollipo S, Charabaty A, de-Madaria E, Rodriguez-Parra AF, Sanchez-Luna SA, Zorniak M, Walsh CM, Grover SC, Siau K (2020) Impact of COVID-19 on endoscopy trainees: an international survey. *Gastrointest Endosc* 92(4):925–935. <https://doi.org/10.1016/j.gie.2020.06.010>
28. Prezotti JA, Henriques JVT, Favorito LA, Canalini AF, Machado MG, Brandao TBV, Barbosa AMV, Moromizato JKM, Anzolic KMJ, Fernandes RC, Rodrigues FRA, Bellucci CHS, Silva CS, Pompeo ACL, de Bessa Jr J, Gomes CM (2021) Impact of COVID-19 on education, health and lifestyle behaviour of Brazilian urology residents. *Int Braz J Urol* 47(4):753–776. <https://doi.org/10.1590/S1677-5538.IBJU.2021.99.09>
29. Segui-Moya E, Gonzalez-Padilla DA, Ortega-Polledo LE, Sanchez-Garcia M, Lopez-Gonzalez JA, Anton-Juanilla M, Mercader-Barrull C, Carrion DM, Gomez Rivas J, Alvarez-Ossorio JL (2020) Impact of COVID-19 in Spanish urology residents: recommendations and perspective. *Arch Esp Urol* 73(5):471–478
30. Adesunkanmi AO, Ubom AE, Olasehinde O, Wuraola FO, Ijarotimi OA, Okon NE, Ikimalo JI, Fasubaa OB, Adesunkanmi ARK (2021) Impact of the COVID-19 pandemic on surgical residency training: perspective from a low-middle income

- country. *World J Surg* 45(1):10–17. <https://doi.org/10.1007/s00268-020-05826-2>
31. Alvarez ML, Waissbluth S, Gonzalez C, Napolitano C, Torrente M, Delano PH, Alarcon R, Fernandez F, Bitran R (2021) How the COVID-19 pandemic affects specialty training: an analysis of a nationwide survey among otolaryngology residents in Chile. *Medwave* 21(1):e8098. <https://doi.org/10.5867/medwave.2021.01.8097>
 32. Aziz H, James T, Remulla D, Sher L, Genyk Y, Sullivan ME, Sheikh MR (2021) Effect of COVID-19 on surgical training across the United States: a national survey of general surgery residents. *J Surg Educ* 78(2):431–439. <https://doi.org/10.1016/j.jsurg.2020.07.037>
 33. Brar B, Bayoumy M, Salama A, Henry A, Chigurupati R (2021) A survey assessing the early effects of COVID-19 pandemic on oral and maxillofacial surgery training programs. *Oral Surg Oral Med Oral Pathol Oral Radiol* 131(1):27–42. <https://doi.org/10.1016/j.oooo.2020.08.012>
 34. Group C-SCS (2021) COVID-19 impact on surgical training and recovery planning (COVID-STAR) - a cross-sectional observational study. *Int J Surg* 88:105903. <https://doi.org/10.1016/j.ijsu.2021.105903>
 35. Meybodi KT, Habibi Z, Nejat F (2021) The effects of COVID-19 pandemic on pediatric neurosurgery practice and training in a developing country. *Childs Nerv Syst* 37(4):1313–1317. <https://doi.org/10.1007/s00381-020-04953-4>
 36. Nicholas C, Hatchell A, Webb C, Temple-Oberle C (2021) COVID-19 and the impact on surgical fellows: uniquely vulnerable learners. *J Surg Educ* 78(2):375–378. <https://doi.org/10.1016/j.jsurg.2020.08.017>
 37. Nnamani Silva ON, Hernandez S, Kim EH, Kim AS, Gosnell J, Roman SA, Lin MYC (2021) Surgery clerkship curriculum changes at an academic institution during the COVID-19 pandemic. *J Surg Educ* 78(1):327–331. <https://doi.org/10.1016/j.jsurg.2020.07.009>
 38. Reed AJM, Chan JKK (2021) Plastic surgery training during COVID-19: challenges and novel learning opportunities. *J Plast Reconstr Aesthet Surg* 74(2):407–447. <https://doi.org/10.1016/j.bjps.2020.08.076>
 39. Saqib SU, Saleem O, Riaz A, Riaz Q, Zafar H (2021) Impact of a global pandemic on surgical education and training—review, response, and reflection. *J Pak Med Assoc* 71(Suppl 1 (1)):S49–S55
 40. Tahir H, Beg MA, Siddiqui F (2021) The reduction in clinical and surgical exposure of trainees during COVID-19 and its impact on their training. *J Pak Med Assoc* 71(Suppl 1 (1)):S18–S22
 41. Vailas M, Sotiropoulou M, Mulita F, Drakos N, Ambalov E, Maroulis I (2021) The impact of COVID-19 on surgical training at a tertiary hospital in Greece: a 'hidden infectious enemy' for junior surgeons? *Eur Surg J* 1–6. <https://doi.org/10.1007/s10353-021-00699-8>
 42. Whelehan DF, Connelly TM, Ridgway PF (2021) COVID-19 and surgery: a thematic analysis of unintended consequences on performance, practice and surgical training. *Surgeon* 19(1):e20–e27. <https://doi.org/10.1016/j.surge.2020.07.006>
 43. White EM, Shaughnessy MP, Esposito AC, Slade MD, Korah M, Yoo PS (2021) Surgical education in the time of COVID: understanding the early response of surgical training programs to the novel coronavirus pandemic. *J Surg Educ* 78(2):412–421. <https://doi.org/10.1016/j.jsurg.2020.07.036>
 44. Al-Jabir A, Kerwan A, Nicola M, Alsafi Z, Khan M, Sohrabi C, O'Neill N, Iosifidis C, Griffin M, Mathew G, Agha R (2020) Impact of the coronavirus (COVID-19) pandemic on surgical practice - part 2 (surgical prioritisation). *Int J Surg* 79:233–248. <https://doi.org/10.1016/j.ijsu.2020.05.002>
 45. Boskoski I, Pecere S, Bove V, Barbaro F, Perri V, Costamagna G (2020) Impact of SARS-CoV-2 on a high volume endoscopy center in Italy. *Dig Liver Dis* 52(8):819–822. <https://doi.org/10.1016/j.dld.2020.06.009>
 46. Caruana EJ, Patel A, Kendall S, Rathinam S (2020) Impact of coronavirus 2019 (COVID-19) on training and well-being in subspecialty surgery: a national survey of cardiothoracic trainees in the United Kingdom. *J Thorac Cardiovasc Surg* 160(4):980–987. <https://doi.org/10.1016/j.jtcvs.2020.05.052>
 47. Crockett M, Skaar C, Oakley J, Malki M, Hussain M, Barber N (2020) Surgical training during COVID: a positive story. *BJU Int* 126(5):641. <https://doi.org/10.1111/bju.15270>
 48. Daodu O, Panda N, Lopushinsky S, Varghese TK Jr, Brindle M (2020) COVID-19 - considerations and implications for surgical learners. *Ann Surg* 272(1):e22–e23. <https://doi.org/10.1097/SLA.0000000000003927>
 49. Johnson J, Chung MT, Stathakios J, Gonik N, Siegel B (2020) The impact of the COVID-19 pandemic on fellowship training: a national survey of pediatric otolaryngology fellowship directors. *Int J Pediatr Otorhinolaryngol* 136:110217. <https://doi.org/10.1016/j.ijporl.2020.110217>
 50. Nesemeier BR, Lebo NL, Schmalbach CE, Barnes KJ, Vernon D, Ting JY, Shipchandler TZ (2020) Impact of the COVID-19 global pandemic on the otolaryngology fellowship application process. *Otolaryngol Head Neck Surg* 163(4):712–713. <https://doi.org/10.1177/0194599820934370>
 51. Perez-Escamirosa F, Medina-Alvarez D, Ruiz-Vereo EA, Ordorica-Flores RM, Minor-Martinez A, Tapia-Jurado J (2020) Immersive virtual operating room simulation for surgical resident education during COVID-19. *Surg Innov* 27(5):549–550. <https://doi.org/10.1177/1553350620952183>
 52. Rana T, Hackett C, Quezada T, Chaturvedi A, Bakalov V, Leonardo J, Rana S (2020) Medicine and surgery residents' perspectives on the impact of COVID-19 on graduate medical education. *Med Educ Online* 25(1):1818439. <https://doi.org/10.1080/10872981.2020.1818439>
 53. Al-Marhoon MS (2020) The impact of COVID-19 on urology practice in Oman. *Sultan Qaboos Univ Med J* 20(4):e404–e405. <https://doi.org/10.18295/squmj.2020.20.04.023>
 54. Cheserem JB, Esene IN, Mahmud MR, Kalangu K, Sanoussi S, Musara A, El-Ghandour NMF, Fieggen G, Qureshi M (2021) A continental survey on the impact of COVID-19 on neurosurgical training in Africa. *World Neurosurg* 147:e8–e15. <https://doi.org/10.1016/j.wneu.2020.11.008>
 55. Rasyid N, Birowo P, Parikesit D, Rahman F (2020) The impact of the COVID-19 pandemic on urology practice in Indonesia: a nationwide survey. *Urol J* 17(6):677–679. <https://doi.org/10.22037/uj.v16i7.6459>
 56. Hennessy O, Fowler AL, Hennessy C, Hogan A, Nugent E, Joyce M (2020) Covid 19 and surgical training: Carpe Diem. *Br J Surg* 107(12):e591. <https://doi.org/10.1002/bjs.12032>
 57. Molina G, Mehtsun WT, Qadan M, Hause KC, Raut CP, Fairweather M (2020) Virtual interviews for the complex general surgical oncology fellowship: the Dana-Farber/Partners Experience. *Ann Surg Oncol* 27(9):3103–3106. <https://doi.org/10.1245/s10434-020-08778-y>
 58. Teo SH, Abd Rahim MR, Nizlan NM (2020) The impact of COVID-19 pandemic on orthopaedic specialty in Malaysia: a cross-sectional survey. *J Orthop Surg (Hong Kong)* 28(2):2309499020938877. <https://doi.org/10.1177/2309499020938877>
 59. Haji FA (2021) Simulation in neurosurgical education during the COVID-19 pandemic and beyond. *Can J Neurol Sci* 48(2):152–154. <https://doi.org/10.1017/cjn.2020.234>

60. Salganik MJ (2012) Commentary: respondent-driven sampling in the real world. *Epidemiology* 23(1):148–150. <https://doi.org/10.1097/EDE.0b013e31823b6979>
61. Vadivoo S, Gupte MD, Adhikary R, Kohli A, Kangusamy B, Joshua V, Mathai AK, Kumar K, Mainkar M, Goswami P, Team IS (2008) Appropriateness and execution challenges of three formal size estimation methods for high-risk populations in India. *AIDS* 22(Suppl 5):S137–148. <https://doi.org/10.1097/01.aids.0000343772.82647.2d>
62. Chabata ST, Fearon E, Webb EL, Weiss HA, Hargreaves JR, Cowan FM (2020) Assessing bias in population size estimates among hidden populations when using the service multiplier method combined with respondent-driven sampling surveys: survey study. *JMIR Public Health Surveill* 6(2):e15044. <https://doi.org/10.2196/15044>
63. Sheridan GA, Hughes AJ, Quinlan JF, Sheehan E, O'Byrne JM (2020) Quantifying the impact of the COVID-19 pandemic on orthopaedic trainees: a national perspective. *Bone Jt Open* 1(10):645–652. <https://doi.org/10.1302/2633-1462.110.BJO-2020-0149.R1>
64. Garcia-Rojo E, Manfredi C, Santos-Perez-de-la-Blanca R, Tejido-Sanchez A, Garcia-Gomez B, Aliaga-Benitez M, Romero-Otero J, Rodriguez-Antolin A (2021) Impact of COVID-19 outbreak on urology surgical waiting lists and waiting lists prioritization strategies in the post-COVID-19 era. *Actas Urol Esp* 45(3):207–214. <https://doi.org/10.1016/j.acuro.2020.11.001>
65. Wang VT, Odani T, Ito M (2021) Considerations and strategies for restarting elective spine surgery in the midst of a pandemic of COVID-19. *Spine Surg Relat Res* 5(2):52–60. <https://doi.org/10.22603/ssrr.2020-0154>
66. Zouari S, Saadi A, Chakroun M, Oueslati A, Fliess M, Bouzouita A, Derouiche A, Slama RB, Ayed H, Chebil M (2020) Urological activity at the time of COVID-19 pandemic: is there any difference between public and private field? *Pan Afr Med J* 37:389. <https://doi.org/10.11604/pamj.2020.37.389.25297>
67. Guadalajara H, Munoz de Nova JL, Fernandez Gonzalez S, Yiasemidou M, Recarte Rico M, Juez LD, Garcia Septiem J, Galindo Jara P, Garcia Virosta M, Lobo Martinez E, Martin-Perez E, Garcia-Olmo D, PC Group (2020) Patterns of acute surgical inflammatory processes presentation of in the COVID-19 outbreak (PIACO Study): surgery may be the best treatment option. *Br J Surg* 107(11):e494–e495. <https://doi.org/10.1002/bjs.11950>
68. Bogani G, Ditto A, De Cecco L, Lopez S, Guerrisi R, Piccioni F, Micali A, Daidone MG, Raspagliesi F (2020) Transmission of SARS-CoV-2 in surgical smoke during laparoscopy: a prospective, proof-of-concept study. *J Minim Invasive Gynecol*. <https://doi.org/10.1016/j.jmig.2020.12.026>
69. Emile SH, Hamid HKS (2020) A critical review of the safety of minimally invasive surgery in the era of COVID-19. *Minim Invasive Ther Allied Technol* 1–7. <https://doi.org/10.1080/13645706.2020.1838549>
70. Ghazali W, Nallaluthan P, Hasan RZ, Adlan AS, Boon NK (2020) Gynecological Endoscopic Society of Malaysia statement and recommendations on gynecological laparoscopic surgery during COVID-19 pandemic. *Gynecol Minim Invasive Ther* 9(4):185–189. https://doi.org/10.4103/GMIT.GMIT_109_20
71. Trivedi PH, Trivedi SP, Ghadge NM, Bajani DP, Trivedi AS (2020) Safe gynecological laparoscopic surgery during COVID times. *J Hum Reprod Sci* 13(4):310–316. https://doi.org/10.4103/jhrs.JHRS_185_20
72. Covidurg Collaborative GC (2021) SARS-CoV-2 vaccination modelling for safe surgery to save lives: data from an international prospective cohort study. *Br J Surg*. <https://doi.org/10.1093/bjs/znab101>
73. Simoes J, Bhangu A, CovidSurg C (2020) Should we be re-starting elective surgery? *Anaesthesia* 75(12):1563–1565. <https://doi.org/10.1111/anae.15296>
74. Glasbey JC, Nepogodiev D, Simoes JFF, Omar O, Li E, Venn ML, Pgdme, Abou Char MK, Capizzi V, Chaudhry D, Desai A, Edwards JG, Evans JP, Fiore M, Videria JF, Ford SJ, Ganly I, Griffiths EA, Gujjuri RR, Koliass AG, Kaafarani HMA, Minaya-Bravo A, McKay SC, Mohan HM, Roberts KJ, San Miguel-Mendez C, Pockney P, Shaw R, Smart NJ, Stewart GD, SundarMrcog S, Vidya R, Bhangu AA, COCollaborative (2021) Elective cancer surgery in COVID-19-free surgical pathways during the SARS-CoV-2 pandemic: an international, multicenter, comparative cohort study. *J Clin Oncol* 39(1):66–78. <https://doi.org/10.1200/JCO.20.01933>
75. Collaborative CO (2020) Elective surgery cancellations due to the COVID-19 pandemic: global predictive modelling to inform surgical recovery plans. *Br J Surg* 107(11):1440–1449. <https://doi.org/10.1002/bjs.11746>
76. Wouters OJ, Shadlen KC, Salcher-Konrad M, Pollard AJ, Larson HJ, Teerawattananon Y, Jit M (2021) Challenges in ensuring global access to COVID-19 vaccines: production, affordability, allocation, and deployment. *Lancet* 397(10278):1023–1034. [https://doi.org/10.1016/S0140-6736\(21\)00306-8](https://doi.org/10.1016/S0140-6736(21)00306-8)
77. Bleier BS, Ramanathan M Jr, Lane AP (2021) COVID-19 vaccines may not prevent nasal SARS-CoV-2 infection and asymptomatic transmission. *Otolaryngol Head Neck Surg* 164(2):305–307. <https://doi.org/10.1177/0194599820982633>
78. de Caro F, Hirschmann TM, Verdonk P (2020) Returning to orthopaedic business as usual after COVID-19: strategies and options. *Knee Surg Sports Traumatol Arthrosc* 28(6):1699–1704. <https://doi.org/10.1007/s00167-020-06031-3>
79. Hoi KK, Curtis SH, Driver L, Wisnosky E, Zopf DA, Bohm LA (2021) Adoption of telemedicine for multidisciplinary care in pediatric otolaryngology. *Ann Otol Rhinol Laryngol*. <https://doi.org/10.1177/0003489421997651>
80. Mencia MM, Goalan R (2021) COVID-19 and its effects upon orthopaedic surgery: the Trinidad and Tobago experience. *World J Orthop* 12(3):94–101. <https://doi.org/10.5312/wjo.v12.i3.94>
81. Stanley C, Kelly M, Elzaki M, Butler A, Condon F, Lenehan B (2021) Lessons from lockdown: virtual clinics and service reorganisation in fracture management during COVID 19 experience of an Irish Regional Trauma Unit. *Surgeon*. <https://doi.org/10.1016/j.surge.2021.02.006>
82. Silva N, Laiginhas R, Meireles A, Barbosa Breda J (2020) Impact of the COVID-19 pandemic on ophthalmology residency training in Portugal. *Acta Med Port* 33(10):640–648. <https://doi.org/10.20344/amp.14341>
83. Van Heest A, Brandt AM, Dyer G, Homer CJ, Murray PM (2021) COVID-19: impact on orthopaedic graduate medical education in the U.S.: AOA Critical Issues Symposium. *J Bone Joint Surg Am*. <https://doi.org/10.2106/JBJS.20.01948>
84. Jenkins PJGA, Murray O, Anthony I (2014) The Glasgow fracture pathway: a virtual clinic. *BJJ News* 2:22–24
85. Kelly MOKN, Francis A (2020) Connolly Hospital Trauma Assessment clinic: a virtual solution to patient flow. *Ir J Med Sci* 189:425–429
86. Chaudhry H, Nadeem S, Mundi R (2021) How satisfied are patients and surgeons with telemedicine in orthopaedic care during the COVID-19 pandemic? A systematic review and meta-analysis. *Clin Orthop Relat Res* 479(1):47–56. <https://doi.org/10.1097/CORR.0000000000001494>
87. Or M, Breathnach O, Conlon B, Kiernan C, Sheehan E (2019) Trauma assessment clinic: virtually a safe and smarter way of managing trauma care in Ireland. *Injury* 50(4):898–902. <https://doi.org/10.1016/j.injury.2019.03.046>

88. Llorens R, Noe E, Colomer C, Alcaniz M (2015) Effectiveness, usability, and cost-benefit of a virtual reality-based telerehabilitation program for balance recovery after stroke: a randomized controlled trial. *Arch Phys Med Rehabil* 96(3):418–425 e412. <https://doi.org/10.1016/j.apmr.2014.10.019>
89. Ottesen TD, Montoya RL, Ogunleye TD, Brown KE, Woolley PM, Dejean J, Qudsi RA, Agarwal-Harding KJ, Dyer GSM (2021) Implementation and impact evaluation of a virtual orthopaedic continuing medical education conference in a low-resource country. *J Surg Educ*. <https://doi.org/10.1016/j.jsurg.2021.01.002>
90. Martin-Gorgojo A, Bernabeu-Wittel J, Linares-Barríos M, Russo-de la Torre F, Garcia-Doval I, Rio-de D, la Torre E (2020) Attendee survey and practical appraisal of a Telegram(R)-based dermatology congress during the COVID-19 confinement. *Actas Dermosifiliogr* 111(10):852–860. <https://doi.org/10.1016/j.ad.2020.08.009>
91. McDowell L, Goode S, Sundaresan P (2020) Adapting to a global pandemic through live virtual delivery of a cancer collaborative trial group conference: the TROG 2020 experience. *J Med Imaging Radiat Oncol* 64(3):414–421. <https://doi.org/10.1111/1754-9485.13047>
92. Terhune KP, Choi JN, Green JM, Hildreth AN, Lipman JM, Aarons CB, Heyduk DA, Misra S, Anand RJ, Fise TF, Thorne CB, Edwards GC, Joshi ART, Clark CE, Nfonsam VN, Chahine A, Smink DS, Jarman BT (2020) Harrington DT (2020) Ad astra per aspera (Through Hardships to the Stars): lessons learned from the First National Virtual APDS Meeting. *J Surg Educ* 77(6):1465–1472. <https://doi.org/10.1016/j.jsurg.2020.06.015>
93. Tuma F, Nituica C, Mansuri O, Kamel MK, McKenna J, Blebea J (2021) The academic experience in distance (virtual) rounding and education of emergency surgery during COVID-19 pandemic. *Surg Open Sci*. <https://doi.org/10.1016/j.sopen.2021.03.001>
94. DePietro DM, Santucci SE, Harrison NE, Kiefer RM, Trerotola SO, Sudheendra D, Shamimi-Noori S (2021) Medical student education during the COVID-19 pandemic: initial experiences implementing a virtual interventional radiology elective course. *Acad Radiol* 28(1):128–135. <https://doi.org/10.1016/j.acra.2020.10.005>
95. Vervoort D, Dearani JA, Starnes VA, Thourani VH, Nguyen TC (2021) Brave new world: virtual conferencing and surgical education in the coronavirus disease 2019 era. *J Thorac Cardiovasc Surg* 161(3):748–752. <https://doi.org/10.1016/j.jtcvs.2020.07.094>
96. Woodruff P, Wallis CJD, Albers P, Klaassen Z (2021) Virtual conferences and the COVID-19 pandemic: are we missing out with an online only platform? *Eur Urol*. <https://doi.org/10.1016/j.eururo.2021.03.019>
97. Repici A, Maselli R, Colombo M, Gabbiadini R, Spadaccini M, Anderloni A, Carrara S, Fugazza A, Di Leo M, Galtieri PA, Pellegatta G, Ferrara EC, Azzolini E, Lagioia M (2020) Coronavirus (COVID-19) outbreak: what the department of endoscopy should know. *Gastrointest Endosc* 92(1):192–197. <https://doi.org/10.1016/j.gie.2020.03.019>
98. Tang JW, Li Y, Eames I, Chan PK, Ridgway GL (2006) Factors involved in the aerosol transmission of infection and control of ventilation in healthcare premises. *J Hosp Infect* 64(2):100–114. <https://doi.org/10.1016/j.jhin.2006.05.022>
99. Johnston ER, Habib-Bein N, Dueker JM, Quiroz B, Corsaro E, Ambrogio M, Kingsley M, Papachristou GI, Kreiss C, Khalid A (2019) Risk of bacterial exposure to the endoscopist's face during endoscopy. *Gastrointest Endosc* 89(4):818–824. <https://doi.org/10.1016/j.gie.2018.10.034>
100. Wong TW, Lee CK, Tam W, Lau JT, Yu TS, Lui SF, Chan PK, Li Y, Bresee JS, Sung JJ, Parashar UD, Outbreak Study G (2004) Cluster of SARS among medical students exposed to single patient. *Hong Kong. Emerg Infect Dis* 10(2):269–276. <https://doi.org/10.3201/eid1002.030452>
101. Siau K, Iacucci M, Duncckley P, Penman I, EndoTrain Survey C (2020) The impact of COVID-19 on gastrointestinal endoscopy training in the United Kingdom. *Gastroenterology*. 159(4):1582–1585 e1583. <https://doi.org/10.1053/j.gastro.2020.06.015>
102. Yiasemidou M, Glassman D, Tomlinson J, Song D, Gough MJ (2017) Perceptions about the present and future of surgical simulation: a national study of mixed qualitative and quantitative methodology. *J Surg Educ* 74(1):108–116. <https://doi.org/10.1016/j.jsurg.2016.07.011>
103. Yiasemidou M, Glassman D, Khan K, Downing J, Sivakumar R, Fawole A, Biyani CS (2020) Validation of a cost-effective appendectomy model for surgical training. *Scott Med J* 65(2):46–51. <https://doi.org/10.1177/0036933019900340>
104. Yiasemidou M, Gkaragkani E, Glassman D, Biyani CS (2018) Cadaveric simulation: a review of reviews. *Ir J Med Sci* 187(3):827–833. <https://doi.org/10.1007/s11845-017-1704-y>
105. Yiasemidou M, de Siqueira J, Tomlinson J, Glassman D, Stock S, Gough M (2017) “Take-home” box trainers are an effective alternative to virtual reality simulators. *J Surg Res* 213:69–74. <https://doi.org/10.1016/j.jss.2017.02.038>
106. Glassman D, Yiasemidou M, Venkateswaran B, Sivakumar R, Majumder S, Biyani CS (2016) A multi-specialty surgical course for residents transitioning from early to intermediate training. *Int J Med Educ* 7:130–131. <https://doi.org/10.5116/ijme.5708.e9ea>
107. Yiasemidou M, Glassman D, Mushtaq F, Athanasiou C, Williams MM, Jayne D, Miskovic D (2017) Mental practice with interactive 3D visual aids enhances surgical performance. *Surg Endosc* 31(10):4111–4117. <https://doi.org/10.1007/s00464-017-5459-3>
108. Yiasemidou M, Glassman D, Jayne D, Miskovic D (2018) Is patient-specific pre-operative preparation feasible in a clinical environment? A systematic review and meta-analysis. *Comput Assist Surg (Abingdon)* 23(1):57–68. <https://doi.org/10.1080/24699322.2018.1495266>
109. Yiasemidou M, Galli R, Glassman D, Tang M, Aziz R, Jayne D, Miskovic D (2018) Patient-specific mental rehearsal with interactive visual aids: a path worth exploring? *Surg Endosc* 32(3):1165–1173. <https://doi.org/10.1007/s00464-017-5788-2>
110. Yiasemidou M, Mushtaq F, Basheer M, Galli R, Panagiotou D, Stock S, Preston N, Mon-Williams M, Jayne DG, Miskovic D (2021) Patient-specific mental rehearsal with three-dimensional models before low anterior resection: randomized clinical trial. *BJS Open* 5:(2)zraa004. <https://doi.org/10.1093/bjsopen/zraa004>
111. Tomlinson JE, Yiasemidou M, Watts AL, Roberts DJ, Timothy J (2016) Cadaveric spinal surgery simulation: a comparison of cadaver types. *Global Spine J* 6(4):357–361. <https://doi.org/10.1055/s-0035-1563724>
112. Pears M, Yiasemidou M, Ismail MA, Veneziano D, Biyani CS (2020) Role of immersive technologies in healthcare education during the COVID-19 epidemic. *Scott Med J* 65(4):112–119. <https://doi.org/10.1177/0036933020956317>
113. Auricchio G, Kaganerh E (2015) How digitalization is changing the way executives learn. *IESE Insight* 26:31–38
114. Dimension Studio. XR content and virtual production www.dimensionstudio.co Accessed 15th May 2020

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.