



Contents lists available at ScienceDirect

Saudi Pharmaceutical Journal

journal homepage: www.sciencedirect.com



Short communication

## Study of DNA damage caused by dipyrone in presence of some transition metal ions



Bruna Corrêa Roriz, Horacio Dorigan Moya\*

Faculdade de Medicina da Fundação do ABC, CEPES (Centro de Estudos, Pesquisa, Prevenção e Tratamento em Saúde), Av. Príncipe de Gales, 821, Príncipe de Gales, Santo André - CEP - 09060-650, Santo André, SP, Brazil

## ARTICLE INFO

## Article history:

Received 26 October 2016

Accepted 24 February 2017

Available online 27 February 2017

## Keywords:

DNA damage

Dipyrone

Cu<sup>2+</sup>Fe<sup>3+</sup>Ni<sup>2+</sup>Mn<sup>3+</sup>

## ABSTRACT

The DNA damage in the presence of dipyrone (used as its sodium salt, NaDip) and some transition metal ions in an air saturated ([O<sub>2</sub>] ≈ 0.25 mM) non-buffered solution at T = (25.0 ± 0.5)°C was investigated by agarose gel electrophoresis. As metal ions Cu<sup>2+</sup>, Fe<sup>3+</sup>, Ni<sup>2+</sup> and Mn<sup>3+</sup> were selected and evaluated in the present study because of the important role they play in a biological system.

pUC19 plasmid DNA damage-induced by NaDip (80–600 μM) was observed in the presence of 100 μM Cu<sup>2+</sup>. The damage was proportional to the NaDip concentration provided that the order of addition of reagents (pUC19 plasmid DNA + Cu<sup>2+</sup> + NaDip) is obeyed. Addition in the reaction medium of ligands for Cu<sup>2+</sup> and Cu<sup>+</sup>, respectively EDTA and neocuproine, promoted total inhibition or reduction of the pUC19 plasmid DNA damage suggesting the involvement of the Cu<sup>2+</sup>/Cu<sup>+</sup> cycle. Besides, the decrease in the pUC19 plasmid DNA damage after addition of catalase (1.0 × 10<sup>-4</sup> mg μL<sup>-1</sup>) in the same reaction medium indicates that H<sub>2</sub>O<sub>2</sub> is also involved in the damage process.

In NaDip concentration range (80–600 μM), and under same the experimental conditions, it was not possible to conclude whether there was pUC19 plasmid DNA damage caused by 10 μM Fe<sup>3+</sup>. No damage was observed in the presence of Mn<sup>3+</sup> or Ni<sup>2+</sup>.

Although the technique used in this study is sensitive to detect the pUC19 plasmid DNA damage it was not possible to identify in which DNA base this damage occurs. Further studies with other techniques should be made to unambiguously identify the oxidative intermediates that are responsible for the DNA damage.

As far as we know, this is the first study dealing with the pUC19 plasmid DNA damage-induced by NaDip in presence of copper, iron, nickel and manganese ions.

© 2017 The Authors. Production and hosting by Elsevier B.V. on behalf of King Saud University. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

### 1. Introduction

Cells must maintain an appropriate balance between the level of free radicals and antioxidants compounds to ensure its biological structural integrity. If the level of these free radicals exceeds the antioxidants compounds, acquired from diet or endogenously produced, biomolecules such as proteins and DNA can be damaged (Kawanishi et al., 2005).

Several *in vitro* studies have shown that some chemical compounds can induce DNA damage (Schweigert et al., 2000; Asad et al., 2002; Zheng et al., 2006). Among these compounds there are some drugs as L-Dopa (Husain and Hadi, 1995), N-acetylcysteine (Oikawa et al., 1999), procabazine (Ogawa et al., 2003) gentamicin (Lesniak et al., 2003), AZT (Iwamoto et al., 2003), and chloramphenicol (Ohnishi et al., 2015) that, under specific conditions and in the presence of a transition metal ion (M<sup>n+</sup>), can also lead to DNA damage. As a common sense it is believed that these drugs can reduce M<sup>n+</sup> to M<sup>(n-1)+</sup>, which in the presence of H<sub>2</sub>O<sub>2</sub> generates free radicals inducing DNA damage as a consequence.

NaDip, also known as dipyrone, novalgine or metamizole (Fig. 1), is a nonsteroidal anti-inflammatory drug indicated for the treatment of painful manifestations and fever with a complex mechanism responsible for the analgesic effect (Shimada et al., 1994). Despite the suspected occurrence of agranulocytosis, a potentially fatal side

\* Corresponding author.

E-mail address: [horacio.moya@fmabc.br](mailto:horacio.moya@fmabc.br) (H.D. Moya).

Peer review under responsibility of King Saud University.



Production and hosting by Elsevier




Fig. 1. Structure of sodium dipyrone (NaDip).

effect (Hedenmalm and Spigset, 2002; Wessel et al., 2006), this drug is still prescribed and used in many countries. Besides, as far as we know, no study has been done to evaluate the possible DNA damage-induced by NaDip in presence of trace elements.

In the present study pUC-19 plasmid DNA was exposed to some transition metal ions ( $\text{Cu}^{2+}$ ,  $\text{Fe}^{3+}$ ,  $\text{Ni}^{2+}$  and  $\text{Mn}^{3+}$ ), which play important roles in a biological system. Copper is an important element for growth and many enzymes require it to function properly. Iron is essential for hemoglobin formation (macronutrient in pregnant women) and most of this metal is ingested as  $\text{Fe}^{3+}$  in human diet. Nickel is a micronutrient able to prevent anemia by aiding the absorption of iron. Manganese, also a trace element, is a cofactor for a number of important enzymes. Although present as  $\text{Mn}^{2+}$  in human diet an *in vitro* study pointed out that the trivalent state ( $\text{Mn}^{3+}$ ) appear to be more cytotoxic (Chen et al., 2001).

Oxidative damage to the pUC-19 plasmid DNA was verified by agarose gel electrophoresis through the conversion of supercoiled pUC-19 plasmid DNA (supercoil, SC, native conformation) to open circular (OC) form, resulting from single break, in solution containing NaDip in an unbuffered and air saturated solution ( $[\text{O}_2] \approx 0.25 \text{ mM}$ ) (Moreno et al., 2007).

In all experiments the quantification of pUC-19 plasmid DNA damage was performed by the ratio between the total amount of OC (normalized with respect to the background produced by pUC-19 plasmid DNA alone) and the total amount of pUC-19 plasmid DNA present. In none of the experiments was observed the linear form (L), which is the result of the double-strand breaks.

Under these experimental conditions a marked damage to DNA was observed in the presence of  $\text{Cu}^{2+}$  and NaDip, being proportional to the drug concentration. Addition of EDTA and neocuproine, chelating agents for  $\text{Cu}^{2+}$  and  $\text{Cu}^+$  respectively, inhibited or diminished the pUC-19 plasmid DNA damage suggesting the participation of the  $\text{Cu}^{2+}/\text{Cu}^+$  redox cycle. Moreover, the decrease in the pUC-19 plasmid DNA damage after the introduction of catalase allows inferring that  $\text{H}_2\text{O}_2$  is also involved in this damage process.

## 2. Materials and methods

### 2.1. Apparatus

Electrophoresis experiments were performed in a mini submerged horizontal gel chamber using a PowerPac™ Basic power supply.

A transilluminator UV ZT-21 (BioGlow® Crystal) coupled with a digital camera PowerShot G10 (Canon®) was used to obtain the photograph of gels.

pUC-19 plasmid DNA bands were quantified with the AlphaEase® FC Software (Pharmacia Biotech, San Francisco, CA, USA).

### 2.2. Reagents

Deionized water was purified with a Milli-Q Plus Water System (Millipore®) and used to prepare all analytical-grade chemicals (unless stated otherwise).

Supercoiled pUC-19 plasmid DNA (native form)  $0.5 \mu\text{g} \mu\text{L}^{-1}$  and agarose electrophoresis were from MBI Fermentas®.

Ethidium bromide ( $\text{C}_{21}\text{H}_{20}\text{BrN}_3$ ,  $394.31 \text{ g mol}^{-1}$ ,  $500 \mu\text{g mL}^{-1}$ ), Ficoll type 400, bromophenol blue ( $\text{C}_{19}\text{H}_{10}\text{Br}_4\text{O}_5\text{S}$ ,  $669.96 \text{ g mol}^{-1}$ ), boric Acid ( $\text{H}_3\text{BO}_3$ ,  $61.82 \text{ g mol}^{-1}$ ,  $\geq 99.5\%$ ), Neocuproine (NC,  $\text{C}_{14}\text{H}_{12}\text{N}_2$ ,  $208.26 \text{ g mol}^{-1}$ ,  $\geq 98\%$ ), sodium dipyrone (NaDip,  $\text{C}_{13}\text{H}_{16}\text{N}_3\text{NaO}_4\text{S}$ ,  $333.34 \text{ g mol}^{-1}$ ,  $>99.9\%$ ), catalase from bovine liver ( $13,500 \text{ U mg}^{-1}$ ) were from Sigma®.

1,10-phenanthroline monohydrate (phen,  $\text{C}_{12}\text{H}_8\text{N}_2\cdot\text{H}_2\text{O}$ ,  $198.26 \text{ g mol}^{-1}$ ,  $99.5\%$ ) and ethanol ( $\text{CH}_3\text{CH}_2\text{OH}$ ,  $46.06 \text{ g mol}^{-1}$ ,  $99.5\%$ ), TRIS (tris(hydroxymethyl)aminomethane,  $\text{C}_4\text{H}_{11}\text{NO}_3$ ,  $121.14 \text{ g mol}^{-1}$ , disodium EDTA ( $\text{Na}_2\text{EDTA}$ ,  $\text{C}_{10}\text{H}_{14}\text{N}_2\text{O}_8\text{Na}_2\cdot 2\text{H}_2\text{O}$ ,  $372.24 \text{ g mol}^{-1}$ ), manganese(III) acetate dihydrate ( $\text{C}_6\text{H}_9\text{MnO}_6\cdot\text{H}_2\text{O}$ ,  $268.10 \text{ g mol}^{-1}$ ,  $>98\%$ ), copper disodium EDTA ( $\text{Na}_2\text{Cu}[\text{EDTA}]$ ,  $397.74 \text{ g mol}^{-1}$ ,  $\geq 99.5\%$ ) were from Merck®.

### 2.3. Solutions

pUC-19 plasmid DNA  $1.875 \times 10^{-2} \mu\text{g} \mu\text{L}^{-1}$  solution was obtained transferring  $6 \mu\text{L}$  pUC-19 plasmid DNA  $0.5 \mu\text{g} \mu\text{L}^{-1}$  (freshly thawed) to a 1.5 mL microtube and adding  $154 \mu\text{L}$  water.

NaDip 7.5 mM stock solution was prepared by transferring 0.2500 g into a 100 mL volumetric flask and completed with water. Working solutions (0.5–800  $\mu\text{M}$ ) were obtained by appropriate dilution.

$\text{Cu}(\text{NO}_3)_2$ ,  $\text{Fe}(\text{ClO}_4)_3$  and  $\text{Ni}(\text{NO}_3)_2$  0.20 M stock solutions (respectively at pH 6.3, 3.5 and pH 4.5) were prepared from reaction of pure electrolytic metal (99.99%, Sigma®) with doubly distilled nitric acid (99.99% Sigma®) as described elsewhere (Moreno et al., 2007; Alipázaga et al., 2009, 2010). A 0.75 mM diluted solution of each ion was daily prepared by diluting 375  $\mu\text{L}$  of 0.20 M stock solution in a 100.0 mL volumetric flask.

$\text{Cu}[\text{EDTA}]^{2-}$  0.75 mM complex solution was prepared by dissolving 0.0140 g  $\text{Na}_2\text{Cu}[\text{EDTA}]$  in a 50.0 mL volumetric flask (pH 7.5).

$\text{Cu}(\text{NC})_2^+$  0.75 mM complex solution was prepared by mixing 0.0156 g NC and 188  $\mu\text{L}$   $\text{Cu}(\text{NO}_3)_2$  0.2 M stock solution in a 50.0 mL volumetric flask (pH 7.5).

$\text{Fe}[\text{EDTA}]^-$  0.75 mM complex solution was prepared by dissolving 0.0140 g  $\text{Na}_2\text{EDTA}$  and mixing with 188  $\mu\text{L}$   $\text{Fe}(\text{ClO}_4)_3$  0.20 M in a 50.0 mL volumetric flask (pH 4.5).

Solution mixture containing  $\text{Fe}^{3+}$  and phen, both 0.75 mM, was prepared by dissolving 0.0223 g phen and mixing with 188  $\mu\text{L}$   $\text{Fe}(\text{ClO}_4)_3$  0.2 M in a 50.0 mL volumetric flask (pH 5.0).

$\text{Mn}(\text{H}_3\text{C-COO})_3$  0.75 mM was prepared just before use transferring 0.020 g manganese(III)-acetate-dihydrate to a 100.0 mL volumetric flask containing 50 mL of water (pH 4.5).

$\text{Cu}^{2+}$  (1.0–100  $\mu\text{M}$ ),  $\text{Fe}^{3+}$  (5.0–100  $\mu\text{M}$ ),  $\text{Ni}^{2+}$  and  $\text{Mn}^{3+}$  (10–100  $\mu\text{M}$ ) working solutions were all obtained by appropriate dilution.

Catalase freshly solution was prepared by dissolving 1.0 mg in 2.0 mL water.

The preparation of TBE buffer (pH 8.0) and agarose gel 0.8% are described in Supplementary Material.

## 3. Procedures

### 3.1. Strategy for preparation of mixtures

In all experiments, pUC-19 plasmid DNA were exposed to a transition metal ion (complexed or not) in air saturated solution ( $[\text{O}_2] \approx 0.25 \text{ mM}$ ) followed by the addition of NaDip solution. The percentages of pUC-19 plasmid DNA strand breaks were observed when the freshly prepared solutions of the reagents were mixed.

Three series of experiments were carried out in the following order:

- In the presence of metal ion and absence of NaDip: 20  $\mu\text{L}$  of pUC-19 plasmid DNA  $1.875 \times 10^{-2} \mu\text{g} \mu\text{L}^{-1}$  were transferred to eight 1.5 mL microtubes. From the second microtube 10  $\mu\text{L}$  a transition metal ion solution of different concentrations were added.
- In the presence of a fixed concentration of metal ion containing NaDip: 20  $\mu\text{L}$  of pUC-19 plasmid DNA  $1.875 \times 10^{-2} \mu\text{g} \mu\text{L}^{-1}$  were transferred to eight 1.5 mL microtubes. From the second microtube 10  $\mu\text{L}$  a transition metal ion solution (complexed or not) were added and from the third microtube 10  $\mu\text{L}$  of NaDip solution of different concentrations were added.
- In the presence of a fixed concentration of NaDip containing metal ion: 20  $\mu\text{L}$  pUC-19 plasmid DNA  $1.875 \times 10^{-2} \mu\text{g} \mu\text{L}^{-1}$  were transferred to eight 1.5 mL microtubes. From the second microtube 10  $\mu\text{L}$  of NaDip solution were added and from the third microtube 10  $\mu\text{L}$  a transition metal ion solution (complexed or not) of different concentrations were added.

All of microtubes were made up to 50  $\mu\text{L}$  with water. The temperature was kept at  $(25.0 \pm 0.5) ^\circ\text{C}$ . The concentrations of the final solutions are indicated in the legends of the figures.

The gel electrophoresis experiments are described in [Supplementary Material](#).

## 4. Results and discussion

### 4.1. Electrophoresis experiments in the presence of $\text{Cu}^{2+}$

Several control experiments were carried out in the present work. No pUC-19 plasmid DNA damage was observed when  $\text{Cu}^{2+}$  ranges from 10 to 120  $\mu\text{M}$  in the absence of NaDip ([Fig. 1, Supplementary Material](#)) under these experimental conditions. Since the  $\text{Cu}^{2+}$  concentration used is not strictly established ([Husain and Hadi, 1995; Ogawa et al., 2003; Iwamoto et al., 2003; Lesniak et al., 2003](#)), in the studies involving DNA damage in the presence of drugs, a  $\text{Cu}^{2+}$  100  $\mu\text{M}$  solution ([Frelon et al., 2003](#)) was chosen to be used in the present study ([Fig. 2C](#)).

Under the experimental conditions used in the present study no pUC-19 plasmid DNA damage was markedly observed when NaDip ranges from 0.5 to 100  $\mu\text{M}$  ([Fig. 2A](#)), however the damage can be clearly noted from 200  $\mu\text{M}$  ([Fig. 2B, lane 4](#)). [Fig. 2C](#) shows the results of an experiment by varying  $\text{Cu}^{2+}$  concentration and main-

taining fixed NaDip at 500  $\mu\text{M}$  in which it is possible to notice that the pUC-19 plasmid DNA damage is complete when  $\text{Cu}^{2+}$  is 100  $\mu\text{M}$  ([Fig. 2C, lane 8](#)). It can be noted from these results that there is a minimum  $[\text{NaDip}:\text{Cu}^{2+}]$  concentration ratio for pUC-19 plasmid DNA damage be initiated (2:1) and then completed (5:1).


Experiments have shown that when  $\text{Cu}^{2+}$  (100  $\mu\text{M}$ ) is added as  $\text{Cu}[\text{EDTA}]^{2-}$  ( $\log \beta = 18.8$ ) ([Smith and Martell, 2004](#)) no pUC-19 plasmid DNA damage was observed ([Fig. 2, Supplementary Material](#)). In addition, when  $\text{Cu}^{2+}$  (100  $\mu\text{M}$ ) is added along with NC (a ligand for  $\text{Cu}^+$ ) the pUC-19 plasmid DNA damage indicated by OC formation is significantly repressed ([Fig. 2D](#)). The decrease of NaDip/ $\text{Cu}^{2+}$ -mediated pUC-19 plasmid DNA strand breaks by NC addition occurs as a result of the  $\text{Cu}(\text{NC})_2^+$  complex formation ( $\log \beta_2 = 19.1$ ) ([Smith and Martell, 2004](#)) removing  $\text{Cu}^+$  from the solution. This corroborates that the  $\text{Cu}^{2+}/\text{Cu}^+$  redox cycle occurs (in the presence of  $\text{O}_2$ ) and is essential for the pUC-19 plasmid DNA damage in presence of NaDip.

In another experiment the enzyme catalase was added in the following order: pUC-19 plasmid DNA, catalase (final concentration  $1 \times 10^{-4} \text{mg} \mu\text{L}^{-1}$ ),  $\text{Cu}^{2+}$  and NaDip (experimental conditions of [Fig. 2](#)). Since the addition of the enzyme catalase inhibited entirely the pUC-19 plasmid DNA damage ([Fig. 3, Supplementary Material](#)) it is possible to infer that the presence of  $\text{H}_2\text{O}_2$  in the reaction medium is necessary to cause this damage.

In the experiments where pUC-19 plasmid DNA damage occurred it was noted that the order of addition of reagents (i.e. pUC-19 plasmid DNA +  $\text{Cu}^{2+}$  + NaDip) is critical. In fact, when it was added pUC-19 plasmid DNA + NaDip +  $\text{Cu}^{2+}$  no damage was observed. This suggests that the copper ions must be linked to the pUC-19 plasmid DNA before the NaDip-mediated damage occurs. In fact, it was reported that  $\text{Cu}^{2+}$  ions are able to interact with DNA in-between the bases ([Lloyd and Phillips, 1999](#)) yielding DNA-copper-hydroperoxo complexes which generates highly reactive species as  $\text{O}_2$  and  $\cdot\text{OH}$  ([Schweigert et al., 2000](#)). The experiments described above seem to show that there is no formation of complexes between NaDip and  $\text{Cu}^{2+}$  under these experimental conditions.




In addition, it has been previously shown that resveratrol ([Win et al., 2002](#)), hydroquinone ([Li and Trush, 1993](#)) and catechol ([Schweigert et al., 2000](#)), all non-drugs, are also capable of inducing DNA strand breaks (OC and L configurations) in the presence of  $\text{Cu}^{2+}$  with effective contribution of  $\text{H}_2\text{O}_2$  and the  $\text{Cu}^{2+}/\text{Cu}^+$  redox cycle.

Based on those reports ([Win et al., 2002; Li and Trush, 1993; Schweigert et al., 2000](#)), and from the observations of the present study, a mechanism for the pUC-19 plasmid DNA damage in the presence of  $\text{Cu}^{2+}$  and NaDip (Eqs. (1) to (8)) is proposed (despite




**Fig. 2.** Effect of  $\text{Cu}^{2+}$ . The pUC19 plasmid DNA is present in two conformations SC (native) and OC (resulting from single strand breaks). Saturated air solution ( $[\text{O}_2] \approx 0.25 \text{mM}$ ), pH 7.5,  $T = (25.0 \pm 0.5) ^\circ\text{C}$ . All lanes (1) contains only pUC19 plasmid DNA  $5 \text{ng} \mu\text{L}^{-1}$ . (A) Lane (2): (1) +  $\text{Cu}^{2+}$  100  $\mu\text{M}$ ; Lanes (3): (2) + NaDip 0.5; 1.0, 5.0; 10; 50 and 100  $\mu\text{M}$  respectively. (B) Lane (2): (1) +  $\text{Cu}^{2+}$  100  $\mu\text{M}$ ; Lanes (3–8): (2) + NaDip 80; 200; 300; 400; 500 and 600  $\mu\text{M}$ , respectively. (C) Lane (2): (1) + NaDip 500  $\mu\text{M}$ ; Lane (3): (2) +  $\text{Cu}^{2+}$  5.0; 10; 25; 50; 75 and 100  $\mu\text{M}$ , respectively. (D) Lane (2): (1) +  $\text{Cu}(\text{NC})_2^+$  100  $\mu\text{M}$ ; Lanes (3–8): (2) + NaDip 80; 200; 300; 400; 500 and 600  $\mu\text{M}$ , respectively.

the fact that the oxidized form of Dip<sup>-</sup> has not yet been properly elucidated and that the O<sub>2</sub><sup>-</sup> and ·OH free radicals were not determined in the present study):




From the data obtained in the *in vitro* study presented here (Fig. 5) it can be seen that NaDip concentration that caused markedly pUC-19 plasmid DNA damage (in presence of Cu<sup>2+</sup>) is about 7500 times smaller than the NaDip concentration usually available




**Fig. 5.** pUC-19 plasmid DNA damage in the presence of NaDip and Cu<sup>2+</sup> or Fe<sup>3+</sup> (both 100 μM). Saturated air solution ([O<sub>2</sub>] ≈ 0.25 mM). pH = 7.5. T = (25.0 ± 0.5) °C. The data points represent the average of three OC% values (error between 1 and 10%). Experimental conditions shown in Figs. 2 and 3.

in most pharmaceutical formulations (500 mg mL<sup>-1</sup> ≈ 1.60 M). Nevertheless, it does not allow inferring that the same damage degree occurs *in vivo* conditions. In fact, in the human body the presence of ceruloplasmin (protein responsible for binding to the free copper) may prevent Cu<sup>2+</sup> from being a catalytic inducing factor in redox reactions.



**Fig. 3.** Effect of Fe<sup>3+</sup>. The pUC19 plasmid DNA is present in two conformations SC (native) and OC (resulting from single strand breaks). Saturated air solution ([O<sub>2</sub>] ≈ 0.25 mM). pH 7.5; T = (25.0 ± 0.5) °C. All lanes (1) contains only pUC19 plasmid DNA 5 ng μL<sup>-1</sup>. (A) Lanes (2) a (8): (1) + Fe<sup>3+</sup> 5.0; 10; 20; 25; 50; 75 and 100 μM, respectively (No NaDip). (B) Lane (2): (1) + NaDip 500 μM; Lanes (3–8): (2) + Fe<sup>3+</sup> 10; 20; 25; 50; 75 and 100 μM, respectively. (C) Lane (2): (1) + Fe<sup>3+</sup> 10 μM; Lanes (3) a (8): (2) + NaDip 300; 400; 500; 600; 700 and 800 μM, respectively. (D) Lane (2): (1) + Fe<sup>3+</sup> and phen (both 10 μM); Lane (3–8): (2) + NaDip 300; 400; 500; 600; 700 and 800 μM, respectively.



**Fig. 4.** Effect of Ni<sup>2+</sup> or Mn<sup>3+</sup>. The pUC19 plasmid DNA is present in two conformations SC (native) and OC (resulting from single strand breaks). Saturated air solution ([O<sub>2</sub>] ≈ 0.25 mM). pH 7.5. T = (25.0 ± 0.5) °C. Solutions containing Ni<sup>2+</sup> (Figures A and C) or Mn<sup>3+</sup> (Figures B and D). All lanes (1) contains only pUC19 plasmid DNA 5 ng μL<sup>-1</sup>. (A) Lanes (2) a (8): (1) + Ni<sup>2+</sup> 10; 20; 30; 40; 50; 75 and 100 μM, respectively (No NaDip). (B) Lane (2): (1) + Mn<sup>3+</sup> 10; 20; 30; 40; 50; 75 and 100 μM, respectively (No NaDip). (C) Lane (2): (1) + Ni<sup>2+</sup> 10 μM; Lanes (3–8): (2) + NaDip 300; 400; 500; 600; 700 and 800 μM, respectively. (D) Lane (2): (1) + Mn<sup>3+</sup> 10 μM; Lane (3–8): (2) + NaDip 300; 400; 500; 600; 700 and 800 μM, respectively.

#### 4.2. Electrophoresis experiments in the presence of Fe<sup>3+</sup>

In the absence of NaDip and in air saturated solution ([O<sub>2</sub>] ≈ 0.25 mM) the pUC-19 plasmid DNA damage occurs from addition of Fe<sup>3+</sup> 10 μM (Fig. 3A, lane 3). An observable smear, a damage noted when DNA is divided into small fragments, begins at Fe<sup>3+</sup> 50 μM and it is complete at 100 μM (Fig. 3A, lane 8). The repetition of the same experiment in a solution NaDip 500 μM does not produce any appreciable effect which seems surprisingly to show that NaDip does not reduce Fe<sup>3+</sup> promoting the Fe<sup>3+</sup>/Fe<sup>2+</sup> redox cycle (Fig. 3B) under these experimental conditions (similar reaction represented by Eq. (1)).

Moreover, keeping Fe<sup>3+</sup> 10 μM, minimum concentration at which the damage was observed (Fig. 3A, lane 3), and NaDip in the range 300–800 μM, the % OC formed remains constant, (15 ± 5)% up to 600 μM (Fig. 3C, lane 6). The damage in the presence of Fe<sup>3+</sup> cannot be surely concluded in these experimental conditions ([O<sub>2</sub>] ≈ 0.25 mM and T = (25.0 ± 0.5) °C) due to small increases of OC (21 and 23%) in presence of NaDip (700 and 800 μM), respectively.

On the other hand, if 10 μM Fe<sup>3+</sup> is added complexed with EDTA (in presence of NaDip 300–800 μM) the Fe[EDTA]<sup>-</sup> complex formed (log β = 25) (Smith and Martell, 2004) totally suppress this small pUC-19 plasmid DNA damage (Fig. 4, Supplementary Material) observed in Fig. 3A and B. Similarly, if a mixture containing 10 μM Fe<sup>3+</sup> and 10 μM phen (which binds stronger with Fe<sup>2+</sup>, log β<sub>3</sub> = 21.5, but also can form complex with Fe<sup>3+</sup>, log β<sub>3</sub> = 14.1) (Smith and Martell, 2004), was added in presence of NaDip (300–800 μM) no pUC-19 plasmid DNA damages is observed (Fig. 3D).

The reagent addition order strategy used in this study, (item 3.1), allows the metal ion to associate with the pUC-19 plasmid DNA before the addition of NaDip. The small pUC-19 plasmid DNA damage in the presence of Fe<sup>3+</sup> might be attributed to the final pH of the mixtures containing pUC-19 plasmid DNA, Fe<sup>3+</sup> and NaDip because in this experimental condition (pH 7.5) ferric-hydroxo complexes should be present (Baes and Mesmer, 1986).

#### 4.3. Electrophoresis experiments in the presence of Ni<sup>2+</sup> and Mn<sup>3+</sup>

Ni<sup>2+</sup> and Mn<sup>3+</sup> cause pUC-19 plasmid DNA damage (in the absence of NaDip) after 20 μM addition of each metal ion (Fig. 4A and B) and smear was observed from Ni<sup>2+</sup> 50 μM (Fig. 4A, lane 5). For Mn<sup>3+</sup> the % OC remained constant (50 ± 6)% up 100 μM and no occurrence of smear was observed (Fig. 4B).

Ni<sup>2+</sup> experiments carried out at 10 μM (fixed concentration) and NaDip from 300 to 800 μM did not form significant amounts of OC form (10 ± 2)% (Fig. 4C). It was pointed out that Ni<sup>2+</sup> complex preferentially with phosphate groups on the DNA backbone (Lloyd and Phillips, 1999). Since Ni<sup>2+</sup> cannot be readily reduced in aqueous solutions, even in presence of suitable complexing agents, it can be inferred that under these experimental conditions the addition of NaDip does not promote the formation of the Ni<sup>2+</sup>/Ni<sup>+</sup> redox cycle. In fact, Ni<sup>2+</sup> was able of causing pUC-19 plasmid DNA damage, induced by autoxidation of S(IV), when complexed with glycylglycylhistidine (GGH), forming the Ni<sup>3+</sup>/GGH complexes with formation of none Ni<sup>+</sup> complexes (Alipázaga et al., 2005).

Despite of the high value of the standard reduction potential of the Mn(III)/Mn(II) couple in aqueous solution (E<sup>0</sup> = 1.51 V vs. NHE) (Moya et al., 1997) the pUC-19 plasmid DNA damage in the presence of the reducing agent such NaDip was not observed.

In fact, for Mn(III) the increase of OC form was also insignificant in presence of NaDip, (10 ± 4)%. Here again the presence of hydrolysed species of Mn<sup>3+</sup> can prevent the formation of Mn<sup>3+</sup>-pUC-19 plasmid DNA interaction since there is considerable evidence that

Mn<sup>3+</sup> hydrolyses even in concentrated acid solution (Baes and Mesmer, 1986). Besides, previous work observed that DNA damage does not occur in a solution containing Mn<sup>3+</sup> and AZT even when exposed to UVB (Iwamoto et al., 2003).

## 5. Conclusions

Supercoiled of pUC 19 plasmid DNA configuration becomes converted into OC after exposure to 100 μM Cu<sup>2+</sup> and (80–600) μM NaDip. There is a critical order in the addition of reagents (pUC-19 plasmid DNA + Cu<sup>2+</sup> + NaDip) for the pUC-19 plasmid DNA damage. No damage was observed when NaDip was added before Cu<sup>2+</sup>.

No marked pUC-19 plasmid DNA damage was observed in the presence of NaDip and Fe<sup>3+</sup>, Ni<sup>2+</sup> or Mn<sup>3+</sup> under the experimental conditions of this study (air saturated solution with [O<sub>2</sub>] ≈ 0.25 mM and 25.0 °C). This may be because in these experimental conditions, and with order of addition of reagents, Fe<sup>2+</sup>, Ni<sup>+</sup> and Mn<sup>2+</sup> are not originated after addition of NaDip and do not return to their oxidized states, which prevents the formation of the M<sup>n+</sup>/M<sup>(n-1)+</sup> redox cycle. In oxygen saturated solutions and in other temperatures these remarks could be different.

In spite of the gel electrophoresis study performed here has been efficacious to detect the pUC-19 plasmid DNA damage it was not possible to find out which DNA base was damaged.

Further studies with other techniques (e.g. HPLC) should be made to identify in which pUC-19 plasmid DNA base this damage occurs since, at this point, it is still difficult to unambiguously identify the oxidative intermediates that are responsible for the damage.

## Author contributions

B.C. Roriz carried out the experiments and wrote a first draft of the paper. H.D. Moya conceived and designed the experiments and wrote the final draft of the paper.

## Conflicts of interest

The authors declare no conflict of interest.

## Acknowledgements

The authors acknowledge the financial support of FAPESP (Fundação de Amparo à Pesquisa do Estado de São Paulo) and NEPAS (Núcleo de Ensino, Pesquisa e Assessoria a Saúde) from FMABC to the scholarship granted to B.C. Roriz.

## Appendix A. Supplementary material

Supplementary data associated with this article can be found, in the online version, at <http://dx.doi.org/10.1016/j.jpsps.2017.02.010>.

## References

- Alipázaga, M.V., Moreno, R.G.M., Linares, E., Medeiros, M.H.G., Coichev, N., 2005. Oxidative DNA damage induced by autoxidation of microquantities of S(IV) in the presence of Ni(II)-gly-gly-his. *Dalton Trans.*, 3738–3744.
- Alipázaga, V.M., Cerchiaro, G., Moya, H.D., Coichev, N., 2009. Oxidative DNA damage mediated by Cu<sup>2+</sup> and Cu<sup>+</sup> complexes in the presence of S(IV). *J. Braz. Chem. Soc.* 20 (7), 1302–1312.
- Alipázaga, V.M., Moya, H.D., Coichev, N., 2010. Effect of some antioxidants on the oxidative DNA damage induced by autoxidation of microquantities of sulfite in the presence of Ni<sup>2+</sup>/Gly-Gly-L-His. *J. Coord. Chem.* 63 (14), 2450–2460.
- Asad, S.F., Singh, S., Ahmad, A., Hadi, S.M., 2002. Bilirubin/biliverdin-Cu<sup>2+</sup> induced DNA breakage; reaction mechanism and biological significance. *Toxicol. Lett.* 131, 181–189.

- Baes, C.F., Mesmer, R.E., 1986. The Hydrolysis of Cations. Wiley, New York, pp. 223–233.
- Chen, J.-Y., Tsao, G.C., Zhao, Q., Zheng, W., 2001. Differential cytotoxicity of Mn(II) and Mn(III): special reference to mitochondrial [Fe-S] containing enzymes. *Toxicol. Appl. Pharmacol.* 175 (2), 160–168.
- Frelon, S., Douki, T., Favier, A., Cadet, J., 2003. Hydroxyl radical is not the main reactive species involved in the degradation of DNA bases by copper in the presence of hydrogen peroxide. *Chem. Res. Toxicol.* 16, 191–197.
- Hedenmalm, K., Spigset, O., 2002. Agranulocytosis and other blood dyscrasias associated with dipyrone (metamizole). *Eur. J. Clin. Pharmacol.* 58 (4), 265–274.
- Husain, S., Hadi, S.M., 1995. Strand scission in DNA induced by L-dopa in the presence of Cu<sup>2+</sup>. *FEBS Lett.* 364 (75), 78.
- Iwamoto, T., Hiraku, Y., Oikawa, S., Mizutani, H., Kojima, M., Kawanishi, S., 2003. Oxidative DNA damage induced by photodegradation products of 3'-azido-3'-deoxythymidine. *Arch. Biochem. Biophys.* 416, 155–163.
- Kawanishi, S., Oikawa, S., Murata, M., 2005. Evaluation for safety of antioxidant chemopreventive agents. *Antioxid. Redox Signal.* 7 (11–12), 1728–1739.
- Lesniak, W., Harris, W.R., Kravitz, J.Y., Schacht, J., Pecoraro, V.L., 2003. Solution chemistry of copper(II)-gentamicin complexes: relevance to metal-related aminoglycoside Toxicity. *Inorg. Chem.* 42, 1420–1429.
- Li, Y., Trush, M.A., 1993. DNA damage resulting from the oxidation of hydroquinone by copper: role for a Cu<sup>2+</sup>/Cu<sup>+</sup> redox cycle and reactive oxygen generation. *Carcinogenesis* 14, 1303–1311.
- Lloyd, D.R., Phillips, D.H., 1999. Oxidative DNA damage mediated by copper(II), iron (II) and nickel(II). Fenton reactions: evidence for site-specific mechanisms in the formation of double-strand breaks, 8-hydroxydeoxyguanosine and putative intrastrand cross-links. *Mutat. Res.* 424, 23–36.
- Moreno, R.G.M., Alipázaga, M.V., Gomes, O.F., Linares, E., Medeiros, M.H.G., Coichev, N., 2007. DNA damage and 2'-deoxyguanosine oxidation induced by S(IV) autoxidation catalyzed by copper(II) tetraglycine complexes: synergistic effect of a second metal ion. *J. Inorg. Biochem.* 101, 866–875.
- Moya, H.D., Neves, A.E., Coichev, N., 1997. The stabilization of manganese (III) by azide ions in aqueous solution. *Talanta* 44 (797), 803.
- Ogawa, K., Hiraku, Y., Oikawa, S., Murata, M., Sugimura, Y., Kawamura, J., Kawanishi, S., 2003. Molecular mechanisms of DNA damage induced by procarbazine in the presence of Cu<sup>2+</sup>. *Mutat. Res.* 539, 145–155.
- Oikawa, S., Yamada, K., Yamashita, N., Tada-Oikawa, S., Kawanishi, S., 1999. N-acetylcysteine, a cancer chemopreventive agent, causes oxidative damage to cellular and isolated DNA. *Carcinogenesis* 20 (8), 1485–1490.
- Ohnishi, S., Murata, M., Ida, N., Oikawa, S., Kawanishi, S., 2015. Oxidative DNA damage induced by metabolites of chloramphenicol, an antibiotic drug. *Free Radical Res.* 49 (9), 1165–1172.
- Schweigert, N., Acero, L., Gunten, U., Canonica, S., Zehnder, A.J.B., Eggen, R.I.L., 2000. DNA degradation by the mixture of copper and catechol is caused by DNA-copper-hydroperoxo complexes, probably DNA-Cu + OOH. *Environ. Mol. Mutagen.* 36 (1), 5–12.
- Shimada, S.G., Otterness, I.G., Stitt, J.T., 1994. A study of the mechanism of action of the mild analgesic dipyrone. *Agents Actions* 41 (3–4), 188–192.
- Smith, R.M., Martell, A.E., 2004. Critically selected stability constants of metal complexes. *Nist 46, Version 8.0.*
- Wessel, J., Matya, M., Neugebauer, M., Kiefer, H., Daldrup, T., Torbah, F., Weber, H., 2006. Characterization of oxalic acid derivatives as new metabolites of metamizol (dipyrone) in incubated hen's egg and human. *Eur. J. Pharm. Sci.* 28, 15–25.
- Win, W., Cao, Z., Peng, X., Trush, M.A., Li, Y., 2002. Different effects of genistein and resveratrol on oxidative DNA damage in vitro. *Mutat. Res.* 513, 113–120.
- Zheng, Li-Fang, Wei, Qing-Yi, Cai, Yu-Jun, Fang, Jian-Guo, Zhou, B., Yang, L., Liu, Zhong-Li, 2006. DNA damage induced by resveratrol and its synthetic analogues in the presence of Cu<sup>2+</sup> ions: mechanism and structure-activity relationship. *Free Radical Biol. Med.* 41, 1807–1816.