

Review



Cite this article: Cuttler K, Hassan M, Carr J, Cloete R, Bardien S. 2021 Emerging evidence implicating a role for neurexins in neurodegenerative and neuropsychiatric disorders. *Open Biol.* **11**: 210091. <https://doi.org/10.1098/rsob.210091>

Received: 13 April 2021
Accepted: 25 August 2021

Subject Area:

genetics/cellular biology/molecular biology

Keywords:

neurexin, neuroligin, synapse, neurodegenerative disorders, neuropsychiatric disorders, protein interactions

Author for correspondence:

Katelyn Cuttler
e-mail: kcuttler@sun.ac.za

Electronic supplementary material is available online at <https://doi.org/10.6084/m9.figshare.c.5628933>.

Emerging evidence implicating a role for neurexins in neurodegenerative and neuropsychiatric disorders

Katelyn Cuttler¹, Maryam Hassan³, Jonathan Carr^{2,4}, Ruben Cloete³ and Soraya Bardien^{1,4}

¹Division of Molecular Biology and Human Genetics, Department of Biomedical Sciences, and ²Division of Neurology, Department of Medicine, Faculty of Medicine and Health Sciences, Stellenbosch University, Cape Town, South Africa

³South African Medical Research Council Bioinformatics Unit, South African National Bioinformatics Institute, University of the Western Cape, Cape Town, South Africa

⁴South African Medical Research Council/Stellenbosch University Genomics of Brain Disorders Research Unit, Cape Town, South Africa

KC, 0000-0003-1833-5426

Synaptopathies are brain disorders characterized by dysfunctional synapses, which are specialized junctions between neurons that are essential for the transmission of information. Synaptic dysfunction can occur due to mutations that alter the structure and function of synaptic components or abnormal expression levels of a synaptic protein. One class of synaptic proteins that are essential to their biology are cell adhesion proteins that connect the pre- and post-synaptic compartments. Neurexins are one type of synaptic cell adhesion molecule that have, recently, gained more pathological interest. Variants in both neurexins and their common binding partners, neuroligins, have been associated with several neuropsychiatric disorders. In this review, we summarize some of the key physiological functions of the neurexin protein family and the protein networks they are involved in. Furthermore, examination of published literature has implicated neurexins in both neuropsychiatric and neurodegenerative disorders. There is a clear link between neurexins and neuropsychiatric disorders, such as autism spectrum disorder and schizophrenia. However, multiple expression studies have also shown changes in neurexin expression in several neurodegenerative disorders, including Alzheimer's disease and Parkinson's disease. Therefore, this review highlights the potential importance of neurexins in brain disorders and the importance of doing more targeted studies on these genes and proteins.

1. Introduction

There is accumulating evidence to suggest that synaptic dysfunction is present in both neuropsychiatric disorders, such as autism spectrum disorders (ASDs), schizophrenia and bipolar disorder (BD), and neurodegenerative disorders, such as Parkinson's disease (PD), Alzheimer's disease (AD) and Huntington's disease (HD) [1]. In fact, involvement of the synapse is such a prominent feature of the pathogenesis of various brain disorders that it has led to the coining of a specific term, 'synaptopathies'. Indeed, in the case of PD, the involvement of synaptopathy as an initial and central event in the disease pathogenesis, which precedes neuronal damage, has been postulated [2]. Synaptic dysfunction can occur due to mutations that alter the structure and function of synaptic components or abnormal expression levels of a synaptic protein.

Synapses are specialized junctions between neurons that transmit information and they connect neurons into millions of 'neural circuits' that underlie all brain functions [3]. The information transmitted allows the nervous system to respond

to external stimuli and controls bodily functions, behaviour, emotions and memories [4]. This system is tightly controlled and regulated, and even slight perturbations can lead to synaptic dysfunction. An important aspect of synapse biology is the cell adhesion molecules that connect pre- and post-synaptic compartments [5]. These interactions in the synaptic cleft help to maintain synapse structure by delineating mutual boundaries [6]. These proteins are also important in synapse plasticity as synaptic cell adhesion is able to regulate the remodelling of synapses [7]. Interestingly, they are also involved in trans-synaptic signalling [5]. Thus, these proteins are highly important in the organization of synaptic junctions and overall brain function.

Neurexins are one type of synaptic cell adhesion molecule. They are pre-synaptically localized and bind to neuroligins and other proteins in the post-synapse (figure 1). Neurexins and their common binding partners, neuroligins, have recently gained more pathological interest as variants in both have been associated with several neuropsychiatric disorders, including autism and schizophrenia [8]. This further suggests that synaptic dysfunction plays a role in the development of these disorders. Synaptic dysfunction is also known to occur in neurodegenerative disorders [9]; however, it was considered an endpoint of these disorders, due to the considerably later onset of clinical symptoms and progressive appearance of cognitive deficits. This dichotomy has, recently, been challenged by the creation of ‘disease-in-a-dish’ models for multiple central nervous system (CNS) pathologies [9]. This research has identified commonalities between developmental and degenerative disorders, at both the cellular and molecular level, with most of these common mechanisms meeting at the synapse level [9]. Indeed, our laboratory has, recently, found a novel variant (p.G849D) in the *NRXN2* gene which may be implicated in PD [10]. Therefore, we believe it is important to investigate the potential role of neurexins in various neuropsychiatric and neurodegenerative disorders.

In this review, we summarize some of the key physiological functions of the neurexin protein family and the protein networks they are involved in. We also examine the available published literature to determine what research has been done on neurexins in neuropsychiatric and neurodegenerative disorders. This analysis provides an overview on what progress has been made in understanding the roles of synaptic functioning in these disorders and reveals the gaps in knowledge in this field.

2. Structure and biological functions of neurexins

Neurexins were first identified using affinity chromatography when neurexin 1 α was found in rat brain extract on a column of α -latrotoxin [11]. α -latrotoxin is a potent neurotoxin from black widow spider venom that stimulates synaptic vesicle exocytosis and induces massive neurotransmitter release [12]. This work has been continued by Südhof and co-workers [13] who have characterized the neurexin proteins and their binding partners, the neuroligins [14].

In mammals, the neurexins are encoded by three *NRXN* genes (*NRXN1-3*), each of which has both an upstream promoter that is used to generate the α -neurexins, and a downstream promoter that is used to generate the shorter β -neurexins [13,15]. Neurexins also undergo extensive alternate splicing




Figure 1. Location of neurexins and their binding partners, neuroligins, in the synapse. Several neurexin–neuroligin pathway proteins are shown as well as synaptic vesicle-binding proteins. NMDAR, *N*-methyl-D-aspartate receptor; mGluR5, metabolic glutamate receptor 5; PSD-95, post-synaptic density protein 95; Shank, SH3 and multiple ankyrin repeat domains protein.




Figure 2. Structural domain organization of the α and β forms of neurexin. (a) α -neurexin. (b) β -neurexin. EGF, epidermal growth factor-like region; LNS, laminin/neurexin/sex hormone-binding domain; PDZ, PSD-95, DLG1, ZO-1 domain; SP, signal peptide; TM, transmembrane domain.

at five splice sites, generating significant diversity of more than 2000 potential variants [13,16]. The fact that neurexin splice insert sequences and their positions are well conserved among neurexin genes and between species supports the idea that alternative splicing has important functional roles.

The neurexins are transmembrane proteins that consist of an extracellular region responsible for trans-synaptic interactions, a transmembrane domain and a smaller cytoplasmic domain named PSD-95, DLG1, ZO-1 binding domain (PDZ) that is involved in intracellular protein interactions and signalling (figure 2) [13]. α -neurexins are composed of six large extracellular laminin/neurexin/sex




Figure 3. A summary of GO terms associated with neurexins 1–3. (a) Biological processes. (b) Cellular components. (c) Molecular functions. All: total number of proteins analysed. The number above each bar indicates the number of proteins assigned to that category. Figure generated by WebGestalt (<http://www.webgestalt.org>) [18].

hormone-binding (LNS) globulin domains with three interspersed epidermal growth factor (EGF)-like regions (figure 2). β -neurexins are shorter and only have the sixth extracellular LNS domain and no EGF-like regions (figure 2). Only neurexin 1 protein structures (both α and β forms) have been solved experimentally in *Mus musculus*, *Rattus norvegicus* and *Bos taurus*. However, these structures have not yet been solved in humans.

Neurexins are localized pre-synaptically and are distributed to both excitatory and inhibitory synapses [8]. Their functions are mediated by their binding to neuroligins (figure 1). Neuroligins have five known isoforms and are expressed post-synaptically [17]. Consequently, neurexins and neuroligins form synaptic complexes in the synaptic cleft and have been found to control synapse formation, maturation, validation and function [17]. Various combinations of the different neurexins and neuroligin binding partners at synapses may also help determine synapse specificity through differential interactions between multiple splice variants and isoforms of these proteins [8].

Primarily, neurexins function to maintain synaptic organization. Gene ontology (GO) analysis by WebGestalt (<http://www.webgestalt.org>) [18] of the three neurexins indicates that they all function in protein binding, ion binding and possess molecular transducer activity (figure 3). They are also involved in cellular component organization, developmental processes, response to stimuli, cell communication and biological regulation. These processes thus demonstrate how neurexins are able to maintain synaptic organization but also show their multi-functional nature. As such, it is conceivable that disruptions in neurexins could be detrimental to their various functions and affect overall neuronal function and integrity.

3. Biological pathways and interacting partners of neurexins

To understand the broader biological pathways that the neurexins are involved in, protein–protein interaction network

analysis and co-expression analysis was performed using the tools, STRING (<https://string-db.org>) [19] and GeneMania (<https://genemania.org>) [20]. STRING finds related genes by accessing the STRING database which contains experimental data and computational predictions. Data in STRING are weighted and integrated and a confidence score of 0–1 is calculated for all interacting protein partners. GeneMANIA finds proteins related to neurexins by leveraging functional association data, such as interactions, pathways, co-expression, co-localization and protein domain similarity. All functional data for the proteins observed in these networks were obtained from UniProt (<https://www.uniprot.org>) [21], unless otherwise stated, while pathway data were obtained from KEGG (<https://www.kegg.jp>) [22].

3.1. String analysis

Weighted string analysis was conducted on neurexin 1, 2 and 3 individually to determine their binding partners (figure 4a–c). Based on this analysis, there is strong evidence that neurexin 1 interacts with 10 proteins including calcium/calmodulin-dependent serine protein kinase (CASK), leucine-rich repeat transmembrane neuronal protein 1 (LRRTM1), LRRTM2, LRRTM3, neuroligin 1, neuroligin 2, neuroligin 3, neuroligin 4X, SH3 and multiple ankyrin repeat domains protein 2 (SHANK2) and synaptotagmin-1 with scores of 0.987, 0.983, 0.985, 0.975, 0.998, 0.997, 0.998, 0.997, 0.975, 0.974, respectively. Similarly, neurexins 2 and 3 also have 10 interactors each. There is strong evidence that neurexin 2 interacts with CASK, discs large homologue 4 (DLG4), LRRTM1, LRRTM2, LRRTM3, neuroligin 1, neuroligin 2, neuroligin 3, neuroligin 4X and SHANK2 with scores of 0.979, 0.977, 0.984, 0.983, 0.972, 0.998, 0.998, 0.998, 0.997 and 0.969, respectively. There is strong evidence that neurexin 3 interacts with CASK, DLG4, LRRTM1, LRRTM2, LRRTM3, neuroligin 1, neuroligin 2, neuroligin 3, neuroligin 4X and SHANK2 with scores of 0.978, 0.971, 0.979, 0.983, 0.971, 0.998, 0.997, 0.997, 0.997 and 0.969, respectively. The STRING analyses performed on the three neurexins identified




Figure 4. Protein interacting partners of neurexins 1–3. (a–c) STRING network of neurexins 1–3. Nodes represent gene-encoded proteins. Edges represent protein–protein associations. Connections between nodes represent the relationship between proteins. A bold line implies a higher confidence level. (d) GeneMANIA network of all three neurexins. Nodes represent gene-encoded proteins. Larger nodes indicate higher protein scores. Edges represent protein–protein associations. Connections between nodes represent the relationship between proteins. A bold line implies a higher confidence level.

interacting proteins with very high confidence scores since the lowest score across the analyses was 0.969. This means that there is strong experimental evidence that these proteins interact with one or more of the neurexins.

Analysis of the identified neurexin binding partners revealed many proteins important in the maintenance and functioning of synapses. Notably, variants in several of these proteins are implicated in neuropsychiatric and developmental disorders. Variants in neuroligin 1 and SHANK2 have been implicated in susceptibility to autism [23,24], while variants in neuroligin 3 have been implicated in Asperger syndrome and autism [25]. Variants in neuroligin 4X have been implicated in X-linked forms of Asperger syndrome, autism susceptibility and mental retardation [25–27]. Variants in CASK have been implicated in FG syndrome 4, an X-linked genetic disorder and mental retardation [28–32], variants in DLG4 have been implicated in intellectual developmental disorder 62 [33,34] and variants in synaptotagmin-1 have been implicated in Baker–Gordon syndrome [35].

Furthermore, binding partners of these proteins as well as the pathways they occur in could also give insight into the development of disease. CASK binds to amyloid precursor protein and neuroligin 1 binds to amyloid- β , both of which are important in AD. LRRTM3 is also a known positive regulator of amyloid- β formation. Notably, *LRRTM3* may be considered a candidate gene for late-onset AD as it promotes the processing of amyloid precursor protein which leads to toxic amyloid- β accumulation [36]. DLG4 is involved in dopamine receptor binding and synaptotagmin-1 regulates dopamine secretion. The loss of dopamine functioning is crucial in PD. Indeed, DLG4 is involved in several pathways of

neurodegeneration (in multiple diseases), the HD pathway as well as cocaine addiction.

3.2. GeneMANIA

GeneMANIA analysis was performed on the neurexins to reveal further potential protein–protein interactions (figure 4d; electronic supplementary material, table S1). We performed the analysis by selecting only proteins with stronger evidence of neurexin interactions, such as interactions with physical evidence, and evidence from co-expression and co-localization studies.

All of the binding partners observed by STRING analysis were still present; however, more interacting proteins were also identified. These proteins have more diverse functions but still function in overall synapse maintenance.

This analysis further identified afadin (AFDN), Rho GTPase activating protein 10 (ARHGAP10), cerebellin 1 (CBLN1), dystroglycan (DAG1), microtubule actin cross-linking factor 1 (MACF1), neurexophilin-2, neurexophilin-3, PDZ domain-containing protein 2 (PDZD2), proteolipid protein 1 (PLP1), syndecan binding protein 1 (SDCBP), SDCBP2, SH3 domain-containing GRB2-like protein 2 (SH3GL2), signal-induced proliferation-associated 1-like protein 1 (SIPA1L1), synaptotagmin-13 (SYT13) Tafa chemokine-like family member 1 (TAF1), TUBB-like protein 1 (TULP1) and XK-related protein 4 (XKR4) as interactors of one or more neurexin proteins. AFDN, ARHGAP10, MACF1 and SIPA1L1 are all involved in actin filament binding/organization, while PDZD2, SDCBP and SDCBP2 are involved in cell binding and cytoskeletal organization.

Dysregulation of any of these proteins could thus affect cell adhesion and binding at the synapse. In addition, variants in *MACF1* have been implicated in lissencephaly 9 with complex brainstem malformation [37]. *CBLN1* is essential for synapse integrity and plasticity and its disruption could lead to synapse dysfunction. *DAG1* has multiple functions, such as laminin and basement membrane assembly, cell survival, peripheral nerve myelination, nodal structure and cell migration. Variants in *DAG1* have been implicated in both type A and type C muscular dystrophy–dystroglycanopathy [38–41]. Muscular dystrophies are genetic disorders characterized by the degeneration of skeletal muscle. Type C muscular dystrophy–dystroglycanopathy affects the limb-girdle area [40], while type C is congenital with brain and eye anomalies [39]. Neurexophilin-2 and neurexophilin-3 are both ligands for α -neurexins and are involved in the neuropeptide signalling pathway. Disruption of these proteins could, therefore, affect neurotransmitter release and the subsequent signalling. *PLP1* is the major myelin protein in the CNS and is important for maintaining the structure of myelin. Disruption of this protein could, therefore, negatively affect the downstream myelination of neurons, as is seen in multiple sclerosis (MS). Interestingly, *PLP1* is also involved in the development of the substantia nigra, the main brain region affected by PD. Therefore, *PLP1* alterations could also lead to disruptions in this brain region. *SH3GL2* has been implicated in synaptic vesicle endocytosis, while synaptotagmin-13 may be involved in transport vesicle docking to the plasma membrane. Dysregulation of these proteins could thus affect neurotransmitter functioning. *TAF1* is involved in the modulation of neural stem cell proliferation and differentiation; therefore, dysregulation of this protein could result in developmental disorders. *TULP1* is required for normal development of photoreceptor synapses. Variants in *TULP1* are associated with Leber congenital amaurosis [42,43] and retinitis pigmentosa [42,44–47]. However, this protein is also involved in actin filament binding, therefore, its dysregulation could also affect cell adhesion and binding at the synapse. Not much is known about *XKR4* except that it is involved in apoptosis during development. Therefore, its dysregulation could also possibly result in developmental disorders.

GO terms and physiological/pathway information on all binding partners identified by STRING and GeneMANIA are available in electronic supplementary material, figure S1 and table S1.

4. Role of neurexins in neuropsychiatric disorders

Literature-based searches using neurexin as a search term identified several studies that reported an association of neurexins in various neuropsychiatric disorders. The main findings of these studies are reported in table 1 and are summarized below.

4.1. Human studies

NRXN1 has been well documented for its association with ASDs [54]. Several genetic analyses of families and populations of people with ASD have shown that copy number variations (CNVs) and de novo mutational events at the *NRXN1* locus are enriched in ASD [48,49,51,52,54]. In one

study, *NRXN1* was sequenced in cases of ASD with mental retardation [50]. Mutations (c.–3G > T in the Kozak region, c.3G > T at the initiation codon (p.M1), p.R375Q and p.G378S) were found in the *NRXN1 β* coding region thereby potentially implicate synapse dysfunction an important determinant in ASD [50].

The first evidence for a potential role of *NRXN2* in ASD was provided by a report of a frameshift mutation within *NRXN2* exon 12 (c.2733delT) in a boy with ASD and his father who had severe language delay [57]. This mutation results in a truncated neurexin 2 α protein that lacks the binding sites for the established post-synaptic binding partners *LRRTM2* and *neuroligin 2* [57]. Subsequently, a 21-year-old man with a clinical phenotype including autistic traits, such as speech and language deficits and pathological insistence on routine, was reported to have a 570 kb de novo deletion of 24 genes at chromosome 11q13.1, including *NRXN2* [58].

Using microarray analyses on RNA extracted from brain tissue, *Mirnic et al.* [69] did not observe a difference in neurexin 1 expression between schizophrenia (SCZ) and control samples. However, since then, a link between neurexin 1 and SCZ has been reported in other studies. One study reported that *NRXN1* deletions are more common in those with SCZ; however, it also found that there was incomplete penetrance of these deletions in families with SCZ [72]. *Kirov et al.* [70] observed a deletion in an SCZ patient at 2p16.3 that disrupts *NRXN1* and predicted that it was highly likely to be pathogenic. Also, *NRXN1* deletions were shown to segregate with several neuropsychiatric disorders in a study of a complex family [78]. The proband had SCZ and other members of his family had mental retardation, schizophreniform disorder and affective disorder [78]. After genotyping the proband and eight family members, they found two rare deletions upstream of the *NRXN1* gene (2p16.3) that co-segregate with these disorders [78]. Notably, this shows that deletions in *NRXN1* may manifest as multiple neuropsychiatric phenotypes.

Angione et al. [64] implicated *NRXN1* in epilepsy. They identified a 2p16.3 deletion in an 8-year-old male patient diagnosed with epilepsy showing symptoms of myoclonic-astonic seizures (EMAS) [64]. This deletion included the first five exons of the *NRXN1* gene [64].

NRXN genes may also be involved in treatment response. In one study, it was found that variants in *NRXN1* may affect the long-term treatment outcome of patients with BD by modulating the effects of antipsychotics [61]. In a study of *Levetiracetam* resistance, an antiepileptic drug, *Grimminger et al.* [63] found that neurexin 1 is differentially expressed in non-responder and responder patients with mesial temporal lobe epilepsy (mTLE), whereby lower levels of neurexin 1 were observed in responder patients.

4.1.1. Association studies

A genome-wide association study (GWAS) by *Liu et al.* [51] specifically examined *NRXN1* in an autism cohort of the Chinese Han population and discovered 22 variants that were associated with ASD. In this cohort, one SNP (rs2303298) was also significantly associated with a risk of developing ASD [51]. Furthermore, a GWAS of SCZ in Spain showed that a *NRXN1* single nucleotide polymorphism (SNP) (rs3850333) was close to the significance threshold [71], while another GWAS in American patients of European or African ancestry showed that *NRXN1* is associated with antipsychotic

Table 1. List of studies that have implicated neurexin genes in neuropsychiatric disorders. AAV, adeno-associated virus; AGRE, Autism Genetic Resource Exchange; Array-CGH, array comparative genomic hybridization; CBDB, Clinical Brain Disorders Branch; CIBERSAM, Centro de Investigación Biomédica en Red de Salud Mental; CNV, copy number variation; EMAS, epilepsy with myoclonic-atonic seizures; GWAS, genome-wide association study; hESC, human embryonic stem cell; iN, induced neuron; iPSC, induced pluripotent stem cell; KO, knockout; LC-MS/MS, liquid chromatography mass spectrometry/mass spectrometry; mESC, mouse embryonic stem cell; mTLE, mesial temporal lobe epilepsy; NGS, next-generation sequencing; NIMH, National Institute of Mental Health; RT-PCR, reverse transcriptase–polymerase chain reaction; SNP, single nucleotide polymorphism; SSC, Simons Simplex Collection; STEP-BD, Systematic Treatment Enhancement Program for Bipolar Disorder; WT, wild-type.

disorder/disease process	neurexin gene	type of study	methods	main finding	reference
autism spectrum disorder (ASD)	NRXN1	genetic analysis	used SSC samples and the SSC database to extract 'trios' consisting of a mother, father and an ASD-affected child	a de novo CNV in <i>NRXN1</i> was discovered in a large cohort of families with a single ASD-affected child and at least one unaffected sibling	[48]
ASD	NRXN1	GWAS	performed genetic analyses to identify CNVs 1174 families from SSC were genotyped identified CNVs and de novo events	rare de novo events/CNVs at <i>NRXN1</i> are strongly associated with autism	[49]
ASD	NRXN1 β	genetic analysis	86 patients with both ASD and mental retardation The coding sequence of the <i>NRXN1β</i> gene was analysed by PCR	four novel mutations in <i>NRXN1β</i> were identified by sequencing the exon of the gene in cases with autism and mental retardation	[50]
ASD	NRXN1	genetic analysis	313 ASD patients and 500 healthy controls from a Chinese autism cohort were recruited performed genomic DNA sequencing	22 variants in the <i>NRXN1</i> gene were discovered in the Chinese Han population; one SNP (rs2303298) was significantly associated with a risk of autism in this cohort	[51]
ASD	NRXN1	genetic analysis	2478 ASD individuals from SSC and 719 ASD individuals from AGRE 580 controls from ClinSeq and NIMH used a custom microarray to analyse CNVs	recurrent CNVs in <i>NRXN1</i> are enriched in autism.	[52]
ASD	NRXN1, 2 and 3	cell culture	iPSCs were produced from probands and unaffected family members iPSCs underwent neuronal differentiation to organoids RNA sequencing was performed on both iPSCs and differentiated organoids	neurexin 1, 2 and 3 mRNA is overexpressed in patient-derived iPSCs and differentiated organoids	[53]
ASD	NRXN1	genetic analysis	2591 families from SSC were genotyped identified CNVs, de novo deletions and ASD risk genes	<i>NRXN1</i> is an ASD risk gene	[54]
ASD	NRXN1 and 2	animal study	RNA was isolated from the whole brain of age-matched monoamine oxidase A KO mice and wild-type mice Microarrays were used to determine gene expression levels	neurexin 1 and 2 are downregulated in monoamine oxidase A KO mice	[55]
ASD	α -NRXNs	animal and cell culture study	transfected <i>C. elegans</i> strains and HEK-293 cells with plasmids expressing NRXN1 α and different α 2 δ subunits performed co-immunoprecipitation and pull-down assays	changes in α -neurexin binding to α 2 δ -3 subunits of N-type calcium channels could be important in some forms of autism spectrum disorders	[56]
ASD	NRXN2	genetic analysis	142 ASD patients and 94 non-syndromic controls sequenced NRXN 1,2 and 3 genes	observed a frameshift mutation in <i>NRXN2</i> exon 12 in a patient with ASD inherited from a father with severe language delay	[57]

(Continued.)

Table 1. (Continued.)

disorder/disease process	neurexin gene	type of study	methods	main finding	reference
ASD	NRXN2	genetic analysis	recruited a patient with speech problems, autistic traits and pancreatic gastrinoma performed array-CGH	a de novo 0.57 Mb microdeletion was observed in chromosome 11q13.1, including NRXN2	[58]
ASD	NRXN2	animal study	used previously collected human faecal samples from typically developing children and children with ASD	mice colonized by microbiota from ASD patients showed differential splicing of NRXN2	[59]
bipolar disorder (BD)	NRXN3	GWAS	C57BL/6 J weanlings were colonized with human faecal samples brain tissue RNA was extracted and sequenced	NRXN3 shows a significant association with bipolar disorder	[60]
BD	NRXN1	genetic analysis	obtained participants from a family study of mood disorders in Taiwan (2008–2012) performed a multi-stage GWAS	NRXN1 may affect the long-term treatment outcome of bipolar disorder	[61]
borderline personality disorder (BPD)	NRXN3	association study	obtained patient genotyping and clinical data from STEP-BD analysed data to determine the effect of individual markers on phenotypes	several NRXN3 SNPs were nominally associated with BPD phenotype in heroin-dependent cases	[62]
epilepsy	NRXN1	microarray analysis	obtained 53 biopsy specimens from mTLE patients	neurexin 1 is differentially expressed in non-responder and responder mTLE patients to the antiepileptic drug Levetiracetam	[63]
epilepsy	NRXN1	genetic testing	performed microarray analysis	lower levels of neurexin 1 are observed in responder patients	[64]
epilepsy/seizures	NRXN2 α	animal study	77 patients were identified at Children's Hospital Colorado various genetic tests were conducted	a 2p16.3 deletion, which includes the first five exons of the NRXN1 gene, was identified in an 8-year-old male EMAS patient	[65]
fragile X syndrome	NRXN3	animal study	treated adult Wistar rats with kainite or pentylenetetrazole to induce seizures	following kainate- and pentylenetetrazole-induced seizures in rats, neurexin 2 α expression increased in the dentate gyrus of the hippocampus	[66]
major depressive disorder (MDD)	NRXN1, 2 and 3	animal study	isolated total RNA from whole-rat brains and hippocampi performed RT-PCR to determine the levels of different NRXNs	there is increased neurexin 3 mRNA in female fragile X mice, but decreased neurexin 3 mRNA in male fragile X mice	[67]
			used male and female WT and FMR1 KO mice (4–6 per experiment) analysed brain sections using riboprobes for NRXN1, 2 and 3 and NLGN 1, 2 and 3	neurexin 1, 2 and 3 were not differentially expressed in a rat chronic mild stress model of depression	
			81 healthy Sprague-Dawley rats were subjected to various mild stress factors		
			extracted proteins from hippocampal post-synaptic density fractions analysis by LC-MS/MS		

(Continued.)

Table 1. (Continued.)

disorder/disease process	neurexin gene	type of study	methods	main finding	reference
neuropsychiatric disorders	NRXN1	cell culture	cultured cortical neurons from NRXN1 α KO mice introduced conditional NRXN1 mutations into hESCs using AAV recombination, and differentiated them into human iNs analysed neuronal development, synapse formation and neurotransmitter release	heterozygous NRXN1 mutations are able to selectively impair neurotransmitter release and increase the levels of the synaptic scaffolding protein, CASK in human iNs but not in the cortical neurons of NRXN1 α KO mice	[68]
schizophrenia (SCZ)	NRXN1	microarray analysis	obtained brain tissue from 12 SCZ patients and 10 controls extracted total RNA performed a microarray analysis	the expression of neurexin 1 was not significantly different between the schizophrenic and control subjects	[69]
SCZ	NRXN1	genetic analysis	selected 45 male and 48 female proband-parent trios from a sample of 600 Bulgarian SCZ trios performed Array-CGH	observed a 0.25 Mb deletion on 2p16.3 in both the proband and affected sibling which disrupts NRXN1	[70]
SCZ	NRXN1	GWAS	3063 SCZ patients and 2847 controls from CIBERSAM performed a GWAS at 95 SNPs	the rs3850333 SNP in the NRXN1 gene was close to the significant threshold in a GWAS of schizophrenia in Spain	[71]
SCZ	NRXN1	genetic analysis	obtained DNA of 635 SCZ patients and 635 controls from the CBDB Sibling Study	NRXN1 deletions are more frequent in schizophrenia patients	[72]
SCZ	NRXN1	genetic analysis	genotyped samples and analysed them for CNVs and deletions data from 572 SCZ patients and 551 controls were used to select genes for sequencing 153 SCZ patients and 153 controls were sequenced for 21 chosen genes using NGS	there is incomplete penetrance of NRXN1 deletions in families with schizophrenia missense variants at NRXN1 may be protective against schizophrenia susceptibility	[73]
SCZ	NRXN1	cell culture	isolated primary rat neurons from hippocampi differentiated human neurons derived from human iPSCs overexpressed Caveolin-1 in both cell types western blotting was used to measure the expression of proteins involved in synaptic plasticity as well as DISC1, an SCZ-associated protein	overexpressing Caveolin-1, a potential therapeutic for schizophrenia, in neurons increased expression of proteins involved in synaptic plasticity (PSD95, synaptobrevin, synaptophysin, neurexin 1 and syntaxin 1) as well as DISC1	[74]

(Continued.)

Table 1. (Continued.)

disorder/disease process	neurexin gene	type of study	methods	main finding	reference
SCZ	NRXN1	animal study	generated iPSCs from 5 childhood-onset SCZ patients and 4 controls differentiated iPSCs into glial cells transplanted glial cells into mice via injection into the corpus colosum performed molecular analyses on both the differentiated glial cells and chimeric mice	neurexin 1 was downregulated in chimeric mice produced from iPSCs derived from patients with childhood-onset schizophrenia	[75]
SCZ	NRXN1	GWAS	obtained genetic data and treatment response data of 302 SCZ patients treated with lurasidone and 117 SCZ patients treated with a placebo from two clinical SCZ trials performed a GWAS	NRXN1 is associated with antipsychotic response to lurasidone in schizophrenia patients	[76]
SCZ	NRXN1	cell culture	generated iPSCs from 3 NRXN1 deletion SCZ patients and 3 controls and differentiated them into human iNs generated mESCs from NRXN1 KO mice analysed neuronal development, synapse formation and neurotransmitter release	heterozygous NRXN1 deletions impair neurotransmitter release and synaptic function, and increase the levels of the synaptic scaffolding protein, CASK in human iNs but not mESCs generated from NRXN1 KO mice	[77]
SCZ and other neuropsychiatric disorders	NRXN1	genetic analysis	recruited a family with multiple neuropsychiatric disorders the proband has SCZ, while other family members have mental retardation, schizophreniform disorder and affective disorder genotyped the proband and eight family members	two rare deletions upstream of the NRXN1 gene (2p16.3) segregate with schizophrenia, schizophreniform disorder, and affective disorder in a family	[78]

response to lurasidone in SCZ patients [76]. Additionally, an association study on Spanish SCZ patients showed that missense mutations in *NRXN1* may actually protect against susceptibility to SCZ [73].

In a Taiwanese GWAS study, a significant association between *NRXN3* and BD was found [60]. And finally, an association study on Australian borderline personality disorder (BPD) patients showed that several *NRXN3* SNPs were nominally associated with BPD in heroin-dependent cases [62].

4.2. *In vitro* and *in vivo* models of disease

Functional *in vitro* and *in vivo* studies have also found evidence for the roles of neurexins in ASD. Monoamine oxidase A knockout (KO) mice, which are an animal model for autism, exhibited downregulated levels of both neurexin 1 and neurexin 2 [55]. Furthermore, mice colonized with the microbiota of ASD patients showed differential splicing of *NRXN2* [59]. Another animal study showed that changes in the binding of α -neurexins to N-type calcium channels could be important for some forms of ASD as it mediates synaptic inhibition [56]. Finally, a study using ASD patient-derived induced pluripotent stem cells (iPSCs) and differentiated organoids showed that neurexin 1, 2 and 3 mRNA is overexpressed in these samples [53].

One study examined neurexins in Fragile X syndrome, a genetic disorder with features similar to ASD, and characterized by the silencing of the *FMR1* gene [79]. Individuals with Fragile X experience a range of neurodevelopmental problems, such as learning disabilities and cognitive impairment, and males are usually more severely affected. Using *FMR1* KO mice, researchers probed brain sections to determine the levels of neurexin 1, 2 and 3 [66]. Interestingly, they found that neurexin 3 mRNA levels are increased in female mice but decreased in male mice and predicted that this may help explain the sex difference observed in this disorder [66].

In an animal study of SCZ, neurexin 1 was found to be downregulated [75]. This study generated iPSCs from patients with childhood-onset SCZ, differentiated them into glial cells and injected the glial cells into mice to form chimeric mice as a model organism [75]. Interestingly, an *in vitro* study of SCZ showed that overexpressing Calveolin-1, a potential therapeutic for SCZ, actually increased the levels of neurexin 1 as well as other proteins involved in synaptic plasticity [74].

Neurexin 2 α has been implicated in epilepsy and, more specifically, in seizures. Researchers observed an increase in neurexin 2 α expression in the dentate gyrus of the hippocampus in an induced-seizure mouse model [65]. Finally, in one study, a rat chronic mild stress model of depression was used to determine if neurexin expression was altered in major depressive disorder; however, no change in neurexin 1, 2 or 3 levels was observed [67].

So far, there have been two studies validating the effect of NRXNs *in vitro*, both by Pak *et al.* [68,77]. These studies cultured human stem cells as well as mice cells generated from *NRXN1* KO mice. The first study introduced two conditional *NRXN1* mutations previously seen in a range of neuropsychiatric disorders, including ASD and SCZ, into human embryonic stem cells (hESCs) using adeno-associated virus recombination and differentiated them into human-induced neurons (iNs) [68]. These cells were compared to cortical

neurons generated from *NRXN1 α KO mice [68]. The second study generated iPSCs from three *NRXN1* deletion SCZ patients and three controls, and again differentiated them into human iNs [77]. These cells were compared to mouse embryonic stem cells (mESCs) from *NRXN1* KO mice [77]. Both studies showed that heterozygous *NRXN1* deletions were able to impair neurotransmitter release and synaptic function, and increase the levels of the synaptic scaffolding protein, CASK, in human iNs but not in mice cells [68,77]. Therefore, these studies provide evidence that *NRXN1* deletions exhibit a major synaptic transmission phenotype in humans and are thus meaningful at a pathophysiological level.*

In summary, these studies demonstrate a link between *NRXNs* and neuropsychiatric disorders such as ASD and SCZ, especially involving full or partial deletions of these genes. *NRXNs* have also been associated with BD and BPD. In addition, protein expression studies have shown changes in neurexin expression in animal models of epilepsy/seizures and Fragile X syndrome.

5. Role of neurexins in neurodegenerative disorders

Additionally, literature-based searches provided proof for the involvement of neurexins in various neurodegenerative disorders, and these studies are listed in table 2 and discussed below.

5.1. Human studies

Studies examining cerebrospinal fluid (CSF) from AD patients have observed lowered expression of neurexin 1 [81], as well as neurexin 2 α and neurexin 3 α [85]. In addition, it was found that these changes precede the neurodegeneration markers as they were observed in the preclinical stage 1 of AD [85]. Moreover, A β ₄₂ fibrils in CSF were found to bind to neurexin 1, 2 and 3 as well as proteoglycans and growth factors [83]. Levels of the synaptic proteins neuronal pentraxin 2 (NPTX2), GluA4-containing glutamate (AMPA4), neuroligin 1 and neurexin 2 α are also declined in plasma neuron-derived exomes and this decline was associated with AD progression [82]. Neurexin 3 protein expression has also been seen to be specifically downregulated in blood samples of AD patients [84].

Another expression analysis on CSF from MS patients identified neurexin 2 α levels as a potential biomarker for the disorder [98], while a genetic analysis found that a mutant miRNA, MIR8485, overexpresses neurexin, which leads to a calcium overload in pre-synapses [99]. It was hypothesized that this could induce neurodegeneration in MS [99].

A study examining gene expression in brain tissue samples of patients with PD found that genes related to nerve function, such as protocadherin-8 (PCDH8) and neurexin 3, were downregulated [109].

Two studies on mild cognitive impairment (MCI) found promising results. MCI is a milder form of dementia that is considered the intermediate state of cognitive decline between normal ageing and dementia [114]. Berchtold *et al.* [95] found that neurexin 1 and neurexin 2 are upregulated in MCI. In addition, neurexin 1 expression was found to be associated with longitudinal phenotypes in MCI, but not in AD [96].

One study examined neurexins in order to identify genes that are differentially regulated by HIV encephalitis [94]. This

Table 2. List of studies that have implicated neurexin genes in neurodegenerative disorders and ageing. 6-OHDA, 6-hydroxydopamine; ACP-RT-PCR, annealing control primer reverse transcriptase-polymerase chain reaction; ADNI, Alzheimer's disease neuroimaging initiative; AMPA4, GluA4-containing glutamate; CSF, cerebrospinal fluid; EAE, experimental autoimmune encephalomyelitis; ELISA, enzyme-linked immunosorbent assay; FTD-GWAS, frontotemporal dementia genome-wide association study; GEO, gene expression omnibus; GWAS, genome-wide association study; HIV, human immunodeficiency virus; HYPERGENES, European Network for Genetic-Epidemiological Studies; LC-MS/MS, liquid chromatography mass spectrometry; LC-SRM, liquid chromatography single reaction monitoring; MAP, Rush Memory and Ageing Project; MR, magnetic resonance; MRI, magnetic resonance imaging; NPTX2, neuronal pentaxin 2; ONIND, other non-inflammatory neurological disease; PCDH8, protocadherin-8; PPM1, Parkinson's Progression Markers Initiative; qRT-PCR, quantitative real-time PCR; RAP-PCR, reverse arbitrarily primed PCR; rMOG, rat myelin oligodendrocyte glycoprotein; RRMS, relapsing-remitting MS; RT-PCR, reverse transcriptase-PCR; SNP, single nucleotide polymorphism; UV-CLIP, ultraviolet cross-linking and immunoprecipitation; WES, whole-exome sequencing.

disorder/disease process	neurexin gene	type of study	methods	main finding	reference
Alzheimer's disease (AD)	NRXN3	GWAS of brain structure	obtained neuroimaging and genetic data from 818 subjects as part of ADNI performed a GWAS with 546,314 SNPs using temporal lobe and hippocampal volume as quantitative phenotypes	NRXN3 (<i>KIAA0743</i>) is associated with temporal lobe structure in AD patients	[80]
AD	NRXN1	protein expression analysis	collected CSF samples from 10 AD patients and 15 healthy controls analysis using LC-MS/MS	the concentrations of the synaptic proteins neurexin 1 and neuronal PTX1, as well as neurofascin, were significantly lowered in AD CSF	[81]
AD	NRXN2 α	protein expression analysis	collected blood and CSF samples from 28 AD patients and 28 controls extracted plasma neuron-derived exomes CD81, NPTX2, AMPA4, neuroligin 1 and neurexin 2 α proteins were quantified using ELISAs	significantly lower levels of the synaptic proteins NPTX2, AMPA4, neuroligin 1 and neurexin 2 α in the plasma neuron-derived exomes correlate with AD progression	[82]
AD	NRXN1, 2 and 3	protein expression analysis	collected CSF samples from six AD patients and five non-AD patients binding assays were performed to determine which proteins in CSF bind to A β ₄₂ fibrils and/or protofibrils	A β ₄₂ fibrils in AD CSF are involved in binding to proteoglycans, growth factors and neuron-associated proteins, such as neurexin 1, 2 and 3	[83]
AD	NRXN3	transcriptome and RNA expression analysis	selected data from 263 AD patients and 151 non-demented controls sampled from the religious orders study performed RNA expression profiling	neurexin 3 expression is downregulated in AD	[84]
AD	NRXN2 α and NRXN3 α	protein expression analysis	collected CSF samples from AD patients and cognitively normal controls (three stage study with different <i>n</i> for each stage) performed LC-MS/MS and LC-SRM	levels of neurexin 2 α and neurexin 3 α , as well as other synaptic proteins are decreased in preclinical AD CSF	[85]
AD and ageing	NRXN1, 2 and 3	microarray analysis	obtained frozen brain samples from 26 AD cases and 55 non-AD controls from National Institute on Ageing Alzheimer's disease brain banks used microarrays to evaluate expression profiles of 340 synaptic genes	SYNAPTIC proteins, including neurexin 1, 2 and 3, undergo altered expression in ageing and AD	[86]
AD and ageing	NRXN3	animal study	mice were divided into four groups, with four mice in each group: memory intact AD-transgenic mice, memory impaired AD-transgenic mice, memory intact aged mice and memory impaired aged mice performed proteomics on the hippocampus of each mouse	neurexin 3 is downregulated in AD-transgenic mice with impaired memory, but not in normal aged mice with impaired memory	[87]

(Continued.)

Table 2. (Continued.)

disorder/disease process	neurexin gene	type of study	methods	main finding	reference
AD and ageing	NRXN1 and 3	microarray analysis	performed a microarray analysis on 47 post-mortem brain samples from cognitively intact aged individuals from the MAP study identified 48 microarrays from the public GEO: 16 young cases, 18 cognitively intact aged cases and 14 AD cases analysed data to identify genes related to physical activity, ageing and AD cerebella were removed from three adult C57BL/6 J mice and three aged C57BL/6Jnia mice	neurexin 1 and 3 have decreased expression in ageing and AD but have increased expression in association with late-life physical activity	[88]
ageing	NRXN3	animal study	RNA was extracted and sequenced monocytes were purified from PBMCs analysis of methylation was performed on genomic DNA from monocytes	neurexin 3 is downregulated in the cerebellum of aged mice	[89]
ageing	NRXN2	methylation analysis	28 Swiss albino mice were divided into four groups by age: young, adult, middle age and old molecular techniques were used to analyse neurexin 1 and neuroigin 3 expression	CpG sites associated with <i>MRP1</i> , <i>NRXN2</i> and <i>miR-29b-2</i> are hypomethylated in monocytes during ageing	[90]
ageing	NRXN1	animal study	collected data of 2202 post-mortem human brain samples of neurologically healthy individuals with different ages Calculated signal expression of genes performed UV-CLIP experiments on SH-SY5Y cells to find TDP-43 targets validated these results on lumbar spinal cords from 4 ALS patients and 4 controls using RT-PCR	neurexin 1 and neuroigin 3 are differentially expressed in cerebral cortex and hippocampus during different stages of ageing, which might be responsible for alterations in synaptic plasticity during ageing	[91]
ageing	NRXN2 and 3	transcriptome analysis	received cortical brain tissue from 13 HIV patients: eight with HIV encephalitis and five without extracted total RNA performed microarray analysis	neurexin 2 and 3 are downregulated in ageing	[92]
amyotrophic lateral sclerosis (ALS)	NRXN1	cell culture and expression analysis	aged frozen brain samples from 16 MCI cases, 25 AD cases and 24 aged controls from National Institute on Aging Alzheimer's Disease brain banks extracted total RNA performed a microarray analysis	neurexin 1 and other TDP-43 targets are dysregulated in ALS	[93]
HIV encephalitis	NRXN1	microarray analysis	obtained frozen brain samples from 16 MCI cases, 25 AD cases and 24 aged controls from National Institute on Aging Alzheimer's Disease brain banks extracted total RNA performed a microarray analysis	neurexin 1 is downregulated in HIV encephalitis	[94]
mild cognitive impairment (MCI)	NRXN1 and 2	microarray analysis		neurexin 1 and 2 are upregulated in MCI	[95]

(Continued.)

Table 2. (Continued.)

disorder/disease process	neurexin gene	type of study	methods	main finding	reference
MCI and AD	NRXN1	association study	obtained brain MR images of 400 MCI subjects, 400 AD subjects and 200 aged controls from the ADNI database obtained genotype data for 510 of these subjects from the ADNI database performed an association study	neurexin 1 expression is associated with longitudinal phenotypes in MCI, but not in AD	[96]
multiple sclerosis (MS)	NRXN3	animal study	induced EAE in 17 rats by injecting rMOG six control rats were treated with saline extracted total RNA used a cDNA expression array	neurexin 3 is downregulated in grey matter of EAE-induced rats	[97]
MS	NRXN2 α	protein expression analysis	collected CSF samples from 37 RRMS patients, 50 patients with ONMIND and patients with non-neurological (orthopaedic) diseases analysis using LC-MS/MS	neurexin 2 α in CSF is a potential biomarker for MS	[98]
MS	NRXN1	genetic analysis	collected blood from a female patient with RRMS	overexpression of neurexin 1 by mutant MIR8485 leads to calcium overload	[99]
MS	NRXN1	cell culture	performed WES and screened for mutations treated THP-1 cells with ceramides to induce hypermethylation of DNA isolated genomic DNA	in pre-synapses. This could induce neurodegeneration in MS ceramide-induced hypermethylation of DNA was associated with decreased transcript levels of neurexin 1 in cultured human monocytes	[100]
neurodegeneration	NRXN3	animal study	measured levels of neurexin 1, FZD7 and TP63 using qRT-PCR 45 DA(RT1 ^{aw}) and 45 PVG(RT1 ^s) adult rats subjected rats to ventral root avulsion extracted total RNA used a cDNA expression assay and performed RT-PCR	neurexin 3 is downregulated in rats with axonal damage caused by ventral root avulsion	[101]
neurodegeneration	NRXN3	animal study	three experimental groups with five ICR mice each injected kainic acid into ICR mice extracted total RNA from the hippocampus performed ACP-RT-PCR and sequenced the PCR products	neurexin 3 is downregulated in the hippocampus of mice treated with kainic acid, an inducer of neurodegeneration	[102]
neurodegeneration	NRXN1	bioinformatics and cell culture	analysed cross-linking, immunoprecipitation and sequencing data from the ArrayExpress archive to identify RNAs bound to TDP-43 in human and mouse brains quantitative RT-PCR was used to measure mRNA expression	a novel TDP-43 binding miRNA, miR-NID1 (miR-8485), represses neurexin 1 expression and may play a role in neurodegeneration	[103]

(Continued.)

Table 2. (Continued.)

disorder/disease process	neurexin gene	type of study	methods	main finding	reference
neurodegeneration	NRXN1 β	cell culture	transfected rat hippocampal neurons to overexpress acetylcholinesterase performed a co-immunoprecipitation assay with neurexin 1 and acetylcholinesterase co-transfected HEK-293 cells to express neurexin 1 β and neuroligin 1 and cultured these cells in acetylcholinesterase conditioned media performed a co-immunoprecipitation assay with neurexin 1 β and neuroligin 1	excessive glycosylated acetylcholinesterase could competitively disrupt neurexin 1 β -neuroligin junctions and impair the integrity of glutamatergic synapses	[104]
neurotoxicity	NRXN3 β	animal study	groups of 3 Sprague–Dawley rats were treated with sarin via intramuscular injection rats were sacrificed 15 min or 3 months after sarin exposure dissected brains and extracted total RNA performed a microarray analysis	sarin exposure causes a persistent downregulation of neurexin 1 β and breakdown of the blood–brain barrier	[105]
neurotoxicity	NRXN2 α	animal study	wild-type zebrafish were repeatedly exposed to domoic acid via intracoelomic injection dissected brains and extracted total RNA performed a microarray analysis	neurexin 2 α was upregulated in zebrafish two weeks after exposure to domoic acid, a neurotoxin	[106]
Parkinson's disease (PD)	NRXN1	cell culture	cultured SH-SY5Y cells and primary mouse mesencephalic cells treated cells with 6-OHDA performed RAP-PCR and analysed the PCR products using RT-PCR and qRT-PCR	downregulation of neurexin 1 mRNA and protein was observed in the 6-OHDA-induced cell culture models of PD	[107]
PD	NRXN2	animal study	transgenic mice were assigned to 4 treatment groups with 20 mice per group cholesterol oximes were administered in food pellets	transgenic mice overexpressing α -synuclein have increased levels of neurexin 2 chronic administration of cholesterol oximes to these mice decreased neurexin 2 levels	[108]
PD	NRXN3	genetics analysis	TH+ neurons were isolated from the substantia nigra and subjected to a transcriptome analysis obtained genomic data of 29 PD samples and 18 controls from the GEO database analysed the data to identify disease-related genes and differential gene expression	genes related to nerve function, such as PCDH8 and neurexin 3, are downregulated in PD brain tissue samples	[109]

(Continued.)

Table 2. (Continued.)

disorder/disease process	neurexin gene	type of study	methods	main finding	reference
PD	NRXN1	animal study	adult Wistar rats were divided into five treatment groups, with 6–8 rats in each group experimental groups had 6-OHDA brain injections with or without different concentrations of allopregnanolone western blots were performed to evaluate the levels of the synaptic proteins PSD95 and neurexin 1 in the striatum	neurexin 1 is significantly decreased in the striatum of 6-OHDA-induced rats treatment with allopregnanolone attenuates this and other molecular changes	[110]
PD	NRXN1	RNA expression analysis	MRI data from 149 PD patients and 64 healthy controls were obtained from the PPMI database 17 genes of interest implicated in PD were selected for whole-brain expression analysis	neurexin 1 does not have an expression pattern that predicts regional atrophy in PD	[111]
PD	NRXN1	animal study	adult Wistar rats were divided into seven treatment groups, with seven rats in each group experimental groups had 6-OHDA brain injections with or without different concentrations of apelin-13 western blots were performed to evaluate the levels of the synaptic proteins PSD95, neurexin 1 and neuroligin in the striatum	neurexin 1 expression is decreased in the striatum of 6-OHDA-induced rats 6-OHDA rats treated with apelin-13 showed increased neurexin 1 expression in the striatum	[112]
spinal muscular atrophy (SMA)	NRXN2 α	animal study	used HB9-D3cpv/MN-transgenic zebrafish and <i>Snn-1</i> ^{-/-} / <i>SMN2</i> mice isolated total RNA from both models performed a microarray analyses and qRT-PCR	<i>SMN</i> -deficiency downregulates neurexin 2 α expression and alters its splicing in zebrafish and mouse models of SMA	[113]

microarray study showed that neurexin 1 is downregulated in HIV encephalitis.

Finally, González-Velasco *et al.* [92] showed that neurexin 2 and neurexin 3 mRNA levels are downregulated in ageing. Another study found that neurexin 1, 2 and 3 underwent altered expression in both AD and ageing [86]. A more recent study from the same group confirmed decreased expression of both neurexin 1 and neurexin 3 in AD and ageing [88]. Interestingly, they also found that late-life physical activity is associated with increased expression of these proteins [88].

5.1.1. Association study

A GWAS performed by Stein *et al.* [80] showed that the SNP rs7155434 within *NRXN3* is associated with temporal lobe structure in AD patients. Temporal lobe volume deficits are a known risk factor for AD; therefore, this study potentially implicates *NRXN3* with AD risk [80].

5.2. *In vitro* and *in vivo* models of disease

Several studies involving cell culture and/or rodent disease models have also shown differences in the expression of neurexin proteins. Three studies showed that neurexin 1 is downregulated in PD. One of these measured neurexin 1 mRNA in two 6-OHDA (6-hydroxydopamine)-induced cell culture models; one using human neuroblastoma (SH-SY5Y) cells and the other using primary mouse mesencephalic cells [107]. The other studies used a 6-OHDA-induced rat model of PD and both saw a decrease in neurexin 1 in the striatum [110,112]. In addition, these studies showed that treatment with apelin-13 [112] or allopregnanolone [110] is able to attenuate this change. Apelin-13 is an endogenous ligand for APJ [115] that has been investigated as a potential protective neuropeptide due to the role of the apelin-APJ system in neuronal survival [116], while allopregnanolone is a reduced metabolite of progesterone [117] and has reduced CSF levels in PD patients [118]. Freeze *et al.* [111], however, noted that the expression pattern of neurexin 1 does not predict regional atrophy in PD. This suggests that neurexin 1 is not a marker for PD; however, it does not exclude it as an important protein in PD pathogenesis. Another study in PD-transgenic mice overexpressing α -synuclein found that neurexin 2 expression was also upregulated [108]. In addition, chronic administration of cholesterol oximes was able to increase the transcription of cytoprotective genes and undo transcriptome alterations, including the alteration of neurexin 2 expression [108].

Two studies using induced models of MS implicated neu-rexins in this disorder. One study induced experimental autoimmune encephalomyelitis (EAE) in rats and observed downregulation of neurexin 3 [97]. This is a commonly used model that mimics certain aspects of MS. The other study used an *in vitro* model of MS, cultured human monocytes, and observed an association between ceramide-induced hypermethylation of DNA and neurexin 1 mRNA [100].

An animal study performed by Neuner *et al.* [87] showed that neurexin 3 is downregulated in AD-transgenic mice, but not in normal aged mice with impaired memory. However, Popesco *et al.* [89] found that neurexin 3 is downregulated in the cerebellum of aged mice. Another study found that levels of both neurexin 1 and neuroligin 3 are differentially expressed in cerebral cortex and hippocampus of mice and

that these expression levels change during different stages of ageing [91]. They predicted that this may be responsible for the changes in synaptic plasticity observed with age [91]. Finally, a DNA methylation study by Tserel *et al.* [90] showed that CpG sites associated with *NRP1*, *NRXN2* and miR-29b-2 are hypomethylated in monocytes during ageing.

To date, only one study has examined neu-rexins in amyotrophic lateral sclerosis (ALS) and spinal muscular atrophy (SMA). In a cell culture model of ALS, neurexin 1 and other RNA targets of TDP-43 were dysregulated [93]. TDP-43 is a component of the cytoplasmic inclusion bodies present in ALS patients [93]. Fragments of TDP-43 are ubiquitinated, hyperphosphorylated and then accumulate in neurons and glia [119]. In zebrafish and mouse KO models of SMA, the *SMN*-deficiency downregulated neurexin 2 α expression and altered its splicing [113]. SMA is associated with mutation or deletions in the *SMN* gene [120] and lack of the *SMN* protein causes degeneration and results in anterior horn cell dysfunction.

5.3. Models of induced neurodegeneration and toxicity

Several studies investigated neu-rexins in models of neurodegeneration or toxicity instead of studying a specific neurodegenerative disease.

Four studies examined the role of neu-rexins in models of induced neurodegeneration. Two of these studies hypothesized that neurexin 1 could play a role in neurodegeneration. The first study showed that a novel TDP-43 binding miRNA, miR-NID1 (miR-8485) is able to repress neurexin 1 and predicted that this could play a role in neurodegeneration [103]. Xiang *et al.* [104] found *in vitro* that excessive glycosylated acetylcholinesterase could competitively disrupt the neurexin 1 β -neuroligin junctions and impair the integrity of glutamatergic synapses, which could lead to neurodegeneration. The other two studies showed that neurexin 3 is downregulated in animal models of neurodegeneration [101,102]. Suh *et al.* [102] saw that neurexin 3 was downregulated in the hippocampus of mice treated with kainic acid, an inducer of neurodegeneration, while Swanberg *et al.* [101] found that neurexin 3 is downregulated in rats with axonal damage caused by ventral root avulsion.

Two studies were conducted in animal models of neuro-toxicity. One study exposed zebrafish to chronic, low levels of the neurotoxin domoic acid and saw an upregulation of neurexin 2 α after two weeks [106]. The other study exposed rats to acute doses of sarin, which caused a persistent down-regulation of neurexin 1 β and breakdown of the blood-brain barrier [105].

In summary, multiple studies have shown changes in neu-rexin expression in AD, ALS, MS, PD and SMA. Many of these studies have observed downregulation of protein expression for neurexin 1, 2 and 3 in these disorders. Similarly, downregulation of neurexin 1, 2 and 3 were observed in disorders such as HIV encephalitis and MCI and in studies on ageing, in models of neuronal toxicity, and animal models of MS and ALS.

6. Concluding remarks

A clear link between synaptic dysfunction and neurodegenerative as well as neuropsychiatric disorders has been established in recent years. Our literature-based searches revealed several

studies that have linked CNVs, deletions or expression changes in neurexins to different disorders. The evidence is most compelling for a role of neurexins in neuropsychiatric disorders, particularly in regard to the involvement of neurexin 1 in ASD and SCZ. Currently, there is comparatively less evidence for the involvement of neurexins in neurodegenerative disorders. Although there have been some studies that have suggested that neurexins may be important in these disorders, at this stage more experimental data are still needed to draw concrete conclusions. Therefore, it is apparent that more targeted studies in various disorders involving these genes as well as the proteins they encode are warranted. In terms of their broader biological and physiological functions, the neurexins function as molecular inducers, are involved in iron and protein binding, and play a role in cell-to-cell communication and response to stimuli, consequently making them critical for normal cell functioning. Furthermore, these proteins interact with various other proteins such as the neuroligins and the LRRTM proteins identified via protein interaction networks. This implicates the neurexins' involvement in synaptic integrity and functioning making them promising candidates as disease genes for a wide range of brain pathologies.

In summary, this review serves to highlight the potential importance of the neurexin genes and proteins in human disease and recommends that more targeted studies on these genes and proteins are warranted. Furthermore, with the wealth of exomic and genomic sequences and genome-wide transcriptomic datasets now available, it has become plausible

to interrogate them for their involvement in various human disorders, on a scale not previously possible. In addition, the human neurexin protein structures urgently need to be solved to understand the function and infer accurate protein–protein interactions as well as to understand the effect of mutations on the protein structure. Ultimately, improved knowledge on synapses and their individual components are necessary to develop novel therapeutic approaches for the emerging and exciting field of synaptopathies.

Data accessibility. This article has no additional data.

Authors' contributions. K.C. performed the research and wrote the first draft of the manuscript. M.H. and R.C. assisted with some of the analysis. K.C. and S.B. conceptualized the study. All authors critically reviewed and edited the manuscript.

Competing interests. The authors declare that there are no competing interests.

Funding. This work is based on the research supported wholly/in part by the National Research Foundation of South Africa (NRF) (grant nos. 106052, 129249); the South African Medical Research Council (SAMRC) (self-initiated research grant); the Harry Crossley Foundation and Stellenbosch University, South Africa. SAMRC and The Higher Education Department, Next Generation of Academic Programme (nGAP), provided support for R.C. in the form of a full-time academic position and salary.

Acknowledgements. We acknowledge Minke Bekker for the drawing of figure 1. We also acknowledge the support of the NRF-DST Centre of Excellence for Biomedical Tuberculosis Research; South African Medical Research Council Centre for Tuberculosis Research; Division of Molecular Biology and Human Genetics, Faculty of Medicine and Health Sciences, Stellenbosch University, Cape Town.

References

- Torres VI, Vallejo D, Inestrosa NC. 2017 Emerging synaptic molecules as candidates in the etiology of neurological disorders. *Neural Plast.* **2017**, 8081758. (doi:10.1155/2017/8081758)
- Imbriani P, Schirizzi T, Meringolo M, Mercuri NB, Pisani A. 2018 Centrality of early synaptopathy in Parkinson's disease. *Front. Neurol.* **9**, 103. (doi:10.3389/fneur.2018.00103)
- Südhof TC. 2017 Synaptic neurexin complexes: a molecular code for the logic of neural circuits. *Cell* **171**, 745–769. (doi:10.1016/j.cell.2017.10.024)
- Lepeta K *et al.* 2016 Synaptopathies: synaptic dysfunction in neurological disorders—a review from students to students. *J. Neurochem.* **138**, 785–805. (doi:10.1111/jnc.13713)
- Missler M, Südhof TC, Biederer T. 2012 Synaptic cell adhesion. *Cold Spring Harb. Perspect. Biol.* **4**, a005694. (doi:10.1101/cshperspect.a005694)
- Schikorski T, Stevens CF. 1997 Quantitative ultrastructural analysis of hippocampal excitatory synapses. *J. Neurosci.* **17**, 5858–5867. (doi:10.1523/jneurosci.17-15-05858.1997)
- Toni N, Buchs PA, Nikonenko I, Bron CR, Müller D. 1999 LTP promotes formation of multiple spine synapses between a single axon terminal and a dendrite. *Nature* **402**, 421–425. (doi:10.1038/46574)
- Krueger DD, Tuffy LP, Papadopoulos T, Brose N. 2012 The role of neurexins and neuroligins in the formation, maturation, and function of vertebrate synapses. *Curr. Opin. Neurobiol.* **22**, 412–422. (doi:10.1016/j.conb.2012.02.012)
- Taoufik E, Kouroupi G, Zygogianni O, Matsas R. 2018 Synaptic dysfunction in neurodegenerative and neurodevelopmental diseases: an overview of induced pluripotent stem-cell-based disease models. *Open Biol.* **8**, 180138. (doi:10.1098/rsob.180138)
- Sebate B, Cuttler K, Cloete R, Britz M, Christoffels A, Williams M, Carr J, Bardien S. 2021 Prioritization of candidate genes for a South African family with Parkinson's disease using in-silico tools. *PLoS ONE* **16**, e0249324. (doi:10.1371/journal.pone.0249324)
- Ushkaryov YA, Petrenko AG, Geppert M, Südhof TC. 1992 Neurexins: synaptic cell surface proteins related to the α -latrotoxin receptor and laminin. *Science* **257**, 50–56. (doi:10.1126/science.1621094)
- Rosenthal L, Meldolesi J. 1989 α -Latrotoxin and related toxins. *Pharmacol. Ther.* **42**, 115–134. (doi:10.1016/0163-7258(89)90024-7)
- Missler M, Südhof TC. 1998 Neurexins: three genes and 1001 products. *Trends Genet.* **14**, 20–26. (doi:10.1016/S0168-9525(97)01324-3)
- Ichtchenko K, Nguyen T, Südhof TC. 1996 Structures, alternative splicing, and neurexin binding of multiple neuroligins. *J. Biol. Chem.* **271**, 2676–2682. (doi:10.1074/jbc.271.5.2676)
- Rudenko G, Nguyen T, Chelliah Y, Su TC, Deisenhofer J, Hughes H. 1999 The structure of the ligand-binding domain of neurexin I β : regulation of LNS domain function. *Cell* **99**, 93–101. (doi:10.1016/S0092-8674(00)80065-3)
- Tabuchi K, Südhof TC. 2002 Structure and evolution of neurexin genes: insight into the mechanism of alternative splicing. *Genomics* **79**, 849–859. (doi:10.1006/geno.2002.6780)
- Craig AM, Kang Y. 2007 Neurexin-neuroligin signaling in synapse development. *Curr. Opin. Neurobiol.* **17**, 43–52. (doi:10.1016/j.conb.2007.01.011)
- Liao Y, Wang J, Jaehnig EJ, Shi Z, Zhang B. 2019 WebGestalt 2019: gene set analysis toolkit with revamped UIs and APIs. *Nucleic Acids Res.* **47**, W199–W205. (doi:10.1093/nar/gkz401)
- Szklarczyk D *et al.* 2019 STRING v11: protein-protein association networks with increased coverage, supporting functional discovery in genome-wide experimental datasets. *Nucleic Acids Res.* **47**, D607–D613. (doi:10.1093/nar/gky1131)
- Warde-Farley D *et al.* 2010 The GeneMANIA prediction server: biological network integration for gene prioritization and predicting gene function. *Nucleic Acids Res.* **38**, W214–W220. (doi:10.1093/nar/gkq537)
- The UniProt Consortium. 2021 UniProt: the universal protein knowledgebase in 2021. *Nucleic Acids Res.* **49**, D480–D489. (doi:10.1093/nar/gkaa1100)
- Kanehisa M, Furumichi M, Sato Y, Ishiguro-Watanabe M, Tanabe M. 2021 KEGG: integrating viruses and cellular organisms. *Nucleic Acids Res.* **49**, D545–D551. (doi:10.1093/nar/gkaa970)

23. Nakanishi M, Nomura J, Ji X, Tamada K, Arai T, Takahashi E, Bucan M, Takumi T. 2017 Functional significance of rare neuroligin 1 variants found in autism. *PLoS Genet.* **13**, e1006940. (doi:10.1371/journal.pgen.1006940)
24. Berkel S *et al.* 2010 Mutations in the SHANK2 synaptic scaffolding gene in autism spectrum disorder and mental retardation. *Nat. Genet.* **42**, 489–491. (doi:10.1038/ng.589)
25. Jamain S *et al.* 2003 Mutations of the X-linked genes encoding neuroligins NLGN3 and NLGN4 are associated with autism. *Nat. Genet.* **34**, 27–29. (doi:10.1038/ng1136)
26. Laumonnier F *et al.* 2004 X-linked mental retardation and autism are associated with a mutation in the NLGN4 gene, a member of the neuroligin family. *Am. J. Hum. Genet.* **74**, 552–557. (doi:10.1086/382137)
27. Lawson-Yuen A, Saldivar JS, Sommer S, Picker J. 2008 Familial deletion within NLGN4 associated with autism and Tourette syndrome. *Eur. J. Hum. Genet.* **16**, 614–618. (doi:10.1038/sj.ejhg.5202006)
28. Najm J *et al.* 2008 Mutations of CASK cause an X-linked brain malformation phenotype with microcephaly and hypoplasia of the brainstem and cerebellum. *Nat. Genet.* **40**, 1065–1067. (doi:10.1038/ng.194)
29. Piluso G *et al.* 2008 A missense mutation in CASK causes FG syndrome in an Italian family. *Am. J. Hum. Genet.* **84**, 162–177. (doi:10.1016/j.ajhg.2008.12.018)
30. Tarpey PS *et al.* 2009 A systematic, large-scale resequencing screen of X-chromosome coding exons in mental retardation. *Nat. Genet.* **41**, 535–543. (doi:10.1038/ng.367)
31. Hackett A *et al.* 2010 CASK mutations are frequent in males and cause X-linked nystagmus and variable XLMR phenotypes. *Eur. J. Hum. Genet.* **18**, 544–552. (doi:10.1038/ejhg.2009.220)
32. Moog U *et al.* 2011 Phenotypic spectrum associated with CASK loss-of-function mutations. *J. Med. Genet.* **48**, 741–751. (doi:10.1136/jmedgenet-2011-100218)
33. Relievelid SH *et al.* 2016 Meta-analysis of 2,104 trios provides support for 10 new genes for intellectual disability. *Nat. Neurosci.* **19**, 1194–1196. (doi:10.1038/nn.4352)
34. Moutton S *et al.* 2018 Truncating variants of the DLG4 gene are responsible for intellectual disability with marfanoid features. *Clin. Genet.* **93**, 1172–1178. (doi:10.1111/cge.13243)
35. Baker K *et al.* 2018 SYT1-associated neurodevelopmental disorder: a case series. *Brain* **141**, 2576–2591. (doi:10.1093/brain/awy209)
36. Majercak J *et al.* 2006 LRRTM3 promotes processing of amyloid-precursor protein by BACE1 and is a positional candidate gene for late-onset Alzheimer's disease. *Proc. Natl Acad. Sci. USA* **103**, 17 967–17 972. (doi:10.1073/pnas.0605461103)
37. Dobyms WB *et al.* 2018 MACF1 mutations encoding highly conserved zinc-binding residues of the GAR domain cause defects in neuronal migration and axon guidance. *Am. J. Hum. Genet.* **103**, 1009–1021. (doi:10.1016/j.ajhg.2018.10.019)
38. Dong M, Noguchi S, Endo Y, Hayashi YK, Yoshida S, Nonaka I, Nishino I. 2015 DAG1 mutations associated with asymptomatic hyperCKemia and hypoglycosylation of α -dystroglycan. *Neurology* **84**, 273–279. (doi:10.1212/WNL.0000000000001162)
39. Geis T, Marquard K, Rödl T, Reihle C, Schirmer S, Von Kalle T, Bornemann A, Hehr U, Blankenburg M. 2013 Homozygous dystroglycan mutation associated with a novel muscle-eye-brain disease-like phenotype with multicystic leucodystrophy. *Neurogenetics* **14**, 205–213. (doi:10.1007/s10048-013-0374-9)
40. Hara Y *et al.* 2011 A dystroglycan mutation associated with limb-girdle muscular dystrophy. *N. Engl. J. Med.* **364**, 939–946. (doi:10.1056/nejmoa1006939)
41. Riemersma M *et al.* 2015 Absence of α - and β -dystroglycan is associated with Walker-Warburg syndrome. *Neurology* **84**, 2177–2182. (doi:10.1212/WNL.0000000000001615)
42. Hanein S *et al.* 2004 Leber congenital amaurosis: comprehensive survey of the genetic heterogeneity, refinement of the clinical definition, and genotype-phenotype correlations as a strategy for molecular diagnosis. *Hum. Mutat.* **23**, 306–317. (doi:10.1002/humu.20010)
43. Mataftsi A, Schorderet DF, Chachoua L, Boussalah M, Nouri MT, Barthelmes D, Borrut FX, Munier FL. 2007 Novel TULP1 mutation causing leber congenital amaurosis or early onset retinal degeneration. *Investig. Ophthalmol. Vis. Sci.* **48**, 5160–5167. (doi:10.1167/iovs.06-1013)
44. Banerjee P *et al.* 1998 TULP1 mutation in two extended Dominican kindreds with autosomal recessive Retinitis pigmentosa. *Nat. Genet.* **18**, 177–179. (doi:10.1038/ng0298-177)
45. Den Hollander AI, Van Lith-Verhoeven JJC, Arends ML, Strom TM, Cremers FPM, Hoyng CB. 2007 Novel compound heterozygous TULP1 mutations in a family with severe early-onset retinitis pigmentosa. *Arch. Ophthalmol.* **125**, 932–935. (doi:10.1001/archoph.125.7.932)
46. Hagstrom SA, North MA, Nishina PM, Berson EL, Dryja TP. 1998 Recessive mutations in the gene encoding the tubby-like protein TULP1 in patients with Retinitis pigmentosa. *Nat. Genet.* **18**, 174–176. (doi:10.1038/ng0298-174)
47. Kondo H, Qin M, Mizota A, Kondo M, Hayashi H, Hayashi K, Oshima K, Tahira T, Hayashi K. 2004 A homozygosity-based search for mutations in patients with autosomal recessive retinitis pigmentosa, using microsatellite markers. *Investig. Ophthalmol. Vis. Sci.* **45**, 4433–4439. (doi:10.1167/iovs.04-0544)
48. Levy D *et al.* 2011 Rare de novo and transmitted copy-number variation in autistic spectrum disorders. *Neuron* **70**, 886–897. (doi:10.1016/j.neuron.2011.05.015)
49. Sanders SJ *et al.* 2011 Multiple recurrent de novo CNVs, including duplications of the 7q11.23 Williams Syndrome region, are strongly associated with autism. *Neuron* **70**, 863–885. (doi:10.1016/j.neuron.2011.05.002)
50. Camacho-Garcia RJ, Planelles MI, Margalef M, Pecero ML, Martínez-Leal R, Aguilera F, Vilella E, Martínez-Mir A, Scholl FG. 2012 Mutations affecting synaptic levels of neurexin-1 β in autism and mental retardation. *Neurobiol. Dis.* **47**, 135–143. (doi:10.1016/j.nbd.2012.03.031)
51. Liu Y *et al.* 2012 Mutation analysis of the NRXN1 gene in a Chinese autism cohort. *J. Psychiatr. Res.* **46**, 630–634. (doi:10.1016/j.jpsychires.2011.10.015)
52. Girirajan S *et al.* 2013 Refinement and discovery of new hotspots of copy-number variation associated with autism spectrum disorder. *Am. J. Hum. Genet.* **92**, 221–237. (doi:10.1016/j.ajhg.2012.12.016)
53. Mariani J *et al.* 2015 FOXG1-dependent dysregulation of GABA/glutamate neuron differentiation in autism spectrum disorders. *Cell* **162**, 375–390. (doi:10.1016/j.cell.2015.06.034)
54. Sanders SJ *et al.* 2015 Insights into autism spectrum disorder genomic architecture and biology from 71 risk loci. *Neuron* **87**, 1215–1233. (doi:10.1016/j.neuron.2015.09.016)
55. Chen K, Kardys A, Chen Y, Flink S, Tabakoff B, Shih JC. 2017 Altered gene expression in early postnatal monoamine oxidase A knockout mice. *Brain Res.* **1669**, 18–26. (doi:10.1016/j.brainres.2017.05.017)
56. Tong XJ, López-Soto EJ, Li L, Liu H, Nedelcu D, Lipscombe D, Hu Z, Kaplan JM. 2017 Retrograde synaptic inhibition is mediated by α -neurexin binding to the α 2 δ subunits of N-type calcium channels. *Neuron* **95**, 326–340. (doi:10.1016/j.neuron.2017.06.018)
57. Gauthier J *et al.* 2011 Truncating mutations in NRXN2 and NRXN1 in autism spectrum disorders and schizophrenia. *Hum. Genet.* **130**, 563–573. (doi:10.1007/s00439-011-0975-z)
58. Mohrmann I, Gillissen-Kaesbach G, Siebert R, Caliebe A, Hellenbroich Y. 2011 A de novo 0.57 Mb microdeletion in chromosome 11q13.1 in a patient with speech problems, autistic traits, dysmorphic features and multiple endocrine neoplasia type 1. *Eur. J. Med. Genet.* **54**, 6. (doi:10.1016/j.ejmg.2011.04.006)
59. Sharon G *et al.* 2019 Human gut microbiota from autism spectrum disorder promote behavioral symptoms in mice. *Cell* **177**, 1600–1618. (doi:10.1016/j.cell.2019.05.004)
60. Kuo PH *et al.* 2014 Identification of novel loci for bipolar I disorder in a multi-stage genome-wide association study. *Prog. Neuro-Psychopharmacol. Biol. Psychiatry* **51**, 58–64. (doi:10.1016/j.pnpb.2014.01.003)
61. Fabbri C, Serretti A. 2016 Genetics of long-term treatment outcome in bipolar disorder. *Prog. Neuro-Psychopharmacol. Biol. Psychiatry* **65**, 17–24. (doi:10.1016/j.pnpb.2015.08.008)
62. Panagopoulos VN *et al.* 2013 Examining the association of NRXN3 SNPs with borderline personality disorder phenotypes in heroin dependent cases and socio-economically disadvantaged controls. *Drug Alcohol Depend.* **128**, 187–193. (doi:10.1016/j.drugalcdep.2012.11.011)

63. Grimming T *et al.* 2013 Levetiracetam resistance: synaptic signatures & corresponding promoter SNPs in epileptic hippocampi. *Neurobiol. Dis.* **60**, 115–125. (doi:10.1016/j.nbd.2013.08.015)
64. Angione K, Eschbach K, Smith G, Joshi C, Demarest S. 2019 Genetic testing in a cohort of patients with potential epilepsy with myoclonic-atonic seizures. *Epilepsy Res.* **150**, 70–77. (doi:10.1016/j.eplepsyres.2019.01.008)
65. Górecki DC, Szkarczyk A, Lukasiuk K, Kaczmarek L, Simons JP. 1999 Differential seizure-induced and developmental changes of neuexin expression. *Mol. Cell. Neurosci.* **13**, 218–227. (doi:10.1006/mcne.1999.0740)
66. Lai JKY, Doering LC, Foster JA. 2016 Developmental expression of the neuroligins and neuexins in fragile X mice. *J. Comp. Neurol.* **524**, 807–828. (doi:10.1002/cne.23868)
67. Han X, Shao W, Liu Z, Fan S, Yu J, Chen J, Qiao R, Zhou J, Xie P. 2015 ITRAQ-based quantitative analysis of hippocampal postsynaptic density-associated proteins in a rat chronic mild stress model of depression. *Neuroscience* **298**, 220–292. (doi:10.1016/j.neuroscience.2015.04.006)
68. Pak CH, Danko T, Zhang Y, Aoto J, Anderson G, Maxeiner S, Yi F, Wernig M, Südhof TC. 2015 Human neuropsychiatric disease modeling using conditional deletion reveals synaptic transmission defects caused by heterozygous mutations in NRXN1. *Cell Stem Cell* **17**, 316–328. (doi:10.1016/j.stem.2015.07.017)
69. Mirnics K, Middleton FA, Marquez A, Lewis DA, Levitt P. 2000 Molecular characterization of schizophrenia viewed by microarray analysis of gene expression in prefrontal cortex. *Neuron* **28**, 53–67. (doi:10.1016/S0896-6273(00)00085-4)
70. Kirov G *et al.* 2008 Comparative genome hybridization suggests a role for NRXN1 and APBA2 in schizophrenia. *Hum. Mol. Genet.* **17**, 458–465. (doi:10.1093/hmg/ddm323)
71. Ivorra JL *et al.* 2014 Replication of previous genome-wide association studies of psychiatric diseases in a large schizophrenia case-control sample from Spain. *Schizophr. Res.* **159**, 107–113. (doi:10.1016/j.schres.2014.07.004)
72. Todarello G, Feng N, Kolachana BS, Li C, Vakkalanka R, Bertolino A, Weinberger DR, Straub RE. 2014 Incomplete penetrance of NRXN1 deletions in families with schizophrenia. *Schizophr. Res.* **155**, 1–7. (doi:10.1016/j.schres.2014.02.023)
73. Suárez-Rama JJ *et al.* 2015 Resequencing and association analysis of coding regions at twenty candidate genes suggest a role for rare risk variation at AKAP9 and protective variation at NRXN1 in schizophrenia susceptibility. *J. Psychiatr. Res.* **66–67**, 38–44. (doi:10.1016/j.jpsychires.2015.04.013)
74. Kassan A *et al.* 2017 Caveolin-1 regulation of disrupted-in-schizophrenia-1 as a potential therapeutic target for schizophrenia. *J. Neurophysiol.* **117**, 436–444. (doi:10.1152/jn.00481.2016)
75. Windrem MS *et al.* 2017 Human iPSC glial mouse chimeras reveal glial contributions to schizophrenia. *Cell Stem Cell* **21**, 195–208. (doi:10.1016/j.stem.2017.06.012)
76. Li J, Yoshikawa A, Brennan MD, Ramsey TL, Meltzer HY. 2018 Genetic predictors of antipsychotic response to lurasidone identified in a genome wide association study and by schizophrenia risk genes. *Schizophr. Res.* **192**, 194–204. (doi:10.1016/j.schres.2017.04.009)
77. Pak CH *et al.* 2021 Cross-platform validation of neurotransmitter release impairments in schizophrenia patient-derived NRXN1-mutant neurons. *Proc. Natl Acad. Sci. USA* **118**, 2021. (doi:10.1073/pnas.2025598118)
78. Duong LTT, Hoeffding LK, Petersen KB, Knudsen CD, Thygesen JH, Klitten LL, Tommerup N, Ingason A, Werge T. 2015 Two rare deletions upstream of the NRXN1 gene (2p16.3) affecting the non-coding mRNA AK127244 segregate with diverse psychopathological phenotypes in a family. *Eur. J. Med. Genet.* **58**, 650–653. (doi:10.1016/j.ejmg.2015.11.004)
79. Verkerk AJMH *et al.* 1991 Identification of a gene (FMR-1) containing a CGG repeat coincident with a breakpoint cluster region exhibiting length variation in fragile X syndrome. *Cell* **65**, 905–914. (doi:10.1016/0092-8674(91)90397-H)
80. Stein JL *et al.* 2010 Genome-wide analysis reveals novel genes influencing temporal lobe structure with relevance to neurodegeneration in Alzheimer's disease. *Neuroimage* **51**, 542–554. (doi:10.1016/j.neuroimage.2010.02.068)
81. Brinkmalm G, Sjödin S, Simonsen AH, Hasselbalch SG, Zetterberg H, Brinkmalm A, Blennow K. 2018 A parallel reaction monitoring mass spectrometric method for analysis of potential CSF biomarkers for Alzheimer's disease. *Proteom. Clin. Appl.* **12**, 1700131. (doi:10.1002/prca.201700131)
82. Goetzl EJ, Abner EL, Jicha GA, Kapogiannis D, Schwartz JB. 2018 Declining levels of functionally specialized synaptic proteins in plasma neuronal exosomes with progression of Alzheimer's disease. *FASEB J.* **32**, 888–893. (doi:10.1096/fj.201700731R)
83. Rahman MM, Westermark GT, Zetterberg H, Härd T, Sandgren M. 2018 Protofibrillar and fibrillar amyloid- β binding proteins in cerebrospinal fluid. *J. Alzheimer's Dis.* **66**, 1053–1064. (doi:10.3233/JAD-180596)
84. Canchi S, Rao B, Masliah D, Rosenthal SB, Sasik R, Fisch KM, De Jager PL, Bennett DA, Rissman RA. 2019 Integrating gene and protein expression reveals perturbed functional networks in Alzheimer's disease. *Cell Rep.* **28**, 1103–1116. (doi:10.1016/j.celrep.2019.06.073)
85. Lleó A *et al.* 2019 Changes in synaptic proteins precede neurodegeneration markers in preclinical Alzheimer's disease cerebrospinal fluid. *Mol. Cell. Proteom.* **18**, 546–560. (doi:10.1074/mcp.RA118.001290)
86. Berchtold NC, Coleman PD, Cribbs DH, Rogers J, Gillen DL, Cotman CW. 2013 Synaptic genes are extensively downregulated across multiple brain regions in normal human aging and Alzheimer's disease. *Neurobiol. Aging* **34**, 1653–1661. (doi:10.1016/j.neurobiolaging.2012.11.024)
87. Neuner SM, Wilmott LA, Hoffmann BR, Mozhui K, Kaczorowski CC. 2017 Hippocampal proteomics defines pathways associated with memory decline and resilience in normal aging and Alzheimer's disease mouse models. *Behav. Brain Res.* **322**, 288–298. (doi:10.1016/j.bbr.2016.06.002)
88. Berchtold NC, Prieto GA, Phelan M, Gillen DL, Baldi P, Bennett DA, Buchman AS, Cotman CW. 2019 Hippocampal gene expression patterns linked to late-life physical activity oppose age and AD-related transcriptional decline. *Neurobiol. Aging* **78**, 142–154. (doi:10.1016/j.neurobiolaging.2019.02.012)
89. Popesco MC, Lin S, Wang Z, Ma ZJ, Friedman L, Frostholm A, Rotter A. 2008 Serial analysis of gene expression profiles of adult and aged mouse cerebellum. *Neurobiol. Aging* **29**, 774–788. (doi:10.1016/j.neurobiolaging.2006.12.006)
90. Tserel L, Limbach M, Saare M, Kisand K, Metspalu A, Milani L, Peterson P. 2014 CpG sites associated with NRP1, NRXN2 and miR-29b-2 are hypomethylated in monocytes during ageing. *Immun. Ageing* **11**, 1. (doi:10.1186/1742-4933-11-1)
91. Kumar D, Thakur MK. 2015 Age-related expression of neuexin1 and neuroligin3 is correlated with presynaptic density in the cerebral cortex and hippocampus of male mice. *Age (Omaha)* **37**, 17. (doi:10.1007/s11357-015-9752-6)
92. González-Velasco O, Papy-García D, Le Douaron G, Sánchez-Santos JM, De Las Rivas J. 2020 Transcriptomic landscape, gene signatures and regulatory profile of aging in the human brain. *Biochim. Biophys. Acta - Gene Regul. Mech.* **1863**, 194491. (doi:10.1016/j.bbagr.2020.194491)
93. Xiao S *et al.* 2011 RNA targets of TDP-43 identified by UV-CLIP are deregulated in ALS. *Mol. Cell. Neurosci.* **47**, 167–180. (doi:10.1016/j.mcn.2011.02.013)
94. Everall I, Salaria S, Roberts E, Corbeil J, Sasik R, Fox H, Grant I, Masliah E. 2005 Methamphetamine stimulates interferon inducible genes in HIV infected brain. *J. Neuroimmunol.* **170**, 158–171. (doi:10.1016/j.jneuroim.2005.09.009)
95. Berchtold NC, Sabbagh MN, Beach TG, Kim RC, Cribbs DH, Cotman CW. 2014 Brain gene expression patterns differentiate mild cognitive impairment from normal aged and Alzheimer's disease. *Neurobiol. Aging* **35**, 1961–1972. (doi:10.1016/j.neurobiolaging.2014.03.031)
96. Vounou M, Janousova E, Wolz R, Stein JL, Thompson PM, Rueckert D, Montana G. 2012 Sparse reduced-rank regression detects genetic associations with voxel-wise longitudinal phenotypes in Alzheimer's disease. *Neuroimage* **60**, 700–716. (doi:10.1016/j.neuroimage.2011.12.029)
97. Zeis T, Kinter J, Herrero-Herranz E, Weissert R, Schaeeren-Wiemers N. 2008 Gene expression analysis of normal appearing brain tissue in an animal model for multiple sclerosis revealed grey matter alterations, but only minor white matter changes.

- J. Neuroimmunol.* **205**, 10–19. (doi:10.1016/j.jneuroim.2008.09.009)
98. Opsahl JA *et al.* 2016 Label-free analysis of human cerebrospinal fluid addressing various normalization strategies and revealing protein groups affected by multiple sclerosis. *Proteomics* **16**, 1154–1165. (doi:10.1002/pmic.201500284)
99. Kattimani Y, Veerappa AM. 2018 Dysregulation of NRXN1 by mutant MIR8485 leads to calcium overload in pre-synapses inducing neurodegeneration in Multiple sclerosis. *Mult. Scler. Relat. Disord.* **22**, 153–156. (doi:10.1016/j.msard.2018.04.005)
100. Castro K *et al.* 2019 Body mass index in multiple sclerosis modulates ceramide-induced DNA methylation and disease course. *EBioMedicine* **43**, 392–410. (doi:10.1016/j.ebiom.2019.03.087)
101. Swanberg M, Duvefelt K, Diez M, Hillert J, Olsson T, Piehl F, Lidman O. 2006 Genetically determined susceptibility to neurodegeneration is associated with expression of inflammatory genes. *Neurobiol. Dis.* **24**, 67–88. (doi:10.1016/j.nbd.2006.05.016)
102. Suh YA, Kwon OM, Yim SY, Lee HJ, Kim SS. 2007 Identification of differentially expressed genes in murine hippocampus by modulation of nitric oxide in kainic acid-induced neurotoxic animal model. *Korean J. Physiol. Pharmacol.* **11**, 149–154.
103. Fan Z, Chen X, Chen R. 2014 Transcriptome-wide analysis of TDP-43 binding small RNAs identifies miR-NID1 (miR-8485), a novel miRNA that represses NRXN1 expression. *Genomics* **103**, 76–82. (doi:10.1016/j.ygeno.2013.06.006)
104. Xiang YY, Dong H, Yang BB, Macdonald JF, Lu WY. 2014 Interaction of acetylcholinesterase with neurexin-1 β regulates glutamatergic synaptic stability in hippocampal neurons. *Mol. Brain* **7**, 1–7. (doi:10.1186/1756-6606-7-15)
105. Damodaran TV, Patel AG, Greenfield ST, Dressman HK, Lin SM, Abou-Donia MB. 2006 Gene expression profiles of the rat brain both immediately and 3 months following acute sarin exposure. *Biochem. Pharmacol.* **71**, 497–520. (doi:10.1016/j.bcp.2005.10.051)
106. Hiolski EM *et al.* 2014 Chronic low-level domoic acid exposure alters gene transcription and impairs mitochondrial function in the CNS. *Aquat. Toxicol.* **155**, 151–159. (doi:10.1016/j.aquatox.2014.06.006)
107. Noelker C *et al.* 2012 Differentially expressed gene profile in the 6-hydroxy-dopamine-induced cell culture model of Parkinson's disease. *Neurosci. Lett.* **507**, 10–15. (doi:10.1016/j.neulet.2011.11.035)
108. Richter F *et al.* 2014 Chronic administration of cholesterol oximes in mice increases transcription of cytoprotective genes and improves transcriptome alterations induced by alpha-synuclein overexpression in nigrostriatal dopaminergic neurons. *Neurobiol. Dis.* **69**, 263–275. (doi:10.1016/j.nbd.2014.05.012)
109. Fu LM, Fu KA. 2015 Analysis of Parkinson's disease pathophysiology using an integrated genomics-bioinformatics approach. *Pathophysiology* **22**, 15–29. (doi:10.1016/j.pathophys.2014.10.002)
110. Nezhadi A, Esmaeili-Mahani S, Sheibani V, Shabani M, Darvishzadeh F. 2017 Neurosteroid allopregnanolone attenuates motor disability and prevents the changes of neurexin 1 and postsynaptic density protein 95 expression in the striatum of 6-OHDA-induced rats' model of Parkinson's disease. *Biomed. Pharmacother.* **88**, 1188–1197. (doi:10.1016/j.biopha.2017.01.159)
111. Freeze B, Acosta D, Pandya S, Zhao Y, Raj A. 2018 Regional expression of genes mediating trans-synaptic alpha-synuclein transfer predicts regional atrophy in Parkinson disease. *NeuroImage Clin.* **18**, 456–466. (doi:10.1016/j.nicl.2018.01.009)
112. Haghparast E, Sheibani V, Abbasnejad M, Esmaeili-Mahani S. 2019 Apelin-13 attenuates motor impairments and prevents the changes in synaptic plasticity-related molecules in the striatum of Parkinsonism rats. *Peptides* **117**, 170091. (doi:10.1016/j.peptides.2019.05.003)
113. See K *et al.* 2014 SMN deficiency alters Nrnx2 expression and splicing in zebrafish and mouse models of spinal muscular atrophy. *Hum. Mol. Genet.* **23**, 1754–1770. (doi:10.1093/hmg/ddt567)
114. Petersen RC. 2011 Mild cognitive impairment. *N. Engl. J. Med.* **364**, 2227–2234. (doi:10.1056/NEJMc0910237)
115. Tatemoto K *et al.* 1998 Isolation and characterization of a novel endogenous peptide ligand for the human APJ receptor. *Biochem. Biophys. Res. Commun.* **251**, 471–476. (doi:10.1006/bbrc.1998.9489)
116. Zhang X *et al.* 2011 Up-regulation of apelin in brain tissue of patients with epilepsy and an epileptic rat model. *Peptides* **32**, 1793–1799. (doi:10.1016/j.peptides.2011.08.006)
117. Baulieu EE, Robel P, Schumacher M. 2001 Neurosteroids: beginning of the story. *Int. Rev. Neurobiol.* **46**, 1–32. (doi:10.1016/S0074-7742(01)46057-0)
118. di Michele F, Longone P, Romeo E, Lucchetti S, Brusa L, Pierantozzi M, Bassi A, Bernardi G, Stanzione P. 2003 Decreased plasma and cerebrospinal fluid content of neuroactive steroids in Parkinson's disease. *Neurol. Sci.* **24**, 172–173. (doi:10.1007/s10072-003-0115-1)
119. Mackenzie IRA, Rademakers R. 2008 The role of transactive response DNA-binding protein-43 in amyotrophic lateral sclerosis and frontotemporal dementia. *Curr. Opin. Neurol.* **21**, 693–700. (doi:10.1097/WCO.0b013e3283168d1d)
120. Lefebvre S *et al.* 1995 Identification and characterization of a spinal muscular atrophy-determining gene. *Cell* **80**, 155–165. (doi:10.1016/0092-8674(95)90460-3)