

Review Article

Omega-3 Fatty Acids and PPAR γ in Cancer

Iris J. Edwards¹ and Joseph T. O'Flaherty²

¹Department of Pathology, Wake Forest University School of Medicine, Winston-Salem, NC 27157, USA

²Department of Internal Medicine, Wake Forest University School of Medicine, Winston-Salem, NC 27157, USA

Correspondence should be addressed to Iris J. Edwards, iedwards@wfubmc.edu

Received 28 February 2008; Revised 30 May 2008; Accepted 24 June 2008

Recommended by Dipak Panigrahy

Omega-3 (or n-3) polyunsaturated fatty acids (PUFAs) and their metabolites are natural ligands for peroxisome proliferator receptor activator (PPAR) γ and, due to the effects of PPAR γ on cell proliferation, survival, and differentiation, are potential anticancer agents. Dietary intake of omega-3 PUFAs has been associated with a reduced risk of certain cancers in human populations and in animal models. In vitro studies have shown that omega-3 PUFAs inhibit cell proliferation and induce apoptosis in cancer cells through various pathways but one of which involves PPAR γ activation. The differential activation of PPAR γ and PPAR γ -regulated genes by specific dietary fatty acids may be central to their distinct roles in cancer. This review summarizes studies relating PUFAs to PPAR γ and cancer and offers a new paradigm relating an n-3 PUFA through PPAR γ to the expression of the cell surface proteoglycan, syndecan-1, and to the death of cancer cells.

Copyright © 2008 I. J. Edwards and J. T. O'Flaherty. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

1. INTRODUCTION

The peroxisome proliferator-activated receptor (PPAR) family of nuclear receptors comprises three distinct gene products, PPAR α , β/δ , and γ , that differ in ligand specificity, tissue distribution, and developmental expression [1–3]. PPARs demonstrate a relatively high level of constitutive transcriptional activity which is further increased upon binding their activating ligands [4–7]. These ligands are primarily long chain unsaturated and polyunsaturated fatty acids (PUFAs) and certain metabolites of these fatty acids [8–10]. Initially, PPARs were thought mainly to govern lipid homeostasis by binding fatty acids and their metabolites to thereby become more active in regulating genes for proteins involved in lipid metabolism [8, 10, 11]. Indeed, PPAR α is expressed predominantly in tissues with high fatty acid requirements such as liver, heart, and kidney, while PPAR γ isoforms γ 1 and γ 2 are highly enriched in adipose tissue to regulate adipocyte differentiation and lipid storage [3]. However, expression of PPAR γ 1, as with PPAR β/δ and PPAR α , has now been extended to most other tissues and regulatory roles for PPARs extended to other systemic functions such as carbohydrate regulation, immune modulation, and the

proliferation, survival and differentiation of cells [3]. The latter effects have led to intense interest in the PPARs in relation to cancer.

PPAR α and its ligand activators regulate fatty acid and lipoprotein metabolism and promote the development of hepatocellular carcinoma in rodents and reduce the metastasis of melanoma in hamsters [12]. These and other of their effects do not, in general, translate to humans. PPAR β/δ plays a key role in lipid metabolism of peripheral tissues. Its high expression in colon has been shown to promote colon cancer [12, 13], in a mechanism that involves the stimulation of PPAR β/δ by arachidonic acid, PPAR β/δ -dependent upregulation of cyclooxygenase (COX)-2 leading to overproduction of prostaglandin (PG)E₂, and PGE₂-induced growth of colon cancer cells. There is relatively little documentation of a role for PPAR β/δ in other cancers [14]. By contrast, PPAR γ has a broad range of effects on cancer. PPAR γ controls fat metabolism by regulating genes involved in lipogenesis, insulin sensitivity, and adipocyte differentiation [3, 15]. These effects underlie the use of thiazolidinediones, which bind and activate PPAR γ , to treat insulin-resistant type II diabetes [3, 15]. Although PPAR γ activators have been widely shown to inhibit growth in

cultured cancer cells, *in vivo* effects have proved to be complex: they inhibit but sometimes promote cancer growth [16] probably due to stimulation of antiproliferative and apoptotic signaling pathways or proliferative and antiapoptotic pathways, depending on cellular conditions [3, 12, 15–18]. These findings led to the idea of selective PPAR γ modulators (SPARMs), drugs analogous to selective estrogen receptor modulators (SERMs) in which distinct actions of the modulator depend on the cellular context [19] and on distinct receptor conformations, and therefore different gene interactions [20]. Fatty acids may be considered as natural SPARMs since their binding does not necessarily lead to PPAR activation and target gene transcription [11].

The considerations discussed above raise a possibility that managed alterations in the type of fatty acids in tissues, can alter the activity of PPARs and thereby the genes they control for therapeutic benefit. The fatty acid content of tissues is dependent mainly on dietary intake. Omega-3 PUFAs, docosahexaenoic acid (DHA), and eicosapentaenoic acid (EPA) are enriched in the diets of many populations that enjoy a low incidence of cancer [21]. These diets also obtain some modest success ameliorating advanced cancer in humans [22] and have been widely used to inhibit carcinogenesis and tumor progression in animal models. The ability of specific fatty acids to differentially activate PPARs and PPAR-regulated genes may be central to their distinct roles in cancer. This review will focus on PPAR γ , its activation by fatty acids, and functional results in cancer cells.

2. FATTY ACID METABOLISM

2.1. Fatty acid types and interconversions

Fatty acids are hydrocarbons with a terminal carboxyl group. The carbons of saturated fatty acids are all connected by single bonds, whereas the chains of monounsaturated and polyunsaturated fatty acids (PUFAs) contain one or more double bonds, respectively. The n-3 and n-6 designation describes the position of the double bond closest to the (omega) carbon at the methyl end of the molecule (Figure 1). Oleic acid (18 : 1) has a single double bond between carbons 9 and 10 from the omega carbon and is designated an n-9 or omega-9 monounsaturated fatty acid. Like the saturated fatty acids, oleic acid can be synthesized *de novo* in mammalian cells. It can also be obtained from the diet through intake of oils such as olive and canola. By contrast, PUFAs cannot be synthesized *de novo* in mammals and must be obtained from the diet. The shortest of the n-6 PUFAs is linoleic acid (LA, 18 : 2, n-6). Its 18 carbon, n-3 counterpart is α -linolenic acid (ALA, 18 : 3, n-3). Both LA and ALA are metabolized through a series of elongation and desaturation steps to longer chain PUFAs: LA to arachidonic acid (AA, 20 : 4, n-6) and ALA to EPA (20 : 5, n-3) and DHA (22 : 6, n-3) (Figure 2). The first and rate limiting step in this pathway is the introduction of a double bond by the Δ 6 desaturase (for review see [23]). For n-3 PUFAs, ALA is converted to stearidonic acid (SDA, 18 : 4, n-3), elongated, and desaturated by Δ 5-desaturase to form EPA. In mammalian cells, the conversion of EPA to DHA follows the Sprecher pathway in which EPA is

elongated to docosapentaenoic acid (DPA, 22 : 5, n-3), then to tetracosapentaenoic acid (TPA, 24 : 5, n-3), and desaturated to tetracosahexaenoic acid (THA, 24 : 6, n-3). THA is translocated from the endoplasmic reticulum to peroxisomes, where β -oxidation results in the loss of 2 carbons to form DHA [24]. The PUFAs are also metabolized, most importantly for this review, to PPAR γ activators (see Section 2.3).

2.2. Dietary fatty acids

The results of both dietary intake and stable isotope studies have shown that the conversion of ALA to DHA in humans is extremely inefficient (for review see [25]). Most of the ingested ALA is an immediate target for β -oxidation to provide energy, leaving an estimated 8–10% to enter the elongation-desaturation pathway [26, 27]. A kinetic analysis of ^2H -labeled fatty acids estimated that conversion of ALA to EPA was only 0.2%, EPA to DPA was 0.13%, and DPA to DHA was 0.05% [28]. There is some evidence of gender-related differences in the activity of the elongation-desaturation pathway that result in a greater efficiency of conversion of ALA to DHA in females than in males [25, 27, 29]. Support for a role of sex hormones in the conversion pathway is provided by data indicating higher DHA in plasma lipids associated with oral contraceptive use [27] as well in males supplemented with estrogen during sex-change procedures [30]. Moreover, testosterone treatment of female-male transsexuals was shown to decrease plasma DHA [30].

Because common enzymes in the elongation-desaturation pathway are responsible for conversion of both n-3 and n-6 PUFAs, background diet is also a factor in efficiency of conversion. LA is the most abundant fatty acid in the Western diet with consumption in US that is ten-fold that of ALA (reviewed in [31]). Studies have shown that a high intake of LA is associated with a low conversion of ALA to EPA [26]. In spite of limited metabolism of ALA to its long chain derivatives in the stable isotope tracer studies, feeding studies have consistently shown that increased consumption of ALA does result in higher levels of EPA in plasma or cell lipids [31]. However, there was no measurable increase in DHA in these pools. Likewise diets supplemented with EPA do not result in a detectable increase in plasma DHA [32]. Thus, the inefficiency of this pathway does not appear limited to one step but rather extends throughout the pathway. The consensus of a number of studies is that the only way to increase plasma and tissue levels of a specific PUFA is to increase the consumption of that fatty acid. This may be of particular importance in the light of recent *in vitro* studies on the antitumor effects of DHA.

2.3. PUFAs metabolism to PPAR γ activators

Tissues metabolize PUFAs to oxygenated products that have quite different impacts on PPAR γ than their parent molecules. Moreover, n-3 PUFA inhibit the metabolism n-6 PUFAs and subplant them from their oxygenation pathways to form products [33–35] that have different effects on PPAR γ than their n-6 PUFAs oxygenated counterparts.




FIGURE 1: Structures of unsaturated fatty acids: oleic acid (n-9 monounsaturated), linoleic acid and arachidonic acid (n-6 polyunsaturated), α -linolenic acid, eicosapentaenoic acid, and docosahexaenoic acid (n-3 polyunsaturated). The “n” numbers are counted from the methyl or omega terminus.




FIGURE 2: The elongation-desaturation pathway for the metabolism of n-6 and n-3 polyunsaturated fatty acids.

It is therefore important to consider PUFAs oxygenation pathways. LA, AA, and DHA require $>10\text{--}30\ \mu\text{M}$ to activate PPAR γ but are commonly converted to stronger ($>0.1\text{--}10\ \mu\text{M}$) activators in cells. LA is metabolized (Figure 3, upper panel) by 15-lipoxygenases (LOX)-1/2 to 9(S)- and 13(S)-HODE (hydroxy-octadecaenoate) and by cyclooxygenases (COX)-1/2 to 9(R)- and 13(S)-HODE. The HODEs can be converted to 13-oxo- and 9-oxo-ODE by a dehydrogenase [36–39]. The hydroxy and to a greater extent oxo LA analogs have greater PPAR γ -activating potency than LA [36, 40–42]. AA is metabolized (Figure 3, center panel) via 5-LOX to 5(S)-HETE (hydroxy-eicosatetra-enoate) and via 15-LOX-1/2 to 15(S)-HETE. These HETEs can be converted to oxo-ETEs and 5-oxo-15(S)-hydroxy-ETE as shown in Figure 3 [39, 43–50]. 15-HETE has weak and 5-HETE essentially no ability to activate PPAR γ . However, their oxo counterparts have appreciable ability to do so with 5-oxo-15(S)-hydroxy-ETE showing the greatest potency in binding and activating PPAR γ [43]. AA is also metabolized (Figure 3, center panel) by COX1/2 to PG (prostaglandin) D₂ which as a consequence of successive dehydrations and an isomerization, perhaps by nonenzymatic routes, convert to PGJ₂, Δ^{12} -PGJ₂, and 15-deoxy $\Delta^{12,14}$ -PGJ₂ (15-d- $\Delta^{12,14}$ -PGJ₂); these PGJ₂'s have greater ability than PGD₂ to activate PPAR γ with 15-d- $\Delta^{12,14}$ -PGJ₂ being a most potent ($>0.1\text{--}1\ \mu\text{M}$) naturally occurring PPAR γ activator [9, 43, 51–56]. In one study, the K_d 's of 15-d- $\Delta^{12,14}$ -PGJ₂, 5-oxo-15-OH-ETE, PGJ₂, 5-oxo-ETE, and 5(S)-HETE in binding to PPAR γ were 1.4, 11, 37, 81, and $>1000\ \mu\text{M}$, respectively; their potency in activating a cell-based PPAR γ reporter paralleled these K_d 's [43]. DHA is metabolized (Figure 3, bottom panel) by 15-LOX or other oxygenase to 17-OH- and 7-OH-DHA, products that activate PPAR γ with greater potency (ED₅₀'s in activating a cell-based PPAR γ reporter of $\sim 5\ \mu\text{M}$) than DHA [57]. 4-OH-, and 4-oxo-DHA [53], while not yet shown to be made by cancer cells, also activate PPAR γ with greater potency (ED₅₀'s of 13.4 and $7.8\ \mu\text{M}$ in activating a cellular PPAR γ reporter, resp.) than DHA (ED₅₀ $> 10\ \mu\text{M}$) [53]. Hence, in this DHA series, similar to the 5-HETE series of AA metabolites, the oxo analog exhibits the greatest potency. We note that the more potent PPAR γ activators, the oxo-PUFAs, form preferentially in cells undergoing excessive oxidation, free radical, and NADPH/NADH-depleting reactions [43, 44, 48, 57, 58]. This suggests that PPAR γ may serve as a sensor for oxo-PUFA thereby monitoring cellular oxidative stress and when this stress is severe, engaging cell death programs [43, 58]. This PPAR γ function, we suggest, could contribute to the necrosis that occurs in tumors particularly after chemical and radiation treatment [59].

Cells process PUFAs in other relevant ways. They convert them to nitrates, probably in nonenzymatic reactions, where the nitric oxide made during cell stimulation attacks the PUFAs. Nitrated LA and AA are stronger PPAR γ activators than their precursors [60–62]. Cells also convert PUFAs to cannabinoids such as anandamide (ethanolamine amide of AA) and arachidonoylglycerol which also activate PPAR γ with greater potency than AA [63–65]. Finally, cells conjugate glutathione to PUFAs that contain an α,β -unsaturated

ketone such as 15-d- $\Delta^{12,14}$ -PGJ₂ and 5-oxo-ETE [66–68]. Since the conjugates are rapidly excreted from cells by multidrug-resistance transporters, conjugation inhibits the ability of α,β -unsaturated ketones to activate PPAR γ [66]. Cancer cells excrete anticancer drugs through these same transporters and become drug-resistant by overexpressing these transporters [69]. Such mutated cells may also be resistant to α,β -unsaturated ketone activators of PPAR γ .

2.4. Low-density lipoproteins (LDL) as deliverers of PPAR γ -activating n-3 PUFA

LDL carry esterified PUFAs in glycerolipids and cholesterol. They bind to cell surface LDL receptors and then internalize in endocytic vesicles which merge with lysosomes to de-esterify and release the PUFAs into the cytosol [70]. This route differs from the direct delivery of PUFA: it bypasses cell surface G protein-coupled fatty acid receptors (GPR 40 and 120; see Section 4.3), deposits PUFA in cells more slowly, and thereby avoids stimulation of G protein-coupled receptors and, perhaps, an array of C domain-bearing proteins which are activated by PUFA. This is also an important pathway for delivering PUFA to tumor cells because of a significant increase in LDL receptor activity in neoplastic tissues [71–73]. We have obtained from monkeys fed special diets, LDL enriched with n-6 PUFA (mostly AA and LA) or n-3 PUFA (mostly DHA and EPA). The n-3 but not n-6 PUFA-rich LDL mimicked thiazolidinediones and DHA in inhibiting cancer cell growth [74] and activating PPAR γ [75, 76].


3. PPAR γ

3.1. Structural considerations


PPAR γ 1 and γ 2 originate from the PPAR γ gene through separate promoters and 5' exons. Compared to the ubiquitously expressed PPAR γ 1, PPAR γ 2, which is limited mainly to adipose tissue, has 30 additional amino acids at its NH₂ terminus and is a more potent transcription activator [77]. Because they appear to have the same targets, however, the two isoforms are here considered together under the term PPAR γ . PPAR γ is comprised of four functional parts: the NH₂-terminal A/B region bears a ligand-independent transcription-activating motif AF-1; C region binds response elements (PPREs with a DR-1 consensus half-sequence of AGGTCA); D region binds various transcription cofactors; and E/F region has an interface for dimerizing with 9-*cis* retinoic acid receptors (RXRs), an AF-2 ligand-dependent transcription-activating motif, and a ligand-binding domain (LBD) [3, 12, 15, 17]. The LBD has a spacious cavity that binds ligands having a polar head group extending from a hydrophobic tail such as diverse PUFAs and PUFA metabolites [7, 77].

3.2. PPAR γ regulation by other signaling pathways


PPAR γ is phosphorylated by extracellular signal-regulated kinases (ERK)-1/2 and C-Jun N-terminal kinase; when so



(a)



(b)



(c)

FIGURE 3: The cellular metabolism of LA, AA, and DHA to more potent activators of PPAR γ . ODE is octadecaenoate; HETE is hydroxy-eicosatetraenoate; ETE is eicosatetraenoate; PG is prostaglandin.

phosphorylated, it has less ligand-binding affinity and gene-regulating activity [3, 78, 79]. The phosphorylation and attendant decrease in activity of PPAR γ occur in cells treated with PPAR γ activators and may cause the activators to show little or no ability to stimulate PPAR γ [3, 79–81]. ERK pathways impact PPAR γ in another way: the ERK-activating enzyme, MEK, when activated, binds with PPAR γ 's AF-2 motif. This causes PPAR γ to release from PPRE complexes and, bound to MEK and directed by MEK's nuclear export signal, to exit the nucleus [81, 82]. It is important to note that PUFAs and PUFA metabolites can activate the MEK/ERK pathway (see Section 4.3) and therefore may have biphasic effects: they not only directly activate PPAR γ but also entrain events inhibiting PPAR γ .

PPAR γ is targeted for degradation by ubiquitylation and sumoylation. Ligand binding, certain protein kinases, and some transcription cofactors (e.g., p300) promote ubiquitin-dependent degradation of PPAR γ in proteasomes [3]. Sumoylation occurs on K107 of PPAR γ 2 in a ligand-independent fashion to inhibit AF-1 function and on K365 of PPAR γ in a ligand-dependent fashion to promote PPAR γ 's binding of nuclear receptor corepressor [83, 84]. Sumoylation of PPAR γ causes its proteasomal degradation. ERK phosphorylation promotes K107 sumoylation. This reaction represents yet another means by which ERKs can inhibit PPAR γ [84].

3.3. PPAR γ transcriptional cofactors

PPARs bind a specific DNA sequence termed peroxisome proliferator response element (PPRE) in the 5'-flanking region of target genes as a heterodimer with RXR. Studies using various techniques [3, 85, 86] suggest the following model: PPAR γ •RXR complexes (the interaction is ligand-independent) exist in nuclei as macrocomplexes associated with various transcription corepressors [3, 87]. Some complexes, ligand-bound or not, may associate with transcription coactivators to control the basal expression of genes. In any event, PPAR γ •RXR complexes are highly mobile, rapidly scanning chromatin, although this scanning does not involve their DNA binding domain [86]. Ligands trigger PPAR γ •RXR to localize at their cognate PPREs and to exchange corepressors for coactivators such as cyclic AMP response element binding protein (CREB) and p300 [3, 16, 87, 88]. At some gene sites, activators cause PPAR γ •RXR to recruit corepressors and thereby cause gene repression [3, 89, 90]. However, the availability of cofactors differs between cell types and within cells over time depending on the cell's history and the association of the cofactors to other genes [3, 15, 16], for example, activation of PPAR γ deprives T cell factor/lymphoid enhancing factor (TCF/LEF) of cofactors to thereby inhibit oncogenic signaling by the Wnt pathway [16]. Thus, the effects of PPAR γ activation vary depending on context and cofactor availability at each genetic site. It seems at least possible that the PUFA ligands for PPAR γ will have differential effects in impacting its interactions with these transcriptional cofactors in a manner similar to the SPARMs model [19].

4. TARGETS OF PPAR γ RELEVANT TO CANCER

4.1. Gene targets of PPAR γ

Most known target genes of PPAR γ regulate lipid metabolism and transport [15] with few cancer-related genes having been confirmed as induced by PPAR γ . PPAR γ does induce G₀/G₁ switch gene 2 whose product causes growth arrest in 3T3-L1 cells [91, 92]. PPAR γ also binds the NF κ B promoter of p53 to stimulate expression of p53 and, in consequence, p21^{WAF1/Cip1}. It also binds to a promoter in the Fas ligand gene to induce the expression of this member of the extrinsic apoptosis pathway. These effects appear responsible for slowing growth and causing apoptosis in MCF7 breast cancer [93], human umbilical vein endothelial [94], and possibly Reh [95] cells. Recent studies have identified the heparan sulfate proteoglycan, syndecan 1, as a target for PPAR γ in human breast [75, 76] and prostate [96] cancer cells. The upregulation of syndecan 1 by PPAR γ resulted in apoptosis induction [76].

4.2. Other targets of PPAR γ

PPAR γ impacts many growth-promoting elements through its secondary actions that, while ligand-dependent, do not directly involve its gene promoters. It interacts with nuclear factor of activated T cells, phosphorylated signal transducer, and activator of transcription (STAT)-3, and nuclear factor κ B (NF κ B) to block signaling through these pathways [3]. It binds transcription cofactors to alter these cofactors' availability to other transcription factors: ligand bound-PPAR γ deprives NF κ B of AP-1; deprives STAT-1 of CREB binding protein; and releases SMRT to render it available to repress STAT-3's transcriptional activity [3, 16, 17, 97]. PPAR γ activation is also associated with the activation of ERK1/2, protein kinases C, protein kinase A, AMP-activated protein kinase α [17]; induction of p16, p18, and p21 cyclin-dependent kinase inhibitors [3, 17, 18]; decreased expression of cyclooxygenase 2, *cmyc*, *cmyb*, D1, and D3 cell cycle control genes, and regenerating gene 1A [17, 18]; decreased secretion of cytokines and growth factors [17, 98]; depression of the Akt survival pathway by upregulating PTEN and inhibiting the phosphorylation of Akt and mTOR [3, 17]; inhibiting retinoblastoma protein (Rb) activity to repress the activities of cyclins D3 and E [3]; and regulating a host of other elements involved in the growth and death of cells [3, 12, 16–18]. It is not clear which if any of these effects are due to the action of PPAR γ or PPAR γ activators. PUFAs impact many of these same targets but can do so not only by PPAR γ -dependent but also PPAR γ -independent routes (see the next section).

4.3. Targets of PPAR γ -activating Ligands

Studies of PPAR γ function depend on challenging cells with PPAR-activating ligands that have numerous side effects impacting cell growth. 15-d- $\Delta^{12,14}$ -PGJ₂ has a reactive α,β -unsaturated ketone (Figure 3) that covalently binds to cysteine sulfur on PPAR γ ; this renders its PPAR γ binding irreversible [58, 68]. 15-d- $\Delta^{1,14}$ -d-PGJ₂ also binds to cysteines

in the IKK β subunit of I κ B kinase, thereby inhibiting NF κ B activation [99, 100]. Other ligands with an α,β unsaturated ketone (e.g., oxo-ODEs and oxo-ETEs; see Figure 3) have this chemical reactivity [58] and along with 15-d- $\Delta^{1,14}$ -d-PGJ₂ may exert anticancer effects by covalently attaching to signal molecules like IKK β [58, 99, 101] or elements needed for expressing the epidermal growth factor receptor (EGFR) and JAK [102, 103].

Naturally occurring ligands have other PPAR γ -independent effects. The D and J series of PGs including 15-d- $\Delta^{12,14}$ -PGJ₂ bind to PGD₂ receptors [104], 5-oxo- and 5-oxo-15-hydroxy-ETE bind to the OXE receptor [105], and AA, EPA, and DHA bind to GPR40 and GPR120 receptors [106, 107]. These G protein-coupled receptors regulate signal pathways that effect cancer cell growth. For example, 5-oxo-15-hydroxy ETE acts on OXE to stimulate cells to activate ERK and Akt and proliferate; this stimulation counters its antigrowth activity in various cancer cell types. Indeed, HEK293 cells lack OXE receptors and in contrast to OXE receptor-bearing breast, prostate and ovarian cancer cell lines respond to 5-oxo-ETE and 5-oxo-15-oxo-ETE only by slowing, not speeding, their proliferation [43]. PUFAs activation of GPR120 also causes ERK and Akt activation to increase the survival of serum-starved STC-1 cells [108]. Finally, PUFAs are also metabolized to products that act on G protein receptors to promote cell growth, for example, prostate cancer cells convert AA to PGE₂, which acts through its receptors to stimulate the NF κ B pathway and thereby the expression of various cytokines and growth factors [109]. The G protein receptor-dependent actions of PPAR γ ligands may explain reports that these ligands have biphasic effects in stimulating proliferation and antiproliferation in cancer cells [110].

Thiazolidinediones stimulate cells to activate ERK1/2, p38, and JNK [111–113] by discharging Ca²⁺ from the ER to evoke an ER stress response; this activates Ca²⁺/calmodulin kinase II, proline-rich tyrosine kinase 2, protein kinases C, *c-Src*, EGFR, the ERK1/2 and JNK pathways, the double stranded RNA-activated protein kinase, and p38 [111]. Double stranded RNA-activated protein kinase inactivates eukaryotic initiation factor-2 to depress protein translation [111, 114]. Since EPA has recently been shown to have similar effects on ER calcium discharge [111, 115], it seems likely that various other PUFAs activate the ER stress pathway. Nonetheless, PPAR γ activators often show very different side effects [42, 103, 116–120]. For example, among three PPAR γ agonists, ciglitazone, 9-HODE, and 13-HODE, only 9-HODE induced apoptosis in U937 cells [38], 15d- $\Delta^{12,14}$ -PGJ₂, but not various other PPAR γ ligands, reduced EGFR expression in squamous carcinoma cells [99], 15d- $\Delta^{12,14}$ -PGJ₂, but not troglitazone, inhibited the stimulated induction of MHC class II molecules in retinal pigmented epithelial cells [112], and DHA, but not EPA, stimulated the target gene, syndecan 1 to inhibit the proliferation and induce apoptosis in breast and prostate cancer cell lines [75, 76, 96]. Numerous other examples of differential effects among PPAR γ agonists exist (e.g., [113–116]), but it is worth stressing that n-3 PUFAs inhibit the metabolism of n-6 PUFAs to products that promote the growth of cancer cells

such as PGE₂, 5-HETE, and leukotriene B₄ [33–35, 45, 113]. This inhibitory effect may make an important contribution to the anticancer effects of n-3 PUFAs.

5. DIETARY FATTY ACIDS AND CANCER

5.1. Human studies

Although there are inconsistencies [121], human population studies have shown that consumption of a diet enriched in n-3 PUFAs may offer protection against a number of cancers including those of breast [122–124], prostate [125, 126], and colon [127–129]. Although many of these studies have relied on dietary intake data from self-reported questionnaires or estimates based on national consumption, a few have used the fatty acid composition of tissues as a measure of exposure to dietary fats. The EURAMIC study is one of the largest to provide evidence that the balance between n-3 and n-6 PUFA may play a role in breast cancer [130]. Adipose tissue aspirates from breast cancer patients and controls demonstrated that the ratio of long chain n-3 to n-6 PUFAs was inversely associated with breast cancer in four of five centers studied. In human prostate tissue, lower EPA and DHA as well as lower n-3 to n-6 PUFAs ratios were associated with cancer compared to benign prostate hyperplasia [131] and with advanced stage compared to organ confined disease [132]. This inverse association of n-3 PUFAs and prostate cancer is supported by analyses of fatty acids in serum and red-cell membranes of patients with prostate disease [133, 134].

5.2. Animal studies

Animal studies provide convincing evidence of a negative relationship with n-3 PUFA diets and a positive relationship with n-6 PUFA diets for breast, prostate, and colon cancer. In studies of breast cancer induced by chemical carcinogens in rats [135–137], and human cancer cell xenografts in nude mice [138–140], tumor growth rate, size, and metastases were all suppressed by dietary n-3 PUFA supplementation. Likewise for colon cancer, antitumor properties of n-3 PUFA diets have been shown in transplantable mouse tumors [141–143] as well as in chemically induced rat tumors [144–151]. Although there have been fewer animal studies of PUFAs in prostate cancer, they are consistent with those in breast and colon cancer. In xenograft models of prostate cancer, n-3 PUFAs enriched diets inhibited tumor growth compared to n-6 PUFA diets [152–154]. Recently, a prostate-specific Pten knockout mouse model was used to demonstrate that a dietary ratio of n-6 to n-3 PUFA lower than 5 was effective in suppressing tumor growth, and extending animal lifespan [155].

5.3. Cell culture studies

Insight into the mechanism(s) responsible for the anticancer properties of n-3 PUFAs have been provided by animal studies as well as by in vitro investigations using human cancer cell lines. A major focus for such studies has been




FIGURE 4: The syndecan-1 pathway for n-3 PUFA induction of apoptosis. Dashed lines indicate that effects may be indirect with involvement of other metabolites and signaling molecules.

the competitive inhibition between n-6 and n-3 PUFAs for the enzymes involved in their metabolism. The desaturation and elongation of LA to AA were shown to be decreased in the presence of high n-3 PUFAs due to enzyme preference for the n-3 substrates [156]. AA and EPA compete for the COX and LOX enzymes, again with preferential n-3 utilization that results in a reduction in the highly reactive eicosanoids generated from AA [157, 158] in favor of less inflammatory n-3 eicosanoids [159]. The decreased growth of prostate xenograft tumors was shown to involve inhibition of COX 2 and PGE₂ in the tissues [154]. Thus, the combined human, animal, and cell culture studies indicate that diet is an important regulator of the levels of n-3 versus n-6 PUFAs in tissues, including those that are cancerous. High levels of n-3 PUFAs may directly evoke antitumor events, become metabolized to products with antitumor activity, or suppress the production of tumor-promoting metabolites such as those formed by n-6 PUFAs.

6. n-3 PUFA REGULATION OF SYNDECAN-1

Increasing evidence implicates PPAR γ in the divergent effects of n-3 and n-6 PUFAs in cancer cells and point to a growth inhibitory role for PPAR γ [160–164]. We recently found that n-3 PUFAs—but not n-6 PUFAs—enriched LDL, inhibited the proliferation, and induced apoptosis in human breast cancer cells [74–76]. The n-3 LDL delivered both EPA and DHA to the cells. When these individual fatty acids were delivered to cells by albumin, DHA but not EPA proved effective in stimulating apoptosis in a pathway that involved activation of PPAR γ [75]. The molecular target for both DHA and PPAR γ in these cells was shown to be the heparan sulfate proteoglycan, syndecan-1. Syndecan-1 itself was effective in apoptosis induction and when syndecan-1 was silenced, the ability of DHA to induce apoptosis was completely blocked as it was in the presence of a dominant negative PPAR γ [76]. Moreover, syndecan-1 siRNA was effective in blocking troglitzone-induced apoptosis. Thus, a novel pathway linking n-3 PUFAs to apoptosis in tumor cells is as follows: DHA activates PPAR γ , which results in transcriptional upregulation of the syndecan-1 target gene, and the syndecan-1 protein induces apoptosis (Figure 4). This novel pathway has been confirmed in human prostate cancer cells [96].

Although PPAR γ was not a target for EPA in breast and prostate cancer cells, a recent report has demonstrated that EPA was an effective PPAR γ transactivator in HT-29 human colon cancer cells [165]. In contrast, both EPA and DHA were shown to reduce PPRE reporter activity in an HCT-116 colon cancer cells [166]. DHA has recently

been shown to reduce the growth of human lung cancer cells in a process that was associated with increased PPAR γ protein [167]. These conflicting reports are consistent with data showing selective modulation of PPAR γ by different ligands in different cells [168]. Several other reasons may be proposed for the differential response to DHA and EPA in the breast and prostate tumor cells including (1) PPAR γ activation may be mediated by a unique DHA metabolite rather than DHA itself; (2) there may be a difference in the bioavailability of the two fatty acids following cellular uptake; (3) EPA may be a ligand for or metabolized to a ligand (e.g., 5(S)-hydroxy-eicosapentaenoic acid) for a G protein-coupled receptor that activates ERK and thereby inactivates or in some other way counteracts PPAR γ ; (4) EPA may directly, or after being metabolized, activate other pathways that counteract PPAR γ signaling.

The identification of syndecan-1 as a target gene for PPAR γ in the breast and prostate cancer cells was a novel but not unexpected finding. The syndecan-1 promoter contains a DR-1 element that is recognized by a several members of the nuclear hormone receptor superfamily including PPAR γ . Although there are conflicting reports of a role for syndecan-1 in cancer, the importance of these studies is the identification of a PPAR γ molecular target that is regulated by PUFAs and results in functional response in the tumor cells. As more such targets emerge, we may be able to understand how different dietary fatty acids play divergent roles in cancer.

ACKNOWLEDGMENTS

This work was supported by National Institutes of Health Grants P01CA106742 (IJE, JO) and R01CA115958 (IJE). The content of this manuscript is solely the responsibility of the authors and does not necessarily represent the official views of the National Cancer Institute or the National Institutes of Health.

REFERENCES

- [1] S. Kersten, B. Desvergne, and W. Wahli, “Roles of PPARs in health and disease,” *Nature*, vol. 405, no. 6785, pp. 421–424, 2000.
- [2] C.-H. Lee, P. Olson, and R. M. Evans, “Minireview: lipid metabolism, metabolic diseases, and peroxisome proliferator-activated receptors,” *Endocrinology*, vol. 144, no. 6, pp. 2201–2207, 2003.
- [3] J. N. Feige, L. Gelman, L. Michalik, B. Desvergne, and W. Wahli, “From molecular action to physiological outputs: peroxisome proliferator-activated receptors are nuclear receptors at the crossroads of key cellular functions,” *Progress in Lipid Research*, vol. 45, no. 2, pp. 120–159, 2006.
- [4] G. Lazennec, L. Canaple, D. Saugy, and W. Wahli, “Activation of peroxisome proliferator-activated receptors (PPARs) by their ligands and protein kinase A activators,” *Molecular Endocrinology*, vol. 14, no. 12, pp. 1962–1975, 2000.
- [5] F. Molnár, M. Matilainen, and C. Carlberg, “Structural determinants of the agonist-independent association of human peroxisome proliferator-activated receptors with coactivators,” *The Journal of Biological Chemistry*, vol. 280, no. 28, pp. 26543–26556, 2005.

- [6] L. Michalik, V. Zoete, G. Krey, et al., "Combined simulation and mutagenesis analyses reveal the involvement of key residues for peroxisome proliferator-activated receptor α helix 12 dynamic behavior," *The Journal of Biological Chemistry*, vol. 282, no. 13, pp. 9666–9677, 2007.
- [7] V. Zoete, A. Grosdidier, and O. Michielin, "Peroxisome proliferator-activated receptor structures: ligand specificity, molecular switch and interactions with regulators," *Biochimica et Biophysica Acta*, vol. 1771, no. 8, pp. 915–925, 2007.
- [8] H. E. Xu, M. H. Lambert, V. G. Montana, et al., "Molecular recognition of fatty acids by peroxisome proliferator-activated receptors," *Molecular Cell*, vol. 3, no. 3, pp. 397–403, 1999.
- [9] B. M. Forman, P. Tontonoz, J. Chen, R. P. Brun, B. M. Spiegelman, and R. M. Evans, "15-deoxy- $\Delta^{12,14}$ -prostaglandin J_2 is a ligand for the adipocyte determination factor PPAR γ ," *Cell*, vol. 83, no. 5, pp. 803–812, 1995.
- [10] S. A. Kliewer, S. S. Sundseth, S. A. Jones, et al., "Fatty acids and eicosanoids regulate gene expression through direct interactions with peroxisome proliferator-activated receptors α and γ ," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 94, no. 9, pp. 4318–4323, 1997.
- [11] B. M. Forman, J. Chen, and R. M. Evans, "Hypolipidemic drugs, polyunsaturated fatty acids, and eicosanoids are ligands for peroxisome proliferator-activated receptors α and δ ," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 94, no. 9, pp. 4312–4317, 1997.
- [12] P. Sertznig, M. Seifert, W. Tilgen, and J. Reichrath, "Present concepts and future outlook: function of peroxisome proliferator-activated receptors (PPARs) for pathogenesis, progression, and therapy of cancer," *Journal of Cellular Physiology*, vol. 212, no. 1, pp. 1–12, 2007.
- [13] L. Michalik, B. Desvergne, and W. Wahli, "Peroxisome-proliferator-activated receptors and cancers: complex stories," *Nature Reviews Cancer*, vol. 4, no. 1, pp. 61–70, 2004.
- [14] L. Xu, C. Han, and T. Wu, "A novel positive feedback loop between peroxisome proliferator-activated receptor- δ and prostaglandin E_2 signaling pathways for human cholangiocarcinoma cell growth," *The Journal of Biological Chemistry*, vol. 281, no. 45, pp. 33982–33996, 2006.
- [15] L. Michalik, J. Auwerx, J. P. Berger, et al., "International union of pharmacology. LXI. Peroxisome proliferator-activated receptors," *Pharmacological Reviews*, vol. 58, no. 4, pp. 726–741, 2006.
- [16] A. Krishnan, S. A. Nair, and M. R. Pillai, "Biology of PPAR γ in cancer: a critical review on existing lacunae," *Current Molecular Medicine*, vol. 7, no. 6, pp. 532–540, 2007.
- [17] S. Han and J. Roman, "Peroxisome proliferator-activated receptor γ : a novel target for cancer therapeutics?" *Anti-Cancer Drugs*, vol. 18, no. 3, pp. 237–244, 2007.
- [18] T. Wang, J. Xu, X. Yu, R. Yang, and Z. C. Han, "Peroxisome proliferator-activated receptor γ in malignant diseases," *Critical Reviews in Oncology/Hematology*, vol. 58, no. 1, pp. 1–14, 2006.
- [19] M. B. Sporn, N. Suh, and D. J. Mangelsdorf, "Prospects for prevention and treatment of cancer with selective PPAR γ modulators (SPARMs)," *Trends in Molecular Medicine*, vol. 7, no. 9, pp. 395–400, 2001.
- [20] J. M. Olefsky, "Treatment of insulin resistance with peroxisome proliferator-activated receptor γ agonists," *The Journal of Clinical Investigation*, vol. 106, no. 4, pp. 467–472, 2000.
- [21] R. A. Siddiqui, S. R. Shaikh, L. A. Sech, H. R. Yount, W. Stillwell, and G. P. Zaloga, "Omega 3-fatty acids: health benefits and cellular mechanisms of action," *Mini Reviews in Medicinal Chemistry*, vol. 4, no. 8, pp. 859–871, 2004.
- [22] R. Colomer, J. M. Moreno-Nogueira, P. P. García-Luna, et al., " n -3 fatty acids, cancer and cachexia: a systematic review of the literature," *British Journal of Nutrition*, vol. 97, no. 5, pp. 823–831, 2007.
- [23] H. Sprecher, "Metabolism of highly unsaturated n -3 and n -6 fatty acids," *Biochimica et Biophysica Acta*, vol. 1486, no. 2-3, pp. 219–231, 2000.
- [24] H. Sprecher, "The roles of anabolic and catabolic reactions in the synthesis and recycling of polyunsaturated fatty acids," *Prostaglandins Leukotrienes and Essential Fatty Acids*, vol. 67, no. 2-3, pp. 79–83, 2002.
- [25] C. M. Williams and G. Burdge, "Long-chain n -3 PUFA: plant v. marine sources," *Proceedings of the Nutrition Society*, vol. 65, no. 1, pp. 42–50, 2006.
- [26] E. A. Emken, R. O. Adlof, and R. M. Gulley, "Dietary linoleic acid influences desaturation and acylation of deuterium-labeled linoleic and linolenic acids in young adult males," *Biochimica et Biophysica Acta*, vol. 1213, no. 3, pp. 277–288, 1994.
- [27] G. C. Burdge and S. A. Wootton, "Conversion of α -linolenic acid to eicosapentaenoic, docosapentaenoic and docosahexaenoic acids in young women," *British Journal of Nutrition*, vol. 88, no. 4, pp. 411–420, 2002.
- [28] R. J. Pawlosky, J. R. Hibbeln, J. A. Novotny, and N. Salem Jr., "Physiological compartmental analysis of α -linolenic acid metabolism in adult humans," *Journal of Lipid Research*, vol. 42, no. 8, pp. 1257–1265, 2001.
- [29] R. J. Pawlosky, J. R. Hibbeln, Y. Lin, et al., "Effects of beef- and fish-based diets on the kinetics of n -3 fatty acid metabolism in human subjects," *The American Journal of Clinical Nutrition*, vol. 77, no. 3, pp. 565–572, 2003.
- [30] E. J. Giltay, L. J. G. Gooren, A. W. F. T. Toorians, M. B. Katan, and P. L. Zock, "Docosahexaenoic acid concentrations are higher in women than in men because of estrogenic effects," *The American Journal of Clinical Nutrition*, vol. 80, no. 5, pp. 1167–1174, 2004.
- [31] L. M. Arterburn, E. B. Hall, and H. Oken, "Distribution, interconversion, and dose response of n -3 fatty acids in humans," *The American Journal of Clinical Nutrition*, vol. 83, no. 6, pp. 1467S–1476S, 2006.
- [32] Y. Park and W. S. Harris, "Omega-3 fatty acid supplementation accelerates chylomicron triglyceride clearance," *Journal of Lipid Research*, vol. 44, no. 3, pp. 455–463, 2003.
- [33] D. G. Payan, M. Y. S. Wong, T. Chernov-Rogan, et al., "Alterations in human leukocyte function induced by ingestion of eicosapentaenoic acid," *Journal of Clinical Immunology*, vol. 6, no. 5, pp. 402–410, 1986.
- [34] M. J. James, R. A. Gibson, and L. G. Cleland, "Dietary polyunsaturated fatty acids and inflammatory mediator production," *The American Journal of Clinical Nutrition*, vol. 71, no. 1, pp. 343S–348S, 2000.
- [35] R. A. Gibson, M. A. Neumann, M. J. James, J. S. Hawkes, C. Hall, and L. G. Cleland, "Effect of n -3 and n -6 dietary fats on the lipoxygenase products from stimulated rat neutrophils," *Prostaglandins Leukotrienes and Essential Fatty Acids*, vol. 46, no. 2, pp. 87–91, 1992.
- [36] R. Altmann, M. Hausmann, T. Spöttl, et al., "13-oxo-ODE is an endogenous ligand for PPAR γ in human colonic epithelial cells," *Biochemical Pharmacology*, vol. 74, no. 4, pp. 612–622, 2007.

- [37] S. M. Earles, J. C. Bronstein, D. L. Winner, and A. W. Bull, "Metabolism of oxidized linoleic acid: characterization of 13-hydroxyoctadecadienoic acid dehydrogenase activity from rat colonic tissue," *Biochimica et Biophysica Acta*, vol. 1081, no. 2, pp. 174–180, 1991.
- [38] M. Hamberg, "Stereochemistry of oxygenation of linoleic acid catalyzed by prostaglandin-endoperoxide H synthase-2," *Archives of Biochemistry and Biophysics*, vol. 349, no. 2, pp. 376–380, 1998.
- [39] A. R. Brash, M. Jisaka, W. E. Boeglin, et al., "Investigation of a second 15S-lipoxygenase in humans and its expression in epithelial tissues," *Advances in Experimental Medicine and Biology*, vol. 469, pp. 83–89, 2000.
- [40] J. T. Huang, J. S. Welch, M. Ricote, et al., "Interleukin-4-dependent production of PPAR- γ ligands in macrophages by 12/15-lipoxygenase," *Nature*, vol. 400, no. 6742, pp. 378–382, 1999.
- [41] L. Nagy, P. Tontonoz, J. G. A. Alvarez, H. Chen, and R. M. Evans, "Oxidized LDL regulates macrophage gene expression through ligand activation of PPAR γ ," *Cell*, vol. 93, no. 2, pp. 229–240, 1998.
- [42] J. K. A. Hampel, L. M. Brownrigg, D. Vignarajah, et al., "Differential modulation of cell cycle, apoptosis and PPAR γ 2 gene expression by PPAR γ agonists ciglitazone and 9-hydroxyoctadecadienoic acid in monocytic cells," *Prostaglandins Leukotrienes and Essential Fatty Acids*, vol. 74, no. 5, pp. 283–293, 2006.
- [43] J. T. O'Flaherty, L. C. Rogers, C. M. Paumi, et al., "5-oxo-ETE analogs and the proliferation of cancer cells," *Biochimica et Biophysica Acta*, vol. 1736, no. 3, pp. 228–236, 2005.
- [44] K.-R. Erlemann, J. Rokach, and W. S. Powell, "Oxidative stress stimulates the synthesis of the eosinophil chemoattractant 5-oxo-6,8,11,14-eicosatetraenoic acid by inflammatory cells," *The Journal of Biological Chemistry*, vol. 279, no. 39, pp. 40376–40384, 2004.
- [45] J. T. O'Flaherty, J. F. Cordes, S. L. Lee, M. Samuel, and M. J. Thomas, "Chemical and biological characterization of oxo-eicosatetraenoic acids," *Biochimica et Biophysica Acta*, vol. 1201, no. 3, pp. 505–515, 1994.
- [46] J. T. O'Flaherty, M. Kuroki, A. B. Nixon, et al., "5-oxo-eicosanoids and hematopoietic cytokines cooperate in stimulating neutrophil function and the mitogen-activated protein kinase pathway," *The Journal of Biological Chemistry*, vol. 271, no. 30, pp. 17821–17828, 1996.
- [47] W. S. Powell, F. Gravelle, and S. Gravel, "Metabolism of 5(S)-hydroxy-6,8,11,14-eicosatetraenoic acid and other 5(S)-hydroxyeicosanoids by a specific dehydrogenase in human polymorphonuclear leukocytes," *The Journal of Biological Chemistry*, vol. 267, no. 27, pp. 19233–19241, 1992.
- [48] W. S. Powell, F. Gravelle, and S. Gravel, "Phorbol myristate acetate stimulates the formation of 5-oxo-6,8,11,14-eicosatetraenoic acid by human neutrophils by activating NADPH oxidase," *The Journal of Biological Chemistry*, vol. 269, no. 41, pp. 25373–25380, 1994.
- [49] S. H. Lee, K. Rangiah, M. V. Williams, A. Y. Wehr, R. N. DuBois, and I. A. Blair, "Cyclooxygenase-2-mediated metabolism of arachidonic acid to 15-oxo-eicosatetraenoic acid by rat intestinal epithelial cells," *Chemical Research in Toxicology*, vol. 20, no. 11, pp. 1665–1675, 2007.
- [50] M. Gulliksson, Å. Brunnström, M. Johannesson, et al., "Expression of 15-lipoxygenase type-1 in human mast cells," *Biochimica et Biophysica Acta*, vol. 1771, no. 9, pp. 1156–1165, 2007.
- [51] S. A. Kliewer, J. M. Lenhard, T. M. Willson, I. Patel, D. C. Morris, and J. M. Lehmann, "A prostaglandin J₂ metabolite binds peroxisome proliferator-activated receptor γ and promotes adipocyte differentiation," *Cell*, vol. 83, no. 5, pp. 813–819, 1995.
- [52] P. Sarraf, E. Mueller, W. M. Smith, et al., "Loss-of-function mutations in PPAR γ associated with human colon cancer," *Molecular Cell*, vol. 3, no. 6, pp. 799–804, 1999.
- [53] T. Itoh and K. Yamamoto, "Peroxisome proliferator activated receptor γ and oxidized docosahexaenoic acids as new class of ligand," *Naunyn-Schmiedeberg's Archives of Pharmacology*, vol. 377, no. 4–6, pp. 541–547, 2008.
- [54] O. Nosjean and J. A. Boutin, "Natural ligands of PPAR γ : are prostaglandin J₂ derivatives really playing the part?" *Cellular Signalling*, vol. 14, no. 7, pp. 573–583, 2002.
- [55] B. Zingarelli and J. A. Cook, "Peroxisome proliferator-activated receptor- γ is a new therapeutic target in sepsis and inflammation," *Shock*, vol. 23, no. 5, pp. 393–399, 2005.
- [56] J. U. Scher and M. H. Pillinger, "15d-PGJ₂: the anti-inflammatory prostaglandin?" *Clinical Immunology*, vol. 114, no. 2, pp. 100–109, 2005.
- [57] A. González-Pérez, A. Planagumà, K. Gronert, et al., "Docosahexaenoic acid (DHA) blunts liver injury by conversion to protective lipid mediators: protectin D1 and 17 S-hydroxy-DHA," *The FASEB Journal*, vol. 20, no. 14, pp. 2537–2539, 2006.
- [58] T. Shiraki, N. Kamiya, S. Shiki, T. S. Kodama, A. Kakizuka, and H. Jingami, " α,β -unsaturated ketone is a core moiety of natural ligands for covalent binding to peroxisome proliferator-activated receptor γ ," *The Journal of Biological Chemistry*, vol. 280, no. 14, pp. 14145–14153, 2005.
- [59] B. Rigas and Y. Sun, "Induction of oxidative stress as a mechanism of action of chemopreventive agents against cancer," *British Journal of Cancer*, vol. 98, no. 7, pp. 1157–1160, 2008.
- [60] B. A. Freeman, P. R. S. Baker, F. J. Schopfer, S. R. Woodcock, A. Napolitano, and M. d'Ischia, "Nitro-fatty acid formation and signaling," *The Journal of Biological Chemistry*, vol. 283, no. 23, pp. 15515–15519, 2008.
- [61] P. R. S. Baker, Y. Lin, F. J. Schopfer, et al., "Fatty acid transduction of nitric oxide signaling: multiple nitrated unsaturated fatty acid derivatives exist in human blood and urine and serve as endogenous peroxisome proliferator-activated receptor ligands," *The Journal of Biological Chemistry*, vol. 280, no. 51, pp. 42464–42475, 2005.
- [62] F. J. Schopfer, Y. Lin, P. R. S. Baker, et al., "Nitrolinoleic acid: an endogenous peroxisome proliferator-activated receptor γ ligand," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 102, no. 7, pp. 2340–2345, 2005.
- [63] S. E. O'Sullivan, "Cannabinoids go nuclear: evidence for activation of peroxisome proliferator-activated receptors," *British Journal of Pharmacology*, vol. 152, no. 5, pp. 576–582, 2007.
- [64] A. Lenman and C. J. Fowler, "Interaction of ligands for the peroxisome proliferator-activated receptor γ with the endocannabinoid system," *British Journal of Pharmacology*, vol. 151, no. 8, pp. 1343–1351, 2007.
- [65] M. Bouaboula, S. Hilairret, J. Marchand, L. Fajas, G. Le Fur, and P. Casellas, "Anandamide induced PPAR γ transcriptional activation and 3T3-L1 preadipocyte differentiation," *European Journal of Pharmacology*, vol. 517, no. 3, pp. 174–181, 2005.

- [66] C. M. Paumi, P. K. Smitherman, A. J. Townsend, and C. S. Morrow, "Glutathione S-transferases (GSTs) inhibit transcriptional activation by the peroxisomal proliferator-activated receptor γ (PPAR γ) ligand, 15-deoxy- $\Delta^{12,14}$ -prostaglandin J₂ (15-d-PGJ₂)," *Biochemistry*, vol. 43, no. 8, pp. 2345–2352, 2004.
- [67] C. M. Paumi, M. Wright, A. J. Townsend, and C. S. Morrow, "Multidrug resistance protein (MRP) 1 and MRP3 attenuate cytotoxic and transactivating effects of the cyclopentenone prostaglandin, 15-deoxy- $\Delta^{12,14}$ -prostaglandin J₂ in MCF7 breast cancer cells," *Biochemistry*, vol. 42, no. 18, pp. 5429–5437, 2003.
- [68] R. C. Murphy and S. Zarini, "Glutathione adducts of oxyeicosanoids," *Prostaglandins & Other Lipid Mediators*, vol. 68–69, pp. 471–482, 2002.
- [69] R. O'Connor, "The pharmacology of cancer resistance," *Anticancer Research*, vol. 27, no. 3A, pp. 1267–1272, 2007.
- [70] R. E. Olson, "Discovery of the lipoproteins, their role in fat transport and their significance as risk factors," *The Journal of Nutrition*, vol. 128, no. 2, pp. 439S–443S, 1998.
- [71] S. Vitols, C. Peterson, O. Larsson, P. Holm, and B. Åberg, "Elevated uptake of low density lipoproteins by human lung cancer tissue in vivo," *Cancer Research*, vol. 52, no. 22, pp. 6244–6247, 1992.
- [72] D. F. Lum, K. R. McQuaid, V. L. Gilbertson, and M. Hughes-Fulford, "Coordinate up-regulation of low-density lipoprotein receptor and cyclo-oxygenase-2 gene expression in human colorectal cells and in colorectal adenocarcinoma biopsies," *International Journal of Cancer*, vol. 83, no. 2, pp. 162–166, 1999.
- [73] Y. Chen and M. Hughes-Fulford, "Human prostate cancer cells lack feedback regulation of low-density lipoprotein receptor and its regulator, SREBP2," *International Journal of Cancer*, vol. 91, no. 1, pp. 41–45, 2000.
- [74] I. J. Edwards, I. M. Berquin, H. Sun, et al., "Differential effects of delivery of omega-3 fatty acids to human cancer cells by low-density lipoproteins versus albumin," *Clinical Cancer Research*, vol. 10, no. 24, pp. 8275–8283, 2004.
- [75] H. Sun, I. M. Berquin, and I. J. Edwards, "Omega-3 polyunsaturated fatty acids regulate syndecan-1 expression in human breast cancer cells," *Cancer Research*, vol. 65, no. 10, pp. 4442–4447, 2005.
- [76] H. Sun, I. M. Berquin, R. T. Owens, J. T. O'Flaherty, and I. J. Edwards, "Peroxisome proliferator-activated receptor γ -mediated up-regulation of syndecan-1 by n-3 fatty acids promotes apoptosis of human breast cancer cells," *Cancer Research*, vol. 68, no. 8, pp. 2912–2919, 2008.
- [77] S. Heikkinen, J. Auwerx, and C. A. Argmann, "PPAR γ in human and mouse physiology," *Biochimica et Biophysica Acta*, vol. 1771, no. 8, pp. 999–1013, 2007.
- [78] D. Shao, S. M. Rangwala, S. T. Bailey, S. L. Krakow, M. J. Reginato, and M. A. Lazar, "Interdomain communication regulating ligand binding by PPAR- γ ," *Nature*, vol. 396, no. 6709, pp. 377–380, 1998.
- [79] H. S. Camp, S. R. Tafuri, and T. Leff, "c-Jun N-terminal kinase phosphorylates peroxisome proliferator-activated receptor- γ 1 and negatively regulates its transcriptional activity," *Endocrinology*, vol. 140, no. 1, pp. 392–397, 1999.
- [80] K. A. Burns and J. P. Vanden Heuvel, "Modulation of PPAR activity via phosphorylation," *Biochimica et Biophysica Acta*, vol. 1771, no. 8, pp. 952–960, 2007.
- [81] E. Papageorgiou, N. Pitulis, P. Msaouel, P. Lembessis, and M. Koutsilieris, "The non-genomic crosstalk between PPAR- γ ligands and ERK1/2 in cancer cell lines," *Expert Opinion on Therapeutic Targets*, vol. 11, no. 8, pp. 1071–1085, 2007.
- [82] E. Burgermeister, D. Chuderland, T. Hanoch, M. Meyer, M. Liscovitch, and R. Seger, "Interaction with MEK causes nuclear export and downregulation of peroxisome proliferator-activated receptor γ ," *Molecular and Cellular Biology*, vol. 27, no. 3, pp. 803–817, 2007.
- [83] G. Pascual, A. L. Fong, S. Ogawa, et al., "A SUMOylation-dependent pathway mediates transrepression of inflammatory response genes by PPAR- γ ," *Nature*, vol. 437, no. 7059, pp. 759–763, 2005.
- [84] M. Shimizu, D. Yamashita, T. Yamaguchi, F. Hirose, and T. Osumi, "Aspects of the regulatory mechanisms of PPAR functions: analysis of a bidirectional response element and regulation by sumoylation," *Molecular and Cellular Biochemistry*, vol. 286, no. 1–2, pp. 33–42, 2006.
- [85] J. N. Feige, L. Gelman, C. Tudor, Y. Engelborghs, W. Wahli, and B. Desvergne, "Fluorescence imaging reveals the nuclear behavior of peroxisome proliferator-activated receptor/retinoid X receptor heterodimers in the absence and presence of ligand," *The Journal of Biological Chemistry*, vol. 280, no. 18, pp. 17880–17890, 2005.
- [86] C. Tudor, J. N. Feige, H. Pingali, et al., "Association with coregulators is the major determinant governing peroxisome proliferator-activated receptor mobility in living cells," *The Journal of Biological Chemistry*, vol. 282, no. 7, pp. 4417–4426, 2007.
- [87] E. Powell, P. Kuhn, and W. Xu, "Nuclear receptor cofactors in PPAR γ -mediated adipogenesis and adipocyte energy metabolism," *PPAR Research*, vol. 2007, Article ID 53843, 11 pages, 2007.
- [88] Z. Gao, Q. He, B. Peng, P. J. Chiao, and J. Ye, "Regulation of nuclear translocation of HDAC3 by I κ B α is required for tumor necrosis factor inhibition of peroxisome proliferator-activated receptor γ function," *The Journal of Biological Chemistry*, vol. 281, no. 7, pp. 4540–4547, 2006.
- [89] E. Burgermeister and R. Seger, "MAPK kinases as nucleocytoplasmic shuttles for PPAR γ ," *Cell Cycle*, vol. 6, no. 13, pp. 1539–1548, 2007.
- [90] R. Kostadinova, W. Wahli, and L. Michalik, "PPARs in diseases: control mechanisms of inflammation," *Current Medicinal Chemistry*, vol. 12, no. 25, pp. 2995–3009, 2005.
- [91] F. Zandbergen, S. Mandard, P. Escher, et al., "The G₀/G₁ switch gene 2 is a novel PPAR target gene," *Biochemical Journal*, vol. 392, no. 2, pp. 313–324, 2005.
- [92] B. E. J. Teunissen, P. J. H. Smeets, P. H. M. Willemsen, L. J. De Windt, G. J. Van der Vusse, and M. Van Bilsen, "Activation of PPAR δ inhibits cardiac fibroblast proliferation and the trans-differentiation into myofibroblasts," *Cardiovascular Research*, vol. 75, no. 3, pp. 519–529, 2007.
- [93] D. Bonfiglio, S. Aquila, S. Catalano, et al., "Peroxisome proliferator-activated receptor- γ activates p53 gene promoter binding to the nuclear factor- κ B sequence in human MCF7 breast cancer cells," *Molecular Endocrinology*, vol. 20, no. 12, pp. 3083–3092, 2006.
- [94] T.-C. Ho, S.-L. Chen, Y.-C. Yang, C.-L. Liao, H.-C. Cheng, and Y.-P. Tsao, "PEDF induces p53-mediated apoptosis through PPAR gamma signaling in human umbilical vein endothelial cells," *Cardiovascular Research*, vol. 76, no. 2, pp. 213–223, 2007.
- [95] H. Zand, A. Rhimipour, M. Bakhshayesh, M. Shafiee, I. Nour Mohammadi, and S. Salimi, "Involvement of PPAR- γ and

- p53 in DHA-induced apoptosis in Reh cells," *Molecular and Cellular Biochemistry*, vol. 304, no. 1-2, pp. 71–77, 2007.
- [96] I. J. Edwards, H. Sun, Y. Hu, et al., "In vivo and in vitro regulation of syndecan 1 in prostate cells by n-3 polyunsaturated fatty acids," *The Journal of Biological Chemistry*, vol. 283, no. 26, pp. 18441–18449, 2008.
- [97] J. Liu, H. Wang, Y. Zuo, and S. R. Farmer, "Functional interaction between peroxisome proliferator-activated receptor γ and β -catenin," *Molecular and Cellular Biology*, vol. 26, no. 15, pp. 5827–5837, 2006.
- [98] I. Takada, M. Suzawa, and S. Kato, "Nuclear receptors as targets for drug development: crosstalk between peroxisome proliferator-activated receptor γ and cytokines in bone marrow-derived mesenchymal stem cells," *Journal of Pharmacological Sciences*, vol. 97, no. 2, pp. 184–189, 2005.
- [99] A. Rossi, P. Kapahi, G. Natoli, et al., "Anti-inflammatory cyclopentenone prostaglandins are direct inhibitors of I κ B kinase," *Nature*, vol. 403, no. 6765, pp. 103–118, 2000.
- [100] W. E. Ackerman IV, X. L. Zhang, B. H. Rovin, and D. A. Kniss, "Modulation of cytokine-induced cyclooxygenase 2 expression by PPAR γ ligands through NF κ B signal disruption in human WISH and amnion cells," *Biology of Reproduction*, vol. 73, no. 3, pp. 527–535, 2005.
- [101] D. S. Straus and C. K. Glass, "Cyclopentenone prostaglandins: new insights on biological activities and cellular targets," *Medicinal Research Reviews*, vol. 21, no. 3, pp. 185–210, 2001.
- [102] H. Siavash, N. G. Nikitakis, and J. J. Sauk, "Abrogation of IL-6-mediated JAK signalling by the cyclopentenone prostaglandin 15d-PG $_2$ in oral squamous carcinoma cells," *British Journal of Cancer*, vol. 91, no. 6, pp. 1074–1080, 2004.
- [103] H. Siavash, N. G. Nikitakis, and J. J. Sauk, "Targeting of epidermal growth factor receptor by cyclopentenone prostaglandin 15-deoxy- $\Delta^{12,14}$ -prostaglandin J_2 in human oral squamous carcinoma cells," *Cancer Letters*, vol. 211, no. 1, pp. 97–103, 2004.
- [104] W. Almishri, C. Cossette, J. Rokach, J. G. Martin, Q. Hamid, and W. S. Powell, "Effects of prostaglandin D_2 , 15-deoxy- $\Delta^{12,14}$ -prostaglandin J_2 , and selective DP $_1$ and DP $_2$ receptor agonists on pulmonary infiltration of eosinophils in Brown Norway rats," *Journal of Pharmacology and Experimental Therapeutics*, vol. 313, no. 1, pp. 64–69, 2005.
- [105] J. T. O'Flaherty, J. S. Taylor, and M. J. Thomas, "Receptors for the 5-oxo class of eicosanoids in neutrophils," *The Journal of Biological Chemistry*, vol. 273, no. 49, pp. 32535–32541, 1998.
- [106] C. P. Briscoe, M. Tadayyon, J. L. Andrews, et al., "The orphan G protein-coupled receptor GPR40 is activated by medium and long chain fatty acids," *The Journal of Biological Chemistry*, vol. 278, no. 13, pp. 11303–11311, 2003.
- [107] A. Hirasawa, K. Tsumaya, T. Awaji, et al., "Free fatty acids regulate gut incretin glucagon-like peptide-1 secretion through GPR120," *Nature Medicine*, vol. 11, no. 1, pp. 90–94, 2005.
- [108] S. Katsuma, N. Hatae, T. Yano, et al., "Free fatty acids inhibit serum deprivation-induced apoptosis through GPR120 in a murine enteroendocrine cell line STC-1," *The Journal of Biological Chemistry*, vol. 280, no. 20, pp. 19507–19515, 2005.
- [109] M. Hughes-Fulford, C.-F. Li, J. Boonyaratankornkit, and S. Sayyah, "Arachidonic acid activates phosphatidylinositol 3-kinase signaling and induces gene expression in prostate cancer," *Cancer Research*, vol. 66, no. 3, pp. 1427–1433, 2006.
- [110] C. E. Clay, A. M. Namen, G.-I. Atsumi, et al., "Magnitude of peroxisome proliferator-activated receptor- γ activation is associated with important and seemingly opposite biological responses in breast cancer cells," *Journal of Investigative Medicine*, vol. 49, no. 5, pp. 413–420, 2001.
- [111] O. S. Gardner, B. J. Dewar, and L. M. Graves, "Activation of mitogen-activated protein kinases by peroxisome proliferator-activated receptor ligands: an example of nongenomic signaling," *Molecular Pharmacology*, vol. 68, no. 4, pp. 933–941, 2005.
- [112] M. Abedin, J. Lim, T. B. Tang, D. Park, L. L. Demer, and Y. Tintut, "N-3 fatty acids inhibit vascular calcification via the p38-mitogen-activated protein kinase and peroxisome proliferator-activated receptor- γ pathways," *Circulation Research*, vol. 98, no. 6, pp. 727–729, 2006.
- [113] J. T. O'Flaherty, L. C. Rogers, B. A. Chadwell, et al., "5(S)-hydroxy-6,8,11,14-E,Z,Z,Z-eicosatetraenoate stimulates PC3 cell signaling and growth by a receptor-dependent mechanism," *Cancer Research*, vol. 62, no. 23, pp. 6817–6819, 2002.
- [114] S. S. Palakurthi, H. Aktas, L. M. Grubbisich, R. M. Mortensen, and J. A. Halperin, "Anticancer effects of thiazolidinediones are independent of peroxisome proliferator-activated receptor γ and mediated by inhibition of translation initiation," *Cancer Research*, vol. 61, no. 16, pp. 6213–6218, 2001.
- [115] S. S. Palakurthi, R. Flückiger, H. Aktas, et al., "Inhibition of translation initiation mediates the anticancer effect of the n-3 polyunsaturated fatty acid eicosapentaenoic acid," *Cancer Research*, vol. 60, no. 11, pp. 2919–2925, 2000.
- [116] F. Willermain, S. Dulku, N. S. Gonzalez, et al., "15-deoxy-12,14-prostaglandin J_2 inhibits interferon gamma induced MHC class II but not class I expression on ARPE cells through a PPAR gamma independent mechanism," *Prostaglandins & Other Lipid Mediators*, vol. 80, no. 3-4, pp. 136–143, 2006.
- [117] W. A. Wilmer, C. Dixon, L. Lu, T. Hilbelink, and B. H. Rovin, "A cyclopentenone prostaglandin activates mesangial MAP kinase independently of PPAR γ ," *Biochemical and Biophysical Research Communications*, vol. 281, no. 1, pp. 57–62, 2001.
- [118] D. L. Feinstein, A. Spagnolo, C. Akar, et al., "Receptor-independent actions of PPAR thiazolidinedione agonists: is mitochondrial function the key?" *Biochemical Pharmacology*, vol. 70, no. 2, pp. 177–188, 2005.
- [119] M. Soller, S. Dröse, U. Brandt, B. Brüne, and A. von Knethen, "Mechanism of thiazolidinedione-dependent cell death in Jurkat T cells," *Molecular Pharmacology*, vol. 71, no. 6, pp. 1535–1544, 2007.
- [120] C. L. Chaffer, D. M. Thomas, E. W. Thompson, and E. D. Williams, "PPAR γ -independent induction of growth arrest and apoptosis in prostate and bladder carcinoma," *BMC Cancer*, vol. 6, article 53, pp. 1–13, 2006.
- [121] C. H. MacLean, S. J. Newberry, W. A. Mojica, et al., "Effects of omega-3 fatty acids on cancer risk: a systematic review," *Journal of the American Medical Association*, vol. 295, no. 4, pp. 403–415, 2006.
- [122] L. Kaizer, N. F. Boyd, V. Kriukov, and D. Tritchler, "Fish consumption and breast cancer risk: an ecological study," *Nutrition and Cancer*, vol. 12, no. 1, pp. 61–68, 1989.
- [123] S. D. Hursting, M. Thornquist, and M. M. Henderson, "Types of dietary fat and the incidence of cancer at five sites," *Preventive Medicine*, vol. 19, no. 3, pp. 242–253, 1990.
- [124] S. Sasaki, M. Horacsek, and H. Kesteloot, "An ecological study of the relationship between dietary fat intake and breast cancer mortality," *Preventive Medicine*, vol. 22, no. 2, pp. 187–202, 1993.

- [125] P. D. Terry, J. B. Terry, and T. E. Rohan, "Long-chain (n-3) fatty acid intake and risk of cancers of the breast and the prostate: recent epidemiological studies, biological mechanisms, and directions for future research," *The Journal of Nutrition*, vol. 134, no. 12, pp. 3412S–3420S, 2004.
- [126] M. F. Leitzmann, M. J. Stampfer, D. S. Michaud, et al., "Dietary intake of n-3 and n-6 fatty acids and the risk of prostate cancer," *The American Journal of Clinical Nutrition*, vol. 80, no. 1, pp. 204–216, 2004.
- [127] C. P. Caygill, A. Charlett, and M. J. Hill, "Fat, fish, fish oil and cancer," *British Journal of Cancer*, vol. 74, no. 1, pp. 159–164, 1996.
- [128] C.-C. Yeh, L.-L. Hsieh, R. Tang, C. R. Chang-Chieh, and F.-C. Sung, "Risk factors for colorectal cancer in Taiwan: a hospital-based case-control study," *Journal of the Formosan Medical Association*, vol. 102, no. 5, pp. 305–312, 2003.
- [129] C.-X. Yang, T. Takezaki, K. Hirose, M. Inoue, X.-E. Huang, and K. Tajima, "Fish consumption and colorectal cancer: a case-reference study in Japan," *European Journal of Cancer Prevention*, vol. 12, no. 2, pp. 109–115, 2003.
- [130] N. Simonsen, P. van't Veer, J. J. Strain, et al., "Adipose tissue omega-3 and omega-6 fatty acid content and breast cancer in the euramic study," *American Journal of Epidemiology*, vol. 147, no. 4, pp. 342–352, 1998.
- [131] G. Mamalakis, A. Kafatos, N. Kalogeropoulos, N. Andrikopoulos, G. Daskalopoulos, and A. Kranidis, "Prostate cancer vs hyperplasia: relationships with prostatic and adipose tissue fatty acid composition," *Prostaglandins Leukotrienes and Essential Fatty Acids*, vol. 66, no. 5-6, pp. 467–477, 2002.
- [132] V. L. Freeman, M. Meydani, K. Hur, and R. C. Flanigan, "Inverse association between prostatic polyunsaturated fatty acid and risk of locally advanced prostate carcinoma," *Cancer*, vol. 101, no. 12, pp. 2744–2754, 2004.
- [133] Y. J. Yang, S. H. Lee, S. J. Hong, and B. C. Chung, "Comparison of fatty acid profiles in the serum of patients with prostate cancer and benign prostatic hyperplasia," *Clinical Biochemistry*, vol. 32, no. 6, pp. 405–409, 1999.
- [134] A. E. Norrish, C. M. Skeaff, G. L. B. Arribas, S. J. Sharpe, and R. T. Jackson, "Prostate cancer risk and consumption of fish oils: a dietary biomarker-based case-control study," *British Journal of Cancer*, vol. 81, no. 7, pp. 1238–1242, 1999.
- [135] B. S. Reddy, L. A. Cohen, G. D. McCoy, P. Hill, J. H. Weisburger, and E. L. Wynder, "Nutrition and its relationship to cancer," *Advances in Cancer Research*, vol. 32, pp. 237–345, 1980.
- [136] J. J. Jurkowski and W. T. Cave Jr., "Dietary effects of menhaden oil on the growth and membrane lipid composition of rat mammary tumors," *Journal of the National Cancer Institute*, vol. 74, no. 5, pp. 1145–1150, 1985.
- [137] L. M. Braden and K. K. Carroll, "Dietary polyunsaturated fat in relation to mammary carcinogenesis in rats," *Lipids*, vol. 21, no. 4, pp. 285–288, 1986.
- [138] D. P. Rose, M. A. Hatala, J. M. Connolly, and J. Rayburn, "Effect of diets containing different levels of linoleic acid on human breast cancer growth and lung metastasis in nude mice," *Cancer Research*, vol. 53, no. 18, pp. 4686–4690, 1993.
- [139] D. P. Rose and J. M. Connolly, "Effects of dietary omega-3 fatty acids on human breast cancer growth and metastases in nude mice," *Journal of the National Cancer Institute*, vol. 85, no. 21, pp. 1743–1747, 1993.
- [140] D. P. Rose, J. M. Connolly, J. Rayburn, and M. Coleman, "Influence of diets containing eicosapentaenoic or docosahexaenoic acid on growth and metastasis of breast cancer cells in nude mice," *Journal of the National Cancer Institute*, vol. 87, no. 8, pp. 587–592, 1995.
- [141] F. Cannizzo Jr. and S. A. Broitman, "Postpromotional effects of dietary marine or safflower oils on large bowel or pulmonary implants of CT-26 in mice," *Cancer Research*, vol. 49, no. 15, pp. 4289–4294, 1989.
- [142] M. Iigo, T. Nakagawa, C. Ishikawa, et al., "Inhibitory effects of docosahexaenoic acid on colon carcinoma 26 metastasis to the lung," *British Journal of Cancer*, vol. 75, no. 5, pp. 650–655, 1997.
- [143] M. D. Boudreau, K. H. Sohn, S. H. Rhee, S. W. Lee, J. D. Hunt, and D. H. Hwang, "Suppression of tumor cell growth both in nude mice and in culture by n-3 polyunsaturated fatty acids: mediation through cyclooxygenase-independent pathways," *Cancer Research*, vol. 61, no. 4, pp. 1386–1391, 2001.
- [144] B. S. Reddy and H. Maruyama, "Effect of dietary fish oil on azoxymethane-induced colon carcinogenesis in male F344 rats," *Cancer Research*, vol. 46, no. 7, pp. 3367–3370, 1986.
- [145] T. Minoura, T. Takata, M. Sakaguchi, et al., "Effect of dietary eicosapentaenoic acid on azoxymethane-induced colon carcinogenesis in rats," *Cancer Research*, vol. 48, no. 17, pp. 4790–4794, 1988.
- [146] B. S. Reddy and S. Sugie, "Effect of different levels of omega-3 and omega-6 fatty acids on azoxymethane-induced colon carcinogenesis in F344 rats," *Cancer Research*, vol. 48, no. 23, pp. 6642–6647, 1988.
- [147] M. Takahashi, T. Minamoto, N. Yamashita, T. Kato, K. Yazawa, and H. Esumi, "Effect of docosahexaenoic acid on azoxymethane-induced colon carcinogenesis in rats," *Cancer Letters*, vol. 83, no. 1-2, pp. 177–184, 1994.
- [148] C. W. Hendrickse, M. R. Keighley, and J. P. Neoptolemos, "Dietary ω -3 fats reduce proliferation and tumor yields at colorectal anastomosis in rats," *Gastroenterology*, vol. 109, no. 2, pp. 431–439, 1995.
- [149] N. Onogi, M. Okuno, C. Komaki, et al., "Suppressing effect of perilla oil on azoxymethane-induced foci of colonic aberrant crypts in rats," *Carcinogenesis*, vol. 17, no. 6, pp. 1291–1296, 1996.
- [150] M. Takahashi, M. Fukutake, T. Isoi, et al., "Suppression of azoxymethane-induced rat colon carcinoma development by a fish oil component, docosahexaenoic acid (DHA)," *Carcinogenesis*, vol. 18, no. 7, pp. 1337–1342, 1997.
- [151] J. E. Paulsen, T. Stamm, and J. Alexander, "A fish oil-derived concentrate enriched in eicosapentaenoic and docosahexaenoic acid as ethyl esters inhibits the formation and growth of aberrant crypt foci in rat colon," *Pharmacology & Toxicology*, vol. 82, no. 1, pp. 28–33, 1998.
- [152] R. A. Karmali, P. Reichel, L. A. Cohen, et al., "The effects of dietary ω -3 fatty acids on the DU-145 transplantable human prostatic tumor," *Anticancer Research*, vol. 7, no. 6, pp. 1173–1179, 1987.
- [153] D. P. Rose and L. A. Cohen, "Effects of dietary menhaden oil and retinyl acetate on the growth of DU 145 human prostatic adenocarcinoma cells transplanted into athymic nude mice," *Carcinogenesis*, vol. 9, no. 4, pp. 603–605, 1988.
- [154] N. Kobayashi, R. J. Barnard, S. M. Henning, et al., "Effect of altering dietary ω -6/ ω -3 fatty acid ratios on prostate cancer membrane composition, cyclooxygenase-2, and prostaglandin E₂," *Clinical Cancer Research*, vol. 12, no. 15, pp. 4662–4670, 2006.

- [155] I. M. Berquin, Y. Min, R. Wu, et al., "Modulation of prostate cancer genetic risk by omega-3 and omega-6 fatty acids," *The Journal of Clinical Investigation*, vol. 117, no. 7, pp. 1866–1875, 2007.
- [156] D. P. Rose and J. M. Connolly, "Omega-3 fatty acids as cancer chemopreventive agents," *Pharmacology & Therapeutics*, vol. 83, no. 3, pp. 217–244, 1999.
- [157] G. E. Caughey, E. Mantzioris, R. A. Gibson, L. G. Cleland, and M. J. James, "The effect on human tumor necrosis factor α and interleukin 1β production of diets enriched in n-3 fatty acids from vegetable oil or fish oil," *The American Journal of Clinical Nutrition*, vol. 63, no. 1, pp. 116–122, 1996.
- [158] R. I. Sperling, A. I. Benincaso, C. T. Knoell, J. K. Larkin, K. F. Austen, and D. R. Robinson, "Dietary ω -3 polyunsaturated fatty acids inhibit phosphoinositide formation and chemotaxis in neutrophils," *The Journal of Clinical Investigation*, vol. 91, no. 2, pp. 651–660, 1993.
- [159] P. Needleman, A. Raz, and M. S. Minkes, "Triene prostaglandins: prostacyclin and thromboxane biosynthesis and unique biological properties," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 76, no. 2, pp. 944–948, 1979.
- [160] E. Mueller, P. Sarraf, P. Tontonoz, et al., "Terminal differentiation of human breast cancer through PPAR γ ," *Molecular Cell*, vol. 1, no. 3, pp. 465–470, 1998.
- [161] E. Mueller, M. Smith, P. Sarraf, et al., "Effects of ligand activation of peroxisome proliferator-activated receptor γ in human prostate cancer," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 97, no. 20, pp. 10990–10995, 2000.
- [162] E. Elstner, C. Müller, K. Koshizuka, et al., "Ligands for peroxisome proliferator-activated receptor and retinoic acid receptor inhibit growth and induce apoptosis of human breast cancer cells in vitro and in BNX mice," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 95, no. 15, pp. 8806–8811, 1998.
- [163] E. Elstner, E. A. Williamson, C. Zang, et al., "Novel therapeutic approach: ligands for PPAR γ and retinoid receptors induce apoptosis in bcl-2-positive human breast cancer cells," *Breast Cancer Research and Treatment*, vol. 74, no. 2, pp. 155–165, 2002.
- [164] T. Takashima, Y. Fujiwara, K. Higuchi, et al., "PPAR-gamma ligands inhibit growth of human esophageal adenocarcinoma cells through induction of apoptosis, cell cycle arrest and reduction of ornithine decarboxylase activity," *International Journal of Oncology*, vol. 19, no. 3, pp. 465–471, 2001.
- [165] C. D. Allred, D. R. Talbert, R. C. Southard, X. Wang, and M. W. Kilgore, "PPAR γ 1 as a molecular target of eicosapentaenoic acid in human colon cancer (HT-29) cells," *The Journal of Nutrition*, vol. 138, no. 2, pp. 250–256, 2008.
- [166] J. Y. Lee and D. H. Hwang, "Docosahexaenoic acid suppresses the activity of peroxisome proliferator-activated receptors in a colon tumor cell line," *Biochemical and Biophysical Research Communications*, vol. 298, no. 5, pp. 667–674, 2002.
- [167] A. Trombetta, M. Maggiora, G. Martinasso, P. Cotogni, R. A. Canuto, and G. Muzio, "Arachidonic and docosahexaenoic acids reduce the growth of A549 human lung-tumor cells increasing lipid peroxidation and PPARs," *Chemico-Biological Interactions*, vol. 165, no. 3, pp. 239–250, 2007.
- [168] C. D. Allred and M. W. Kilgore, "Selective activation of PPAR γ in breast, colon, and lung cancer cell lines," *Molecular and Cellular Endocrinology*, vol. 235, no. 1-2, pp. 21–29, 2005.