

BMJ Open Systematic review: unmet supportive care needs in people diagnosed with chronic liver disease

Patricia C Valery,^{1,2} Elizabeth Powell,^{3,4} Neta Moses,⁵ Michael L Volk,⁶ Steven M McPhail,^{7,8} Paul J Clark,^{2,4,9} Jennifer Martin^{5,10}

To cite: Valery PC, Powell E, Moses N, *et al.* Systematic review: unmet supportive care needs in people diagnosed with chronic liver disease. *BMJ Open* 2015;**5**: e007451. doi:10.1136/bmjopen-2014-007451

► Prepublication history for this paper is available online. To view these files please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2014-007451>).

Received 15 December 2014
Revised 22 February 2015
Accepted 9 March 2015

ABSTRACT

Objective: People with chronic liver disease, particularly those with decompensated cirrhosis, experience several potentially debilitating complications that can have a significant impact on activities of daily living and quality of life. These impairments combined with the associated complex treatment mean that they are faced with specific and high levels of supportive care needs. We aimed to review reported perspectives, experiences and concerns of people with chronic liver disease worldwide. This information is necessary to guide development of policies around supportive needs screening tools and to enable prioritisation of support services for these patients.

Design: Systematic searches of PubMed, MEDLINE, CINAHL and PsycINFO from the earliest records until 19 September 2014. Data were extracted using standardised forms. A qualitative, descriptive approach was utilised to analyse and synthesise data.

Results: The initial search yielded 2598 reports: 26 studies reporting supportive care needs among patients with chronic liver disease were included, but few of them were *patient-reported* needs, none used a validated liver disease-specific supportive care need assessment instrument, and only three included patients with cirrhosis. Five key domains of supportive care needs were identified: informational or educational (eg, educational material, educational sessions), practical (eg, daily living), physical (eg, controlling pruritus and fatigue), patient care and support (eg, support groups), and psychological (eg, anxiety, sadness).

Conclusions: While several key domains of supportive care needs were identified, most studies included hepatitis patients. There is a paucity of literature describing the supportive care needs of the chronic liver disease population likely to have the most needs—namely those with cirrhosis. Assessing the supportive care needs of people with chronic liver disease have potential utility in clinical practice for facilitating timely referrals to support services.

INTRODUCTION

Chronic liver disease (CLD) is a major global cause of morbidity and mortality. The prevalence of CLD differs between countries,

Strengths and limitations of this study

- This systematic review comprehensively summarises reported perspectives, experiences and concerns of people who have been diagnosed with chronic liver disease (CLD) worldwide.
- The data presented highlight the shortage of information regarding unmet needs of patients with CLD.
- While several key domains of supportive care needs were identified, most studies included patients with hepatitis, few of them reported *patient* needs, and none used a validated, liver disease-specific supportive care needs assessment instrument. There is a paucity of literature describing the supportive care needs of the CLD population likely to have the most needs—namely those with cirrhosis.
- Assessing the supportive care needs of people with CLD have potential utility in clinical practice for facilitating timely referrals to support services.

affecting approximately 300 million people in China,¹ 29 million in the European Union² and more than 8 million cases in Australia.³ The leading causes of CLD are viral hepatitis B and C (HCV), harmful alcohol consumption and metabolic fatty liver disease associated with obesity and type 2 diabetes.^{1–7} Regardless of aetiology, most of the morbidity and mortality from CLD occurs among individuals with cirrhosis, who are at risk of developing complications including ascites, hepatic encephalopathy, variceal haemorrhage and liver cancer. Liver disease is the fifth greatest cause of death in the UK, where the average age of death from liver disease is 59 years, with large impacts on loss of quality adjusted life years.⁸ The number of individuals with advanced liver disease, liver-related deaths and healthcare costs are predicted to increase over the next decade.^{9–11} Despite these alarming predictions, there is inadequate awareness of the



CrossMark

For numbered affiliations see end of article.

Correspondence to

Dr Patricia C Valery;
Patricia.Valery@menzies.edu.au

disease among the general public and health professionals, and many healthcare systems lack regional or national strategies to address or prevent the increasing burden from complicated CLD.^{3 8}

People with cirrhosis must follow a complex and variable regimen of dietary restrictions, medications, laboratory testing and clinic visits. In addition, patients with decompensated cirrhosis frequently suffer debilitating complications that impact on an individual's quality of life (QoL) and activities of daily living. These impairments combined with the complex management of advanced liver disease are likely to mean that patients are faced with specific and high levels of supportive care needs.¹² In contrast to other advanced end-organ disease, such as heart failure or chronic obstructive pulmonary disease, the potential of devolved models of supportive care in the community or home for patients with CLD is yet to be established. Because hospital care traditionally focuses on medical management of the major complications of portal hypertension such as ascites and variceal bleeding, it is likely that many patients' supportive care needs remain unmet.

The term supportive care needs encompasses the physical, informational, emotional, practical, social and spiritual needs of an individual with chronic disease.¹³ Health needs assessment instruments are increasingly being developed to evaluate specific areas and magnitude of need as a means of improving provision of patient care and outcomes, particularly in the arena of chronic diseases such as cancer and cardiac failure.^{14–18} Advances in medical care have resulted in people with CLD living longer, through better management of disease complications.¹⁹ An imperative exists for a valid and reliable measure that can provide an accurate supportive care needs assessment for people with CLD.

This systematic review addresses the following questions: (1) What are the supportive care needs of people who have been diagnosed with CLD? (2) What are the domains and specific items of need most frequently reported as unmet by patients with CLD, and what is the extent of these needs? and (3) What are the measures for assessment of unmet supportive care needs of people who have been diagnosed with CLD available in the literature? This information is necessary to guide development of policies around supportive needs screening tools and to enable prioritisation of support services for patients with CLD.

METHODS

A systematic review was undertaken to review and synthesise studies investigating the supportive care needs of people diagnosed with CLD. One author (PCV) searched online peer-reviewed journal articles indexed in PubMed, MEDLINE, CINAHL and PsycINFO from the earliest records until 19 September 2014. Titles and abstracts were searched for possible combinations of the terms including chronic liver disease, or chronic

hepatitis, hepatitis, non-alcoholic fatty liver disease, or NAFLD, or cirrhosis, alcoholic liver disease, and unmet need, or support needs, or supportive care needs, or perceived needs, or supportive care, or needs assessment. The search was complemented by manually reviewing the references of retrieved articles for other articles of potential relevance to the research aims.

Two investigators (PCV and JM) independently reviewed all titles; those judged to be potentially helpful were examined. Data were extracted using standardised data collection forms on to a Microsoft Excel spreadsheet (Microsoft Corp, Redmond, Washington, USA). The form included record number, title, year of publication, and abstract for each study, and outcome (inclusion/exclusion) and reason for exclusion were extracted. The following selection criteria were then applied: availability of an abstract; use of primary data; published in English, Spanish or Portuguese; reporting patients' views, perspectives, experiences, concerns; patients were adults or children; and, if reporting on patients with cystic fibrosis, the article had to focus primarily on liver disease. We excluded reviews and editorials and reports for hepatitis A, liver transplant, hepatocellular carcinoma and studies among people with cystic fibrosis that primarily examined respiratory or pancreatic disease. In particular, hepatocellular carcinoma was excluded as a diagnosis of cancer engenders specific supportive care needs around the cancer treatments and side effects. We also excluded papers where patient-reported supportive care needs (as opposed to doctor or carer-reported patient needs) were not investigated and those solely focusing on QoL.

As seen in other disease settings (eg, cancer), although QoL measures provide important insights into the problems experienced by patients, they do not reveal what patients ideally want from the healthcare system or the extent to which their needs are being satisfied.²⁰ Furthermore, they fail to link patient's experience directly with their service desires.

Abstracts with relevant content were selected for full manuscript review. PCV and JM independently reviewed manuscripts; two other investigators (EP and NM) were available to adjudicate disagreements. Data were extracted using predesigned forms on to a Microsoft Excel spreadsheet. The following data were extracted for each study: author, year published, country, disease group, study design, data collection method, survey used for data collection (Was it validated? If yes, the details about validation were collected), number of study participants, inclusion criteria, response rate, summary of findings (key points with regard to perspectives, experiences and concerns of people with CLD), main focus of the paper (eg, needs assessment, patients' experiences) and study limitations. Data extraction was conducted by a single reviewer (PCV), and independently verified by a second reviewer (JM; outcome (inclusion/exclusion) and reason for exclusion were recorded). Study quality was assessed against the following quality criteria: aims

and/or research question stated clearly, study design, recruitment strategy stated clearly, data collection methods, analysis (sufficient presentation of data to permit assessment of analysis), and response rate and assessment of confounding for quantitative studies only. We decided to exclude from our review studies not describing at least three of these criteria. Failure to mention the quality measure in the articles was considered a failure to fulfil the criterion. A qualitative, descriptive approach was utilised to analyse and synthesise data with the reporting of most of the aforementioned details in [tables 1](#) and [2](#).

RESULTS

The initial search yielded 2598 reports: 17 were relevant for our overview, 9 additional studies were found after reviewing the reference lists of relevant articles ([figure 1](#); [tables 1](#) and [2](#)), with a total of 26 articles included in the review. Most studies included patients with hepatitis C (n=19), three included patients with cirrhosis, two with hepatitis B or C, and two included patients with hepatitis or CLD. Of studies including patients with hepatitis, in addition to the confirmation of diagnosis, in three studies participants had to be on combination therapy (interferon and ribavirin).^{21–23} Of the aforementioned studies, five included a mixed group of participants (not only patients). Jessop *et al*²⁴ included members of hepatitis and/or liver support groups; these groups included not just patients with hepatitis, but families, friends and, in some cases, patients with other liver diseases. Rakoski *et al*¹² included elderly people diagnosed with cirrhosis and an age-matched comparison group. Sgorbini *et al*²³ and Bajaj *et al*²⁵ included patients (hepatitis C and cirrhosis, respectively), and their partners or carers. Jennings²⁶ included people who had abnormal laboratory tests and who were referred for further testing for diagnosis of hepatitis C. Rakoski *et al*¹² was the only group reporting information separately for cases and controls.

Fourteen studies used quantitative methodology ([table 1](#)), 12 were cross-sectional, 1 was longitudinal and 1 was a quasi-experimental study. Sample size varied between studies with a range from 36 to 462 patients. Twelve studies (sample size range 5–70 patients) used qualitative methodology ([table 2](#)) to describe patients' experiences, concerns, supportive care needs, perspectives of care, and information and knowledge about their disease. Four used focus groups to collect data, seven used semistructured or unstructured/in-depth interviews, and one used both focus groups and individual in-depth interviews. Many of the qualitative studies began the interview with an open-ended general question about the patients' experience with their disease, followed up by questions or prompts addressing specific areas or topics of interest (eg, stigma, treatment).^{23 27–30}

In reviewing the literature, five common key domains of supportive care needs were identified: 'informational or educational' (eg, educational material, educational

sessions), 'practical' (eg, daily living, financial support), 'physical' (eg, reducing abdominal distention, controlling pruritus and fatigue), 'patient care and support' (eg, support group at the clinic, caregiver support group) and 'psychological' (eg, anxiety, sadness; [table 3](#)).

Most studies reported unmet needs or concerns in the 'informational or educational' domain (15 out of 26 studies). Need for or lack of information about their disease, treatment and tests, controlling symptoms, and disease transmission was among commonly reported concerns reported by 11.^{24 26 28–36} Of the five reports using a supportive care needs assessment tool to collect the data ([table 3](#)), information needs was a common concern in four studies.^{21 26 31 37 38} Only two studies^{21 38} reported that patients perceived themselves as having good support with regard to informational needs. Temple-Smith *et al*³⁹ report mixed results for men and women, with the former denying need for information while the latter were willing to seek health information to better manage their disease.

Other domains of unmet needs or concerns were also reported. Items pertained in the 'patient care and support' domain were reported by eight studies, including access to a pharmacist, a nutritionist, support groups.^{29 31 33 36 38–41} 'Practical needs' including financial stress (cost of care, assistance for obtaining drug coverage plans for medication, worried about being able to provide for their family impact on activities of daily living (eg, dressing, bathing)) were reported by seven studies.^{12 21 23 25 27 31 34 42} Concerns about symptoms, treatment and prognosis, disease transmission (routes of infection, infecting others) were reported by seven studies (here grouped as 'physical' domain).^{23 37 40 42–44} In seven studies, patients reported fear, anxiety, sadness, feelings of isolation, or reported desire for access to psychological counselling ('psychological' domain).^{22 23 30 34 37 38 45}

Of the 14 quantitative studies, 5 used a supportive care needs assessment tool to collect the data.^{21 26 31 37} Balfour *et al*³¹ used the Hepatitis C Needs Assessment Scale (HCNAS), an 11-item self-reported tool developed for their study. For the HCNAS, patients were asked to rank the importance of their healthcare needs on a five-point scale. Zandi *et al*³⁷ used a needs assessment tool that consisted of a list of 20 questions related to common symptoms and management (eg, fatigue, itching, dry mouth, muscular cramps, dietary regimen). Chang *et al* used the Inventory of Socially Supportive Behaviours (ISSB),⁴⁶ a 15-item questionnaire that asks patients to rate on a four-point scale four types of support: emotional, appraisal, informational and tangible.²¹ Jennings²⁶ used a survey consisting of 13 questions including items on the educational needs of patients with HCV (educational delivery methods, interest in support groups, topics of interest related to HCV and preferred services relating to HCV). There was no evidence on literature review of further validation of any of the other needs assessment instruments. Grogan and Timmins³⁸ used a validated survey tool (a 59-item

Table 1 Basic characteristics (setting, disease group, study design, participant characteristics and data collection methods) and main findings including domains of commonly reported supportive care needs items of quantitative studies (n=14) included in the literature review

Author (year published) and country	Study design and study participants	Disease group	Data collection methods and survey used	Focus of the paper, summary of findings and domains of supportive care needs
Gifford <i>et al</i> (2003) ⁴⁰ Australia	Cross-sectional study (462 female patients)	Hepatitis C	Self-completed structured questionnaire on diagnosis, use of health/social services, social support, medical care/treatment, disclosure and discrimination, sexual health and reproductive, caring for children, SF-12 health-related QoL, 3 items from the Hepatitis QoL instrument and general health. Overall response rate was 75%	<i>Patients experiences</i> —48% of the women reported having received less favourable treatment by health professionals because of their HCV status, 58% reported ever having experienced symptoms, 48% rated their health as 'fair-poor'. At diagnosis, 39% of the women were referred to a specialist and 28% were offered information about support groups. At the time of the survey, 56% of the women were currently seeing a doctor specifically for their HCV, but only half were satisfied with their current level of medical care <i>Domains: physical, patient care and support Supportive and information needs</i> —The main motivation for joining the group was to get information or support (all respondents received support from the group, most received support 'routinely' or 'often'). Most respondents were somewhat satisfied with the information received from their doctor, but reported receiving more 'useful' information from the support group. Information needs included available treatments, diet and doctor-patient communication <i>Domains: informational/educational Needs assessment</i> —Over half the patients reported that their knowledge of HCV was inadequate. Receiving information about HCV was regarded as 'important-very important' for most patients. Having access to a pharmacist, a nutritionist, psychological counselling and assistance for obtaining drug coverage plans for antiviral therapy were also regarded as 'important'/'very important' for over two-third of the patients <i>Domains: informational/educational, patient care and support</i>
Jessop <i>et al</i> (2004) ²⁴ USA	Cross-sectional study (88 support group members—patients, family and friends)	Hepatitis or chronic liver disease	Self-completed structured questionnaire motivation for joining the support group, duration of participation, supportive and informational needs, supportive and informational aspects of the group, and lifestyle or treatment changes made as result of membership. The response rates were: 73% for the in-person support group, and unknown for the online group	<i>Patients experiences</i> —Half the men reported experiencing HCV-related symptoms, 35%
Balfour <i>et al</i> (2004) ³¹ Canada	Cross-sectional study (111 patients)	Hepatitis C	Self-completed structured questionnaire. Questions included the Hepatitis C Needs Assessment Scale (developed for this study) and Patient Satisfaction Questionnaires. Patients completed interviews at enrolment and 10 months later. 111 out of 148 eligible patients responded to the questionnaire	<i>Patients experiences</i> —Half the men reported experiencing HCV-related symptoms, 35%
Gifford <i>et al</i> (2005) ⁴² Australia	Cross-sectional study (312 male patients)	Hepatitis C	Self-completed structured questionnaire on diagnosis, use of health and social services,	<i>Patients experiences</i> —Half the men reported experiencing HCV-related symptoms, 35%

Continued

Table 1 Continued

Author (year and country)	Study design and study participants	Disease group	Data collection methods and survey used	Focus of the paper, summary of findings and domains of supportive care needs
Zandi <i>et al</i> (2005) ³⁷ Iran	Quasi-experimental study (44 patients)	Cirrhosis	social support, medical care/treatment, disclosure and discrimination, sexual relationships, SF-12 health-related QoL, 3 items from the Hepatitis QoL instrument and general health. Response rate was 54%	rated their health as 'fair/poor', 40% believed they received less favourable treatment from healthcare professionals than those without HCV, 34% were satisfied with the way they were told about their diagnosis, 26% were offered information about HCV support groups, 36% were worried often/very often about being able to provide for their family, and 44% reported being concerned about feeling physically unable to work <i>Domains:</i> physical, practical <i>Needs assessment</i> —The most commonly reported needs were: controlling/reducing abdominal distention, curative ways in cirrhosis (being treatable/not treatable), ways of controlling symptoms (eg, fatigue, pruritus), principles of care and proper medications, worry, patterns of activity, rest and sleep; routes of transmission as well as diagnostic tests and procedures <i>Domains:</i> physical, informational/educational, psychological <i>Patient concerns</i> —The most common volunteered concerns were disease progression (27%) followed by premature death (19%), infecting family members (13%) and side effects of treatment (11%). From the list of potential concerns, the highest priority was given to infecting family members followed by developing liver cancer, infecting others and developing cirrhosis <i>Domains:</i> physical <i>Need for treatment and information</i> —The need for specific treatment was reported by 60% of patients. A demand for more detailed information about HCV was expressed by 90% of the patients. The amount of knowledge they possessed was directly proportional to their schooling <i>Domains:</i> patient care and support, informational/educational
Minuk <i>et al</i> (2005) ⁴⁴ Canada	Cross-sectional study (185 patients)	Hepatitis C	<i>Face-to-face semistructured interviews.</i> Initial open-ended question about their principal concern regarding their HCV infection (volunteered concern). Patients were then asked to prioritise from a list of 7 potential concerns. No patient refused to be interviewed	
Fabris <i>et al</i> (2006) ³³ Italy	Cross-sectional study (364 patients)	Hepatitis C	Multiple-choice <i>self-completed structured questionnaire</i> on source of infection, impact on family, sex life, diet and alcohol consumption, and psychological status, and the need for treatment and information about HCV. Response rate not reported	

Continued

Table 1 Continued

Author (year published) and country	Study design and study participants	Disease group	Data collection methods and survey used	Focus of the paper, summary of findings and domains of supportive care needs
Alizadeh <i>et al</i> (2008) ⁴³ Iran	Cross-sectional study (36 patients)	Hepatitis B and C	Face-to-face semistructured interviews. Initial open-ended question about patients' main concerns regarding their disease. Patients were then asked to rank 8 potential concerns (eg, liver cancer/cirrhosis) No patient refused to be interviewed Self-completed structured questionnaire. Questions included the HQLQ and ISSB questionnaire. The response rate was 86%	<i>Patient concerns</i> —Concerns about viral transmission, side effects of treatment and disease progression to cirrhosis were the most commonly reported <i>Domains</i> : physical <i>QoL and social support</i> —In general, patients included perceived themselves as having relatively good support (appraisal, emotional, informational and tangible). Almost half the patients reported moderate to severe financial stress, and they were found to have significantly poorer QoL during treatment compared with those without financial stress <i>Domains</i> : practical <i>Supportive and information needs</i> —Overall, patients were very satisfied with support received from the nurse specialist; 57% reported that their needs were met and 76% that support received helped them cope with their treatment better. Most patients agreed that the nurse provided informational support. Items fewer patients felt supported with included counselling related to mood disorders and depression, sleep management, information about support groups, and ongoing support postcompletion of treatment <i>Domains</i> : psychological, patient care and support
Chang <i>et al</i> (2008) ²¹ Taiwan	Cross-sectional study (115 patients)	Hepatitis C	Self-completed structured questionnaire on means of contracting HCV, symptoms and treatment success. Using a six-point Likert scale patients rated their level of satisfaction with information and psychological support received from the nurse specialist. The response rate was 53%	<i>Emotional and financial burden on patients and caregivers</i> —The effect of the financial burden was seen on medical adherence (missed appointments or procedures, did not take or took less prescribed medications) and was associated with the severity of liver disease <i>Domains</i> : practical <i>Patients understanding and needs</i> —7% of the patients had not understood their clinicians' explanation of their diagnosis, 26% had not been counselled about avoiding alcohol, 28%
Grogan <i>et al</i> (2010) ³⁸ Ireland	Cross-sectional study	Hepatitis C	Structured interviews sociodemographic, and financial questions as well as a cognitive battery of tests. 142 patients were approached: 13 refused participation, 25 were not eligible and 104 were included	
Bajaj <i>et al</i> (2011) ²⁵ USA	Cross-sectional study (104 patients and 104 caregivers)	Cirrhosis	Structured interviews and medical chart review. Close-ended questions about the patients' understanding of their clinicians' explanation of their diagnosis, counselling about alcohol,	
Bornschelegel <i>et al</i> (2011) ⁴¹ USA	Cross-sectional study (180 patient interviews—145 charts reviews)	Hepatitis C		

Continued

Table 1 Continued

Author (year published) and country	Study design and study participants	Disease group	Data collection methods and survey used	Focus of the paper, summary of findings and domains of supportive care needs
Jennings (2011) ²⁶ USA	Cross-sectional study (152 patients)	Hepatitis C	Information about support group attendance, vaccination against hepatitis, health status and treatment. The response rate was 47% <i>Structured interviews</i> on the educational needs (delivery methods, interest in support groups, topics of interest and preferred services relating to hepatitis C). The response rate was 100%	had not been counselled about preventing spreading hepatitis C to others, and most (90%) had not attended a hepatitis support group (31% were interested in attending) <i>Domains:</i> informational/educational, patient care and support <i>Educational needs</i> —Most respondents did not agree that there was an adequate amount of educational material about hepatitis C in the clinic, most thought that their support person was interested in receiving educational materials about hepatitis C and in participating in educational sessions <i>Domains:</i> informational/educational
Rakoski <i>et al</i> (2012) ¹² USA	Longitudinal study (317 patients and 951 comparison group)	Elderly patients with cirrhosis	<i>Structured interviews</i> on participants' ability to perform tasks of daily living. Two domains were assessed: ADL (eg, dress oneself), and IADL (eg, manage shopping or housework). Response rate not reported	<i>Patient abilities to perform tasks</i> —38% of patients indicated that they had at least one impaired ADL. Commonly reported ADL included 'dressing', 'bathing' and 'walking across room'. The most IADL impairments among those with cirrhosis were 'grocery shopping', 'cooking' and 'managing money'. 10% of individuals with cirrhosis reported 4–5 impaired IADLs <i>Domains:</i> practical

ADL, activities of daily living; HCV, hepatitis C virus; HQLQ, Hepatitis Quality of Life Questionnaire; IADL, instrumental activities of daily living; ISSB, Inventory of Socially Supportive Behaviours; QoL, quality of life.

Table 2 Basic characteristics (setting, disease group, participant characteristics, methodology and research questions) and main findings including domains of commonly reported supportive care needs items of qualitative studies (n=12) included in the literature review

Author (year published) and country	Disease group and study participants	Methodology	Focus of the paper, summary of findings and domains of supportive care needs
Temple-Smith <i>et al</i> (2004) ³⁹ Australia	Hepatitis C (32 patients)	<i>In-depth interviews</i> —Theme list included circumstances surrounding diagnosis, disclosure of HCV status to others, feelings towards self/others after diagnosis, seeking information and knowledge about HCV, barriers to maintaining health living, access/need for support networks, impact of HCV on view of health/lifestyle, thoughts of the future, and knowledge/concerns about transmission	<i>Patients' experiences</i> —There were gender-related differences in relation to symptom recognition, health-seeking attitudes and notions of social support. Men tended to dismiss the impact of their illness, deny needing social support to help them cope with HCV, and did not see the need for information about self-care. In contrast, women talked about the benefits and the desire for social support, and were more willing to seek health information to better manage their HCV. <i>Domains:</i> informational/educational, patient care and support
Harris <i>et al</i> (2005) ²⁸ New Zealand	Hepatitis C (20 patients)	<i>Semistructured in-depth interviews</i> —Each interview (typically) began with the interviewer asking patients 'how they first found out about their hepatitis C', with subsequent areas of discussion addressing experiences of disclosure, stigma, the impact of hepatitis C on relationships, medical encounters, and views on and experiences of HCV treatment. <i>Narrative theory was used*</i>	<i>Patients' experiences</i> —Participants consistently expressed a desire for information regarding the maintenance of their health, and a frustration with the inability of the medical profession to provide this information <i>Domains:</i> informational/educational
Sutton <i>et al</i> (2007) ³⁵ Australia	Hepatitis C (32 patients)	<i>Semistructured interviews</i> —Participants were asked to discuss the impact of the hepatitis C diagnosis and their experience of clinical markers and understandings and perceptions of test results. <i>The illness trajectories and shifting perspectives of wellness and illness models</i> were used	<i>Information and knowledge</i> —The social consequences of living with HCV (eg, social limitations/isolation) were more significant and had greater impact than clinical markers of disease progress. Participants experienced many negative feelings following diagnosis (eg, shock, fear and denial). Misinformation and lack of information was a major problem experienced by this group, some people were exposed to contradictory information. The belief that HCV is always a terminal illness, in particular, was an impression still held by some participants <i>Domains:</i> informational/educational
Grossl <i>et al</i> (2008) ³⁴ USA	Hepatitis C (22 patients)	<i>Semistructured interviews</i> —Questions focused on three areas: medical history, experiences of contracting and being diagnosed with HCV, and the ways in which HCV affected their lives. <i>The theoretical construct of health-related QoL</i> was used	<i>Patients' experiences</i> —Most patients received HCV education from trained professionals but some reported misconceptions and would appear to benefit from additional education. Participants had difficulty with social and occupational functioning, while some received valued support from others for their condition. Living with chronic HCV often had a psychological impact <i>Domains:</i> informational/educational, practical, psychological
Sgorbini <i>et al</i> (2009) ²³ Australia	Hepatitis C (5 patients and their partners)	<i>Semistructured interviews</i> —Interviews began with an open-ended question: 'Living with hepatitis C and undergoing	<i>Patients' experiences</i> —Chronic hepatitis C and combination therapy had an enormous impact on the lives of the patients, their partners and families. The illness and

Continued

Table 2 Continued

Author (year published) and country	Disease group and study participants	Methodology	Focus of the paper, summary of findings and domains of supportive care needs
		combination therapy, what is it like for you?' <i>Heideggerian phenomenology</i> was used	treatment had significant physiological effects that had an impact on QoL; however, the social and psychological consequences of living with a highly stigmatised disease with an unknown course and outcome cannot be underestimated. Patients reported enduring struggles with their finances, lack of a holistic care approach from services, and fears of a threatened future (risk of complications of hepatitis C and lingering effects of treatment) <i>Domains:</i> practical, physical, psychological
Janke <i>et al</i> (2008) ²² USA	Hepatitis C (40 patients)	<i>Focus groups</i> —A structured discussion guide covered questions about treatment, and the decision-making process. Demographic and self-reported health status were collected via self-reported questionnaire	<i>Patients' concerns</i> —Patients reported a remarkable level of emotional volatility that was associated with reduced impulse control, anger, and feelings of sadness and depression. Participants also reported stigmatisation, communication difficulties related to their HCV diagnosis from providers/social peers, and feelings of isolation <i>Domains:</i> psychological
Ng <i>et al</i> (2013) ³⁰ Malaysia	Hepatitis B (44 patients)	<i>Focus groups</i> —Open-ended questions about healthcare experiences (at diagnosis/follow-up), physical, psychological and social impact of hepatitis B; and healthcare needs	<i>Patients' experiences and needs</i> —Patients' needs at diagnosis include desire for emotional support and information. Patients wanted to know more about the disease (transmission, complications, treatment and prognosis); however, information was not easily accessible. They also reported a wide variation in how patients were counselled and managed <i>Domains:</i> psychological, informational/educational
Treloar <i>et al</i> (2010) ³⁶ Australia	Hepatitis C (24 patients)	<i>Semistructured interviews</i> —Themes included HCV and tests, timing/reason for HCV testing, explanation at diagnosis/referrals, attitudes, awareness, readiness/willingness to undertake treatment, and changes in risk practices	<i>Patients' experiences</i> —Overall, the HCV diagnosis experience of participants was poor. Participants reported confusion in relation to HCV tests performed and the implications of test results. Post-test discussions were inadequate—there was a reported lack of information, support and referral provided to participants <i>Domains:</i> informational/educational, patient care and support
Brunings <i>et al</i> (2013) ³² Canada	Hepatitis C (21 patients)	<i>Focus groups</i> —Questions not described. <i>Concept mapping methodology</i> was used	<i>Patients' perspectives of care</i> —Four main themes accounted for most of the statements by participants: communication, professional competence, education/information and continuity of care. Participants emphasised the importance of being treated holistically, receiving HCV education and information was critical to their ability to manage disease <i>Domains:</i> informational/educational


Continued

Table 2 Continued

Author (year and country)	Disease group and study participants	Methodology	Focus of the paper, summary of findings and domains of supportive care needs
Hill <i>et al</i> (2014) ²⁹ England	Hepatitis C (23 patients)	<i>Unstructured interviews</i> —Interviews began with an open-ended question ‘Can you tell me how it has been for you, living with hepatitis C?’ <i>Descriptive phenomenological methodology</i> was used	<i>Patients’ experiences</i> —Experiences of diagnosis were frequently disappointing. Many were told insensitively and felt poorly prepared; this lack of support, advice and information negatively impacted on how they felt about themselves. A lack of practical advice, information and support dominated people’s experiences. This created difficulty when trying to achieve recommended lifestyle changes or gain any sense of control, knowledge or understanding of the condition. Most felt uncertain about how to manage the disease and feared disease progression/health decline <i>Domains:</i> informational/educational, patient care and support
Burnham <i>et al</i> (2014) ²⁷ USA	CLD (13 patients)	<i>Focus groups</i> —Open-ended questions regarding patients’ beliefs of causes of CLD, risk beliefs, attitude towards themselves, towards patients with CLD and towards prevention, perceived benefits of healthy behaviours and risk of liver disease, and barriers to care. The <i>Health Belief Model</i> constructs were used	<i>Knowledge and beliefs</i> —General lack of knowledge about CLD (causes and risks, screening, disease symptoms, and available treatments) was the most common response. When asked to share how they felt about having CLD, patients reported realistic attitudes and emotions, discriminate sharing of their diagnosis, negative medical side effects, fatalism and general negativity. The most commonly reported barriers to treatment included: lack of or inadequate health insurance, cost of care, general lack of knowledge (about symptoms and outcomes), negative attitudes and emotions, and low economic status <i>Domains:</i> informational/educational, practical
Conrad <i>et al</i> (2006) ⁴⁵ Australia	Hepatitis C (70 patients)	<i>Focus groups</i> and individual in-depth interview—13 guiding questions were use in semistructured interviews to elicit open-ended discussion. <i>Grounded-theory</i> methodology was used	<i>Psychosocial factors and QoL</i> —Uncertainty related to disease progression and transmission of the HCV, as well as fear and anxiety about stigma and discrimination were common experiences among patients with HCV <i>Domains:</i> psychological

*Some details about study methods were obtained from M Harris thesis.⁶² CLD, chronic liver disease; HCV, hepatitis C virus; QoL, quality of life.

Figure 1 Summary of the eligibility criteria for inclusion into the review.



questionnaire designed for the study) to collect data on informational and psychological support; however, the questionnaire was designed to explore patients' level of satisfaction with support from the nurse specialist. Minuk *et al*⁴⁴ and Alizadeh *et al*'s⁴³ approach for data collection was the use of an open-ended question to elicit the patient's principal concern about their disease ('volunteered concern'), then patients were asked to rank a list of seven or eight other potential concerns. The other six studies included a mix of questions in their data collection tool, including some specific items on support needs, information needs or ability of patients to perform daily living tasks. Four studies also included questions about QoL.^{21 37 40 42}

DISCUSSION

This systematic review comprehensively summarises the available literature on reported perspectives, experiences and concerns of people who have been diagnosed with CLD. The number of studies collecting patient-reported data is small, compared with either the number of reports of doctors and carer-reported supportive care needs, or those reporting QoL. In particular, there is a paucity of data on the supportive care needs of patients with advanced liver disease and cirrhosis.

Even though the number of studies was small, this review highlights some of the unmet needs of patients with CLD. Most studies were descriptive and used qualitative methodology. However, only five studies used a supportive care needs assessment tool to describe unmet needs of patients with CLD with just two having been validated. One, the ISSB, has shown to have adequate test-retest and internal consistency, although it is not a disease-specific (liver disease) needs assessment tool,^{46 47} and the other was designed to specifically assess patients' satisfaction with information and psychological support received from the nurse specialist.²⁶

Prior to using a health status questionnaire, it is important that the instrument is validated and is suitable for the population under study (eg, translation or rewording may be necessary).⁴⁸ The Scientific Advisory Committee of the Medical Outcomes Trust⁴⁹ developed comprehensive criteria to evaluate the measurement properties of a questionnaire. Eight attributes of an instrument properties to be considered when evaluating a questionnaire assessment tool include: (1) its conceptual and measurement model, (2) validity, (3) reliability, (4) responsiveness, (5) interpretability, (6) respondent and administrative burden, (7) alternative forms and (8) cultural and language adaptations (translations). Although the ISSB⁴⁶ is not a liver disease-specific tool,

Table 3 Domains of supportive care needs and commonly reported specific need items reported by patients included four studies which used a needs assessment tool to collect data

Author	Diseased group	Domains	Need items commonly reported by study participants
Balfour <i>et al</i> ³¹	Hepatitis C	Information/ education needs	<ul style="list-style-type: none"> ▶ 52% rated their current knowledge of liver disease as being inadequate ▶ 91% regarded receiving information about HCV as 'important/very important'
		Patient care and support	<ul style="list-style-type: none"> ▶ 31% were very dissatisfied/dissatisfied with access to specialists services (eg, pharmacist, psychologists) ▶ Percentages regarding access to services as 'important-very/important': 76% HCV drug plans, 68% psychological counselling, 66% nutritionist, 63% pharmacist, 63% support for family/partners, 48% support groups
Zandi <i>et al</i> ³⁷	Cirrhosis	Information/ education needs	<p>The study assessed patients' educational needs. Below is the percentage of patients reporting need for:</p> <ul style="list-style-type: none"> ▶ 65% curative ways in cirrhosis (being treatable/not treatable) ▶ 45% routes of transmission as well as diagnostic tests
		Physical needs	<ul style="list-style-type: none"> ▶ 70% controlling or reducing abdominal distention ▶ 65% ways of controlling fatigue ▶ 60% principles of care and proper medications ▶ 50% controlling pruritus and fatigue
		Psychological Practical support	<ul style="list-style-type: none"> ▶ 55% worry ▶ 47.8% reported moderate to severe financial stress
Chang <i>et al</i> ²¹	Hepatitis C	Psychological Practical support	<ul style="list-style-type: none"> ▶ 55% worry ▶ 47.8% reported moderate to severe financial stress
Jennings ²⁶	Hepatitis C	Information/ education needs	<ul style="list-style-type: none"> ▶ 71% disagreed that there was an adequate amount of educational material about hepatitis C in the clinic ▶ 67% thought that their support person was interested in receiving educational materials about hepatitis C ▶ 78% thought that their support person would be interested in participating in educational sessions
		Patient support	<ul style="list-style-type: none"> ▶ 61% would be interested in joining a regular support group at the clinic ▶ 42% thought that it would be beneficial for their support person to join a caregiver support group
Grogan <i>et al</i> ³⁸	Hepatitis C	Information/ education needs	<p>The study assessed patients' perceptions of support received from the nurse specialist during HCV treatment. There were low levels of disagreement that 'The nurse provided ...':</p> <ul style="list-style-type: none"> ▶ 17% advice on how to maintain a healthy balanced diet ▶ 14% advice on sleep management ▶ 12% advices on energy conservation ▶ 12% advice on physical exercise
		Patient support	<ul style="list-style-type: none"> ▶ 14% information on support groups that were available to me ▶ 18% ongoing support postcompletion of treatment

HCV, hepatitis C virus.

it has adequate test-retest and internal consistency (reliability coefficient of internal consistency for the total scale was 0.89).^{46 47} The questionnaire used in the Grogan and Timmins³⁸ study had its content validity confirmed by a panel of experts, and had an adequate reliability score (Cronbach's $\alpha=0.85$).

Education is a critical component of any healthcare intervention; it has been found to improve treatment

adherence, facilitate effective decision-making, reduce healthcare costs and improve health outcomes. Research shows that people diagnosed with CLD (eg, cirrhosis,⁵⁰ hepatitis C⁵¹) have a poor understanding of their disease and lack adequate knowledge about important information needed to self-manage their disease. Furthermore, participation in an HCV education class has been shown to increase patients' understanding

of disease symptoms, transmission and treatment.⁵² A recent study has shown that a simple educational intervention (providing a concise booklet about cirrhosis and emphasising its importance) for patients with cirrhosis was associated with a 26% improvement in patient knowledge about their disease.⁵⁰ Practical, physical, patient care and support, and psychological needs were also reported as important.

The role of patient education/knowledge has even more importance with the recognition that modifiable host factors can have a substantial impact on liver disease progression and treatment outcomes. In most patients with chronic hepatitis C, fibrosis progression to cirrhosis typically requires decades. However, host risk factors such as heavy alcohol consumption (>50 g/day) or increased body mass index can lead to more rapid liver disease progression.^{53 54} Similarly, in alcoholic liver disease, individuals with fibrosis who continue to drink alcohol have a high risk of disease progression.^{55 56} Improved education about risks of alcohol, obesity and physical inactivity may reduce the impact of comorbidities on disease progression. Patient education is also essential to ensure compliance with prescribed medications and continued follow-up.⁵⁷ Thus, an increase in knowledge around CLD has the potential to affect behavioural change, enhance patient self-efficacy and, in turn, improve both QoL and disease progression.⁵⁸

Although this review aimed at describing the unmet supportive care needs of people diagnosed with CLD, five reports included a mixed group of study participants. Nevertheless, these reports were included because they provide an insight about the complex array of concerns people living with CLD may have. Additional information could also potentially be found from patients' Blog, or Facebook page. However, as these are not standardised they were not included in this review.

Unmet supportive care needs are those needs which lack the level of service or support an individual perceives is necessary to achieve optimal well-being. One criterion for inclusion of articles was that it had to report the *patients'* views, perspectives, experiences or concerns. In most articles included in this review, the reported unmet supportive care needs of patients are the *investigators'* interpretations of patients' needs, as patients were not specifically asked to report their perceived unmet supportive care needs or concerns, rather asked whether they had been counselled about not drinking alcohol and how to avoid transmitting the virus to other people.⁴¹ Those patients who had not been counselled may not necessarily perceive this as an unmet need or concern. In Gifford *et al's*⁴⁰ study, a large number of women reported ever having symptoms and rated their health as 'fair-poor', yet some women may have received outside help or support to deal with these issues. Similarly, Rakoski *et al*¹² reported over one-third of people diagnosed with cirrhosis had at least one impaired activity of daily living such as dressing or bathing, yet it is possible that they may also have

received help and support (formal or informal care) with these activities, and therefore their supportive care needs are not unmet.

Some clinicians may perceive the social and financial domains of supportive care needs to be outside their realm of practice, but such factors can significantly complicate treatment, reduce adherence to treatment or lifestyle modification, and create management challenges. Poor understanding of medications is one key area known to increase hospitalisation. Clinicians inconsistently ask patients about their unmet supportive care needs and concerns, typically operating in a 'reactive mode' (eg, acting in response to patients' pressing or self-reported problems).⁵⁹ In the USA, for example, each State has a Department of Health and Human Services, where patients can access county resources (eg, case managers who can assist patients with navigating the health system). Eligible Medicare/Medicaid patients can seek public assistance or transportation to medical visits by county transportation.⁶⁰ In Australia, similar arrangements are in place to defray cost for patients in rural or remote areas who are required to travel for their healthcare. It may be that having a structured, validated supportive care needs instrument specific for the assessment of supportive care needs of patients with CLD may allow clinicians and other healthcare workers (eg, nurses, dieticians) to better address deficiencies in patients' support needs. Disease-specific supportive care needs assessments have been used widely in the heart failure setting.⁶¹ Future research could investigate the potential for a validated liver disease-specific supportive care needs assessment instrument that can potentially be administered quickly by clinical staff (eg, nursing) or self-administered by patients in the waiting room. Use of such an instrument could then prompt clinicians to be proactive in addressing patients' unmet supportive care needs and, where appropriate, refer to support services to enhance their QoL.

While a systematic review was undertaken, using pre-specified criteria, it is possible that some relevant publications were missed. Unpublished articles or non-indexed articles may have been missed. The studies included in this review varied substantially with regard to methodology. About half the articles used qualitative methodology based on smaller numbers of patients, while half used quantitative methodology. Some studies focused specifically on needs assessment^{21 26 30 31 37} or patients concerns,^{22 43 44} and the reported findings were clearly the patients' views and perspectives of their supportive care needs. Some articles focused mostly on the patients' health status, QoL or changes in lifestyle. These were included in this review because they reported some information about patient's unmet supportive care needs or concerns, for instance Fabris *et al*³³ reported that most patients wanted more detailed information about HCV and its transmission. Despite the fact that much of the burden of clinical care occurs in patients with cirrhosis, only three of the papers focused specifically on that population's supportive care needs.

Lastly, this review is limited by the varying quality and rigour of the included studies, which had significant heterogeneity in terms of the population group (disease, age, gender and concurrent comorbidity), numbers and types of domains interrogated.

In conclusion, this systematic review found 26 articles reporting supportive care needs among patients with liver disease, but few of them are *patient-reported* needs and none used a validated, liver disease-specific supportive care needs assessment instrument. Furthermore, most studies did not focus on the CLD population likely to have the most needs—namely those with cirrhosis. Development of a validated supportive care needs assessment instrument for people with CLD would not only advance understanding of patients' unmet needs, but have potential utility in clinical practice for facilitating timely referrals to support services. Support for areas raised in this review around knowledge and information are important for both chronic disease management, and for end of life planning for patients with liver failure from CLD.

Author affiliations

¹Menzies School of Health Research, Charles Darwin University, Darwin, Northern Territory, Australia

²School of Medicine, University of Queensland, Brisbane, Queensland, Australia

³Centre for Liver Disease Research, School of Medicine, The University of Queensland, Brisbane, Queensland, Australia

⁴Department of Gastroenterology and Hepatology, Princess Alexandra Hospital, Brisbane, Queensland, Australia

⁵School of Medicine and Public Health, University of Newcastle, Newcastle, New South Wales, Australia

⁶Division of Gastroenterology and Hepatology, University of Michigan Health System, Ann Arbor, Michigan, USA

⁷Institute of Health and Biomedical Innovation and School of Public Health & Social Work, Queensland University of Technology, Brisbane, Queensland, Australia

⁸Centre for Functioning and Health Research, Metro South Health, Brisbane, Queensland, Australia

⁹Cancer Control Group, QIMR Berghofer Medical Research Institute, Brisbane, Queensland, Australia

¹⁰Diamantina Institute, University of Queensland, Brisbane, Queensland, Australia

Acknowledgements We thank D Radford-Smith who provided administrative assistance with this manuscript.

Contributors PCV and JM independently reviewed all manuscripts, and EP and NM adjudicated disagreements. PCV was responsible for data acquisition, and takes responsibility for the integrity and the accuracy of the data. PCV drafted the report, all authors have read and contributed to this manuscript's editing, have agreed to the submission to *BMJ Open*, and have agreed with the content and presentation of the paper.

Funding This work was supported by the National Health and Medical Research Council of Australia and The Queensland Government's Smart State Health and Medical Research Fund. PCV was supported by an Australian Research Council Future Fellowship (#FT100100511). SMM is supported by a National Health and Medical Research Council of Australia fellowship (#1016432).

Competing interests None.

Provenance and peer review Not commissioned; externally peer reviewed.

Data sharing statement No additional data are available.

Open Access This is an Open Access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

REFERENCES

1. Wang FS, Fan JG, Zhang Z, *et al*. The global burden of liver disease: the major impact of China. *Hepatology* 2014;60:2099–108.
2. Blachier M, Leleu H, Peck-Radosavljevic M, *et al*. The burden of liver disease in Europe: a review of available epidemiological data. *J Hepatol* 2013;58:593–608.
3. Deloitte Access Economics. *The economic cost and health burden of liver diseases in Australia*. Sydney, Australia: The Gastroenterological Society of Australia/Australian Liver Association, 2013.
4. Lazo M, Hernaez R, Eberhardt MS, *et al*. Prevalence of nonalcoholic fatty liver disease in the United States: the Third National Health and Nutrition Examination Survey, 1988–1994. *Am J Epidemiol* 2013;178:38–45.
5. MacLachlan JH, Allard N, Towell V, *et al*. The burden of chronic hepatitis B virus infection in Australia, 2011. *Aust N Z J Public Health* 2013;37:416–22.
6. Rehm J, Samokhvalov AV, Shield KD. Global burden of alcoholic liver diseases. *J Hepatol* 2013;59:160–8.
7. Roberts HW, Utuama OA, Klevens M, *et al*. The contribution of viral hepatitis to the burden of chronic liver disease in the United States. *Am J Gastroenterol* 2014;109:387–93; quiz 6, 94.
8. All-Party Parliamentary Hepatology Group (APPHG). *Liver disease: today's complacency, tomorrow's catastrophe. The All-Party Parliamentary Hepatology Group (APPHG) inquiry into improving outcomes in liver disease*. UK: All-Party Parliamentary Hepatology Group (APPHG), 2014.
9. Razavi H, Elkhoury AC, Elbasha E, *et al*. Chronic hepatitis C virus (HCV) disease burden and cost in the United States. *Hepatology* 2013;57:2164–70.
10. MacLachlan JH, Cowie BC. Liver cancer is the fastest increasing cause of cancer death in Australians. *Med J Aust* 2012;197:492–3.
11. Australian Institute of Health and Welfare. *Cancer incidence projections Australia, 2011–2020 (Cat no CAN 62)*. Canberra, Australia: AIHW.
12. Rakoski MO, McCammon RJ, Piette JD, *et al*. Burden of cirrhosis on older Americans and their families: analysis of the health and retirement study. *Hepatology* 2012;55:184–91.
13. Fitch M. Supportive care for cancer patients. *Hosp Q* 2000;3:39–46.
14. Sanson-Fisher R, Giris A, Boyes A, *et al*. The unmet supportive care needs of patients with cancer. Supportive Care Review Group. *Cancer* 2000;88:226–37.
15. Giris A, Boyes A, Sanson-Fisher RW, *et al*. Perceived needs of women diagnosed with breast cancer: rural versus urban location. *Aust N Z J Public Health* 2000;24:166–73.
16. Garvey G, Beesley VL, Janda M, *et al*. The development of a supportive care needs assessment tool for Indigenous people with cancer. *BMC Cancer* 2012;12:300.
17. Davidson P, Cockburn J, Daly J, *et al*. Patient-centered needs assessment: rationale for a psychometric measure for assessing needs in heart failure. *J Cardiovasc Nurs* 2004;19:164–71.
18. Garvey G, Beesley V, Janda M, *et al*. Psychometric properties of an Australian Supportive Care Needs Assessment Tool for Indigenous People (SCNAT-IP) with cancer. *Cancer* 2015. In press.
19. Carbonell N, Pauwels A, Serfaty L, *et al*. Improved survival after variceal bleeding in patients with cirrhosis over the past two decades. *Hepatology* 2004;40:652–9.
20. Bonevski B, Sanson Fisher R, Hersey P, *et al*. Assessing the perceived needs of patients attending an outpatient melanoma clinic. *J Psychosoc Oncol* 1999;17:101–18.
21. Chang SC, Ko WS, Wu SS, *et al*. Factors associated with quality of life in chronic hepatitis C patients who received interferon plus ribavirin therapy. *J Formos Med Assoc* 2008;107:454–62.
22. Janke EA, McGraw S, Garcia-Tsao G, *et al*. Psychosocial issues in hepatitis C: a qualitative analysis. *Psychosomatics* 2008;49:494–501.
23. Sgorbini M, O'Brien L, Jackson D. Living with hepatitis C and treatment: the personal experiences of patients. *J Clin Nurs* 2009;18:2282–91.

24. Jessop AB, Cohen C, Burke MM, *et al.* Hepatitis support groups: meeting the information and support needs of hepatitis patients. *Gastroenterol Nurs* 2004;27:163–9.
25. Bajaj JS, Wade JB, Gibson DP, *et al.* The multi-dimensional burden of cirrhosis and hepatic encephalopathy on patients and caregivers. *Am J Gastroenterol* 2011;106:1646–53.
26. Jennings AR. Assessing educational needs of potential hepatitis C patients at a Veterans Affairs medical center. *Educ Health (Abingdon)* 2011;24:532.
27. Burnham B, Wallington S, Jillson IA, *et al.* Knowledge, attitudes, and beliefs of patients with chronic liver disease. *Am J Health Behav* 2014;38:737–44.
28. Harris M. Living with hepatitis C: the medical encounter. *N Z Sociol* 2005;20:4–19.
29. Hill R, Pfeil M, Moore J, *et al.* Living with hepatitis C: a phenomenological study. *J Clin Nurs* 2015;24:428–38.
30. Ng CJ, Low WY, Wong LP, *et al.* Uncovering the experiences and needs of patients with chronic hepatitis B infection at diagnosis: a qualitative study. *Asia Pac J Public Health* 2013;25:32–40.
31. Balfour L, Cooper C, Tasca GA, *et al.* Evaluation of health care needs and patient satisfaction among hepatitis C patients treated at a hospital-based, viral hepatitis clinic. *Can J Public Health* 2004;95:272–7.
32. Brunings P, Klar S, Butt G, *et al.* "It's a big part of our lives": a qualitative study defining quality of hepatitis C care from the patient's perspective. *Gastroenterol Nurs* 2013;36:249–57.
33. Fabris P, Tositti G, Giordani MT, *et al.* Assessing patients' understanding of hepatitis C virus infection and its impact on their lifestyle. *Aliment Pharmacol Ther* 2006;23:1161–70.
34. Groessl EJ, Weingart KR, Kaplan RM, *et al.* Living with hepatitis C: qualitative interviews with hepatitis C-infected veterans. *J Gen Intern Med* 2008;23:1959–65.
35. Sutton R, Treloar C. Chronic illness experiences, clinical markers and living with hepatitis C. *J Health Psychol* 2007;12:330–40.
36. Treloar C, Newland J, Harris M, *et al.* A diagnosis of hepatitis C—insights from a study on patients' experiences. *Aust Fam Physician* 2010;39:589–92.
37. Zandi M, Adib-Hajbagheri M, Memarian R, *et al.* Effects of a self-care program on quality of life of cirrhotic patients referring to Tehran Hepatitis Center. *Health Qual Life Outcomes* 2005;3:35.
38. Grogan A, Timmins F. Patients' perceptions of information and support received from the nurse specialist during HCV treatment. *J Clin Nurs* 2010;19:2869–78.
39. Temple-Smith M, Gifford S, Stoov Eacute M. The lived experience of men and women with hepatitis C: implications for support needs and health information. *Aust Health Rev* 2004;27:46–56.
40. Gifford SM, O'Brien ML, Bammer G, *et al.* Australian women's experiences of living with hepatitis C virus: results from a cross-sectional survey. *J Gastroenterol Hepatol* 2003;18:841–50.
41. Bomschlegel K, Crotty KJ, Sahl S, *et al.* Unmet needs among people reported with hepatitis C, New York City. *J Public Health Manag Pract* 2011;17:E9–17.
42. Gifford SM, O'Brien ML, Smith A, *et al.* Australian men's experiences of living with hepatitis C virus: results from a cross-sectional survey. *J Gastroenterol Hepatol* 2005;20:79–86.
43. Alizadeh AH, Ranjbar M, Yadollahzadeh M. Patient concerns regarding chronic hepatitis B and C infection. *East Mediterr Health J* 2008;14:1142–7.
44. Minuk GY, Gutkin A, Wong SG, *et al.* Patient concerns regarding chronic hepatitis C infections. *J Viral Hepat* 2005;12:51–7.
45. Conrad S, Garrett LE, Cooksley WG, *et al.* Living with chronic hepatitis C means 'you just haven't got a normal life any more'. *Chronic Illn* 2006;2:121–31.
46. Barrera M, Sandler I, Ramsay T. Preliminary development of a scale of social support: studies on college students. *Am J Community Psychol* 1981;9:435–47.
47. Barrera M. Social support in the adjustment of pregnant adolescents: assessment issues. In: Gottlieb BH, ed. *Social networks and social support in community mental health*. Beverly Hills: Sage, 1981:69–96.
48. Terwee CB, Bot SD, de Boer MR, *et al.* Quality criteria were proposed for measurement properties of health status questionnaires. *J Clin Epidemiol* 2007;60:34–42.
49. Scientific Advisory Committee of the Medical Outcomes Trust. Assessing health status and quality-of-life instruments: attributes and review criteria. *Qual Life Res* 2002;11:193–205.
50. Volk ML, Fisher N, Fontana RJ. Patient knowledge about disease self-management in cirrhosis. *Am J Gastroenterol* 2013;108:302–5.
51. Stein MD, Maksad J, Clarke J. Hepatitis C disease among injection drug users: knowledge, perceived risk and willingness to receive treatment. *Drug Alcohol Depend* 2001;61:211–15.
52. Kizer E, Whitehead A, Indest D, *et al.* Efficacy of group education in veterans with hepatitis C. *Fed Pract* 2006;23:50–7.
53. Hourigan LF, Macdonald GA, Purdie D, *et al.* Fibrosis in chronic hepatitis C correlates significantly with body mass index and steatosis. *Hepatology* 1999;29:1215–19.
54. Hutchinson SJ, Bird SM, Goldberg DJ. Influence of alcohol on the progression of hepatitis C virus infection: a meta-analysis. *Clin Gastroenterol Hepatol* 2005;3:1150–9.
55. Pares A, Caballeria J, Bruguera M, *et al.* Histological course of alcoholic hepatitis. Influence of abstinence, sex and extent of hepatic damage. *J Hepatol* 1986;2:33–42.
56. Worner TM, Lieber CS. Perivascular fibrosis as precursor lesion of cirrhosis. *JAMA* 1985;254:627–30.
57. Devins GM, Mendelsohn DC, Barre PE, *et al.* Predialysis psychoeducational intervention extends survival in CKD: a 20-year follow-up. *Am J Kidney Dis* 2005;46:1088–98.
58. Singal AG, Volk ML, Rakoski MO, *et al.* Patient involvement in healthcare is associated with higher rates of surveillance for hepatocellular carcinoma. *J Clin Gastroenterol* 2011;45:727–32.
59. Horsfall L, Macdonald G, Scott I, *et al.* Use of standardised assessment forms in referrals to hepatology outpatient services: implications for accurate triaging of patients with chronic hepatitis C. *Aust Health Rev* 2013;37:218–22.
60. Bonner JE, Barritt AS IV, Fried MW, *et al.* Tangible resources for preparing patients for antiviral therapy for chronic hepatitis C. *Dig Dis Sci* 2012;57:1439–44.
61. Low J, Pattenden J, Candy B, *et al.* Palliative care in advanced heart failure: an international review of the perspectives of recipients and health professionals on care provision. *J Card Fail* 2011;17:231–52.
62. Harris M. *Negotiating the pull of the normal: embodied narratives of living with hepatitis C in New Zealand and Australia*. NSW, Australia: University of New South Wales, 2010.