

# Immune Checkpoint Inhibitor Therapy in Neuroendocrine Tumors



## Authors

Sriram Gubbi<sup>1</sup> , Namrata Vijayvergia<sup>2</sup>, Jian Q Yu<sup>3</sup>, Joanna Klubo-Gwiezdzinska<sup>4\*</sup> , Christian A. Koch<sup>5, 6\*</sup>

## Affiliations

- 1 Endocrinology, National Institutes of Health Clinical Center, Bethesda, United States
- 2 Oncology, Fox Chase Cancer Center, Philadelphia, United States
- 3 Nuclear Medicine, Fox Chase Cancer Center, Philadelphia, United States
- 4 National Institute of Diabetes and Digestive and Kidney Diseases, National Institutes of Health, Bethesda, United States
- 5 Medicine/Endocrinology, The University of Tennessee Health Science Center, Memphis, United States
- 6 Medicine, Fox Chase Cancer Center, Philadelphia, United States

## Key words

neuroendocrine tumor, immunotherapy, somatostatin analogues, medullary thyroid carcinoma; MTC, pheochromocytoma, paraganglioma

received 30.06.2022

accepted after revision 25.07.2022

published online 02.09.2022

## Bibliography

Horm Metab Res 2022; 54: 795–812

DOI 10.1055/a-1908-7790

ISSN 0018-5043

© 2022. The Author(s).

This is an open access article published by Thieme under the terms of the Creative Commons Attribution-NonDerivative-NonCommercial-License, permitting copying and reproduction so long as the original work is given appropriate credit. Contents may not be used for commercial purposes, or adapted, remixed, transformed or built upon. (<https://creativecommons.org/licenses/by-nc-nd/4.0/>)

Georg Thieme Verlag KG, Rüdigerstraße 14,  
70469 Stuttgart, Germany

## Correspondence

Prof. Christian A. Koch, FACP, MACE  
Fox Chase Cancer Center  
Medicine, 333 Cottman Ave  
Philadelphia  
19111-2497  
United States  
Tel.: 215 728 2713  
[Christian.koch65@gmail.com](mailto:Christian.koch65@gmail.com)

## ABSTRACT

Neuroendocrine tumors (NETs) occur in various regions of the body and present with complex clinical and biochemical phenotypes. The molecular underpinnings that give rise to such varied manifestations have not been completely deciphered. The management of neuroendocrine tumors (NETs) involves surgery, locoregional therapy, and/or systemic therapy. Several forms of systemic therapy, including platinum-based chemotherapy, temozolomide/capecitabine, tyrosine kinase inhibitors, mTOR inhibitors, and peptide receptor radionuclide therapy have been extensively studied and implemented in the treatment of NETs. However, the potential of immune checkpoint inhibitor (ICI) therapy as an option in the management of NETs has only recently garnered attention. Till date, it is not clear whether ICI therapy holds any distinctive advantage in terms of efficacy or safety when compared to other available systemic therapies for NETs. Identifying the characteristics of NETs that would make them (better) respond to ICIs has been challenging. This review provides a summary of the current evidence on the value of ICI therapy in the management of ICIs and discusses the potential areas for future research.

## Introduction

The incidence of neuroendocrine tumors (NETs) is increasing and is estimated to be around 1.01–5.25 cases per 100 000 population based on data from the United States, Japan, and European regis-

tries [1–3]. Several forms of locoregional and systemic medical and surgical therapies have been used with varying efficacies. Immunotherapy, particularly with immune checkpoint inhibitors (ICIs) has only recently emerged as one of the options for systemic therapy for NETs. To date, the role of ICI therapy and other forms of immunotherapy in the management of NETs is not well established.

\* Co-senior authors


The various molecular and cellular mechanisms as well as the tumor microenvironmental factors that may regulate the response of NETs to ICI therapy are yet to be completely understood. In this review, we discuss the current knowledge on the efficacy of ICI therapy in the treatment of NETs and discuss the potential areas for future research.

## Neuroendocrine tumor biology and current standard of care

NETs are benign or malignant tumors that demonstrate variable rates of growth and have the ability to store and secrete biologically active peptides and amine compounds [4]. These tumors arise from the so-called diffuse neuroendocrine system of the body, the embryologic origins of which thought to be either from the neural crest or from the gut endoderm [5]. NETs arise from the various neuroendocrine cell types, including the ganglionic cells of the nervous system, paraganglionic chromaffin cells, pancreatic islets, adrenal medulla, and thyroid c-cells [5]. These tumors also demonstrate immunohistochemical reactivity to neuronal markers such as chromogranin A, synaptophysin, or neuron-specific enolase [4]. NETs can occur in various regions of the body, including but not limited to the gastrointestinal tract, pancreas, lungs, bronchus, thymus, skin, cervix, prostate, and the thyroid [6–10]. While most of these tumors are slow growing, some tumors such as high-grade NETs, neuroendocrine carcinomas (NECs), medullary thyroid cancer (MTC), Merkel cell carcinoma (MCC), and certain forms of pheochromocytomas and paragangliomas (PPGLs) tend to be more aggressive and can metastasize to distant sites [6, 11, 12].

As per the 2019 World Health Organization guidelines, NETs are classified into different grades depending on the mitotic rate or Ki-67 proliferative index: G1 (< 2 mitoses/10 high power field (HPF), < 3% Ki-67 index), G2 (2–20 mitoses/10 HPF, 3–20% Ki-67 index), and G3 (> 20 mitoses/10 HPF, > 20% Ki-67 index) [13]. NECs are now a distinct subtype from NETs and are further classified into small-cell NECs (SCNECs) and large-cell NECs (LCNECs). While both G3 NETs and NECs have a high mitotic rate/Ki-67 index, the main difference lies in their differentiation and clinical response to treatment: NECs tend to be poorly differentiated and respond to platinum-based chemotherapy while G3 NETs are well-differentiated and relatively resistant to platinum-based chemotherapy but can be less aggressive than NECs [13, 14]. Apart from these, mixed NET-non-NET neoplasms are also a part of this classification where the differentiation and histopathology can be variable. The course of management of NETs is determined by the anatomical location of the primary tumor, biochemical phenotype, histological grade of the tumor, and staging of the disease [12, 15–17]. Surgery is the preferred modality of treatment for localized non-metastatic NETs, and a cytoreductive surgical approach could be utilized for the treatment of metastatic disease [18]. Locoregional therapies such as radiofrequency ablation, transarterial embolization, stereotactic radiotherapy, and chemoembolization are also feasible options for liver metastases, as well as palliative radiation to bone metastases [18–20].

Such a multimodality approach is displayed in the illustrative history of a 48-year-old man who was found to have an elevated serum calcium up to 14 mg/dl (normal, 8.6–10.0), alkaline phosphatase and gamma glutamyl transferase during routine blood



**Fig. 1** a:  $^{68}\text{Ga}$ -DOTATATE PET/CT scan in Maximum Intensity Projection (MIP) display. Scan was on 09/2020 with multiple liver lesions (hot tumors) and primary pancreatic NET (blue arrow). b:  $^{68}\text{Ga}$ -DOTATATE PET/CT scan in Maximum Intensity Projection (MIP) display. Scan was on 02/2022 with multiple liver lesions again visualized (hot tumors) and primary pancreatic NET (blue arrow). The size is smaller for some of the liver lesions.

work in 2018. Computed tomography (CT) of the abdomen revealed a pancreatic mass and liver tumors. Biopsy of the pancreatic mass showed a low-grade neuroendocrine tumor. Hypercalcemia was mediated by parathyroid hormone-related peptide. He received monthly long-acting octreotide (Sandostatin LAR) 30 mg and was started on denosumab and switched to zoledronic acid with slight decline but without normalization of serum calcium. In 3/2020, he underwent transarterial chemoembolization of multiple metastases within the right and left hepatic lobe. Hypercalcemia continued. He then had a gallium-68 ( $^{68}\text{Ga}$ )-DOTATATE positron emission tomography computed tomography (PET/CT) imaging (► Fig. 1a, ► 2a) and underwent therapy with 4 cycles of lutetium-177 ( $^{177}\text{Lu}$ )-DOTATATE therapy in 09/2020, 11/2020, 1/2021, 3/2021. The lesions were stable on 02/2022  $^{68}\text{Ga}$ -DOTATATE PET/CT imaging (► Fig. 1b, ► 2b). Several months later his serum Ca has dropped but remains above normal range and in 06/2022 was measured at 10.6 mg/dl (8.6–10.0) while continuing monthly Sandostatin LAR 30 mg. Zoledronic acid was stopped in June 2021.

In patients with metastatic disease but with low tumor burden, frequent follow-up with or without treatment with somatostatin analogs (SSAs) such as octreotide [21, 22], and lanreotide [23, 24], especially in patients with somatostatin receptor-positive (SSTR+) NETs and/or hormonally active NETs such as carcinoids, insulinomas, glucagonomas, and others [20] is needed. Several systemic therapy options are available for the treatment of advanced, metastatic NETs [6, 12, 17, 20, 25]. Some of the systemic therapy options include mechanistic target of rapamycin (mTOR) inhibitor everolimus [26, 27], tyrosine kinase inhibitor (TKI) sunitinib [28, 29], vascular endothelial growth factor (VEGF) inhibitor bevacizumab [30], interferon  $\alpha$  [30, 31], and for the treatment of SSTR+ NETs,

peptide receptor radionuclide therapy (PRRT) with radiolabeled SSAs such as lutetium-177 ( $^{177}\text{Lu}$ )-DOTA-Tyr3-octreotate (DOTATATE) [32]. A combination of these therapies is also utilized for the treatment of NETs [27, 30, 32]. Chemotherapeutic regimens are also utilized in the treatment of more aggressive forms of NETs/neuroendocrine carcinomas (NECs), some of which include cisplatin/etoposide [33], carboplatin/etoposide [20], oxaliplatin-based therapy (FOLFOX, CAPEOX) [20], irinotecan-based therapy (cisplatin/etoposide, FOLFIRI, FOLFIRINOX) [20], and temozolomide, either as a single agent or with capecitabine [34, 35].

Such a treatment approach including immunotherapy is shown in the history of a 43-year-old man who presented with abdominal pain due to small bowel obstruction in 2015. He was previously worked up for irritable bowel syndrome in 2013 and had ongoing diarrhea before 2013. After small bowel resection and ongoing abdominal discomfort and diarrhea, he presented to a NET center in 2016 and underwent an indium-111 ( $^{111}\text{In}$ )-octreoscan showing evidence of mesenteric lymphadenopathy ▶ Fig. 3a, b). Bulky mesenteric lymph nodes, small bowel containing the primary G3 NET (Ki-67 index 25%), and the gallbladder were resected, and monthly octreotide LAR was started in an adjuvant setting. Four months later, imaging studies suggested tumor recurrence in the liver and peritoneum and the patient started systemic therapy with capecitabine and temozolomide, however, developed progressive disease ▶ Fig. 4a). Therapy was changed to carboplatin/etoposide for 8 months which was ineffective, and then the patient received 3 cycles of pembrolizumab, unfortunately with disease progression ▶ Fig. 4b).

A meta-iodobenzylguanidine (MIBG) scan in 6/2016 was not avid for liver lesions that were seen on magnetic resonance imaging (MRI). Molecular profiling did not find disease associated mutations or variants of uncertain significance. Microsatellite instability (MSI) testing with immunohistochemical stains for MLH1, MSH2, PMS2, and MSH6 showed continued nuclear expression of all four proteins


in the poorly differentiated NEC, implying low probability of a MSI high tumor. PD-L1 expression was negative.

The National Comprehensive Cancer Network (NCCN) 2021 guidelines endorse the use of all of these locoregional and systemic therapy for the management of NETs as category 2A (low-level evidence and uniform NCCN consensus that a therapy is appropriate) recommendations [20]. The North American Neuroendocrine Tumor Society (NANETS) released its compendium guidelines in 2021 in partnership with the Commonwealth Neuroendocrine Tumor Research Collaboration (CommNETs) comprising Canada, Australia, and New Zealand, along with endorsements and updating of the 2015 European Neuroendocrine Society (ENETS) guidelines [36]. In the NANETS guidelines, most of these systemic therapies were endorsed as grade B or C recommendations as per the Oxford Centre for Evidence-Based Medicine [36].

### Role of immune checkpoint inhibitor therapy in neuroendocrine tumors

ICIs are monoclonal antibodies that target the immune co-inhibitory receptors as well as their respective ligands, including programmed cell death protein-1 (PD-1), PD-ligand 1 (PD-L1), and cytotoxic T-lymphocyte antigen 4 (CTLA-4) [37, 38]. The efficacy of ICIs has been well-demonstrated in the management of various cancers [39–43]. In addition to these proteins, the T-cell immunoglobulin and mucin-domain containing-3 (TIM-3), lymphocyte activation gene-3 (LAG-3), and T-cell immunoglobulin and ITIM domain (TIGIT) are additional co-inhibitory proteins that could serve as potential targets for the next generation of ICIs [44, 45].


Several phase Ib and phase II studies as well as retrospective studies have evaluated the efficacy of ICIs for the treatment of a variety of NETs [46–62]. A summary of data published on ICI therapy in NETs is provided in ▶ Table 1. As evident in this table, full-length articles related to ICI therapy in NETs, except for MCC, have been published only since as recently as 2020. However, the objective response rates (ORR) have been low to modest, with most trial



▶ Fig. 2 a:  $^{68}\text{Ga}$ -DOTATATE PET/CT scan in transaxial fused image. Scan was on 09/2020 with multiple liver lesions (hot tumors) with central necrosis and primary pancreatic NET (blue arrow). Same patient as ▶ Fig. 1. b:  $^{68}\text{Ga}$ -DOTATATE PET/CT scan in transaxial fused image. Scan was on 02/2022 with multiple liver lesions (hot tumors) with smaller size and less central necrosis post therapy with full dose of peptide receptor radionuclide therapy (PRRT). The primary pancreatic NET (blue arrow) is again seen with minimally higher intensity. The overall findings are stable disease. Same patient as ▶ Fig. 1.



► **Fig. 3** **a:**  $^{111}\text{In}$ -Octreoscan in whole body display. Scan was on 06/2016 without significant focal lesions (cold tumors). **b:**  $^{111}\text{In}$ -Octreoscan in transaxial fused image focusing on liver. Scan was on same day of whole-body scan with small liver lesions but no activity above liver background (cold tumors).



► **Fig. 4** **a:** CT of abdomen in trans axial display. Scan was on 12/2016 with more small liver lesions. Same patient as ► **Fig. 3** and therapy changed from temozolomide/capecitabine to carboplatin/etoposide. **b:** CT of abdomen in trans axial display. Scan was on 11/2017. There are more liver lesions (in large circle in center, also posteriorly) and new peritoneal metastases (in smaller circle to right). The overall findings were suggestive of disease progression.

studies demonstrating ORRs of < 10%. In the phase II KEYNOTE-158 study which investigated the safety and efficacy of pembrolizumab across multiple malignancies, the drug's utility in 107 patients with advanced, well-differentiated NETs was also evaluated [50]. After a median follow-up of 24.2 months, ORR was seen in 3.7% (4/107) patients, with partial response (PR) noted in all four and with none achieving complete response (CR). Moreover, all four patients had no PD-L1 expression in the NETs while 17 patients from the remainder of the study population demonstrated positive

PD-L1 expression, therefore suggesting poor correlation between PD-L1 tumor expression and likelihood of response to ICI [63]. Similarly, in another study evaluating pembrolizumab therapy in 29 patients with G3 extrapulmonary NETs, ORR was seen in one (3.4%) patient with esophageal NEC [51]. Moreover, there were no differences in the disease control rate, overall survival (OS), or progression-free survival (PFS) between patients with PD-L1-positive and PD-L1-negative NETs. While there are individual reports of combination of PD-1 therapy and chemotherapy being effective in tum-

► **Table 1** List of studies evaluating the efficacy of ICI in NETs.

Study	Year	Phase	Number of patients	Types of NET	Type of ICI (target protein)	Duration of follow-up in months	Outcomes
Mehnert et al. (KEYNOTE-028) [46]	2020	I	41	25 carcinoid	pembrolizumab (PD1) 16 pNETs	carcinoid: median: 20 (range: 2–35) pNET: median: 21 (range: 5–32)	carcinoid: ORR: 12% (95% CI: 2.5–31.2%) PR: 12% (3/25 patients) SD at ≥6 months: 32% (8/25 patients) PD: 28% (7/25 patients) Median PFS in months: 5.6 (range: 3.5–10.7) Median OS in months: 21.1 (95% CI: 9.1–2.4) AEs: 17 (68%) patients SAEs: 8 (32%) patients pNET: ORR: 6.3% (0.2–30.2%) PR: 6.3% (1/16 patients) SD at ≥6 months: 31.3 (5/16 patients) PD: 6.3% (1/16 patients) Median PFS in months: 4.5 (95% CI: 3.6–8.3) Median OS in months: 21 (95% CI: 20.2–not reached) AEs: 11 (68.8%) patients SAEs: 1 (6.3%) patient
Patel et al. (DART SWOG 1609) [48] 6 lung NETs 11 NETs at other sites	2020	II	32	Non-pancreatic NETs: 15 GI NETs	ipilimumab (CTLA4) + nivolumab (PD1)	Up to 15	ORR: 25% (95% CI: 13–42%) CR: 3% (1/32 patients) PR: 22% (7/32 patients) PD: 34% (11/32 patients) Median PFS in months: 4 Median OS in months: 11 AEs: 27 (84.4%) patients SAEs: 16 (50%) patients
Lu et al. [49] 8 NETs at other sites	2020	Ib	40	32 GEP-NETs	toripalimab (PD1)	Up to 24	ORR: 20% (95% CI: 9.1–35.7%) PR: 20% (8/40 patients) SD: 15% (6/40 patients) PD: 80% (24/40 patients) Median PFS in months: 2.5 (95% CI: 1.9–3.1) Median OS in months: 7.8 (95% CI: 5.0–10.8) AEs: 38 (95%) patients SAEs: 11 (27.5%) patients

▶ **Table 1** List of studies evaluating the efficacy of ICI in NETs.

Study	Year	Phase	Number of patients	Types of NET	Type of ICI (target protein)	Duration of follow-up in months	Outcomes
Klein et al. (CA209-538) [58]	2020	II	29	10 bronchial carcinoids 1 lung LCNEC 7 GEP-NETs 3 GEP-NECs 5 thymic NETs 1 prostate NET 1 cervical NET	ipilimumab (CTLA4) + nivolumab (PD1)	26+	ORR: 24% SD: 48.3% (14/29 patients) PD: 10% (3/29 patients) Median PFS in months: 4.82 (95% CI: 2.71–10.53) Median OS in months: 4.78 (95% CI: 4.07–21.25) AEs: 19 (66%) patients SAEs: 10 (34%) patients
Strosberg et al. (KEYNOTE-158) [50]	2020	II	107	83 GEP-NETs 14 lung NETs 10 NETs at other sites	pembrolizumab (PD1)	Median: 24.2 (range: 0.6–33.4)	ORR: 3.7% (95% CI: 1.0–9.3) CR: 0/107 patients PR: 4/107 patients Median PFS in months: 4.1 (95% CI: 3.5–5.4) Median OS in months: 24.2 (95% CI, 15.8–32.5) AEs: 81 (75.7%) patients SAEs: 23 (21.5%) patients
Vijayvergia et al. [51]	2020	II	29	24 GEP-NETs 5 non-GEP-NETs	pembrolizumab (PD1)	Up to 36	ORR: 3.4% (1/29 patients) SD: 20.7% (6/29 patients) DCR: 24.1% (7/29 patients) PD: 58.6% (17/29 patients) Median PFS in months: 2 (95% CI: 6–9.43) Median OS in months: 4.7 (95% CI: 12.86–not estimated) AEs: 36 events SAEs: 9 events
Frumovitz et al. [56]	2020	II	7	6 cervical SCNEC 1 vulvar SCNEC	pembrolizumab (PD1)	Up to 27	ORR: 0% PD: 100% (7/7 patients) at 27 weeks Median PFS in months: 2.1 (range: 0.8–3.3) AEs: 7 events SAEs: 2 events
Özdirik et al. [66]	2020	Retro-spective study	8	1 larynx NET 1 kidney NET 6 GEP-NETs, including a mixed NET	Monotherapy: pembrolizumab (PD1) avelumab (PD-L1) Combination therapy: ipilimumab (CTLA4) + pembrolizumab (PD1)	3–32	PR: 37.5% (3/8 patients) SD: 12.5% (1/8 patients) PD: 50% (4/8 patients)
Sherman et al. [62]	2020	Retro-spective study	23	lung LCNEC	Monotherapy: pembrolizumab (PD1) nivolumab (PD1), atezolizumab (PD-L1) ipilimumab (CTLA4) + pembrolizumab (PD1)	Median: 6.2 (IQR 2.2–12.1)	ORR: 33% CR: 11% (2/23 patients) PR: 22% (4/23 patients) SD: 6% (1/23 patients) PD: 61% (11/23 patients) Median PFS in months: 4.2 (95% CI: 2.4–8.1) Median OS in months: 11.8 (95% CI: 3.7–not reached)

► **Table 1** List of studies evaluating the efficacy of ICI in NETs.

Study	Year	Phase	Number of patients	Types of NET	Type of ICI (target protein)	Duration of follow-up in months	Outcomes
Shirasawa et al. [61]	2020	Retrospective study	13	lung LCNEC	nivolumab (PD1) or pembrolizumab (PD1)	NA	PR: 39% (5/13 patients) SD: 16% (2/13 patients) PD: 45% (6/13 patients) Median PFS in months: 4.2 (95% CI: 2.7–5.7) Median OS in months: 25.2 (95% CI: 21.3–29.1)
Yao et al. [47]	2021	II	116	95 thoracic NETs and GEP-NETs 21 GEP-NECs	spartalizumab (PD1)	Median: 13.4 (range 11–17)	NETs: ORR: 7.4% (95% CI: 3%–14.6%) PR: 7.4% (7/95 patients) SD: 55.8% (55/95 patients) PD: 30.5% (29/95 patients) Median PFS in months: 3.8 Median OS in months: not estimated AEs: 91 (95.8%) patients SAEs: 46 (48.4%) patients NECs: ORR: 4.8% (95% CI: 0.1% - 23.8%) PR: 4.8% (1/21 patients) SD: 14.3% (3/21 patients) PD: 66.7% (14/21 patients) Median PFS in months: 1.8 Median OS in months: 6.8 AEs: 19 (90.5%) patients SAEs: 12 (57.1%) patients
Gile et al. [63]	2021	Retrospective study	57	11 NETs: 8 GEP-NETs 3 NETs at other sites (6 G1 + G2 NETs, and 5 G3 NETs) 46 NECs: 18 GEP-NECs 28 NECs at other sites	Monotherapy: pembrolizumab (PD1) nivolumab (PD1), atezolizumab (PD-L1) Combination therapy: ipilimumab (CTLA4) + pembrolizumab (PD1) ICIs + platinum-based chemotherapy	NETs (all received monotherapy): Median: 79.8 (95% CI: 7.9–251) NECs: Median: 10.7 (95% CI: 1.7–97.9)	G1 + G2 NETs: ORR: 25% CR: 0% PR: 17% (1/6 patients) SD: 17% (1/6 patients) PD: 33% (2/6 patients) Median PFS in months: not reached (95% CI: 1.5–not reached) Median OS in months: 25.2 (95% CI: 2.1–not reached) G3 NETs: ORR: 0% CR: 0% PR: 0% SD: 20% (1/5 patients) PD: 40% (2/5 patients) Median PFS in months: 2.9 (95% CI: 1.4–4.2) Median OS in months: 15.4 (95% CI: 13.7–17.2) NECs: ORR: 19% CR: 2% (1/46 patients) PR: 11% (5/46 patients) SD: 17% (8/46 patients) PD: 39% (18/46 patients) Median PFS in months: NA Median OS in months: 7.2 (95% CI: 3.9–10.8)

► **Table 1** List of studies evaluating the efficacy of ICI in NETs.

Study	Year	Phase	Number of patients	Types of NET	Type of ICI (target protein)	Duration of follow-up in months	Outcomes
Zhang et al. * [53]	2018	Ib	23	5 NETs 15 NECs 3 mixed adenoNECs Sites: GEP and non-GEP	toripalimab (PD1)	NA	ORR: overall: 28.6 % NETs: 40 % NECs: 25 % Median PFS in months: 2.8 (95 % CI: 1.6–4) SAEs 2 (9 %) patients PR: 7 % (2/29 patients)
Fottner et al. * (AVENEC) [54]	2019	II	29	11 G3 NET 16 G3 NEC Sites: both GEP and non-GEP	avelumab (PD-L1)	NA	SD: 14 % (4/29 patients) Median OS in months: 4.2 (1 - > 12) AEs: 11 (38 %) patients
Mulvey et al. * [55]	2019	II	14	extra-pulmonary NETs 11 SCNECs 1 LCNECs 2 not specified	pembrolizumab (PD1)	NA	ORR: 7 % CR: 7 % (1/14 patients) PR: 0 % (0/14 patients) SD: 14 % (2/14 patients) PD: 71 % (10/14 patients) Median PFS in months: 2
Halperin et al. * [52]	2020	II	40	20 pNETs 20 epNETs	atezolizumab (PD-L1) + bevacizumab (VEGF)	NA	pNETs: ORR: 20 % (95 % CI: 6–44 %) Median PFS in months: 19.6 (95 % CI: 10.6–not reached) epNETs: ORR: 15 % (95 % CI: 3–38 %) Median PFS in months: 14.9 (95 % CI: 6.1–not reached)
Rodriguez-Freixinos et al. * [57]	2020	IIb	9	epNECs	avelumab (PD-L1)	NA	ORR: 0 % Median PFS in months: 3 (1–10) Median OS in months: 5 (2–15)
Capdevila et al. * [59]	2020	II	123	cohort (C) 1 lung carcinoids C2. G1/G2 G-HNETs C3. G1/G2 pNETs C4. G3 GEP-NETs	durvalumab (PD-L1) + tremelimumab (CTLA4)	10.8	ORR: C1/2/3/4: 7.4/0%/6.3%/9.1 % respectively. CBR: C1/2/3: 7.4%/32.3%/25 % respectively. OS rate for C4 at 9 months: 36.1 %
Chan et al. * [60]	2021	II	22	extra-pulmonary NETs 8 SCNECs 6 LCNECs 8 not specified	pembrolizumab + irinotecan or paclitaxel	NA	ORR: 9 % PR 9 % (2/22 patients) SD 14 % (3/22 patients) PD 60 % (13/22 patients) Median PFS in months: 2 Median OS in months: 4 SAEs: 7 (32 %) patients
Jimenez et al. [87]	2020	II	11	metastatic PPGL	pembrolizumab (PD1)	Median: 17.9 (IQR: 9.3–20.2)	PR: 9 % (1/11 patients) SD: 64 % (7/11 patients) PD: 27 % (3/11 patients) Median PFS in months: 5.7 (95 % CI: 4.37–not reached) Median OS in months: 19 (95 % CI: 9.9–not reached) AEs: 26 events SAEs: 4 events



► **Table 1** List of studies evaluating the efficacy of ICI in NETs.

Study	Year	Phase	Number of patients	Types of NET	Type of ICI (target protein)	Duration of follow-up in months	Outcomes
Kaufman et al. [102]	2016	II	88	MCC	avelumab (PD-L1)	Median: 10.4 (IQR 8.6–13.1)	ORR: 31.8% (95% CI: 42.3%–79.3%) CR: 9% (8/88 patients) PR: 23% (20/88 patients) SD: 10% (9/88 patients) PD: 36% (32/88 patients) Median PFS: 2.7 months (95% CI 1.4–6.9) Median OS: 11.3 months (7.5–14.0) AEs: 62 (70%) patients SAEs: 4 (5%) patients
D'Angelo et al. [103]	2018	II	29	MCC	avelumab (PD-L1)	Median: 5.1 (range: 0.3–11.3)	3-month ORR 62.1% (95.9% CI: 21.9%–43.1%) CR: 13.8% (4/29 patients) PR: 48.3% (14/29 patients) SD: 10.3% (3/29 patients) PD: 24.1% (7/29 patients) AEs: 28 (71.8%) patients SAEs: 8 (20.5%) patients
Nghiem et al. [101]	2019	II	50	MCC	pembrolizumab (PD1)	Median: 14.9 (range: 0.4–36.4+)	ORR: 56% (95% CI: 41.3% - 70%) CR: 24% (12/50 patients) PR: 32% (16/50 patients) SD: 10% (5/50 patients) PD: 32% (16/50 patients) Median PFS: 16.8 months (95% CI: 4.6 months–not estimable) Median OS: not reached AEs: 50 (100%) patients SAEs: 14 (28%) patients
Topalian et al. (CheckMate 358) [100]	2020	I/II	36	MCC	neoadjuvant nivolumab (PD1) 4 weeks prior to surgery	Median: 20.3 (range: 0.5–39.7)	CR: 47.2% (17/36 patients) PR: 54.5% (18/33 patients) AEs: 18 (46.2%) patients SAEs: 3 (7.7%) patients

NET: Neuroendocrine tumor; GEP: Gastroenteropancreatic; p: Pancreatic; ep: Extrapancreatic; NEC: Neuroendocrine carcinoma; SCNEC: Small cell NEC; LCNEC: Large cell NEC; PPGL: Pheochromocytoma and paraganglioma; MCC: Merkel cell carcinoma; PD-1: Programmed cell death protein-1; PD-L1: PD-ligand 1; CTLA-4: Cytotoxic T-lymphocyte antigen 4; VEGF: Vascular endothelial growth factor; ORR: Objective response rate; PR: Partial response; SD: Stable disease; PD: progressive disease; TTP: Time-to-progression; OS: Overall survival; PFS: Progression-free survival; DCR: Disease control rate; CBR: Clinical benefit rate; CI: Confidence interval; IQR: Interquartile range; AE: Adverse event; SAE: Severe adverse event; NA: Not available; \* These data are currently only published in the form of abstracts.

ors with high tumor mutational burden (TMB) such as pancreatic NECs, a strong correlation between TMB and efficacy of ICI therapy in cancers in general has not been identified to date [64]. In a retrospective study on LCNEC patients, those on anti-PD1 therapy with dense tumoral CD8 + T lymphocyte infiltration ( $> 38$  cells/mm<sup>2</sup>) had a significantly better PFS and OS than patients with lower infiltration density [60]. In this study, 90% of the 13 patients who received anti-PD1 therapy had PD-L1-negative tumors, yet they responded to anti-PD1 therapy. Moreover, three patients with a pathogenic *TP53* variant along with other co-occurring variants such as *PIK3CA* and *RB1* responded well to anti-PD1 therapy, suggesting a probable correlation between presence of certain pathogenic gene variants and response to ICI therapy.

The PD1 inhibitor spartalizumab was utilized to treat 95 patients with thoracic and GEP-NETs as well as 21 GEP-NECs in a phase II study [47]. After a median follow-up of 13.4 months, the ORR in the NET group was 7.4% in the NET patients and 4.8% in the NEC patients, while the median PFS was 3.8 months and 1.8 months in the NET and NEC groups, respectively, with the 12-month Kaplan-Meier estimated median PFS rate of 19.5% in the NET group and 0% in the NEC group. The therapy was also associated with SAEs in close to half the patients in both NET and NEC cohorts, and a quality-of-life assessment did not reveal any meaningful improvement from the PD1 therapy in the global health status or functional scales in these patients. Forty patients with recurrent/metastatic GEP and non-GEP-NETs were treated with the PD1 inhibitor toripalimab in a phase Ib study [49]. The ORR was 20% and the disease control rate (DCR) was 35%, with a median duration of response of 15.2 months. Patients with high PD-L1 expression ( $> 10\%$ ) and a high TMB had better ORRs compared to patients with low PD-L1 expression and a low TMB. One patient who responded to therapy had multiple genomic rearrangement with high prediction score for neoantigens, but interestingly had low TMB, negative PD-L1 expression, and without microsatellite instability. SAEs were noted in  $< 30\%$  of the patients.

PD-L1 inhibitor monotherapy has also been evaluated with avelumab, and atezolizumab [53, 56, 61, 62, 65], although most data are from retrospective studies in which other ICIs including PD1 inhibitors or combination ICI therapies were utilized (► **Table 1**). In a phase II study, combination therapy with the PD-L1 inhibitor atezolizumab + bevacizumab in 40 pancreatic and extrapancreatic NETs led to an ORR of 20 and 15%, and a median PFS of 19.6 months and 14.9 months, respectively [66]. A combination of PD1 or PD-L1 + CTLA4 inhibitors have also been utilized in various studies, including the combinations of nivolumab and ipilimumab [48, 57], pembrolizumab and ipilimumab [61, 62, 65], and durvalumab and tremelimumab [58]. The ORRs with combination therapies have been better compared to monotherapy with PD1 inhibitors, ranging from 24% to 33%, with a comparable AE profile (► **Table 1**).

A meta-analysis on 14 phase I/II studies was performed by Bongiovanni et al. to evaluate the safety and efficacy of ICI therapy in NETs [67]. The efficacy data were available from 636 patients. The pooled ORR was 10% (95% CI: 6–15%;  $I^2 = 67\%$ ), with a DCR of 42% (95% CI: 28–56%,  $I^2 = 93\%$ ). The highest ORR was noted with toripalimab, and combination regimens of PD1 + CTLA4 inhibitors (nivolumab + ipilimumab) or PD-L1 + VEGF inhibitors (atezolizumab + bevacizumab) were superior compared to PD1 inhibitor mon-

otherapy. The DCR was better for G1/G2 NETs as compared to G3 NETs and NECs, but the DCR was not significantly different based on the site of origin of the tumors. Most common AEs noted were dermatologic conditions (rash, pruritis, dermatitis), fatigue, gastrointestinal symptoms, transaminase elevation, hypothyroidism, and loss of appetite. SAEs were noted at a rate of 22% for treatment-related AEs and 18% for immune-related AEs. The median PFS was 4.1 months (95% CI: 2.6–5.4;  $I^2 = 96\%$ ), and the median OS was 11 months (95% CI: 4.8–21.1;  $I^2 = 98\%$ ). A sub-analysis of studies in which PD-L1 expression available [49, 50, 52, 58] revealed that patients with tumors positive for PD-L1 expression had a better ORR compared to patients without PD-L1 tumor expression. Another meta-analysis by Park et al. compared 10 studies comprised of 464 patients with advanced/metastatic NETs [68]. The pooled ORR was 15.5% (95% CI: 9.5–24.3%;  $I^2 = 72\%$ ). The ORR was better with thoracic NETs (24.7%) compared to GEP-NETs (9.5%). Interestingly, poorly-differentiated NET group had better ORR (22.7%) compared to the well-differentiated NET group (10.4%), with the probable explanation being that the poorly-differentiated NETs may have a higher PD-L1 expression and a higher TMB [69]. The median PFS was 3.8 months (95% CI: 3.5–4.1), and the median OS was 22.7 months (95% CI: 20.1–25.9), with the shortest median OS noted with poorly-differentiated GEP-NETs and longest OS with well-differentiated thoracic NETs. Similar to the first meta-analysis, combination therapy resulted in better ORRs compared to monotherapy. The differences noted with ORRs between this meta-analysis and the meta-analysis by Bongiovanni et al. was due to different set of studies included: the Park et al. study included only full-length studies (phase I/II studies and retrospective studies), while the Bongiovanni et al. study included phase I/II full-length studies and abstracts but did not include retrospective studies. In comparison, the ORRs and survival outcomes, and AE profiles have been somewhat better with some of the other established systemic therapies for NETs as summarized in ► **Table 2**. However, it must be noted that these are not head-to-head comparisons, and the type of study is not the same across the articles described in ► **Table 2**.

## Medullary thyroid cancer

While surgery and systemic chemotherapy were the main treatment options available for MTC for several years [11], the management of advanced/metastatic MTC has dramatically changed over the past decade, mainly due to the advent of the tyrosine kinase inhibitors (TKIs) such as vandetanib and cabozantinib [70, 71], and of selective rearranged during transfection (RET) inhibitors, including selpercatinib and pralsetinib [72, 73]. As with other NETs, several studies have utilized PRRT to treat MTC with reasonable success [74–79]. As with other NETs, the utility of ICI in MTC is yet to be thoroughly investigated.

The immune landscape of MTC has been described in a few studies, although the implications of the findings in the clinical management of these tumors needs to be established. PD1/PD-L1 expression has been identified in certain MTC post-surgical specimens, and PD-L1 expression in the tumor cells and the associated immune cells has been shown to be associated with distant metastases at the time of surgery and co-expression of PD1/PD-L1 has shown significant association with advanced stages (III/IV), worse OS, but not with worse PFS [80]. Other immune checkpoint-relat-

► **Table 2** A comparison of outcomes of ICI therapy and other systemic therapies in NETs.

Study	Year	Study type	Number of patients	Types of NET	Type of treatment	Outcomes
Bongiovanni et al. [67]	2021	Meta-analysis	636	Multiple	Various ICIs (monotherapy and combination)	ORR: 10% (95% CI: 6–15%) DCR: 42% (95% CI: 28–56%) Median PFS in months: 4.1 (95% CI: 2.6–5.4) Median OS in months: 11 (95% CI: 4.8–21.1) trSAEs: 22% (95% CI: 13–32%)
Park et al. [68]	2022	Meta-analysis	464	Multiple	Various ICIs (monotherapy and combination)	ORR: 15.5% (95% CI: 9.5–24.3%) Median PFS in months: 3.8 (95% CI: 3.5–4.1) Median OS in months: 22.7 (95% CI: 20.1–25.9) SAEs: NA
Strosberg et al. (NETTER-1) [32]	2017	Phase III RCT, LAR octreotide alone as control group	116 in treatment group	Midgut NETs	<sup>177</sup> Lu-DOTATATE (with LAR octreotide)	ORR: 18% Estimated PFS: 65% trSAEs: 9%
Yao et al. (RADIANT-4) [26]	2015	Phase III, placebo-controlled RCT	203 in treatment group	WD lung and GEP-NETs	everolimus	Median PFS: 11 months (95% CI: 9.2–13.3) trSAEs: <1–9%
Raymond et al. [28]	2011	Phase III, placebo-controlled RCT	86 in treatment group	WD pNETs	sunitinib	ORR: 9.3% Median PFS: 11.4 months SAEs: 1–12%
Caplin et al. [23]	2014	Phase III, placebo-controlled RCT	101 in treatment group	EP-NETs	lanreotide	Median PFS: not reached SAE: 26%
Rinke et al. (PROMID) [21]	2009	Phase IIIb, placebo-controlled RCT	42 in treatment group	midgut NETs	octreotide LAR	Median TTP: 14.3 months SD: 66.7% at 6 months serious AEs: 11 patients
Al-Toubah et al. [35]	2021	Retrospective analysis	462	WD and PD NETs from multiple sites	capecitabine + temozolomide	ORR: 46% DCR: 81% Median PFS in months: 18 (95% CI: 14.0–21.9) Median OS in months: 51 (95% CI: 42.8–59.2) trSAEs: 0.2–8.4%

NETs: Neuroendocrine tumors; ICI: Immune checkpoint inhibitor; RCT: Randomized controlled trial; GEP: Gastroenteropancreatic; EP: Enteropancreatic; p: Pancreatic; ORR: Objective response rate; DCR: Disease control rate; PFS: Progression-free survival; AE: Adverse events; SAEs: Severe adverse events; tr: Treatment-related; WD: Well-differentiated; PD: Poorly-differentiated; LAR: Long-acting repeatable; NA: Not available.

ed candidate antigens such as CD276 have been also identified to be overexpressed in MTC cells as compared to normal thyroid tissue [81]. In a study evaluating tissue microarray expression of immune inhibitory receptors expression comprised of CTLA-4, PD1, TIM-3, LAG-3, and TIGIT in MTC surgical specimens from 200 patients, positive expression was identified at variable levels, ranging from 48% of patients for TIM-3 positivity to 3% for LAG-3 and TIGIT [82]. In this study, CTLA-expression, PD-1/PD-L1 co-expression, and TIM-3 expression were associated with worse recurrence-free survival, and moderate to strong CTLA-4, PD-1, or PD-L1 expression along with consistent TIM-3 expression was noted in MTC of patients who developed advanced disease. Currently, data on ICI therapy in thyroid cancer, including MTC, is extremely sparse [83, 84]. In one report, a patient with advanced, metastatic MTC demonstrated substantial improvement in calcitonin doubling time and tumor burden following yeast-CEA vaccine, followed by surgery and then ICI therapy with avelumab under a phase I trial [85]. Further studies are needed to establish the role of ICI therapy in MTC. For instance, a phase 2 clinical trial (NCT03246958) is evaluating the safety and efficacy of the combination of nivolumab and ipilimumab in the treatment of aggressive thyroid cancer, including cohorts of MTC.

### Pheochromocytomas and paragangliomas

Current standard of care of PPGL includes surgery, locoregional interventions such as radiofrequency or cryoablation and chemoembolization, temozolomide, TKIs, and PRRT [17, 86]. PPGLs with germline pathogenic variants can have varying biochemical and phenotypic presentations (clusters 1, 2, and 3), and the current management of PPGL has been steering towards personalized/targeted therapy, for instance, hypoxia pathway-targeting agents for treating cluster 1 tumors and TKIs for treating cluster 2 tumors [86]. Similar to MTC and other NETs, the utility of ICI therapy has been understudied in PPGL. In a phase II study, pembrolizumab was utilized in 11 patients with progressive, metastatic PPGL, eight of whom at least had prior surgery with or without other systemic therapy (▶ **Table 1**) [87]. The primary endpoint of non-progression at 27 weeks was observed in four patients, while the ORR was 9%, and a clinical benefit rate of 73%. Grade 3 adverse events were noted in four patients while none had Grade 4 or 5 adverse events. However, the favorable treatment responses did not correlate with primary tumor PD-L1 positivity, hormonal status, hereditary syndrome status, or infiltrating mononuclear cells in the primary tumor. A combination therapy with ipilimumab and nivolumab used off-label in a 60-year-old patient with sporadic, metastatic, inoperable pheochromocytoma resulted in substantial reduction in tumor burden after close to 20 months of therapy [88].

### Pituitary tumors

Pituitary tumors are considered as a subtype of NETs and the potential for the utility of ICI in the treatment of these tumors has recently gained some interest [89–91]. While most pituitary tumors are adenomas and can be cured with surgery, the gross tumor resection rate is about 66.4–74% [92]. Also, the so-called refractory adenomas which tend to be more invasive with a high Ki-67 index as well as pituitary carcinomas can cause substantial morbidity and

mortality and are challenging to treat [93]. Temozolomide has been used to treat aggressive forms of pituitary tumors but only about 60% of the tumors respond to this treatment [91]. The immune cell population seems to be different between normal pituitary gland and pituitary adenomas, and among different pituitary adenomas, 3 distinct immunophenotypic clusters of pituitary adenomas have been identified with each cluster comprising a different set of immune checkpoint molecular expression [94]. Later studies have also identified increased PD-L1 expression in pituitary tumors invading the cavernous sinus and in functional and more aggressive adenomas [89, 95]. Anti-PD-L1 therapy has demonstrated reduction in tumor growth and in ACTH levels, and improved survival in murine models of Cushing's disease [96]. In a recent case report, ipilimumab and nivolumab combination therapy followed by maintenance therapy with nivolumab in a 41-year-old patient with recurrent, invasive adrenocorticotropic hormone (ACTH)-producing pituitary adenoma (refractory to bilateral adrenalectomy and temozolomide therapy) led to biochemical response with reduction in ACTH and cortisol levels, and radiographically stable disease 12 months into ICI therapy [97]. In another report, immunotherapy with autoantigens along with a T helper 1 adjuvant for 24 consecutive weeks resulted in substantial biochemical, radiographic, and clinical response in a 31-year-old lady with refractory macroprolactinoma [98]. Further studies on a larger cohort are needed to establish the role of ICIs and other immunotherapies in the management of aggressive pituitary adenomas and pituitary carcinomas.

### Merkel cell carcinoma

MCC is a rare, aggressive form of non-melanoma skin cancer predominantly occurring in the sun-exposed areas in older, fair-skinned individuals [99]. Due to its immunogenic nature, these tumors can be targeted with ICIs. While CTLA-4 inhibitors are not well-studied in this condition, the PD-L1 inhibitor, avelumab and PD-1 inhibitors pembrolizumab and nivolumab have been studied in advanced MCCs in various trials, including KEYNOTE-017, CheckMate 358, and JAVELIN Merkel 200-part A and part B trials [100–103] (▶ **Table 1**). ICI therapy in MCC is associated with higher ORRs (31.8–62.1%) compared to conventional NETs [101–103], and a neoadjuvant approach has also been utilized with tumor regression in close to 50% of the patients [100]. Further optimization of ICI-based treatment is being evaluated in early phase trials targeting other immune checkpoint markers such as TIM-3, LAG-3, TIGIT in advanced cancers [99].

### Summary of the current knowledge

In general, the role of ICI in the treatment of NETs has not been well-established, and the currently available data demonstrate only modest efficacy. Combination ICI therapy is superior to monotherapy. Some of the factors that determine favorable response to ICI therapy include aggressive tumor biology, high TMB, higher T lymphocyte and other inflammatory cell infiltration into the tumor microenvironment, and presence of certain additional pathogenic gene variants. On the other hand, the extent of PD-L1 expression has not shown clear correlation with response to ICI therapy. The NCCN 2021 guidelines recommend the use of pembrolizumab as a monotherapy as a category 2B recommendation (low-level evi-

dence and NCCN consensus that a therapy is appropriate) in patients with NETs harboring high-TMB (> 10 mutations/Megabase) as confirmed by an FDA-approved test, that have progressed on prior therapies and there are no alternative satisfactory therapies available [20]. The combination of ipilimumab and nivolumab is also recommended (category 2B) by the NCCN guidelines for patients with locally advanced or metastatic NETs with unfavorable tumor biology as an alternative to clinical trials [20]. The NANETS 2021 guidelines comment on ICI therapy particularly for pancreatic NETs and endorse the minimal treatment benefit noted in NET patients with single agent PD-1/PD-L1 therapy and recommend the use of immunotherapy for pancreatic NECs in a clinical trial setting [36, 104]. Similarly, NANETS guidelines on PPGL also acknowledge the lack of knowledge on the mechanisms that determine favorable outcomes with ICI therapy in PPGL and suggest the use of ICI to be limited to clinical trials [17, 36].


### Future directions

In addition to ICI therapy *per se*, targeting additional immunotherapeutic mechanisms may enhance the anti-tumor activity of ICIs, especially in NETs which are particularly indolent and relatively less sensitive to ICI therapy compared to other tumors. Some of these targets include alteration of the tumor microenvironment and tumor vasculature, T cell homing, prevention of T cell exhaustion, enhancement of metabolic pathways or cytokines that sustain a robust CD8+ lymphocyte response, and vaccination of a given patient with the antigens derived from their tumor cells [105]. These strategies may hold potential as combination therapies along with ICIs. The intratumoral heterogeneity exhibited at cellular and genomic levels also contributes towards variable immune response towards these tumors and the relative resistance towards several targeted therapies and may partly explain the biology of treatment-refractory NETs [105, 106]. Understanding the immune-microenvironmental mechanisms driving the intratumoral heterogeneity and identifying potential treatment targets may allow for further optimization of immunotherapy in NETs. Some progress has been made in the description of the NET immune-microenvironment. The immune-microenvironment seems to be higher in pancreatic NETs compared to midgut NETs but without any clear association with expression of immune checkpoint markers or mutational profile [107], and a higher density of T cell infiltration in pancreatic NET primary tumors has been associated with a higher recurrence-free survival meanwhile a high regulatory T cell infiltration has been associated with a lower OS among patients with liver NET metastases [108].

In general, NETs are described as having a 'cold' tumor microenvironment which is thought to be the reason for the modest efficacy of ICI therapy. One of the main areas of further research lies in exploring the mechanisms that can turn these immunologically 'cold' tumors into 'hot' (more immunogenic) tumors. The three major immunologically 'cold' cancer phenotypes described include: 1) the immune desert phenotype which comprises tumors that lack T lymphocyte priming, suboptimal antigen processing and presentation, and lack of antigen-presenting cell – T lymphocyte interaction, 2) the immune excluded phenotype in which the T lymphocytes do not effectively infiltrate the tumor, and 3) the immune inflamed phenotype in which the T lymphocytes infiltrate the tumor but these cells are rendered ineffective either due to T

cell exhaustion or due to checkpoint activation [109]. Several mechanisms underlie the 'cold' tumor phenotype, including low multiple histocompatibility complex I (MHC I) expression, low TMB, activation of certain oncogenic pathways, epigenetic modifications, altered tumor vasculature, tumor hypoxia, tumor microbiome, immunosuppressive tumor microenvironment, among others [109]. The molecular landscape of both sporadic and familial NETs demonstrates involvement of protooncogenes as well as tumor suppressor genes, several of which are involved in the development of one or more of the above-described immune evasion phenotypes of tumors [109, 110]. Certain molecular phenotypes such as metastasis-like primary-1 (MLP-1) subtype of pancreatic NETs are associated with worse prognosis, increased levels of immune-related genes expression including T cell-inflamed-related genes, immune checkpoint antigens, and other immune evasion mechanisms, and such enhanced immune-related gene expressions are associated with hypoxia and necroptosis in pancreatic NETs [111]. Several approaches have been attempted to convert the immunologically 'cold' tumors into 'hot' tumors. Some of these mechanisms include promoting T cell priming (immune adjuvants, oncolytic viruses, chemotherapy/radiation mediating an 'abscopal effect', local ablative therapies), antigen-specific T cell expansion (adoptive cellular therapy such as CAR-T cells, anticancer vaccines), and improving T cell trafficking and infiltration (oncogenic pathway inhibitors, epigenetic modifier inhibitors, antiangiogenic therapies, TGF $\beta$  inhibitors, CXCR4 inhibitors) [112–121]. Some of these mechanisms may hold the key to enhancing the response of NETs to ICI or other forms of immunotherapy (► Fig. 5). Further details on the mechanisms on converting immunologically 'cold' into 'hot' tumors are described elsewhere [109].

Apart from the canonical targets of CTLA-4, PD-1, or PD-L1, targeting other components related to immunoregulation may serve as alternative therapy or augment the clinical efficacy of ICIs. For instance, targeting indoleamine 2,3-dioxygenase, an enzyme that plays a role in immune evasion in cancers may potentiate the effects of ICIs [122]. Other proteins involved in immune checkpoint cascade, including TIM-3, LAG-3, and TIGIT also serve as potential targets for novel therapy in the management of NETs [82, 99]. Other strategies such as targeted arterial injection of recombinant viruses or vaccination against anti-apoptotic molecules such as surviving combined with immunogenic adjuvants are being evaluated for the treatment of NETs [122]. It is possible that those NETs that are deemed unlikely to respond to PRRT due to lack of avidity on diagnostic SSA-based imaging, may in fact be candidates for ICI therapy. The reason for this 'flip-flop' phenomenon could be because the less-avid NET lesions tend to be dedifferentiated, which may in turn translate to increased TMB and immunogenicity leading to increased susceptibility towards ICI therapy. This mechanism is probably analogous to the flip-flop phenomenon observed with differentiated thyroid cancers, in which tumors that are radioiodine non-avid tend to be avid on fluorodeoxyglucose (FDG)-PET/CT scan [123]. In this context, it is important to be aware of the distribution of SSTRs 1–5 in normal human tissue and a normative database [124]. The expression profile of neuropeptide receptors can vary across different types of immune cells [125]. Human monocytes express SSTR2A and SSTR1 when induced to differentiate into macrophages or dendritic cells. NETs can be infiltrated by lymphocytes,



► **Fig. 5** Various mechanisms through which immunologically ‘cold’ NETs can be potentially converted into immunologically ‘hot’ NETs. The data are partially adapted from Liu et al. (2021) [109] (Created with BioRender.com). [rerif]

as shown by immunohistochemistry for CD3, CD4, and CD8. Heavier tumor infiltration by T regulatory cells is associated with weaker anti-tumor immunity [126–128]. A high-density SSTR expression occurs not only on tumor cells but also on peritumoral vessels, activated lymphocytes and monocytes. Somatostatin can inhibit inflammation both locally and distant from the site of release [129]. Certain drugs, for instance, valproic acid which can inhibit the histone deacetylase, can elicit an upregulation of *SSTR2* mRNA and protein expression in human NET cells [130, 131]. Glucocorticoids are anti-inflammatory and in patients with ectopic ACTH secretion and cortisol excess the use of selective non-steroidal glucocorticoid receptor antagonist/modulator mifepristone and relacorilant can lead to an upregulation of *SSTR2* expression in ACTH-secreting neuroendocrine tumors [132]. Inhibition of proprotein convertase subtilisin/kexin type 9 (PCSK9) can lead to increased MHC I expression on tumor cells leading to augmented intratumoral CD8 + lymphocyte infiltration [133]. Whether these mechanisms would turn “cold” NETs into “hot” tumors with regards to improving T-cell infiltration and thereby making such NET more responsive to ICI therapy needs to be shown.

Studies in mouse models have revealed that the effects of ICIs are potentially modulated by the gut microbiome, which is in part mediated through certain microbiome-derived metabolites such as inositol [134]. Studies of the human microbiome and its impact on the efficacy of immunotherapy on NETs needs further investigation. Although ICI therapy is technically effective in treating tumors with high-TMB [68], particularly with NECs [135], the cut-offs associated with TMBs have thus far been inconsistent with the predictability of response to ICI therapy, and on some other cancers, ICI therapy has not resulted in improved ORR among patients with high-TMB as compared to patients with low-TMB [64]. The current FDA-approved indication for the use of pembrolizumab on the basis of high-TMB may be too broad and further tailoring of indications based on other factors such as environmental carcinogen exposure are being suggested for consideration [136].

Several clinical trials are ongoing to investigate the role of ICI therapy (clinicaltrials.gov), particularly in conjunction with other therapies such as VEGF-inhibitors (NCT05000294), TKIs (NCT04197310), platinum-based chemotherapy (NCT03980925), stereotactic radiation (NCT03110978), and <sup>177</sup>Lu-DOTATATE (NCT04525638), for the treatment of NETs and NECs. Deciphering the molecular mechanisms and extraneous factors that modulate the immunogenicity of NETs, and further research on systemic therapies or other agents that could potentially enhance the effects of ICIs hold the key to progressing the field of immunotherapy in the management of NETs.

## Acknowledgements

This work was funded in part by the intramural program of the National Institute of Diabetes and Digestive and Kidney Diseases of the National Institutes of Health.

## Conflict of Interest

Dr. Vijayvergia has a consulting/advisory role for Novartis, Tersera, Ipsen, Astra Zeneca, ITM, Taiho, Halio Dx, and there are stock and other ownership interests: Pfizer. Research Funding: Bayer, Puma.

## References

- [1] Yao JC, Hassan M, Phan A et al. One hundred years after “carcinoid”: epidemiology of and prognostic factors for neuroendocrine tumors in 35,825 cases in the United States. *J Clin Oncol* 2008; 26: 3063–3072
- [2] Ito T, Sasano H, Tanaka M et al. Epidemiological study of gastroenteropancreatic neuroendocrine tumors in Japan. *J Gastroenterol* 2010; 45: 234–243

- [3] Fraenkel M, Kim MK, Faggiano A et al. Epidemiology of gastroenteropancreatic neuroendocrine tumours. *Best Pract Res Clin Gastroenterol* 2012; 26: 691–703
- [4] Vinik AI, Woltering EA, Warner RR et al. NANETS consensus guidelines for the diagnosis of neuroendocrine tumor. *Pancreas* 2010; 39: 713–734
- [5] Rosai J. The origin of neuroendocrine tumors and the neural crest saga. *Mod Pathol* 2011; 24: S53–S57
- [6] Kulke MH, Shah MH, Benson AB 3rd et al. Neuroendocrine tumors, version 1.2015. *J Natl Compr Canc Netw* 2015; 13: 78–108
- [7] Koch CA, Petersenn S. Neuroendocrine neoplasms - think about it and choose the most appropriate diagnostic and therapeutic steps. *Rev Endocr Metab Disord* 2018; 19: 107–109
- [8] Koch CA, Petersenn S. Black swans - neuroendocrine tumors of rare locations. *Rev Endocr Metab Disord* 2018; 19: 111–121
- [9] Koch CA, Azumi N, Furlong MA et al. Carcinoid syndrome caused by an atypical carcinoid of the uterine cervix. *J Clin Endocrinol Metab* 1999; 84: 4209–4213
- [10] Singer J, Werner F, Koch CA et al. Ectopic Cushing's syndrome caused by a well differentiated ACTH-secreting neuroendocrine carcinoma of the ileum. *Exp Clin Endocrinol Diabetes* 2010; 118: 524–529
- [11] Wells SA Jr., Asa SL, Dralle H et al. Revised American thyroid association guidelines for the management of medullary thyroid carcinoma. *Thyroid* 2015; 25: 567–610
- [12] Chen H, Sippel RS, O'Dorisio MS et al. The north American neuroendocrine tumor society consensus guideline for the diagnosis and management of neuroendocrine tumors: pheochromocytoma, paraganglioma, and medullary thyroid cancer. *Pancreas* 2010; 39: 775–783
- [13] Assarzadegan N, Montgomery E. What is new in the 2019 World Health Organization (WHO) classification of tumors of the digestive system: review of selected updates on neuroendocrine neoplasms, appendiceal tumors, and molecular testing. *Arch Pathol Lab Med* 2021; 145: 664–677
- [14] Velayoudom-Céphise FL, Duvillard P, Foucan L et al. Are G3 ENETS neuroendocrine neoplasms heterogeneous? *Endocr Relat Cancer* 2013; 20: 649–657
- [15] Strosberg JR, Halfdanarson TR, Bellizzi AM et al. The north American neuroendocrine tumor society consensus guidelines for surveillance and medical management of midgut neuroendocrine tumors. *Pancreas* 2017; 46: 707–714
- [16] Singh S, Bergsland EK, Card CM et al. Commonwealth neuroendocrine tumour research collaboration and the north American neuroendocrine tumor society guidelines for the diagnosis and management of patients with lung neuroendocrine tumors: an international collaborative endorsement and update of the 2015 European neuroendocrine tumor society expert consensus guidelines. *J Thorac Oncol* 2020; 15: 1577–1598
- [17] Fishbein L, Del Rivero J, Else T et al. The north American neuroendocrine tumor society consensus guidelines for surveillance and management of metastatic and/or unresectable pheochromocytoma and paraganglioma. *Pancreas* 2021; 50: 469–493
- [18] Grandhi MS, Lafaro KJ, Pawlik TM. Role of locoregional and systemic approaches for the treatment of patients with metastatic neuroendocrine tumors. *J Gastrointest Surg* 2015; 19: 2273–2282
- [19] Kennedy A, Bester L, Salem R et al. Role of hepatic intra-arterial therapies in metastatic neuroendocrine tumours (NET): guidelines from the NET-liver-metastases consensus conference. *HPB (Oxford)* 2015; 17: 29–37
- [20] Shah MH, Goldner WS, Benson AB et al. Neuroendocrine and adrenal tumors, version 2.2021, NCCN clinical practice guidelines in oncology. *J Natl Compr Canc Netw* 2021; 19: 839–868
- [21] Rinke A, Müller HH, Schade-Brittinger C et al. Placebo-controlled, double-blind, prospective, randomized study on the effect of octreotide LAR in the control of tumor growth in patients with metastatic neuroendocrine midgut tumors: a report from the PROMID Study Group. *J Clin Oncol* 2009; 27: 4656–4663
- [22] Rinke A, Wittenberg M, Schade-Brittinger C et al. Placebo-controlled, double-blind, prospective, randomized study on the effect of octreotide LAR in the control of tumor growth in patients with metastatic neuroendocrine midgut tumors (PROMID): results of long-term survival. *Neuroendocrinology* 2017; 104: 2632
- [23] Caplin ME, Pavel M, Ćwikła JB et al. Lanreotide in metastatic enteropancreatic neuroendocrine tumors. *N Engl J Med* 2014; 371: 224–233
- [24] Caplin ME, Pavel M, Ćwikła JB et al. Anti-tumour effects of lanreotide for pancreatic and intestinal neuroendocrine tumours: the CLARINET open-label extension study. *Endocr Relat Cancer* 2016; 23: 191–199
- [25] Kaderli RM, Spanjol M, Kollár A et al. Therapeutic options for neuroendocrine tumors: a systematic review and network meta-analysis. *JAMA Oncol* 2019; 5: 480–489
- [26] Yao JC, Fazio N, Singh S et al. Everolimus for the treatment of advanced, non-functional neuroendocrine tumours of the lung or gastrointestinal tract (RADIANT-4): a randomised, placebo-controlled, phase 3 study. *Lancet* 2016; 387: 968–977
- [27] Pavel ME, Hainsworth JD, Baudin E et al. Everolimus plus octreotide long-acting repeatable for the treatment of advanced neuroendocrine tumours associated with carcinoid syndrome (RADIANT-2): a randomised, placebo-controlled, phase 3 study. *Lancet* 2011; 378: 2005–2012
- [28] Raymond E, Dahan L, Raoul JL et al. Sunitinib malate for the treatment of pancreatic neuroendocrine tumors. *N Engl J Med* 2011; 364: 501–513
- [29] Raymond E, Kulke MH, Qin S et al. Efficacy and safety of sunitinib in patients with well-differentiated pancreatic neuroendocrine tumours. *Neuroendocrinology* 2018; 107: 237–245
- [30] Yao JC, Guthrie KA, Moran C et al. Phase III prospective randomized comparison trial of depot octreotide plus interferon alfa-2b versus depot octreotide plus bevacizumab in patients with advanced carcinoid tumors: SWOG S0518. *J Clin Oncol* 2017; 35: 1695–1703
- [31] Kölbl L, Persson G, Franzén S et al. Randomized clinical trial of the effect of interferon alpha on survival in patients with disseminated midgut carcinoid tumours. *Br J Surg* 2003; 90: 687–693
- [32] Strosberg J, El-Haddad G, Wolin E et al. Phase 3 trial of (177) Lu-dotatate for midgut neuroendocrine tumors. *N Engl J Med* 2017; 376: 125–135
- [33] Mityr E, Baudin E, Ducreux M et al. Treatment of poorly differentiated neuroendocrine tumours with etoposide and cisplatin. *Br J Cancer* 1999; 81: 1351–1355
- [34] Welin S, Sorbye H, Sebjornsen S et al. Clinical effect of temozolomide-based chemotherapy in poorly differentiated endocrine carcinoma after progression on first-line chemotherapy. *Cancer* 2011; 117: 4617–4622
- [35] Al-Toubah T, Pelle E, Valone T et al. Efficacy and toxicity analysis of capecitabine and temozolomide in neuroendocrine neoplasms. *J Natl Compr Canc Netw* 2021; 20: 29–36
- [36] NANETS. 2021 NANETS guidelines compendium [https://nanets.net/images/guidelines/2021\\_NANETS\\_Guidelines\\_Compendium.pdf](https://nanets.net/images/guidelines/2021_NANETS_Guidelines_Compendium.pdf) accessed on April 5th, 2022
- [37] Sharma P, Siddiqui BA, Anandhan S et al. The next decade of immune checkpoint therapy. *Cancer Discov* 2021; 11: 838–857
- [38] Sharma P, Allison JP. Immune checkpoint targeting in cancer therapy: toward combination strategies with curative potential. *Cell* 2015; 161: 205–214

- [39] Overman MJ, Lonardi S, Wong KYM et al. Durable clinical benefit with nivolumab plus ipilimumab in DNA mismatch repair-deficient/microsatellite instability-high metastatic colorectal cancer. *J Clin Oncol* 2018; 36: 773–779
- [40] Hellmann MD, Paz-Ares L, Bernabe Caro R et al. Nivolumab plus ipilimumab in advanced non-small-cell lung cancer. *N Engl J Med* 2019; 381: 2020–2031
- [41] Larkin J, Chiarion-Sileni V, Gonzalez R et al. Five-year survival with combined nivolumab and ipilimumab in advanced melanoma. *N Engl J Med* 2019; 381: 1535–1546
- [42] Schoenfeld JD, Hanna GJ, Jo VY et al. Neoadjuvant nivolumab or nivolumab plus ipilimumab in untreated oral cavity squamous cell carcinoma: a phase 2 open-label randomized clinical trial. *JAMA Oncol* 2020; 6: 1563–1570
- [43] Gettinger SN, Redman MW, Bazhenova L et al. Nivolumab plus ipilimumab vs nivolumab for previously treated patients with stage IV squamous cell lung cancer: the lung-MAP S1400I phase 3 randomized clinical trial. *JAMA Oncol* 2021; 7: 1368–1377
- [44] Anderson AC, Joller N, Kuchroo VK. Lag-3, Tim-3, and TIGIT: co-inhibitory receptors with specialized functions in immune regulation. *Immunity* 2016; 44: 989–1004
- [45] Qin S, Xu L, Yi M et al. Novel immune checkpoint targets: moving beyond PD-1 and CTLA-4. *Mol Cancer* 2019; 18: 155
- [46] Mehnert JM, Bergsland E, O'Neil BH et al. Pembrolizumab for the treatment of programmed death-ligand 1-positive advanced carcinoid or pancreatic neuroendocrine tumors: Results from the KEYNOTE-028 study. *Cancer* 2020; 126: 3021–3030
- [47] Yao JC, Strosberg J, Fazio N et al. Spartalizumab in metastatic, well/poorly-differentiated neuroendocrine neoplasms. *Endocr Relat Cancer* 2021 ERC-20-0382.R1. doi:10.1530/ERC-20-0382 Online ahead of print
- [48] Patel SP, Othus M, Chae YK et al. A phase II basket trial of dual anti-CTLA-4 and anti-PD-1 blockade in rare tumors (DART SWOG 1609) in patients with nonpancreatic neuroendocrine tumors. *Clin Cancer Res* 2020; 26: 2290–2296
- [49] Lu M, Zhang P, Zhang Y et al. Efficacy, safety, and biomarkers of toripalimab in patients with recurrent or metastatic neuroendocrine neoplasms: a multiple-center phase Ib trial. *Clin Cancer Res* 2020; 26: 2337–2345
- [50] Strosberg J, Mizuno N, Doi T et al. Efficacy and safety of pembrolizumab in previously treated advanced neuroendocrine tumors: results from the phase II KEYNOTE-158 study. *Clin Cancer Res* 2020; 26: 2124–2130
- [51] Vijayvergia N, Dasari A, Deng M et al. Pembrolizumab monotherapy in patients with previously treated metastatic high-grade neuroendocrine neoplasms: joint analysis of two prospective, non-randomised trials. *Br J Cancer* 2020; 122: 1309–1314
- [52] Zhang P, Lu M, Li J et al. Efficacy and safety of PD-1 blockade with JS001 in patients with advanced neuroendocrine neoplasms: A non-randomized, open-label, phase Ib trial. *Ann Oncol* 2018; 29: viii468
- [53] Fottner C, Apostolidis L, Ferrata M et al. A phase II, open label, multicenter trial of avelumab in patients with advanced, metastatic high-grade neuroendocrine carcinomas NEC G3 (WHO 2010) progressive after first-line chemotherapy (AVENEC). *J Clin Oncol* 2019; 37: 4103–4103
- [54] Mulvey C, Raj NP, Chan JA et al. Phase II study of pembrolizumab-based therapy in previously treated extrapulmonary poorly differentiated neuroendocrine carcinomas: results of part A (pembrolizumab alone). *J Clin Oncol* 2019; 37: 363–363
- [55] Frumovitz M, Westin SN, Salvo G et al. Phase II study of pembrolizumab efficacy and safety in women with recurrent small cell neuroendocrine carcinoma of the lower genital tract. *Gynecol Oncol* 2020; 158: 570–575
- [56] Rodriguez-Freixinos V, Chan D, Doherty M et al. Avelumab in unresectable/metastatic, progressive, poorly differentiated, grade 3 neuroendocrine carcinomas (NECs). *Neuroendocrinology* 2020; 217–217
- [57] Klein O, Kee D, Markman B et al. Immunotherapy of ipilimumab and nivolumab in patients with advanced neuroendocrine tumors: a subgroup analysis of the CA209-538 clinical trial for rare cancers. *Clin Cancer Res* 2020; 26: 4454–4459
- [58] Capdevila J, Teule A, López C et al. 11570 A multi-cohort phase II study of durvalumab plus tremelimumab for the treatment of patients (pts) with advanced neuroendocrine neoplasms (NENs) of gastroenteropancreatic or lung origin: The DUNE trial (GETNE 1601). *Ann Oncol* 2020; 31: S770–S771
- [59] Chan JA, Raj NP, Aggarwal RR et al. Phase II study of pembrolizumab-based therapy in previously treated extrapulmonary poorly differentiated neuroendocrine carcinomas: Results of Part B (pembrolizumab + chemotherapy). *J Clin Oncol* 2021; 39: 4148–4148
- [60] Shirasawa M, Yoshida T, Takayanagi D et al. Activity and immune correlates of programmed death-1 blockade therapy in patients with advanced large cell neuroendocrine carcinoma. *Clin Lung Cancer* 2021; 22: 282–291. e286
- [61] Sherman S, Rotem O, Shochat T et al. Efficacy of immune check-point inhibitors (ICPi) in large cell neuroendocrine tumors of lung (LCNEC). *Lung Cancer* 2020; 143: 40–46
- [62] Gile JJ, Liu AJ, McGarrah PW et al. Efficacy of checkpoint inhibitors in neuroendocrine neoplasms: Mayo clinic experience. *Pancreas* 2021; 50: 500–505
- [63] Al-Toubah T, Pelle E, Strosberg J. What is the role of checkpoint inhibitors in neuroendocrine neoplasms? *Oncotarget* 2020; 11: 3751–3752
- [64] McGrail DJ, Pilié PG, Rashid NU et al. High tumor mutation burden fails to predict immune checkpoint blockade response across all cancer types. *Ann Oncol* 2021; 32: 661–672
- [65] Özdirik B, Jann H, Bischoff P et al. PD-L1 - inhibitors in neuroendocrine neoplasia: Results from a real-life study. *Medicine (Baltimore)* 2021; 100: e23835
- [66] Halperin DM, Liu S, Dasari A et al. Assessment of clinical response following atezolizumab and bevacizumab treatment in patients with neuroendocrine tumors: a nonrandomized clinical trial. *JAMA Oncol* 2022; 8: 904–909
- [67] Bongiovanni A, Maiorano BA, Azzali I et al. Activity and safety of immune checkpoint inhibitors in neuroendocrine neoplasms: a systematic review and meta-analysis. *Pharmaceuticals (Basel)* 2021; 14: 476
- [68] Park EJ, Park HJ, Kim KW et al. Efficacy of immune checkpoint inhibitors against advanced or metastatic neuroendocrine neoplasms: a systematic review and meta-analysis. *Cancers (Basel)* 2022; 14: 794
- [69] Samstein RM, Lee CH, Shoushtari AN et al. Tumor mutational load predicts survival after immunotherapy across multiple cancer types. *Nat Genet* 2019; 51: 202–206
- [70] Wells SA Jr., Robinson BG, Gagel RF et al. Vandetanib in patients with locally advanced or metastatic medullary thyroid cancer: a randomized, double-blind phase III trial. *J Clin Oncol* 2012; 30: 134–141
- [71] Elisei R, Schlumberger MJ, Müller SP et al. Cabozantinib in progressive medullary thyroid cancer. *J Clin Oncol* 2013; 31: 3639–3646
- [72] Wirth LJ, Sherman E, Robinson B et al. Efficacy of selpercatinib in RET-altered thyroid cancers. *N Engl J Med* 2020; 383: 825–835
- [73] Subbiah V, Hu MI, Wirth LJ et al. Pralsetinib for patients with advanced or metastatic RET-altered thyroid cancer (ARROW): a multi-cohort, open-label, registrational, phase 1/2 study. *Lancet Diabetes Endocrinol* 2021; 9: 491–501



- [74] Satapathy S, Mittal BR, Sood A et al. Efficacy and safety of concomitant <sup>177</sup>Lu-DOTATATE and low-dose capecitabine in advanced medullary thyroid carcinoma: a single-centre experience. *Nucl Med Commun* 2020; 41: 629–635
- [75] Parghane RV, Naik C, Talole S et al. Clinical utility of (177) Lu-DOTATATE PRRT in somatostatin receptor-positive metastatic medullary carcinoma of thyroid patients with assessment of efficacy, survival analysis, prognostic variables, and toxicity. *Head Neck* 2020; 42: 401–416
- [76] Iten F, Müller B, Schindler C et al. Response to [90yttrium-DOTA]-TOC treatment is associated with long-term survival benefit in metastasized medullary thyroid cancer: a phase II clinical trial. *Clin Cancer Res* 2007; 13: 6696–6702
- [77] Hayes AR, Crawford A, Al Riyami K et al. Metastatic medullary thyroid cancer: the Role of <sup>68</sup>Gallium-DOTA-somatostatin analogue PET/CT and peptide receptor radionuclide therapy. *J Clin Endocrinol Metab* 2021; 106: e4903–e4916
- [78] Bodei L, Handkiewicz-Junak D, Grana C et al. Receptor radionuclide therapy with <sup>90</sup>Y-DOTATOC in patients with medullary thyroid carcinomas. *Cancer Biother Radiopharm* 2004; 19: 65–71
- [79] Beukhof CM, Brabander T, van Nederveen FH et al. Peptide receptor radionuclide therapy in patients with medullary thyroid carcinoma: predictors and pitfalls. *BMC Cancer* 2019; 19: 325
- [80] Bi Y, Ren X, Bai X et al. PD-1/PD-L1 expressions in medullary thyroid carcinoma: clinicopathologic and prognostic analysis of Chinese population. *Eur J Surg Oncol* 2019; 45: 353–358
- [81] Hińcza-Nowak K, Kowalik A, Walczyk A et al. Immune profiling of medullary thyroid cancer – an opportunity for immunotherapy. *Genes (Basel)* 2021; 12: 1534
- [82] Shi X, Li CW, Tan LC et al. Immune co-inhibitory receptors PD-1, CTLA-4, TIM-3, LAG-3, and TIGIT in medullary thyroid cancers: a large cohort study. *J Clin Endocrinol Metab* 2021; 106: 120–132
- [83] Di Molfetta S, Dotto A, Fanciulli G et al. Immune checkpoint inhibitors: new weapons against medullary thyroid cancer? *Front Endocrinol (Lausanne)* 2021; 12: 667784
- [84] Antonelli A, Ferrari SM, Fallahi P. Current and future immunotherapies for thyroid cancer. *Expert Rev Anticancer Ther* 2018; 18: 149–159
- [85] Del Rivero J, Donahue RN, Martí JL et al. A case report of sequential use of a yeast-CEA therapeutic cancer vaccine and anti-PD-L1 inhibitor in metastatic medullary thyroid cancer. *Front Endocrinol (Lausanne)* 2020; 11: 490
- [86] Nölting S, Bechmann N, Taieb D et al. Personalized management of pheochromocytoma and paraganglioma. *Endocr Rev* 2022; 43: 199–239
- [87] Jimenez C, Subbiah V, Stephen B et al. Phase II clinical trial of pembrolizumab in patients with progressive metastatic pheochromocytomas and paragangliomas. *Cancers (Basel)* 2020; 12: 2307
- [88] Rodriguez RR, Rizwan S, Alhamad K et al. The use of immunotherapy treatment in malignant pheochromocytomas/paragangliomas: a case report. *J Med Case Rep* 2021; 15: 172
- [89] Wang Z, Guo X, Gao L et al. The immune profile of pituitary adenomas and a novel immune classification for predicting immunotherapy responsiveness. *J Clin Endocrinol Metab* 2020; 105: e3207–e3223
- [90] Nie D, Fang Q, Li B et al. Research advances on the immune research and prospect of immunotherapy in pituitary adenomas. *World J Surg Oncol* 2021; 19: 162
- [91] Dai C, Liang S, Sun B et al. The progress of immunotherapy in refractory pituitary adenomas and pituitary carcinomas. *Front Endocrinol (Lausanne)* 2020; 11: 608422
- [92] Almutairi RD, Muskens IS, Cote DJ et al. Gross total resection of pituitary adenomas after endoscopic vs. microscopic transsphenoidal surgery: a meta-analysis. *Acta Neurochir (Wien)* 2018; 160: 1005–1021
- [93] Ilie MD, Jouanneau E, Raverot G. Aggressive pituitary adenomas and carcinomas. *Endocrinol Metab Clin North Am* 2020; 49: 505–515
- [94] Wang PF, Wang TJ, Yang YK et al. The expression profile of PD-L1 and CD8(+) lymphocyte in pituitary adenomas indicating for immunotherapy. *J Neurooncol* 2018; 139: 89–95
- [95] Sato M, Tamura R, Tamura H et al. Analysis of tumor angiogenesis and immune microenvironment in non-functional pituitary endocrine tumors. *J Clin Med* 2019; 8: 695
- [96] Kemeny HR, Elsamadicy AA, Farber SH et al. Targeting PD-L1 initiates effective antitumor immunity in a murine model of Cushing disease. *Clin Cancer Res* 2020; 26: 1141–1151
- [97] Sol B, de Filette JMK, Awada G et al. Immune checkpoint inhibitor therapy for ACTH-secreting pituitary carcinoma: a new emerging treatment? *Eur J Endocrinol* 2021; 184: K1–K5
- [98] Hazrati SM, Aghazadeh J, Mohtarami F et al. Immunotherapy of prolactinoma with a T helper 1 activator adjuvant and autoantigens: a case report. *Neuroimmunomodulation* 2006; 13: 205–208
- [99] Angeles CV, Sabel MS. Immunotherapy for Merkel cell carcinoma. *J Surg Oncol* 2021; 123: 775–781
- [100] Topalian SL, Bhatia S, Amin A et al. Neoadjuvant nivolumab for patients with resectable Merkel cell carcinoma in the CheckMate 358 trial. *J Clin Oncol* 2020; 38: 2476–2487
- [101] Nghiem P, Bhatia S, Lipson EJ et al. Durable tumor regression and overall survival in patients with advanced Merkel cell carcinoma receiving pembrolizumab as first-line therapy. *J Clin Oncol* 2019; 37: 693–702
- [102] Kaufman HL, Russell J, Hamid O et al. Avelumab in patients with chemotherapy-refractory metastatic Merkel cell carcinoma: a multicentre, single-group, open-label, phase 2 trial. *Lancet Oncol* 2016; 17: 1374–1385
- [103] D'Angelo SP, Russell J, Lebbé C et al. Efficacy and safety of first-line avelumab treatment in patients with stage IV metastatic Merkel cell carcinoma: a preplanned interim analysis of a clinical trial. *JAMA Oncol* 2018; 4: e180077
- [104] Halfdanarson TR, Strosberg JR, Tang L et al. The north American neuroendocrine tumor society consensus guidelines for surveillance and medical management of pancreatic neuroendocrine tumors. *Pancreas* 2020; 49: 863–881
- [105] Kubli SP, Berger T, Araujo DV et al. Beyond immune checkpoint blockade: emerging immunological strategies. *Nat Rev Drug Discov* 2021; 20: 899–919
- [106] McGranahan N, Swanton C. Clonal heterogeneity and tumor evolution: past, present, and the future. *Cell* 2017; 168: 613–628
- [107] da Silva A, Bowden M, Zhang S et al. Characterization of the neuroendocrine tumor immune microenvironment. *Pancreas* 2018; 47: 1123–1129
- [108] Katz SC, Donkor C, Glasgow K et al. T cell infiltrate and outcome following resection of intermediate-grade primary neuroendocrine tumours and liver metastases. *HPB (Oxford)* 2010; 12: 674–683
- [109] Liu YT, Sun ZJ. Turning cold tumors into hot tumors by improving T-cell infiltration. *Theranostics* 2021; 11: 5365–5386
- [110] Maharjan CK, Ear PH, Tran CG et al. Pancreatic neuroendocrine tumors: molecular mechanisms and therapeutic targets. *Cancers (Basel)* 2021; 13: 5117
- [111] Young K, Lawlor RT, Ragulan C et al. Immune landscape, evolution, hypoxia-mediated viral mimicry pathways and therapeutic potential in molecular subtypes of pancreatic neuroendocrine tumours. *Gut* 2021; 70: 1904–1913

- [112] Topper MJ, Vaz M, Marrone KA et al. The emerging role of epigenetic therapeutics in immuno-oncology. *Nat Rev Clin Oncol* 2020; 17: 75–90
- [113] Sheybani ND, Price RJ. Perspectives on recent progress in focused ultrasound immunotherapy. *Theranostics* 2019; 9: 7749–7758
- [114] Mariathasan S, Turley SJ, Nickles D et al. TGF $\beta$  attenuates tumour response to PD-L1 blockade by contributing to exclusion of T cells. *Nature* 2018; 554: 544–548
- [115] Ribas A, Dummer R, Puzanov I et al. Oncolytic virotherapy promotes intratumoral T cell infiltration and improves anti-PD-1 immunotherapy. *Cell* 2017; 170: 1109–1119. e1110
- [116] Ngwa W, Irabor OC, Schoenfeld JD et al. Using immunotherapy to boost the abscopal effect. *Nat Rev Cancer* 2018; 18: 313–322
- [117] Peng W, Chen JQ, Liu C et al. Loss of PTEN promotes resistance to T cell-mediated immunotherapy. *Cancer Discov* 2016; 6: 202–216
- [118] June CH, O'Connor RS, Kawalekar OU et al. CAR T cell immunotherapy for human cancer. *Science* 2018; 359: 1361–1365
- [119] Finn RS, Qin S, Ikeda M et al. Atezolizumab plus bevacizumab in unresectable hepatocellular carcinoma. *N Engl J Med* 2020; 382: 1894–1905
- [120] Feig C, Jones JO, Kraman M et al. Targeting CXCL12 from FAP-expressing carcinoma-associated fibroblasts synergizes with anti-PD-L1 immunotherapy in pancreatic cancer. *Proc Natl Acad Sci U S A* 2013; 110: 20212–20217
- [121] Abril-Rodriguez G, Torrejon DY, Liu W et al. PAK4 inhibition improves PD-1 blockade immunotherapy. *Nat Cancer* 2020; 1: 46–58
- [122] Rindi G, Wiedenmann B. Neuroendocrine neoplasia of the gastrointestinal tract revisited: towards precision medicine. *Nat Rev Endocrinol* 2020; 16: 590–607
- [123] Gubbi S, Koch CA, Klubo-Gwiezdzinska J. Peptide receptor radionuclide therapy in thyroid cancer. *Front Endocrinol (Lausanne)* 2022; 13: 896287
- [124] Boy C, Heusner TA, Poeppel TD et al. <sup>68</sup>Ga-DOTATOC PET/CT and somatostatin receptor (sst1-sst5) expression in normal human tissue: correlation of sst2 mRNA and SUVmax. *Eur J Nucl Med Mol Imaging* 2011; 38: 1224–1236
- [125] Dalm VA, van Hagen PM, van Koetsveld PM et al. Expression of somatostatin, cortistatin, and somatostatin receptors in human monocytes, macrophages, and dendritic cells. *Am J Physiol Endocrinol Metab* 2003; 285: E344–E353
- [126] Ameri P, Ferone D. Diffuse endocrine system, neuroendocrine tumors and immunity: what's new? *Neuroendocrinology* 2012; 95: 267–276
- [127] Nishikawa H, Sakaguchi S. Regulatory T cells in tumor immunity. *Int J Cancer* 2010; 127: 759–767
- [128] Cascini GL, Cuccurullo V, Tamburrini O et al. Peptide imaging with somatostatin analogues: more than cancer probes. *Curr Radiopharm* 2013; 6: 36–40
- [129] Pintér E, Helyes Z, Szolcsányi J. Inhibitory effect of somatostatin on inflammation and nociception. *Pharmacol Ther* 2006; 112: 440–456
- [130] Veenstra MJ, van Koetsveld PM, Dogan F et al. Epidrug-induced upregulation of functional somatostatin type 2 receptors in human pancreatic neuroendocrine tumor cells. *Oncotarget* 2018; 9: 14791–14802
- [131] Jin XF, Auernhammer CJ, Ilhan H et al. Combination of 5-fluorouracil with epigenetic modifiers induces radiosensitization, somatostatin receptor 2 expression, and radioligand binding in neuroendocrine tumor cells in vitro. *J Nucl Med* 2019; 60: 1240–1246
- [132] Pivonello R, Munster PN, Terzolo M et al. Glucocorticoid receptor antagonism upregulates somatostatin receptor subtype 2 expression in ACTH-producing neuroendocrine tumors: new insight based on the selective glucocorticoid receptor modulator relacorilant. *Front Endocrinol (Lausanne)* 2021; 12: 793262
- [133] Liu X, Bao X, Hu M et al. Inhibition of PCSK9 potentiates immune checkpoint therapy for cancer. *Nature* 2020; 588: 693–698
- [134] Mager LF, Burkhard R, Pett N et al. Microbiome-derived inosine modulates response to checkpoint inhibitor immunotherapy. *Science* 2020; 369: 1481–1489
- [135] Kang NW, Tan KT, Li CF et al. Complete and durable response to nivolumab in recurrent poorly differentiated pancreatic neuroendocrine carcinoma with high tumor mutational burden. *Curr Oncol* 2021; 28: 4587–4596
- [136] Rousseau B, Foote MB, Maron SB et al. The spectrum of benefit from checkpoint blockade in hypermutated tumors. *N Engl J Med* 2021; 384: 1168–1170