

The Bostrichidae of the Maltese Islands (Coleoptera)

Gianluca Nardi¹, David Mifsud²

1 Centro Nazionale per lo Studio e la Conservazione della Biodiversità Forestale “Bosco Fontana”, Sede di Bosco Fontana – Corpo Forestale dello Stato, Strada Mantova 29, I-46045 Marmirolo (MN), Italy **2** Institute of Earth Systems, Division of Rural Sciences and Food Systems, University of Malta, Msida MSD 2080, Malta

Corresponding author: Gianluca Nardi (l_nardi@hotmail.com)

Academic editor: C. Majka | Received 17 June 2014 | Accepted 6 January 2015 | Published 4 February 2015

<http://zoobank.org/4AB90367-FE56-41C0-8825-16E953E46CEC>

Citation: Nardi G, Mifsud D (2015) The Bostrichidae of the Maltese Islands (Coleoptera). ZooKeys 481: 69–108. doi: 10.3897/zookeys.481.8294

Abstract

The Bostrichidae of the Maltese Islands are reviewed. Ten species are recorded with certainty from this Archipelago, of which 6 namely, *Trogoxylon impressum* (Comolli, 1837), *Amphicerus bimaculatus* (A.G. Olivier, 1790), *Heterobostrychus aequalis* (Waterhouse, 1884), *Sinoxylon unidentatum* (Fabricius, 1801), *Xyloperthella picea* (A.G. Olivier, 1790) and *Apate monachus* Fabricius, 1775 are recorded for the first time. Two of the mentioned species (*H. aequalis* and *S. unidentatum*) are alien and recorded only on the basis of single captures and the possible establishment of these species is discussed. Earlier records of *Scobicia pustulata* (Fabricius, 1801) from Malta are incorrect and should be attributed to *S. chevrieri* (A. Villa & J.B. Villa, 1835). A zoogeographical analysis and an updated checklist of the 12 species of Bostrichidae recorded from the Maltese Islands and neighbouring Sicilian islands (Pantelleria, Linosa and Lampedusa) are also provided.

Rhizopertha dominica (Fabricius, 1792) form *granulipennis* Lesne in Beeson & Bhatia, 1937 from Uttarakhand (northern India) was overlooked by almost all subsequent authors. Its history is summarized and the following new synonymy is established: *Rhizopertha dominica* (Fabricius, 1792) form *granulipennis* Lesne in Beeson & Bhatia, 1937 = *Rhyzopertha dominica* (Fabricius, 1792), **syn. n.**

Finally, records of *Amphicerus bimaculatus* from Azerbaijan, of *Bostrichus capucinus* (Linnaeus, 1758) from Jordan and Syria, of *Scobicia chevrieri* from Jordan and Italy, of *Xyloperthella picea* from Italy, and of *Apate monachus* from Corsica (France) and Italy, are also provided.

Keywords

Bostrichidae, new records, new synonym, alien species, Malta, Italy

Introduction

The larvae of most species of this family are wood borers, and as other saproxylic organisms they play an important role in the decomposition processes. They are thus significant for nutrient cycling in natural and semi-natural ecosystems, particularly forests (cf. Stokland et al. 2012). Several species of this family are of great economic importance since they can cause extensive damage to dry and dead wood, to seasoned sapwood timber, to bamboo, and to wooden or bamboo artifacts through the boring behavior of both adults and larvae. Moreover, several other species are pests of living trees and vines, a few species are store product pests attacking commodities such as grain and tubers. These beetles are frequently transported between countries, especially in wood packing materials such as crating and dunnage, and are often intercepted at ports and cargo distribution centres (cf. Haack and Slansky 1987, Geis 2002, Ivie 2002, Haack 2006, Liu et al. 2006, Bahillo de la Puebla et al. 2007, Liu et al. 2008, Lawrence 2010).

The earliest record of the family comes from the mid Cretaceous (Peris et al. 2014). This family is now mainly represented in subtropical and tropical regions, and currently about 600 described species accommodated in 90 genera are known to occur worldwide (cf. Lawrence and Slipinski 2013, Zahradník and Háva 2014).

The knowledge on Maltese Bostrichidae was very limited, with only five previously recorded species. New data on Maltese Bostrichidae emerged from recent studies carried out on collections made from the Maltese Islands, the results of which are included in the present work.

Material and methods

Study area

The study area comprises all of the Maltese islands (c. 316 km²). Chetcuti et al. (1992), Schembri (1993, 1997) and Giusti et al. (1995) can be consulted for general environmental information on this Archipelago situated in the centre of the Mediterranean basin.

Nomenclature and classification

The suprageneric classification of the family and the nomenclature (family-group names and genus-group names) adopted in the present work follow Zahradník and Háva (2014), but the subfamilies follow Liu and Schönitzer (2011). The species are listed alphabetically as in Borowski and Węgrzynowicz (2007) and Ivie (2010).

Faunistic list

For each species, the following information is provided: nomenclatural combinations (listed chronologically-alphabetically) of the Maltese records found in the literature, literature records on the Maltese Islands, material examined, chorotype, data on ecology, and notes.

When possible, the following data was also provided for each record: island, municipality, locality of collection, altitude, date of collection, collector/s, collecting method, number of specimens, and, in parenthesis, literature reference or abbreviation of the depository. The records are listed in alphabetic order with respect to localities of collection. A semicolon separates different records; if these are from the same site, the name of the site is listed only at the beginning with the older record. When deemed useful for the discussion of some species, material examined from other countries (“Other material examined”) is also provided.

Possible interpolations are given in square brackets; collecting data of Italian specimens that is originally written in Italian, is hereunder provided in English. Information on very old specimens is written in double quotation marks.

Regarding literature records, Luigioni (1929) listed the two previously recorded bostrichids from Malta (cf. Cameron and Caruana Gatto 1907), but both Porta (1929, 1934, 1949, 1959) and Borowski (2007) overlooked these records. Records from “Malta” of Nardi (2004a, 2004b) are based only on previous literature. Finally, Schembri and Lanfranco (1996: 6) recorded an unidentified bostrichid collected in a consignment of tropical logs imported from Africa, but the material was presumably lost (D. Mifsud, unpublished data).

The identifications of the material examined were made by one of us (G. Nardi), and were based on Lesne (1899, 1901b, 1904, 1906, 1909, 1924), Español (1955, 1956a), Cymorek (1961), Liu et al. (2006), Bahillo de la Puebla et al. (2007), Sitticaya et al. (2009) and Beiriger (2010). References, including illustrations of the genitalia are listed when available. This was done since so far little attention was given to the study of the genitalia of these beetles.

Zoogeography

Chorotypes, which were also used in the zoogeographical analysis, were assigned according to Vigna Taglianti et al. (1993, 1999) and are based on the distributions provided by the authors cited for each species. Moreover, the chorotypes of many species show extensions or more confined distributions when compared to the standard ones; these differences were mentioned only when the change is significant. For further information on global distributions, references cited in the text may be consulted.

Acronyms

Specimen depositories

BMNH	The Natural History Museum, London, UK;
CCI	private collection P. Cornacchia, Porto Mantovano (Mantua), Italy;
CMM	private collection D. Mifsud, Malta;
CNBFVR	Centro Nazionale per lo Studio e la Conservazione della Biodiversità Forestale “Bosco Fontana” di Verona. Sede di Bosco Fontana. Marmirolo (Mantua), Italy;
CNI	private collection G. Nardi, Cisterna di Latina (Latina), Italy;
MCSV	Museo civico di Storia Naturale, Verona, Italy;
MCSVA	F. Angelini collection c/o Museo civico di Storia Naturale, Verona, Italy;
MCSVD	G. Dellabeffa collection c/o Museo civico di Storia Naturale, Verona, Italy;
MCZRD	E. De Maggi collection c/o Museo civico di Zoologia, Rome, Italy;
MCZRL	P. Luigioni collection c/o Museo civico di Zoologia, Rome, Italy;
MZUF	Museo di Storia Naturale dell’Università degli Studi di Firenze, Sezione di Zoologia “La Specola”, Florence, Italy;
MZUR	Museo di Zoologia, Università degli Studi di Roma “Sapienza”, Rome, Italy.

Main collectors

AF	A. Falzon;
DD	D. Dandria;
D.P.I.T.	Dana Project Invertebrate Team = [S. De Felici, T. Di Micco De Santo, M. Shubat, F. Al-Eman Al-Husseini & A. Abu Hantash];
DM	D. Mifsud;
GN	G. Nardi;
HB	H. Borg Barthet;
LC	L.F. Cassar;
LF	L. Fancello;
PS	P. Sammut.

Other abbreviations

ex = specimen/s;
Fraz. = Frazione = Hamlet;
prov. = province;
pt = pitfall trap.

Results

Subfamily Lyctinae Billberg, 1820, Tribe Lyctini Billberg, 1820

Lyctus brunneus (Stephens, 1830)

Lyctus brunneus Steph.: Cameron and Caruana Gatto 1907: 398.

Lyctus (Xylotrogus) brunneus Steph.: Luigioni 1929: 642.

Lyctus brunneus (Stephens, 1830): Nardi 2004a, Denux and Zagatti 2010: 366.

Literature records. Malta: Valletta (Cameron and Caruana Gatto 1907); “Mal.” [= Malta] (Luigioni 1929); Malta (Nardi 2004a, Denux and Zagatti 2010).

Material examined. Malta: Naxxar, 7.IX.1997, DD, [in human abitation], 2♂♂ (CMM); Rabat, Wied ta l-Isqof, 28.VI.2002, PS, [in an agricultural area], 1♀ (CNI).

Chorotype. Cosmopolitan (Borowski 2007, as *L. (X.) brunneus*). This species is widespread in the Palaearctic region (cf. Nardi 2004a, Borowski 2007), but the followings countries were overlooked in the species distribution summarized by Nardi (2004a) and/or Borowski (2007): Algeria (Peyerimhoff 1919, as *L. (X.) brunneus*, Lesne 1924), Austria (Horion 1961, Lucht 1987, Adlbauer 1998, Nardi 2004a, Denux and Zagatti 2010, Querner et al. 2011), Azores (cf. Borges 1990, Nardi 2004a, Oromí et al. 2010), Belarus, Bulgaria (Denux and Zagatti 2010), Canary Islands (cf. Machado and Oromí 2000, Nardi 2004a, Oromí et al. 2009), Corsica (cf. Becker 1969, Geis 2002), Croatia (Damoiseau 1966, Nardi 2004a), Cyprus (Baudi di Selve 1873, 1874, Georghiou 1977, Nardi 2004a), Egypt (Kaszab 1959, Attia and Kamel 1965, in both cases as *L. (X.) brunneus*, Hanna 1970, Alfieri 1976, Hamad and Aly 1985), Fujian (southeastern China) (Vrydagh 1960), Greece (Damoiseau 1966, Nardi 2004a, Denux and Zagatti 2010), Iran (Adeli 1972, Niloufari 1985), Israel (cf. Halperin and Geis 1999, Chikatunov et al. 2004b, 2006), Latvia (Denux and Zagatti 2010), Poland (cf. Burakowski et al. 1986, Nardi 2004a, Jabłoński et al. 2007, Krajewski and Mazurek 2009), Serbia and Montenegro (Glavendekic et al. 2005, Denux and Zagatti 2010) and Turkey (cf. Gerberg 1957, Akbulut et al. 2008).

Ecology. In tropical areas this species develops in the wood of a large number of unrelated plants, whereas in temperate regions (where the species is considered as an established alien) it develops mainly in hardwood timber (e.g. *Castanea sativa*, *Fraxinus excelsior*, *Junglans regia*, *Quercus* spp., *Ulmus* spp., etc.) primarily in synanthropical environments (workshops, plywood industries, private houses). In the West Palaearctic region, the species was also found in local trees (*Alnus*, *Eucalyptus*, *Ulmus*, etc.) which were in poor health conditions. The passive transport of this beetle has been documented with both wooden products (timbers, tables, furniture, ethnographic collection, briars, etc.) and manioc (cf. Lesne 1924, Lepesme 1944, Español 1956a, Gerberg 1957, Cymorek 1961, Burlini 1972, Aitken 1975, Cavalloro and Ratti 1978,

Gambetta and Orlandi 1982a, Gambetta 1983, Wang et al. 1996, Halperin and Geis 1999, as *L. (X.) brunneus*, Geis 2002, Peters et al. 2002, Chikatunov et al. 2004, Bahillo de la Puebla et al. 2007, as *L. (X.) brunneus*, Mattson et al. 2007, Krajewski and Mazurek 2009, Denux and Zagatti 2010, Geis 2012). According to Gambetta and Orlandi (1982b: 55), *L. brunneus* and *Trogoxylon impressum* (Comolli, 1837) are the two species of Lyctinae which are the most widespread in woods in Italian storage situations.

Notes. A species native to Asia which has been established throughout Europe for more than 150 years (Denux and Zagatti 2010). The recently collected material confirm its presence in Malta. In the nearby countries, the species is recorded from Tunisia (Lesne 1924, Normand 1936, Borowski 2007) but not from Sicily (cf. Audisio et al. 1995, Sparacio 1997, Chiappini et al. 2001, Nardi 2004a, Denux and Zagatti 2010). Accurate information and illustrations on the morphology of all stages of this species were provided by Iwata and Nishimoto (1981, 1982) and Kucerová and Stesjkal (2008). Moreover the genitalia of both sexes was illustrated by different authors (e.g. Altson 1924: pls. 31–34, Gerberg 1957: pl. II, figs 15–16, Cymorek 1961: 81, fig. 3c, Iwata and Nishimoto 1982: 19, figs 42–44).

Subfamily Lyctinae Billberg, 1820, Tribe Trogoxylini Lesne, 1921

Trogoxylon impressum (Comolli, 1837)

Material examined. Malta: Busket, 15.VI.2003, DM, 1 ex (CMM); Mistra Valley, 21.IV.1994, DM, under leves of *Ficus carica*, 1 ex (CMM); Qormi, Hal-Farrug, 5.V.2003, DM, [on] *Ceratonia siliqua*, 1 ex (CNI); 5.V.2003, DM, 2 ex (CMM; CNI); Rabat, Ta Koronja, 14.VI.2002, PS, 1 ex (CMM); Wied Badu, 3.VII.2002, DM, 1 ex (CMM).

Chorotype. Turanic-European-Mediterranean. This species is widespread from the Canary Islands to Turkmenistan; to the north it reaches Austria, Czech Republic, Germany, Hungary, Slovakia and Switzerland, moreover it has been introduced in some northern European states (Denmark, Finland, Norway and Sweden), into Argentina, Chile and USA (cf. Borowski 2007, Borowski and Węgrzynowicz 2007, Barriga and Cepeda 2009), and it is considered as an established alien in the latter two mentioned territories (Gerberg 1957). The species is possibly also establishment in China (cf. Peters et al. 2002), South Africa and Australia (Geis 2002), but for these territories this was not confirmed (Borowski and Węgrzynowicz 2007). This species was also recorded from Iraq (Knopf 1972) and was introduced to Benelux (Lucht 1987, Drumont and Grootaert 2011) and both countries must be added to its distribution which was summarized by Borowski (2007). Hopkins (1911: 137, as *Lyctus impressus*) in an Appendix to Kraus (1911), recorded this species also for “Cordova, Mex. [= Mexico]”; this record was overlooked since Lesne (1938) probably because this Appendix, which also included data on additional specimens, was added after the submission

of the paper of Kraus (1911) (cf. Hopkins 1911: 130, footnote) and Mexico was not included in the mentioned work.

Ecology. *Trogoxylon impressum* is a polyphagous species; in the Mediterranean area it has been reared from many local and exotic trees and shrubs (e.g. *Ceratonia siliqua*, *Cercis siliquastrum*, *Eucalyptus* spp., *Ficus carica*, *Pistacia lentiscus*, *P. vera*, *Punica granatum*, *Quercus* spp., *Tamarix* sp., *Vitis vinifera*), and also from timber (cf. Nardi and Ratti 1995, Halperin and Geis 1999, Geis 2002, Liberto and Audisio 2005, Bahillo et al. 2007, Baena and Zuzarte 2013).

Notes. This species is a new record for the Maltese Islands; it is recorded also from the nearby Island of Pantelleria (Nardi and Ratti 1995) and from the Balearic Islands (cf. Schaufuss 1882, as *L. impressus capitalis* n. var., Bahillo de la Puebla and López-Colón 2001, Nardi 2004a).

Its male genitalia was figured by Gerberg (1957: pl. XI, figs 15–16) and Iablokoff-Khnzorian (1980: 284, fig. 12.9).

Subfamily Dinoderinae C.G. Thomson, 1863

Rhyzopertha dominica (Fabricius, 1792)

Rhyzopertha dominica: Hyde and Daubney 1960: 120.

Rhyzopertha dominica (Fabricius, 1792): Denux and Zagatti 2010: 348.

Literature records. Malta: Floriana, St. Publius site, 1958, on grain, 1 ex (Hyde and Daubney 1960); “MT” [= Malta] (Denux and Zagatti 2010).

Material examined. Malta: St. Thomas Bay, 27.VI.1990, DM, 1 ex (CMM); Zejtun, 20.VII.1989, DM, attracted to light, 1 ex (CMM); 15.IX.1989, DM, 1 ex (CNI); 21.IX.1989, DM, 1 ex (CMM); 28.IX.1989, DM, 1 ex (CMM); 10.IX.2001, DM, 1 ex (CMM); 24.XI.2001, DM, 2 ex (CMM; CNI); 3.VII.2002 DM, 1 ex (CMM).

Chorotype. Cosmopolitan (Potter 1935, as *Rhizopertha dominica*, Borowski 2007). This species is widespread in the Palaearctic region (cf. Nardi 2004b, Borowski 2007). The followings territories were not included in its distribution provided by Borowski (2007): Bhutan (Taylor and Halliday 1986), Bulgaria (Zidan and Obretenchev 2009), China (Anhui, Chongqing, Guangdong, Guizhou, Hebei, Henan, Hubei, Hunan, Jiangxi, Shandong, Sichuan) (Lesne 1904, as *Rhi. dominica*, Song et al. 2012), Iran (cf. Abivari 2001, as *Rhi. dominica*, Ziae et al. 2006, Ashouri and Shayesteh 2009), Israel (Halperin and Damoiseau 1980, as *Rhi. dominica*, Chikatunov et al. 2004, as *Rhyzoperta* [sic!] *dominica*, Chikatunov et al. 2006, as *Rhi. dominica*), Jordan (Sharaf et al. 1983), Libya (cf. Zanon 1922, Gridelli 1930, Zavattari 1934, in all cases as *Rhi. dominica*, Champ 1986), Lithuania (Ivinskis et al. 2009, as *Rhi. dominica*), Morocco (Martínez de la Escalera 1914, Rungs 1946, Español 1956b, Kocher 1956, in all cases as *Rhi. dominica*, Bartali et al. 1990, López-Colon 2000, as *Rhi. dominica*, Benhalima et al. 2004), Nepal (Taylor and Halliday 1986), Pakistan (cf. Quddus and Qayyum

1982, Ishaque et al. 1982, in both cases as *Rhi. dominica*, Taylor and Halliday 1986, Sardar Alam et al. 1999, Anwar et al. 2005, in both cases as *Rhi. dominica*, Wakil et al. 2006, Ahmed et al. 2008, as *Rhi. dominica*, Wakil et al. 2013), Portugal (Seabra 1943, as *Rhi. dominica*, Nardi 2004b, Bahillo de la Puebla et al. 2007, Baena and Zuzarte 2013), Romania (Ghizdavu and Deac 1995, Nardi 2004b), Saudi Arabia (Damoiseau 1979, Amin et al. 1986, in both cases as *Rhi. dominica*, Mostafa et al. 1981, Rostom 1993, Ahmed 1996), Switzerland (Hoppe 1981, Buchi 1993, Kenis 2005, as *Rhyzopertha* [sic!] *dominica*, Denux and Zagatti 2010), Turkey (Gerini 1971, Aydin and Soran 1987, Yucel 1988, Emekçi and Ferizli 2000), Ukraine (Podobivskii 1991) and Uzbekistan (Asanov 1980).

Ecology. Originally, this species was only associated with wood but is now considered as a primary pest of all kinds of stored grains. Both larvae and adults are able to attack whole grain, causing considerable damage. The species is thought to have originated from the Indian subcontinent, and was introduced worldwide by commerce. It is an economically important pest since it can cause significant economic losses in terms of grain mass and nutrient depletion, and pose a public health risk from contamination by allergens, such as uric acid (cf. Potter 1935, Beeson and Bhatia 1937, Lepesme 1944, in all cases as *Rhi. dominica*, Fisher 1950, Aitken 1975, Maes 1995, Pollini 1998, in both cases as *Rhi. dominica*, Geis 2002, Gelosi and Süss 2001, Nguyen 2006, Borowski 2007, Liu et al. 2008, Denux and Zagatti 2010, Kenis and Branco 2010, Arthur et al. 2012, Edde 2012). This species can also damage books (Hoffman 1933, as *Rhi. dominica*).

Notes. The record from Malta by Hyde and Daubney (1960) was later overlooked by all authors (cf. Saliba 1963, Nardi 2004b, Borowski 2007). Thus this beetle must have been introduced in Malta at least since 1958 where it is now a well established species.

In Italy, *Rhyzopertha dominica* was first collected in Sicily during the nineteenth century (cf. Bertolini 1874, as *Rhi. pusilla* Fabr. [= (Fabricius, 1798)]; Ragusa 1896, as *Rhi. pusilla*), and since the 1950's it was widespread in all regions (cf. Norato 1957, Dal Monte 1958, as *Rhi. dominica*, Genduso 1963). According to Denux and Zagatti (2010: 348) the first European record of this species was from Czech Republic and is dated 1900, but the above Sicilian records, as well as those from other countries (cf. Reitter 1883, as *Rhi. pusilla*, Lesne 1901b), are older. Moreover, this species has been present in Europe at least since 3500–2551 BP, as testified by its archelogical presence in Spain and Greece (cf. King 2009). So the species was introduced and established before 1492 A.D. at least in Spain and Greece and must be considered as parautocthonous (cf. Zapparoli 2008: 98). Illustrations of its male and female genitalia were provided by Potter (1935: 474–475, figs 21–25), Lesne (1945: 149, figs 9–13, as *Rhi. dominica*) and Surtees (1961: 149, fig. 11, as *Rhi. dominica*).

Rhizopertha dominica granulipennis Lesne in Beeson & Bhatia, 1937 from northern India (Uttarakhand, Chandi Randge) (Beeson and Bhatia 1937: 283, Lesne 1945: 146, as *Rhi. dominica granulipennis*) was overlooked in recent catalogues (cf. Borowski 2007, Borowski and Węgrzynowicz 2007, Ivie 2010). In its original description – “A large form with exceptionally strongly developed granulation of the elytral declivity was

bred from *Shorea robusta* [(Dipterocarpaceae)] and labelled *R. dominica granulipennis* by Lesne” (Beeson and Bhatia 1937: 283) – its rank is not unambiguously given, so, according to the Code (ICZN 1999, art. 45.6.4), it is an available subspecific name. Its authorship is here attributed to Lesne in Beeson and Bhatia (1937: 283), since Lesne (1945: 146) has reaffirmed the authorship of this name: “J’ai donné le nom de *R. dominica granulipennis* (1 [= footnote: Cf. Beeson et Bhatia, mémoire cité, p. 283.]) à cette forme remarquable” [= I named *R. dominica granulipennis* (1 [= footnote: Cf. Beeson et Bhatia, memory cited, p. 283.]) this remarkable form]. Unfortunately, it was not possible to study the type material of this taxon, that, according to Lesne (1945: 146), is housed in the Muséum National d’Histoire Naturelle of Paris (France) and in the Forest Research Institute of Derha Dun (India). However, on the basis of numerous specimens of this species from different territories (Nardi, unpublished data) which show a significant range of variability in the granulation of the elytral declivity, the following new synonymy, is here established: *Rhizopertha dominica* form *granulipennis* Lesne, in Beeson and Bhatia 1937: 283 = *Rhyzopertha dominica* (Fabricius, 1792: 359), **syn. n.**

Subfamily Bostrichinae Latreille, 1802, Tribe Bostrichini Latreille, 1802

Amphicerus bimaculatus (A.G. Olivier, 1790)

Material examined. Malta: Grgenti, 9.XII.2002, DM, 3 ♂♂ 4 ♀♀ (CMM; CNI); Zabbar, 29.IX.1995, DM, 1 ♀ (CMM); Zejtun, 29.V.1991, DM, 1 ♂ (CMM).

Other material examined. [Azerbaijan:] Caucase, Elisabetpol [= Ganja], [no date], Babadjanides [leg.], 1 ♂ 1 ♀ (MCSVD).

Chorotype. Turanic-Mediterranean except for Libya and Egypt, with extension into Portugal, Hungary, Tajikistan and Kyrgyzstan (cf. Borowski 2007, as *A. (Caenophrada) bimaculata* [sic!]). The occurrence of this species in Azerbaijan (Lesne 1899, 1905, Clermont 1909, Khalilov 1972, in all cases as *Schistoceros bimaculatus*, Ciampolini et al. 1989) is here confirmed. In fact, the distributional record of this species by Borowski (2007: 321), as “E [= Europe]: AF [= Afghanistan]” is incorrect and should be “E: AB [=Azerbaijan]” (cf. Borowski and Węgrzynowicz 2007, as *A. (C.) bimaculatus*). A record from Iraq (Derwesh 1965, as *S. bimaculatus*) was later overlooked, while those from Germany (cf. Soro 1964, Zocchi 1971, in both cases as *S. bimaculatus*, Ciampolini et al. 1989, Pollini 1998) were never confirmed (cf. Lucht 1987, Köhler and Klausnitzer 1998, Geis 2002, Nardi 2004b, as *S. bimaculatus*, Borowski 2007, Borowski and Węgrzynowicz 2007). *Amphicerus bimaculatus* was intercepted at US ports, but it is not an established species in North America (Fisher 1950, as *A. (S.) bimaculatus*, Ivie 2002, Borowski and Węgrzynowicz 2007), while its establishment in Uruguay (cf. Soro 1964, Zocchi 1971, Pollini 1998), was not reported in recent publications (Borowski 2007, Borowski and Węgrzynowicz 2007, Barriga and Cepeda 2009).

Ecology. Larval development of this species takes place mainly in dead wood of *Vitis* spp. and *Tamarix* spp., but other host plants are also recorded including: *Acacia* sp., *Annona cherimola*, *Cerasus* sp., *Citrus* sp., *Delonix regia*, *Ficus carica*, *Lycium* sp., *Malus communis*, *Olea europaea*, *Prunus amygdalus*, *Punica granatum*, *Pyrus malus* and *Tamarindus indica* (cf. Lesne 1901b, as *S. bimaculatus*, Fisher 1950, Novak 1952, Caillol 1954, in both cases as *S. bimaculatus*, Español 1955, Kocher 1956, as *S. bimaculatus*, Soro 1964, Zocchi 1971, Halperin and Damoiseau 1980, Lundberg et al. 1987, in both cases as *S. bimaculatus*, Molesas 1988, Ciampolini et al. 1989, Ragusa and Russo 1989, fig. 8, as [sic!] *Apate monachus*, Pollini 1998, Aksit et al. 2005, Lliberto and Audisio 2005, in both cases as *S. bimaculatus*, Bahillo de la Puebla et al. 2007, as *A. (C.) bimaculata* [sic!], Tezcan 2008, as *S. bimaculatus*).

Notes. First record for Malta. *Amphicerus bimaculatus* is not recorded from the neighbouring Sicilian islands (Tab. 1), but is known from mainland Sicily (cf. Audisio et al. 1995, Sparacio 1997, Nardi 2004b, in all cases as *S. bimaculatus*) and Tunisia (Borowski 2007). It is recorded also from two circumsardinian islands (Piras and Pisano 1972, as *S. bimaculatus*). Its aedeagus was figured by Iablokoff-Khnzorian (1976: 232, fig. 5, as *S. bimaculatus*).

The above specimens collected during 2002 are almost entirely black, probably for a *post mortem* colouration.

The correct grammatical gender for the specific name of this taxon is *bimaculatus*, since it was described as *Bostrichus bimaculatus* from “Provence” (southern France) (Olivier 1790: 109) and the above mentioned usage of *bimaculata* (Bahillo de la Puebla et al. 2007, Borowski 2007) is a subsequent incorrect spelling of this taxon (cf. Borowski 2013: 3).

The nomenclatorial problems for *Schistoceros* Lesne, 1899 were discussed by Ivie (2010).

***Bostrichus capucinus* (Linnaeus, 1758)**

Bostrychus capucinus (L.): Aitken 1975: 8.

Bostrichus capucinus (Linnaeus, 1758): Nardi 2004b.

Literature records. Malta: “Malta: wooden ornament” (Aitken 1975); “Malta” (Nardi 2004b).

Other material examined. Jordan: Dana Reserve, El-Barrah, 1150 m, NE slope, 36R YU 517 926, 23.IV.–8.V.1995, D.P.I.T., Mediterranean environment, pt, 1 ex (MZUR); ditto, Wadi Araba Camp Site, 15–20.IV.1995, D.P.I.T., pt, 1 ex (MZUR). **Syria:** Palmyra, 15.VII.2003, G. Serra leg., 1 ex (MZUF).

Chorotype. Centralasiatic-Mediterranean, including parts of Northern Europe (cf. Nardi and Ratti 1995, as *Bostry. capucinus*, Borowski 2007), the Algerian Sahara (Lesne 1899, 1901b, both as *Bostry. capucinus*) and coastal Sudan (Cloudsley-Thompson 1962, as *Bostry. capucinus*). This species reaches Altai mountains (Lesne 1901b,

Borowski and Węgrzynowicz 2007), Asian Kazakhstan (Borowski 2007), Kyrgyzstan (Vrydagh 1956, Ovtchinnikov 1996, in both cases as *Bostry. capucinus*), Tajikistan (Lesne 1901b), Northwest China (Lesne 1904, Horion 1961, in both cases as *Bostry. capucinus*, Borowski 2007), China (without further details) (Borowski 2007, Yan et al. 2010, as *Bostry. capucinus* and *Bostry. capucinus* var. *rubriventris* [sic!] Zouf [= *rubriventris* Zoufal, 1894]) and eastern Siberia (Borowski 2007). It was intercepted numerous times at US ports, but is not yet established in North America (cf. Fisher 1950, Ivie 2002). The above record from Jordan is the first for this country (cf. Sharaf et al. 1983, Borowski 2007), even though the presence of this species in Jordan was expected because of its occurrence in neighbouring countries (e.g. Israel, Syria, etc.); in Israel the species was probably introduced with timber from Europe (Bytinsky-Salz 1966, Bytinski-Salz and Sternlicht 1967, Halperin and Damoiseau 1980, in all cases as *Bostry. capucinus*, Borowski 2007).

Ecology. This species develops chiefly in the wood of oaks (*Quercus ilex*, *Q. robur*, *Q. toza*, etc.), but is also recorded from many other broadleaves trees and scrubs such as *Pinus*, and timber (cf. Nardi and Ratti 1995, Sparacio 1997, Liberto and Audisio 2005, Bahillo de La Puebla et al. 2007, Baena and Zuzarte 2013). Occasionally it can produce house infestations (Saccà 1940, as *Bostry. capucinus* Geoffroy [sic!], Cavalloro and Ratti 1978, Lodos 1985, Hellrigl 2006, in all cases as *Bostry. capucinus*).

Notes. The above record of Aitken (1975) was based on material collected in UK from cargo originating from Malta. The presence of this species in Malta needs to be confirmed. Although *Quercus ilex* is present in Malta, its abundance is very scarce on the archipelago (cf. Haslam et al. 1977, Schembri 1993, 1997). *Bostrichus capucinus* is known from other similarly small Mediterranean islands, such as the nearby Pantelleria Island, three circumsardinian islands (Piras and Pisano 1972, as *Bostry. capucinus*, Nardi and Ratti 1995), two northern Adriatic islands (Müller 1923: 28, Schatzmayr and Müller 1925: 74, Luigioni 1929: 611, in all cases as *Bostry. capucinus*), and the Balearic Islands (cf. Español 1955, as *Bostry. capucinus*).

Heterobostrychus aequalis (Waterhouse, 1884)

Material examined. Malta: Rabat, 21.IX.2001, PS, 1 ♂ (CMM).

Chorotype. A cosmopolitan species of Indo-Malaysian origins. It is mainly distributed in tropical and sub-tropical regions and restricted to 40° north and south of the equator (cf. Borowski 2007, Borowski and Węgrzynowicz 2007, Azmi et al. 2011: 500, fig. 2). In the Mediterranean, it was intercepted in Israel, Italy and Spain (cf. Gambetta 1983, Geis 2002, Ratti 2002, 2004b, Bahillo de la Puebla et al. 2007, Ratti 2007, Azmi et al. 2011). Ireland (O'Mahony 1949, as *Heterobostrychus* [sic!] *aequalis*), France (cf. Brustel and Aberlenc 2014) and Oregon (Westcott et al. 2006) must be added to the countries in which this species was intercepted (cf. Azmi et al. 2011), even though the former record was based on collection of death specimens (larvae and

adults), which was later ignored (cf. Anderson et al. 1997, Geis 2002, Nardi 2004b, Borowski 2007, Denux and Zagatti 2010, Alexander and Anderson 2012).

Ecology. Polyphagous species attacking some 36 unrelated host-plant genera; this species breeds not only in logs, but also in planks, furniture, plywood and roots of manioc (cf. Fisher 1950, Kalshoven 1963a, 1963b, Horion 1972, Gambetta 1983, Wang et al. 1996, as *Heterobothrachus* [sic!] *aqualis*, Geis 2002, Maes 2005, Aguilera 2006, Bahillo de la Puebla et al. 2007, Sitticaya et al. 2009, Robinson 2013).

Notes. First record for Malta. This species become established in some countries where it was accidentally introduced. Temperatures of 17 °C and below are said to be unsuitable for the species to breed (cf. Ivie 2002, Azmi et al. 2011). Thus, considering the warm climate of the Maltese Islands, (Chetcuti et al. 1992), it is highly likely that the species is already an established one. The above specimen was collected with a light trap on a terrace, mainly surrounded by agricultural land (P. Sammut, pers. comm., 2002).

Subfamily Bostrichinae Latreille, 1802, Tribe Sinoxylini Marseul, 1857

Sinoxylon unidentatum (Fabricius, 1801)

Material examined. Malta: Malta: Marsa, Ghommieri, 24.I.2007, DM, taken from wood packaging material originating from India, 1 ex (CMM).

Chorotype. A cosmopolitan species native to tropical eastern Asia, and widespread in the intertropical regions of the world. In Europe, it was intercepted in France, Germany, Great Britain, Italy, Poland, Russia, Spain and Ukraine, but its establishment was never confirmed. Moreover, the species is not reported from North Africa and the Middle East (except from Yemen) (cf. Vrydag 1955, Poggi et al. 1994, Peck et al. 1998, Geis 2002, Ratti 2004, Maes 2005, Iwata and Nakano 2006, Liu et al. 2006, Peres Filho et al. 2006, Westcott et al. 2006, in all cases as *S. conigerum* Gerstaecker 1855, Bahillo de la Puebla et al. 2007, Borowski 2007, Borowski and Węgrzynowicz 2007, Barriga and Cepeda 2009, as *S. conigerum*, Savoldelli and Regalin 2009, Price et al. 2011, Brustel and Aberlenc 2014).

Ecology. In its place of origin, *S. unidentatum* develops in the wood of many unrelated plant families mainly: Anacardiaceae, Combretaceae, Dipterocarpaceae, Euphorbiaceae, Lamiaceae, Lauraceae, Leguminosae, Mimosaceae, Myrtaceae, Rubiaceae, Tiliaceae, Ulmaceae, etc. (cf. Poggi et al. 1994, Peres Filho et al. 2006, as *S. conigerum*, Bahillo de la Puebla et al. 2007, Savoldelli and Regalin 2009).

Notes. This alien species is a new record for the central Mediterranean area. Due to its polyphagy, it is frequently exported with wooden packing material (cf. Bahillo de la Puebla et al. 2007). The Maltese climate (Chetcuti et al. 1992, Schembri 1997) is unsuitable for its establishment (J. Borowski, pers. comm. 2012), but in an indoor site of northern Italy, this species was able to complete its development and to spread the infestation (Savoldelli and Regalin 2009).

Subfamily Bostrichinae Latreille, 1802, Tribe Xyloperthini Lesne, 1921

Scobicia chevrieri (A. Villa & J.B. Villa, 1835)

Xyloperta pustulata F.: Cameron and Caruana Gatto 1907: 398.

Scobicia pustulata Fabr.: Luigioni 1929: 641.

Scobicia pustulata (Fabricius, 1801): Nardi 2004b.

Scobicia chevrieri (A. Villa & J.B. Villa, 1835): Mifsud et al. 2012: 9.

Literature records. Malta: [Malta,] “Coll. Gatto” (Cameron and Caruana Gatto 1907); “Mal.” [= Malta] (Luigioni 1929); “Malta” (Nardi 2004b); Buskett, adults emerged between 5–25.X.2011 from dead twigs of *Ficus carica* collected on 8.VII.2011, AF & DM leg., 56 ex (Mifsud et al. 2012).

Material examined. Gozo: Marsalforn Valley, 6.VI.1990, DM, 1 ex (CMM). **Malta:** Bahrija, 5.VIII.1992, LC, 1 ex (CMM); Bingemma, 10.IX.2001, DM, attracted to light, 8 ex (CMM); Buskett, 24.VI.2003, DM, attracted to light, mixed woodland *Pinus/Cupressus*, 5 ex (CMM); Marsa, Ghommieri, 24.III.2002, DM, 1 ex (CMM); “Malta, 9/[= IX.]1901”, “Xylopertha”, “pustulata”, “M. Cameron Coll. / B.M. 1936-555”, “5777” [= *Xylopertha pustulata* F. Marsa Scirocco [= Marsaxlokk]/id. EAN [= ?; maybe identified by E. A. Newbery (cf. Cameron and Caruana Gatto 1907: 383)]], 10 ex (BMNH); Rabat, 14.VI.2002, PS, 1 ex (CMM); 3.VIII.2002, PS, 3 ex (CMM); Rabat, Dwejra, 21.VI.2002, PS, 1 ex (CMM); Rabat, Ta Koronja, 6.VI.2002, PS, 2 ex (CMM); Wied Badu, 3.VII.2002, DM, 5 ex (CMM); Wied tal-Isqof, 16.VII.2002, DM, 1 ex (CMM); 2.VIII.2002, DM, 2 ex (CMM); Zejtun, 10.XI.1989, DM, 1 ex (CMM); 29.IX.1990, DM, development of larvae took place in dead branches of vines, 2 ex (CMM); 27.V.2002, DM, 1 ex (CMM).

Other material examined. Italy: Marche region, Ancona prov., Gole di Frasassi, 20.VI.2001, A. B. Biscaccianti leg., ex [larvae from] *Corylus avellana*, 2 ex (CNI); ditto, Pesaro e Urbino prov., Foce Fiume Metauro, area goleale [= Mouth of Metauro River, floodplain area], 2.VI.1999, A. B. Biscaccianti leg., ex larvae from *Salix* sp., 4 ex (CNBFVR; CNI). Latiun region, Rome prov., Tenuta Presidenziale di Castelporziano, Ponte della Focetta, 10.IX.1997, A. B. Di Giulio leg., hygrophilous wood, light trap 15 W, 1 ex (CNI); ditto, ditto, ditto, Villa di Capocotta, 21.VI.2000, P. Maltzeff leg., mixed light trap 160 W, 2 ex (CNI); ditto, Latina prov., Cisterna di Latina, [33T 319824.15 E 4606546.61 N], 29.IX.1987, GN, night, in a garden, at light, 2 ex (CNI); ditto, ditto, ditto, Fraz. Cerciabella, [33T 319479.96 E 4605030.39 N], 17.VIII.1998, GN, in a garden, at light, 20–21 hours, 2 ex (CNI). **Jordan:** Dana Reserve, Acacia Area, 17.IV.1995, D.P.I.T., night catch, 1 ex (MZUR); ditto, El-Barrah, 1150 m, NE slope, 36R YU 517 926, 23.IV.–8.V.1995, D.P.I.T., Mediterranean environment, pt, 1 ex (MZUR); ditto, Irano Turanian Area 1, 18.IV.1995, D.P.I.T., 1 ex (MZUR).

Chorotype. Mediterranean (northward upto Austria, French Alps, Hungary and Switzerland), with extension westward upto Portugal, and eastward upto Azerbaijan,

Georgia, Iran and southern Russia; this species was intercepted in the USA and Canada, but so far it has not established itself (cf. Fisher 1950, as *S. chevrieri* (Villa) [sic!], Vrydagh 1952, as *S. Chevrieri* (Villa) [sic!], Ivie 2002, Borowski 2007, Borowski and Węgrzynowicz 2007, McCaffrey 2011, as *S. chevrieri* Villa & Villa, 1835 [sic!]). It is known also from Romania (Lesne 1904, Vrydagh 1956, in both cases as *S. Chevrieri* Villa [sic!], Nardi 2004b) and Sinai (cf. Alfieri 1976, as *S. chevrieri* Villa [sic!]), but these two regions were overlooked by Borowski (2007). The above record from Jordan is the first for this country (cf. Sharaf et al. 1983, Borowski 2007).

Ecology. Polyphagous species, with development taking place in death or debilitated branches of several woody plants. The following are plants known to be infested by this species and are present in Malta (cf. Haslam et al. 1977): *Acacia* sp., *Amygdalus communis*, *Arundo* sp., *Bambusa* sp., *Ceratonia siliqua*, *Cercis* sp., *Citrus* sp., *Eucalyptus* sp., *Ficus carica*, *Hibiscus sabdariffa*, *Laurus nobilis*, *Morus alba*, *Olea* sp., *Pinus halepensis*, *Pistacia lentiscus*, *P. vera*, *Prunus avium*, *P. dulcis*, *Punica granatum*, *Quercus* spp., *Rhamnus alaternus*, *Ulmus* sp. and *Vitis* spp. (cf. Lesne 1901b, as *S. Chevrieri* Villa [sic!], Peyerimhoff 1919, Novak 1952, both as *S. Chevrieri* (Villa) [sic!], Caillol 1954, Español 1955, as *S. chevrieri* Vill. [sic!], Bytinski-Salz and Sternlicht 1967, as *S. chevrieri* Villa [sic!], Compte 1970, as *S. chevrieri* (Villa [sic!], 1835), Georghiou 1977, as *S. chevrieri* Villa [sic!], Halperin and Damoiseau 1980, as *S. chevrieri* (Villa) [sic!], Lundberg et al. 1987, as *S. chevrieri* Villa [sic!], Nardi and Ratti 1995, Borowski and Mazur 2001, Nardi and Zahradník 2004, Akşit et al. 2005, as *S. chevrieri* Villa [sic!], Baena and Zuzarte 2013).

In central Italy (Marche region), this species (see above) was reared also from wood of *Salix* sp. and *Corylus avellana*, that represent new host-plant records for this Bostrichid (see above listed literature).

This species is often collected at light (Angelini 1996a, 1998, Chikatunov et al. 2006, Baena and Zuzarte 2013) and by window flight traps. Using these traps, large number of specimens were collected in forests of *Quercus calliprinos*, *Pinus halepensis* and *P. brutia* from northern Israel (Buse et al. 2010), in *Q. suber* forests from southern France (Brin et al. 2005, Brin and Brustel 2006), in an oak-hornbeam forest (*Quercocarpinetum boreoitalicum*) from northern Italy (Nardi and Zahradník 2004), in *Quercus ilex* forests of Sardinia, in a floodplain remnant of northern Italy (Nardi, unpublished data) and in mixed beechwoods of central Italy (Redolfi De Zan et al. 2014).

This species was also recorded from urban areas (Nardi 1997, Inglebert 2004).

Notes. *Scobicia pustulata* (Fabricius, 1801), a closely related Mediterranean species (cf. Borowski 2007), is here excluded from the Maltese fauna, since the record by Cameron and Caruana Gatto (1907) should refer to *S. chevrieri* as established by the examination of the above mentioned historical material. This is not a case of misidentification by Cameron and Caruana Gatto (1907), since they (Cameron and Caruana Gatto 1907: 383) based the nomenclature of this species on Heyden et al. (1891: 467) who listed “*Xyloperta pustulata* F.Kiesw.”, and “*Xyloperta pustulata* Kiesenw. (non F.)” is *S. chevrieri* (Lesne 1938: 57, as *S. Chevrieri* Villa [sic!], 1835). Only on the basis of this old litterature record, *S. pustulata* was erroneously listed from Malta by the above subsequent authors (Luigioni 1929, Nardi 2004b).

Scobicia chevrieri is a good colonizer of Mediterranean islands, since it is recorded also from other islands such as Montecristo (Tuscan Archipelago), Pantelleria, Lampedusa (cf. Nardi and Ratti 1995), Balearic and Columbretes Islands (cf. Lesne 1901b, as *S. Chevrieri* Villa [sic!], 1835, Español 1955, Vrydagh 1960b, as *S. chevrieri* Villa [sic!], 1835, Compte 1970, Nardi 2004b).

Xyloperthella picea (A.G. Olivier, 1790)

Material examined. Malta: Qormi, Hal-Farrug, 5.V.2003, DM [under bark of *Ceratonia siliqua* in an agricultural environment], 1 ♀ (CMM); Rabat, 4.VI.1989, PS, attracted to light [on the roof of private residence], 1 ex (CMM); 18.VI.1992, PS, [attracted to light, in an agricultural environment], 2 ex (CMM); 4.VI.1999, PS, attracted to light [in an agricultural environment], 1 ♂, (CMM); 28.VI.2001, PS, [attracted to light, in an agricultural environment], 1 ♀ (CMM); 21.VI.2002, PS, [ditto], 2 ♂♂ (CNI); 23.VI.2002, PS, [ditto], 2 ♀♀ (CMM; CNI); 1.VII.2002, PS, [ditto], 1 ♂ 1 ♀ (CMM); 3.VI.2003, PS, [ditto] 1 ♀ (CNI); Rabat, Ta Koronja, 21.VI.2002, PS, [attracted to light], 1 ♀ (CMM).

Other material examined. Italy: Apulia region, Lecce prov., S. Cataldo, Ris. [= Riserva = Reserve] WWF Le Cesine, 11–21.VI.1995, F. Angelini, 2 ex (MCSVA). Sardinia region, Sassari prov., Berehidda, 15.VII.1985, M. Daccordi leg., 1 ex (MCSV). Sicily region, Siracusa prov., Noto, Oasi di Vendicari, Cala Mosche, 10 m, N°36 49,066' E15°5,834', 3.VII.2011, D. Birtele & P. Birtele leg., 1 ♀ (CCI).

Chorotype. Afrotropical-Mediterranean species which was also intercepted in Germany, Great Britain and The Netherlands (cf. Lesne 1924, 1938, in both cases as *Xylopertha picea*, Aitken 1975, Akşit et al. 2005, Borowski 2007, in both cases as *Xyloperthella p. picea*, Bahillo de la Puebla et al. 2007, Borowski and Węgrzynowicz 2007, Baena and Zuzarte 2013). According to several authors (cf. Lesne 1901a, 1924, 1938, Gridelli 1939, 1940, Blackwelder 1945, Da Costa Lima 1953, in all cases as *Xylopertha picea*, Vrydagh 1958, 1960a, 1960b, 1961, 1962, Geis 2002, Ivie 2002, Borowski 2007, Peres Filho et al. 2007, Barriga and Cepeda 2009, Baena and Zuzarte 2013), it became established in parts of the Neotropical region (Argentina, Brazil, Colombia, French Guyana, Jamaica, Paraguay, Perù) since a long time but its presence in this region was recently reported as doubtful (Borowski and Węgrzynowicz 2007).

Notes. First record for Malta and the Apulia region; the species is not known from neighbouring Sicilian Islands (Tab. 1) but is widespread in Africa. Whether this species is autochthonous or has been introduced into Malta might be therefore open to debate. It is known from Tunisia and from all other mainland countries of North Africa (Lesne 1901b, Normand 1936, in both cases as *Xylopertha picea*, Vrydagh 1956, Borowski 2007), while from southern Europe it is recorded only from Portugal (cf. Serrano 1981, as *X. picea*, Baena and Zuzarte 2013), Greece (Samos Island) (Vrydagh 1962, Nardi 2004b, as *Xyloperthella p. picea*), southern mainland Italy (Basilicata region), Sardinia,

Sicily, Spain and the Balearic Islands (cf. Lesne 1901a, 1901b, 1905, Winkler 1927, Lui-gioni 1929, Porta 1929, in all cases as *Xylopertha picea*, Cobos 1950, as *Xyloperthe* [sic!] *picea*, Español 1955, 1974, Angelini and Montemurro 1986, Audisio et al. 1995, Sparacio 1997, Angelini 1998, Nardi 2004b, Bahillo de la Puebla et al. 2007, Borowski 2007).

According to some authors (cf. Lesne 1924, Español 1955), the Mediterranean populations may belong to a distinct subspecies, *Xyloperthella picea heydeni* (Schilsky, 1899), nevertheless the species is currently monotypic since this subspecies and also *X. picea plumbeipennis* Lesne, 1924 from Gabon and Democratic Republic of Congo (Lesne 1924), were recently listed as a synonym of *X. picea* (Borowski and Węgrzynowicz 2007). The species was described from Cape Verde Islands (Olivier 1790, as *Bostrihus piceus*) and not from Senegal as indicated by some authors (López-Colón 2000, López-Colón et al. 2001, Bahillo de la Puebla et al. 2007); in this African archipelago it is probably autochthonous (Oromí et al. 2005).

Subfamily Apatinae Billberg, 1820, Tribe Apatini Billberg, 1820

Apate monachus Fabricius, 1775

Material examined. Malta: Ghargħur, 18.VII.2013, DM, on living branchs of *Ficus carica*, 1 ♂ (CMM); Manikata, VII.2012, DM, in healthy branch (3–5 cm in diameter) of *Ceratonia siliqua* tunneled by this beetle, 1 ♀ death (CMM). Mellieħa, Kortin, 24.VII.2004, HB, U.V. light trap, 1 ♀ (CNI); 28.VII.2004, HB, U.V. light trap, 2 ♀♀ (CMM); 29.VII.2004, HB, U.V. light trap, 1 ♂ (CMM); 15.VIII.2004, HB, U.V. light trap, 1 ♂ (CMM); 28.VI.2005, HB, U.V. light trap, 1 ♀ (CMM); 3.VII.2005, HB, U.V. light trap, 1 ♂ (CNI); 5.VII.2005, HB, UV lights, 1 ♀ (CNI); 16.VII.2005, HB, U.V. light trap, 2 ♀♀ (CMM; CNI); 18.VI.2006, HB, UV lights, 1 ♂ 1 ♀ (CMM); 20.VI.2006, HB, UV lights, 1 ♀ (CNI); 2.VII.2006, HB, UV lights, 1 ♀ (CMM); 19.VI.2009, HB, UV lights, 1 ♂ (CMM); 5.VII.2009, HB, UV lights, 3 ♂♂ 1 ♀ (CMM; CNI); 11.VIII.2009, HB, UV lights, 2 ♂♂ (CMM); 13.VIII.2009, HB, UV lights, 1 ♂ (CNI); 22.VIII.2009, HB, UV lights, 1 ♂ (CMM); Mellieħa, Santa Maria Estate, 24.VII.2004, HB, 1 ♀ (CMM).

Other material examined. Italy: Calabria region, [Cosenza prov.,] Sibari, VII.1924, G. Leoni leg., 1 ex (MCZRL) (cf. Luigioni 1929: 642). Sardinia region, Cagliari prov., Geremeas, 18.VIII.2001, LF, 1 ♀ (CNI); ditto, ditto, [Island of Sant'Antioco,] Sant'Antioco, Torre Canai, 25.VIII.[19]79, Ferrara leg., 1 ♂ (MZUR); ditto, Oristano prov., Arborea, S. Anna, 10.VI.2004, LF, 6 ♂♂ 1 ♀ (CNI); ditto, Nuoro prov., Orosei, VI.1956, E. De Maggi leg., 1 ex (MCZRD); ditto, [Nuoro prov., Sinscola,] Capo Comino, 31.VIII.1973, L.G. Donadini leg., 1 ♀ (MCSV); ditto, [Sassari prov.,] Stintino, Punta Negra, 15.VII.1998, G. Mambrini leg., 1 ♂ (CCI). **France:** Corsica, Bastia, Pineto, 3.VIII.[19]80, A. Sette leg., 1 ♀ (MCSV).

Chorotype. Afrotropical-Mediterranean (northward upto Corsica and northern Spain). This species is established in the Neotropical region (Greater Antilles, Brazil,

etc.), and was intercepted in southern France, in central European countries and USA (cf. Vrydagħ 1960b, Reichardt 1964, Horion 1972, Aitken 1975, Geis 2002, fig. 31, Ivie 2002, Nardi 2004b, Bahillo de la Puebla et al. 2007, Borowski 2007, Barriga and Cepeda 2009, Ciesla 2011, Brustel and Aberlenc 2014).

Ecology. The genus *Apate* Fabricius, 1775 is one of the most notorious and troublesome forest pests in Africa (cf. Schabel 2006). *Apate monachus* is a polyphagous beetle, with over 80 host-plants used for larval development (cf. Lesne 1924, Rungs 1946, Boselli 1959, as *A. monachus* var. *rufiventris* P.H. Lucas, 1843, Peretz and Cohen 1961, Boselli 1962, as *Apate* [sic!] *monachus*, Chararas and Balakowski 1962, Prota 1963, Zanardi et al. 1969, Zocchi 1971, Halperin and Damoiseau 1980, Luciano 1982, Benfatto and Longo 1985, Sadok and Gerini 1988, Borowski and Mazur 2001, Gobbi 2003, Pisano et al. 2003, Schabel 2006, Bahillo de la Puebla et al. 2007, Di Franco and Benfatto 2008, Bonsignore et al. 2011, Ciesla 2011, Bonsignore 2012, Cillo and Bazzato 2012) of which, the following occur also in Malta (E. Lanfranco, pers. comm., 2014): *Acacia* spp., *Ailanthus glandulosa*, *Amigdalais communis*, *Arbutus unedo*, *Armeniaca vulgaris*, *Ceratonia siliqua*, *Citrus bigaradia*, *C. limon*, *C. limonia*, *C. nobilis*, *C. sinensis*, *Cupressus* sp., *Erica* sp., *Erythrina* sp., *Eucalyptus* spp., *Grevillea* sp., *Malus communis*, *M. domestica*, *Melia azedarach*, *Myrtus communis*, *Olea europaea*, *O. europaea* var. *oleaster*, *Persica vulgaris*, *Phoenix dactylifera*, *Pinus pinea*, *Pisidium guajava*, *Pistacia lentiscus*, *Pyrus amigdaliformis*, *P. dulcis*, *P. communis*, *P. communis* var. *piraster*, *Prunus amygdalus*, *P. armeniaca*, *P. domestica*, *P. persica*, *P. spinosa*, *Punica granatum*, *Quercus ilex*, *Schinus* sp., *Tamarix* sp., and *Vitis* spp. Adults are nocturnal and as observed also in Malta, they are frequently collected at light (cf. Lesne 1924, Sparacio 1997, Angelini 1998, Ragusa and Russo 1989, Chikatunov et al. 2006, Bahillo de la Puebla et al. 2007). In the Afrotropical region, *Lyctoderma africanum* (Grouvelle, 1900) and *L. testaceum* Lesne, 1913 (Bostrichidae, Lyctinae) are associated with adults of *A. monachus*, usually living under the abdomen and between the legs of these beetles (Lesne 1932, Paulian 1988: 501).

Notes. First record for Malta and it is not known from neighbouring Sicilian Islands (Tab. 1). Most of the Maltese specimens were collected at Mellieħa, Kortin using U.V. lights, with sex ratio of 1:1. This habitat can be best described as garigue but with pockets of low lying *Ceratonia siliqua* and *Pistacia lentiscus*. At Manikata, several healthy branches (3–5 cm in diameter) of *Ceratonia siliqua* were found tunneled by this beetle; the wood was drilled in the late summer, and the above mentioned specimen was found death in one of these holes. This area was recently converted into an agritouristic area and the owners were very concerned when they found the healthy branches of *Ceratonia siliqua* damaged by this beetle. According to the above records, the earliest captures of this species from Malta was 2004. This beetle is very conspicuous (10–19 mm of length: Bahillo de la Puebla et al. 2007) and it may be hypothesised that it is a recently established species in Malta. Having said this however, it is also worth mentioning that very few people in Malta were interested in collecting and studying these beetles in recent years. *Apate monachus* is also recorded from Tunisia (Lesne 1924, Borowski 2007) and Italy. In Italy it is known from three southern mainland

regions (Apulia, Basilicata and Calabria), Sicily and Sardinia (including some small circumsardinian islands) (cf. Dodero 1908, as *A. monachus* ab. *rufiventris*, Luigioni 1929, Porta 1929, as *A. monachus* a. *rufiventris*, Boselli 1959, 1962, Chararas and Balachowsky 1962, Prota 1963, Tassi 1967, Zanardi et al. 1969, Zocchi 1971, Luciano 1982, Benfatto and Longo 1985, Ragusa and Russo 1989, Audisio et al. 1995, Angelini 1996b, Sparacio 1997, Angelini 1998, Gobbi 2003, Pisano et al. 2003, Bonsignore et al. 2011, Bonsignore 2012, Cillo and Bazzato 2012). Its doubtfull presence “N?”, in northern Italy (Audisio et al. 1995) was later confirmed by records from South Tyrol (Kahlen and Hellrighl 1996) that are probably based only on interceptions, since its establismnt in the mentioned Alpine region is climatically improbable (Nardi, unpublished data).

The aedeagus of this species was figured by Jeannel and Paulian (1944: 91, figs 75, 87) and by Jeannel (1955: 57, fig. 32c).

The nomenclatorial problems for *Apate* Fabricius, 1775 were discussed by Borowski and Węgrzynowicz (2009).

Discussion

Table 1 summarized the species of Bostrichidae recorded from the Maltese Archipelago and neighbouring Sicilian Islands with each assigned to a chorotype. Altogether 12 species are known: 4 from Pantelleria, 1 from Linosa, 1 from Gozo, 9 from Malta and 1 from Lampedusa; and only *Scobia chevrieri* is recorded from four islands (except from Linosa).

For Sicily and Tunisia, 22 (Catara and Barbagallo 1980, Benfatto and Longo 1985, Audisio et al. 1995, Nardi 2004b, 2004b, Suma and Russo 2005, Muscarella et al. 2013) and 30 species respectively (Borowski 2007) of Bostrichidae are recorded. The relatively small number of species recorded from the Sicilian Channel Islands (Tab. 1) is probably due to the limited surface area, by the absence of many host plants, and the absence of certain habitats such as moutain and sub-mountain ranges.

Excluding *Sinoxylon sexdentatum* (A.G. Olivier, 1790) (= *muricatum* (Linnaeus, 1767) *nomen oblitum*) and *Xylopertha praeusta* (Germar, 1817), Malta hosts all species recorded from the Sicilian Channel Islands (Tab. 1). Some of the host plants (e.g. *Vitis* spp., *Pistacia* spp., *Ficus carica*, *Malus domestica*, *Olea europaea*, *Prunus persica*, *Pyrus communis*, *Quercus ilex*, *Ceratonia siliqua*, *Cercis siliquastrum*, *Citrus* spp.) of the former species (cf. Lesne 1901b, 1906, Peyerimhoff 1919, Halperin and Damoiseau 1980, Gobbi 1984, Benfatti and Longo 1985, Moleas 1988, Mart et al. 1995, Ratti and Nardi 1995, Mourikis et al. 1998, Pollini 1998, Aksit et al. 2005) are present also in Malta (cf. Borg 1962, Saliba 1963, Haslam et al. 1977), so this polyphagous beetle could be eventually found in Malta or may establish itself. The occurrence of *X. praeusta*, which develops chiefly in the wood of oak trees (*Quercus ilex*, *Q. suber*, *Q. robur*, *Q. mirbecki*) (Lesne 1901b, as *Xylonites praeustus* Germar, Rungs 1946, as *X. praeustus* Germ., Español 1955, Bahillo de la Puebla et al. 2007), is less probable since oak trees are rare in the Maltese Islands.

Table 1. Bostrichidae recorded from Maltese Archipelago and neighbouring islands with respective chorotype codes. Abbreviations: A = Aitken 1975; AFM = Afrotopical-Mediterranean; C = Cameron and Caruana Gatto 1907; CEM = Centralasiatic-European-Mediterranean; COS = Cosmopolitan; D = Denux and Zagatti 2010; EUR = European; F = Falzon et al. 2012; G = Goggi 2004; H = Hyde and Daubney 1960; L = Luigioni 1929; LAMP. = Lampedusa; LINO. = Linosa; MED = Mediterranean; Na = Nardi 2004a; Nb = Nardi 2004b; NR = Nardi and Ratti 1995; PANTEL. = Pantelleria; SEU = S-European; TEM = Turanic-European-Mediterranean; TUM = Turanic-Mediterranean; WME = W-Mediterranean; ZGG = Chorotype; ! = this paper; () = misinterpretation.

Species	PANTEL.	LINO.	GOZO	MALTA	LAMP.	ZGG
<i>Lycus brunneus</i>				C L Na D !		COS
<i>Trogoxylon impressum</i>	NR			!		TEM
<i>Rhyzopertha dominica</i>				H D !		COS
<i>Amphicerus bimaculatus</i>				!		TUM
<i>Bostrychus capucinus</i>	NR			A Nb		CEM
<i>Heterobostrychus aequalis</i>				!		COS
<i>Scobicia chevrieri</i>	NR		!	(C L Nb) F !	NR	MED
<i>Sinoxylon sexdentatum</i>	NR					MED
<i>Sinoxylon unidentatum</i>				!		COS
<i>Xylopertha praeusta</i>		G				WME
<i>Xyloperthella picea</i>				!		AFM
<i>Apate monachus</i>				!		AFM
TOTAL/ISLANDS	4	1	1	10	1	

With the current available data, no conclusions can be drawn on how the bostrichid fauna colonised the Maltese Islands, but we can say that almost all species are also known from Sicily and North Africa, and are normally widely spread in the Mediterranean area. In fact, from a zoogeographical point of view (cf. Tab. 2) wide chorotypes prevails: 40% Cosmopolitan, 30% Palaearctic, 20% Afrotopical-Mediterranean and 10% Mediterranean. European chorotypes are absent; this situation was observed in other families of Coleoptera (Mifsud and Bílý 2002, Nardi and Mifsud 2003, Háva and Mifsud 2006), whereas they are represented in two other families of saproxyllic beetles (Mifsud 2002, Mifsud and Knížek 2009).

Species of bostrichids are easily transported with infested woods and cereal, and are frequently observed in harbour areas (cf. Aitken 1975, Contessi 1991, Contessi and Mucciolini 1993, 1998, Peck et al. 1998, Haack and Cavey 2000, Geis 2002, Nicoli Aldini 2003, 2004, Ratti 2004, Borges et al. 2006, Haack 2006, Liu et al. 2006, Majka 2007, Ratti 2007, Beiriger 2010, Humble 2010, Leal et al. 2010, Price et al. 2011). The interception in Malta of other alien species from warm climatic regions is highly likely, and their possible establishment could also be promoted by global warming (cf. Parmesan 2006, Franceschini et al. 2009, Walther et al. 2009, Fronzek et al. 2010, Robinet and Roques 2010, Roques 2010, Thuiller et al. 2011).

Table 2. Chorotypes of the Bostrichidae from Malta Archipelago and neighbouring islands, their distribution and percents (abbreviations as in Table 1).

Chorotypes	Number of species (%)	Pantelleria	Linosa	Maltese Islands	Lampedusa
COS	4 (33.33)			4 (40)	
AFM	2 (16.66)			2 (20)	
CEM	1 (8.33)	1 (25)		1 (10)	
TEM	1 (8.33)	1 (25)		1 (10)	
TUM	1 (8.33)			1 (10)	
MED	2 (16.66)	2 (50)		1 (10)	1 (100)
WME	1 (8.33)		1 (100)		
Total	12	4	1	10	1

From a faunistic point of view, six species are here recorded for the first time, namely: *Trogoxylon impressum*, *Amphicerus bimaculatus*, *Heterobostrychus aequalis*, *Sinoxylon unidentatum*, *Xyloperthella picea* and *Apate monachus*.

The Mediterranean area is one of the most significantly altered biodiversity hot-spots on Earth, since it has been intensively affected by human activity for millennia. As a result, only 4.7% of its primary vegetation remained unaltered and the landscape has been repeatedly transformed (cf. Myers et al. 2000, Cuttelod et al. 2008, Geri et al. 2010, Carroll et al. 2012). This transformation has negatively influenced xylophagous insect populations, especially in small and isolated environments, like those of islands (cf. Becker 1975, Howart and Ramsay 1991, Becker 1992, Crisp et al. 1998, Mantisi 2001, Pasta and La Mantia 2002). In this framework, half of bostrichid species recorded from the Maltese Islands (*Amphicerus bimaculatus*, *Bostrichus capucinus*, *Scobicia chevrieri*, *Xyloperthella picea*, *Apate monachus*), are included in the European Red list of saproxylic beetles (Nieto and Alexander 2010: 29), but luckily all belong only to the IUCN category “LC” (Least Concern).

Acknowledgements

We thank Mr Henry Borg Barthet (Malta), Mr Alessandro B. Biscaccianti (Rome, Italy), Mr Daniele Baiocchi (Rome, Italy), Dr Louis F. Cassar (University of Malta, Malta), Mr David Dandria (Malta), Dr Andrea Di Giulio (Università Roma Tre, Rome, Italy), Dr Stefano De Felici (Università Tor Vergata, Rome, Italy), Mr Charles Farrugia (Malta), Dr Paolo Maltzeff (Rome, Italy), Dr Lara Redolfi De Zan (Università Roma Tre, Rome, Italy) and Mr Paul Sammut (Malta), who provided specimens of some species mentioned in the present work; Dr Jerzy Borowski (Institute of Forest Sciences, University of Łódź, Branch in Tomaszów Mazowiecki, Poland) for the determination of *Sinoxylon unidentatum* and for relevant information; Mr Paolo Cornacchia (Porto Mantovano, Italy), Dr Luca Bartolozzi (Museo di Storia Naturale dell’Università degli Studi di Firenze, Sezione di Zoologia “La Specola”, Florence, Italy), Dr Roberto Casalini, Dr Vincenzo Vomero and Dr Alberto Zilli (Museo civico di Zoologia, Rome, Ita-

ly), Dr Leonardo Latella (Museo civico di Storia Naturale, Verona, Italy), Mr Maurizio Mei, Dr Emanuele Piattella and Prof. Augusto Vigna Taglianti (Museo di Zoologia, Università degli Studi di Roma “Sapienza”, Rome, Italy) for help during the examination of the collections under their care; Mr Edwin Lanfranco (University of Malta) for information on Maltese vegetation; Dr Gianluca Scaglioni (Porto Mantovano, Italy) and Dr Laura Spada (Centro Nazionale per lo Studio e la Conservazione della Biodiversità Forestale “Bosco Fontana”, Italy), for their kind help in the lab; Dr Piero Leo (Cagliari, Italy), Dr Andrew W. Ridley (Department of Agriculture, Fisheries and Forestry, Brisbane, Australia) and Dr Marco Uliana (Museo di Storia Naturale di Venezia, Venice, Italy) who graciously provided copies of three papers.

References

- Abivardi C (2001) Iranian Entomology. Vol. 2. Applied entomology. Springer Verlag, Berlin-Heidelberg-New York, I-VIII + 445–1033.
- Adeli E (1972) Beitrag zur Kenntnis der Forst schädlichen Insekten des Iran. I. Coleoptera. Zeitschrift für Angewandte Entomologie 70: 8–14. doi: 10.1111/j.1439-0418.1972.tb02144.x
- Adlbauer K (1998) Die Splintholzkäferfauna der Steiermark (Coleoptera, Lyctidae). Mitteilungen Landesmuseum Joanneum, Zoolgie 51: 55–58.
- Aguilera AO (2006) Analisis de riesgo para *Heterobostrychus aequalis*. <http://www.semarnat.gob.mx/temas/gestionambiental/forestalsuelos/Paginas/saludforestal.aspx> [accessed 21 October 2013]
- Ahmed A, Ahmad T, Arian MA, Ahmed M (2008) Management of Bagged Wheat godowns to control the stored grain insect pests. Pakistan Entomologist 30(1): 31–36.
- Ahmed KS (1996) Studies on the Ectoparasitoid, *Anisopteromalus calandrae* Howard (Hymenoptera: Pteromalidae) as a Biocontrol Agent against the Lesser Grain Borer, *Rhyzopertha dominica* (Fab.) in Saudi Arabia. Journal of Stored Products Research 32(2): 137–140. doi: 10.1016/0022-474X(96)00005-7
- Aitken AD (1975) Insect Travellers, 1, Coleoptera. Technical Bulletin Ministry of Agriculture Fisheries and Food, HMSO, London, 31, I–XVI + 1–191.
- Akbulut S, Keten A, Yüksel B (2008) Wood Destroying Insects in Düzce Province. Turkish Journal of Zoology 32: 343–350.
- Akşit T, Çakmak U, Özsemerci F (2005) Some New Xylophagous Species on Fig Trees (*Ficus carica* cv. *Calymirna* L.) in Aydin, Turkey. Turkish Journal of Zoology 29: 211–259.
- Alexander KNA, Anderson R (2012) The beetles of decaying wood in Ireland. A provisional annotated checklist of saproxylic Coleoptera. Irish Wildlife Manuals, No. 65. National Parks and Wildlife Service, Department of the Arts, Heritage and the Gaeltacht, Dublin, Ireland, 161 pp.
- Alfieri A (1976) The Coleoptera of Egypt. Memoires de la Societe entomologique d’Egypte 5: I–XVI + 1–362.

- Altson AM (1924) On the genital system of *Lyctus brunneus* Steph., with a note on *Lyctus linearis* Goeze (Coleoptera). The Journal of the Linnean Society of London, Zoology 35: 581–597 + pls. 31–34.
- Amin AH, Assaggaf AI, Al-Robai AAS (1986) Survey and relative abundance of some Coleopterous Insects attracted to a light trap in Jeddah, Saudi Arabia. Bulletin de la Société entomologique d’Egypte 66: 299–317.
- Anderson R, Nash R, O’Conor JP (1997) Irish Coleoptera. A revised and annotated list. Special Entomological Supplement to The Irish Naturalists’ Journal, 81 pp.
- Angelini F (1996a) Coleotterofauna della riserva naturale WWF Lago di Pignola. Basilicata, Potenza. Alfagrafica Volonnino, Lavello, 135 pp.
- Angelini F (1996b) I Coleotteri in Basilicata. Risorsa Natura in Basilicata. Le Aree protette regionali. Ufficio Stampa Consiglio Regionale Basilicata, Potenza 9(5/6): 165–170.
- Angelini F (1998) Coleotterofauna reperita mediante trappola luminosa in due stazioni umide della Basilicata (Italia meridionale). Quaderni della Stazione di Ecologia del civico Museo di Storia naturale di Ferrara 11: 7–37.
- Angelini F, Montemurro F (1986) Coleotterofauna del bosco di Policoro (Matera). Biogeographia, Lavori della Società italiana di Biogeografia 10 (1984): 545–604.
- Anwar M, Ashfaq M, Hasan M, Anjum FM (2005) Efficacy of *Azadirachta indica* L. oil on bagging material against some insect pests of wheat stored in warehouses at Faisalabad. Pakistan Entomologist 27(1): 89–94.
- Arthur FH, Ondier GO, Siebenmorgen TJ (2012) Impact of *Rhyzopertha dominica* (F.) on quality parameters of milled rice. Journal of Stored Products Research 48: 137–142. doi: 10.1016/j.jspr.2011.10.010
- Asanov K (1980) Predators and parasites of the lesser grain borer. Zashchita Rastenii 5: 23.
- Ashouri S, Shayesteh N (2009) Insecticidal activities of Black Pepper and Red Pepper in powder form on adults of *Rhyzopertha dominica* (F.) and *Sitophilus granarius* (L.). Pakistan Entomologist 31(2): 122–127.
- Attia R, Kamel AH (1965) The fauna of stored products in U.A.R. Bulletin de la Société entomologique d’Egypte 49: 221–232.
- Audisio P, Gobbi G, Liberti G, Nardi G (1995) Coleoptera Polyphaga IX (Bostrichoidea, Cleroidea, Lymexyloidea). In: Minelli A, Ruffo S, La Posta S (Eds) Checklist delle specie della fauna italiana, 54. Calderini, Bologna, 1–29.
- Aydin N, Soran H (1987) The pests and contaminators of stored wheat and flour mills in the Thrace region. Turkiye I. Entomoloji Kongresi Bildirileri, 13–16 Ekim 1987, Ege Universitesi, Bornova, Izmir, 717–726.
- Azmi MSMd, Abood F, Razi NA (2011) World Distribution of *Heterobostrychus aequalis* Waterhouse (Coleoptera: Bostrichidae). Journal of Entomology 8(6): 497–511. doi: 10.3923/je.2011.497.511
- Baena M, Zuzarte AJ (2013) Contribución al estudio de los Bostríquidos de Portugal y actualización del catálogo de la fauna ibérica (Coleoptera, Bostrichidae). Zoologica baetica 24: 25–51.
- Bartali H, Dunkel FV, Said A, Sterling RL (1990) Performance of plastic lining for storage of barley in traditional underground structures (Matmora) in Morocco. Journal of Agricultural Engineering Research 47(4): 297–314. doi: 10.1016/0021-8634(90)80049-Z

- Bahillo de la Puebla P, López-Colón JI (2001) Los Bostrichidae Latreille, 1802 de la Comunidad Autónoma Vasca y áreas limítrofes (Coleoptera). *Heteropterus Revista de Entomología* 1: 25–40.
- Bahillo de la Puebla P, López-Colón JI, Baena M (2007) Los Bostrichidae Latreille, 1802 de la fauna de ibero-balear (Coleoptera). *Heteropterus Revista de Entomología* 7(2): 147–227.
- Barriga JE, Cepeda DE (2009) Bostrichidae (incl. Lyctinae). In: Barriga JE (Ed.) Coleoptera Neotropical. <http://coleoptera-neotropical.org/paginaprincipalhome.html> [accessed 21 October 2013]
- Baudi [di Selve] F (1873) Catalogo dei Dascillidi, Malacodermi e Teredili della Fauna europea e circummediterranea appartenenti alle collezioni del Museo Civico di Genova. Annali del Museo Civico di Storia Naturale di Genova, sezione 1, 4: 226–268.
- Baudi di Selve F (1874) Coleopterorum messis in insula Cypro et Asia minore ab Eugenio Truqui congregatae recensitio: de Europaeis notis quibusdam additis. Pars quinta. *Berliner Entomologische Zeitschrift* 17 (1873): 317–338.
- Becker G (1969) Über einige Funde holzzerstörender Insekten auf Korsika. *Zeitschrift für Angewandte Entomologie* 63: 93–98. doi: 10.1111/j.1439-0418.1969.tb04365.x
- Becker P (1975) Island Colonization by Carnivorous and Herbivorous Coleoptera. *Journal of Animal Ecology* 44(3): 893–906. doi: 10.2307/3726
- Becker P (1992) Colonization of Islands by Carnivorous and Herbivorous Heteroptera and Coleoptera: Effects of Island Area, Plant Species Richness, and ‘Extinction’ Rates. *Journal of Biogeography* 19(2): 163–171. doi: 10.2307/2845502
- Beiriger R (2010) *Heterobostrychus hamatipennis* Lesne (Coleoptera: Bostrichidae) new to Florida. *Insecta Mundi* 138: 1–5.
- Benfatto D, Longo S (1985) I coleotteri fitofagi degli agrumi in Italia. *Informatore fitopatologico* 35(2): 15–22.
- Benhalima H, Chaudhry MQ, Mills KA, Price NR (2004) Phosphine resistance in stored-product insects collected from various grain storage facilities in Morocco. *Journal of Stored Products Research* 40: 241–249. doi: 10.1016/S0022-474X(03)00012-2
- Beeson CF, Bhatia BM (1937) On the biology of the Bostrichidae (Coleoptera). *Indian Forest Records* 2: 223–320.
- Bertolini S (1874) Catalogo sinonimico e topografico dei Coleotteri d’Italia. *Supplemento del Bullettino della Società entomologica italiana* 7: 157–204.
- Blackwelder RE (1945) Checklist of the coleopterous insects of Mexico, Central America, the West Indies, and South America. Part 3. United States National Museum Bulletin 185: I–IV + 341–550.
- Bonsignore CP (2012) *Apate monachus* (Fabricius, 1775) [sic!], a Bostrichid Pest of Pomegranate and Carob Trees in Nurseries – Short Communication. *Plant Protection Science* 48 (2): 94–97.
- Bonsignore CP, Manti F, Castiglione E (2011) *Apate monachus*, un Bostrichidae dannoso al Melograno e al Carrubo in vivaio. *Atti XXIII Congresso Nazionale Italiano di Entomologia*, Genova, 13–16 giugno 2011: 208.
- Borg J (1962) Cultivation and Diseases of Fruit Trees in the Maltese Islands. Government Printig Office, Malta, I–VII + 622 pp.

- Borges PAV (1990) A checklist of the Coleoptera from the Azores with some systematic and biogeographic comments. *Boletim do Museu Municipal do Funchal* 42(220): 87–136.
- Borges PAV, Lobo JM, de Azevedo EB, Gaspar CS, Melo C, Nunes LV (2006) Invasibility and species richness of island endemic arthropods: a general model of endemic vs. exotic species. *Journal of Biogeography* 33(1): 169–187. doi: 10.1111/j.1365-2699.2005.01324.x
- Borowski J (2007) Bostrichidae. In: Löbl I, Smetana A (Eds) Catalogue of Palaearctic Coleoptera. Volume 4. Elateroidea – Derodontoidea – Bostrichoidea – Lymexyloidea – Cleroidea – Cucujoidea. Apollo Books, Stenstrup, 320–328.
- Borowski J (2013) Errata for volume 4 [Bostrichidae]. In: Löbl I, Smetana A (Eds) Catalogue of Palaearctic Coleoptera. Volume 8. Curculionoidea II. Apollo Books, Stenstrup, 33–34.
- Borowski J, Mazur S (2001) Contribution to the knowledge of the Bostrichidae and associated Histeridae of Morocco (Insecta, Coleoptera). *Il Naturalista valtellinese – Atti Museo civico di Storia naturale di Morbegno* 12: 69–75.
- Borowski J, Węgrzynowicz P (2007) World Catalogue of Bostrichidae (Coleoptera). Wydawnictwo Mantis, Olsztyn, 248 pp.
- Borowski J, Węgrzynowicz P (2009) *Apate* Fabricius, 1775 (Bostrichidae: Coleoptera): a protected name. *Annales Zoologici* 59(2): 189–191. doi: 10.3161/000345409X464001
- Boselli F (1959) *Apate monachus* Fabr. var. *rufiventris* Lucas (Coleoptera Bostrychidae) nuovo parassita del mandorlo e del pesco, limone e dell’arancio in Sardegna. *Annali della Sperimentazione agraria* (n.s.) 13(1): 35–46.
- Boselli F (1962) La voce degli osservatori delle malattie delle piante. *Osservatorio della Sardegna. Informatore fitopatologico* 12(4): 271–273.
- Brin A, Brustel H (2006) Réponse des Coléoptères saproxyliques à l’hétérogénéité des subéraies dans le Massif des Maures (France). *Revue d’Ecologie (Terre et Vie)* 61: 327–342.
- Brin A, Brustel H, Valladares L (2005) Inventaire de coléoptères saproxyliques de la suberaie de Saint-Daumas (Maures, Var). WWF-France, CEEP, Ecole Supérieure d’Agriculture de Purpan (ESAP), 27 pp.
- Brustel H, Aberlenc H-P (2014) Les Bostrichidae Latreille, 1802 de la faune de France: espèces autochtones, interceptées, introduites ou susceptibles de l’être (Coleoptera). *R.A.R.E.* 23(2): 65–68.
- Buchi R (1993) Inquiry on the incidence of pests and the use of insecticides in granaries, feed stocks and mills. *Landwirtschaft Schweiz* 6(1): 10–12.
- Burakowski B, Mroczkowski M, Stefańska J (1986) Katalog Fauny Polski – Catalogus faunae Poloniae Część XXIII, Tom 11 Chrząszcze Coleoptera. Dermestoidea, Bostrychoidea, Cleroidea et Lymexyloidea. PNW, Muz. Inst. Zool. PAN Warszawa, 243 pp.
- Burlini M (1972) Una infestazione di *Lyctus brunneus* Steph. sui pavimenti a parchetti (Coleoptera Lyctidae). *Bollettino della Società entomologica italiana* 104(8): 176.
- Buse J, Levanony T, Timm A, Dayan T, Assmann T (2010) Saproxylic beetle assemblages in the Mediterranean region: Impact of forest management on richness and structure. *Forest Ecology and Management* 259: 1376–1384. doi: 10.1016/j.foreco.2010.01.004
- Bytinsky-Salz H (1966) An annotated list of the insects and mites introduced into Israel. *Israel Journal of Entomology* 1: 15–48.

- Bytinski-Salz H, Sternlicht M (1967) Insects associated with Oaks (*Quercus*) in Israel. Israel Journal of Entomology 2: 107–143.
- Cameron M, Caruana Gatto A (1907) A list of the Coleoptera of the Maltese Islands. Transactions of the Entomological Society of London 59(3): 383–403.
- Carroll FA, Hunt CO, Schembri PJ, Bonanno A (2012) Holocene climate change, vegetation history and human impact in the Central Mediterranean: evidence from the Maltese Islands. Quaternary Science Reviews 52: 24–40. doi: 10.1016/j.quascirev.2012.07.010
- Catara A, Barbagallo S (1980) Difesa (Cap. 5). In: Baldini E, Scaramuzzi F (Eds) Gli Agrumi. Manuale REDA, Roma, 167–248.
- Cavalloro F, Ratti E (1978) Una infestazione da *Lyctus brunneus* (Steph.) e *Bostrychus capucinus* (L.) su radica di erica (*Erica arborea* L.) (Coleoptera Lyctidae e Bostrichidae). Bollettino della Società entomologica italiana 110(4/6): 69–71.
- Champ BR (1986) Occurrence of resistance to pesticides in grain storage pests. In: Champ BR, Highley E (Eds) Pesticides and Humid Tropical Grain Storage Systems. Proceedings of an international Seminar, Manila, Philippines, 27–30 May 1985: 229–255.
- Chararas C, Balachowsky R (1962) Famille des Bostrichidae. In: Balachowsky R (Ed.) Entomologie appliquée à l'agriculture. Tome I. Coléoptères premier volume. Mason et Cie., Paris, 304–315.
- Chetcuti D, Buhagair A, Schemnri PJ, Ventura F (1992) The climate of the Maltese Islands: a review. Malta Univesity Press, Msida, I–VI + 108 pp.
- Chiappini E, Liotta G, Reguzzi MC, Battisti A (2001) Insetti e restauro. Legno, carta, tessuti, pellame e altri materiali. Calderini Edagricole, Bologna, 260 pp.
- Chikatunov V, Pavlíček T, Nevo E (2004) Coleoptera of “Evolution Canyon” Lower Nahal Oren, Mount Carmel, Israel Part II. Pentsoft, Sofia–Moscow, Series Faunistica 31, 192 pp.
- Chikatunov V, Kravchenko VD, Dorchin Y, Müller G (2006) The xylophagous beetles (Buprestidae, Lyctidae, Bostrichidae, Anobiidae, Cerambycidae, Scolytidae, Platypodidae) (Coleoptera) collected in the Israeli light trap survey and their association with the major phytogeographical zones of Israel. Esperiana. Buchreiche zur Entomologie 12: 313–320.
- Ciampolini M, Corazia F, Lunghini D (1989) Danni da *Amphicerus bimaculatus* alla vite nel Lazio. L’Informatore agrario 45(3): 93–95.
- Ciesla W (2011) Forest Entomology. Wiley-Blackwell, I–X + 81 pls. + 416 pp.
- Cillo D, Bazzato E (2012) Segnalazioni faunistiche italiane 533 – *Ptinomorphus regalis* (Duftschmid, 1825) (Coleoptera Anobiidae). Bollettino della Società entomologica italiana 144(1): 45.
- Cobos A (1950) Datos para el catalogo de los Coléopteros de España – Especies de los alrededores de Málaga. Boletín de la Real Sociedad Española de Historia Natural 47 (1949): 563–609.
- Cola L (1971) Mit fremden Hölzern eingeschleppte Insekten, insbesondere Scolytidae und Platypodidae. Anzeiger für Schädlingskunde und Pflanzenschutz 44(5): 65–68.
- Compte A (1970) Los coleópteros de las Islas Columbretes. Eos: Revista española de Entomología 45: 97–137.
- Contessi A (1991) *Prostephanus truncatus* Horn, un nuovo potenziale infestante delle derrate in magazzino. Informatore fitopatologico 41(11): 21–24.

- Contessi A, Mucciolini G (1993) Considerazioni sugli insetti infestanti le derrate in importazione attraverso il porto di Ravenna nel periodo 1987–1991. In: Domenichini G (Ed.) Atti del 5° Simposio “La difesa antiparassitaria nelle industrie alimentari e la protezione degli alimenti”, Piacenza, 23–25 settembre 1992. Chiriotti Editori, Pinerolo, 163–173.
- Contessi A, Mucciolini G (1998) Considerazioni sugli Insetti infestanti le derrate in importazione attraverso il porto di Ravenna nel quinquennio 1992–1996. In: Cravedi P (Ed.) Atti del 6° Simposio “La difesa antiparassitaria nelle industrie alimentari e la protezione degli alimenti”, Piacenza, 24–26 settembre 1997. Chiriotti Editori, Pinerolo, 75–95.
- Clermont J (1909) Liste de Coléoptères récoltés en Trancaucasie par M. Louis Mesmin. *Miscellanea entomologica* 17(1): 1–6.
- Cloudsley-Thompson JL (1962) Some aspects of the fauna of the Red Sea hills and coastal plain. *Entomologist's monthly Magazine* 98: 159–161.
- Crisp PN, Dickinson KJM, Gibbs GW (1998) Does native invertebrate diversity reflect native plant diversity? A case study from New Zealand and implications for conservation. *Biological Conservation* 83(2): 209–220. doi: 10.1016/S0006-3207(97)00053-0
- Cuttelod A, García N, Abdul Malak D, Temple H, Katariya V (2008) The Mediterranean: a biodiversity hotspot under threat. In: Vié J-C, Hilton-Taylor C, Stuart SN (Eds) The 2008 Review of The IUCN Red List of Threatened Species. IUCN, Gland, Switzerland, 1–13.
- Cymorek S (1961) Die in Mitteleuropa einheimischen und eingeschleppten Splinthokafer aus der Familie Lyctidae. Ihre Systematik und Notizen zur Verbreitung, zum Schasaufreten und zur Biologie. *Entomologische Blätter für Biologie und Systematik Käfer* 57(2): 76–102.
- Da Costa Lima AM (1953) Insetos do Brasil. 8.º Tomo. Coleópteros. 2.ª parte. Escola Nacional de Agronomia, Série Didática 10, 1–323.
- Dal Monte G (1958) Il Cappuccino dei cereali. *Informatore fitopatologico* 12: 216–220.
- Damoiseau R (1966) Les Bostrichidae du Musée de Moravie à Brno (Coleoptera - Cucujidea). *Casopis Moravskeho Musea, Vedy přírodní – Acta Musei Moraviae, Scientiae naturales* 51: 291–300.
- Damoiseau R (1979) Insects of Saudi Arabia Coleoptera: Fam. Bostrichidae. In: Wittmer W, Büttiker W (Eds) Fauna of Arabia. Pro Entomologica, Natural History Museum, Basel and Ciba-Geigy Ltd., Basel 1, 249–250.
- Denux O, Zagatti P (2010) Coleoptera families other than Cerambycidae, Curculionidae sensu lato, Chrysomelidae sensu lato and Coccinellidae. Chapter 8.5. In: Roques A, Kenis M, Lees D, Lopez-Vaamonde C, Rabitsch W, Rasplus J-Y, Roy DB (Eds) Alien terrestrial arthropods of Europe. *BioRisk* 4(1): 315–406.
- Derwesh AI (1965) A preliminary list of identified Insects and some Arachnids from Iraq. Ministry of Agriculture, Directorate general of agricultural Research and Projects, Bagdad, Iraq, *Bulletin* 112: 1–123.
- Di Franco F, Benfatto D (2008) Nuove infestazioni di coleotteri Bostrichidi nei limoneti siciliani. *Informatore fitopatologico*, 50 supplemento a Terra e Vita 26: 26–27.
- Dodero A (1908) Appunti coleotteroologici. *Rivista coleotteroologica italiana* 6(5): 93–102.
- Drumont A, Grootaert P (2011) Saproxyllic beetles from Belgium, online distribution maps of species (Coleoptera). <http://projects.biodiversity.be/beetles/> [accessed 7 August 2013]

- Edde PA (2012) A review of the biology and control of *Rhyzopertha dominica* (F.) the lesser grain borer. *Journal of Stored Products Research* 48: 1–18. doi: 10.1016/j.jspr.2011.08.007
- Emekçi M, Ferizli AG (2000) Current status of stored products protection in Turkey. In: Adler C, Schoeller M (Eds) IOBC / WPRS Working Group “Integrated Protection in Stored Products”. IOBC wprs Bulletin 23(10): 39–46.
- Español F (1949) Datos para el conocimiento de los tenebriónidos del Mediterráneo occidental. *Graellsia* 7: 43–48.
- Español F (1955) Los bostríquidos de Cataluña y Baleares (Col. Cucujoidea). *Publicaciones del Instituto de Biología Aplicada* 21: 107–135.
- Español F (1956a) Los Lícidos de Cataluña (Col. Cucujoidea). *Publicaciones del Instituto de Biología Aplicada* 23: 123–138.
- Español F (1956b) Lícidos y bostríquidos de la zona mediterránea de Marruecos. *Publicaciones del Instituto de Biología Aplicada* 24: 73–75.
- Español F (1971) Sobre algunos coleópteros asociados a bostríquidos, lícidos y anbidos. *Boletín Servicio Plagas Forestales* 14(27): 17–21.
- Español F (1974) Nuevos datos sobre los Bostrychidae de la fauna española (Col. Cucujoidea). *Graellsia* 28 (1972): 37–44.
- Español F (1979) Los Ulomini de la fauna española (Col. Tenebrionidae). *Memorias de la Real Academia de Ciencias y Artes de Barcelona* 44(15): 413–432.
- Fabricius JC (1792) *Entomologia Systematica emendata et aucta. Secundum classes, ordines, genera, species adjectis synonymis, locis, observationibus, descriptionibus. Tom I. Pars II. C. G. Proft, Hafniae*, 538 pp. doi: 10.5962/bhl.title.36532
- Fisher WS (1950) A revision of the North American Species of Beetles belonging to the family Bostrichidae. *United States Department of Agriculture Miscellaneous Publications* 698: 1–157.
- Franceschini A, Longo S, Moricca S (2009) Avversità biotiche e mutamenti climatici in ambienti forestali. In: Ciancio O (Ed.) *Atti del Terzo Congresso Nazionale di Selvicoltura per il miglioramento e la conservazione dei boschi italiani*, 16–19 Ottobre 2008, Taormina (Messina), volume secondo. Accademia Italiana di Scienze Forestali, 605–610.
- Fronzek S, Carter TR, Jylhä K (2010) Scenarios of Climate Change for Europe. In: Settele J, Penev L, Georgiev T, Grabaum R, Grobelnik V, Hammer V, Klotz S, Kühn I (Eds) *Atlas of Biodiversity Risks*. Pensoft, Sofia–Moscow, 68–71.
- Gambetta A (1983) List of insects intercepted in imported timbers in Italy. Document – The International Research Group on Wood Preservation 1217: 1–3.
- Gambetta A, Orlandi E (1982a) Su alcuni insetti reperiti nel legname nei depositi. I° Lictidi e Bostrichidi. Contributi scientifico-pratici per una migliore conoscenza ed utilizzazione del legno 30(77): 8–29.
- Gambetta A, Orlandi E (1982b) Durabilità naturale di 100 legni indigeni e di importazione a funghi, insetti e organismi marini. Contributi scientifico-pratici per una migliore conoscenza ed utilizzazione del legno 30(79): 45–71.
- Geis K-U (2002) Gebietsfremde Splintholz- und Bohrkäfer, nach Mitteleuropa mit Importholz und anderen Gütern eingeschleppt. – Eine Bestandsaufnahme (Coleoptera: Lyctidae, Bostrichidae). *Mitteilungen des Internationalen Entomologischen Vereins*, Supplement 10: 1–100.

- Geis K-U (2012) Eine neozoische Splintholzkäferart unter Einfluß der Klimaerwärmung: Erster Nachweis mehrjähriger Überwinterungen von *Lyctus brunneus* (Steph.) in einem naturnahen Habitat Südwestdeutschlands (Coleoptera: Bostrichidae: Lyctinae). <http://kaeferklaus.de>
- Genduso P (1963) Terzo decennio di osservazioni entomologiche sui principali insetti nocivi della Sicilia. Bollettino dell'Istituto di Entomologia agraria e dell'Osservatorio di Fitopatologia di Palermo 4 (1960–1961–1962): 159–202.
- Gelosi A, Süss L (1991) Insetti e Acari dei cereali in magazzino. Edagricole, Bologna, 108 pp.
- Georghiou GP (1977) The Insects and Mites of Cyprus with emphasis on species of economic importance to agriculture, forestry, man and domestic animals. Kiphissia, Athens, 347 pp.
- Gerberg EJ (1957) A Revision of the New World Species of Powder-Post Beetles Belonging to the Family Lyctidae. Technical Bulletin. United States Department of Agriculture 1157: 1–55 + pls. I–XIV.
- Geri F, Amici V, Rocchini D (2010) Human activity impact on the heterogeneity of a Mediterranean landscape. Applied Geography 30 (2009): 370–379.
- Gerini V (1971) Contributo alla conoscenza della entomofauna nociva della Turchia. Rivista di agricoltura subtropicale e tropicale 65: 330–368.
- Ghizdavu I, Deac VA (1995) Investigations on the arthropod fauna, harmful to agricultural stored products, in the central area of the western plain of Romania. Buletinul Universității de Științe Cluj-Napoca. Seria Agricultura și Horticultură 48(2) (1994): 119–126.
- Giusti F, Manganelli G, Schembri PJ (1995) The non marine molluscs of the Maltese islands. Monografie - Museo Regionale di Scienze Naturali, Torino 15: 1–607.
- Glavendekic M, Mihajlovic L, Petanovic R (2005) Introduction and spread of invasive mites and insects in Serbia and Montenegro. Plant protection and plant health in Europe: introduction and spread of invasive species, held at Humboldt University, Germany, Berlin, 9–11 June 2005, 229–230.
- Gobbi G (1984) Appunti sui Coleotteri Cleroidei predatori di xilofagi (Coleoptera, Trogositidae, Cleridae, Melyridae). Bollettino dell'Associazione Romana di Entomologia 38(1/4) (1983): 51–62.
- Gobbi G (2003) Nuovi reperti di Coleotteri nel biotopo di Policoro (Matera) ed in altre aree protette della Lucania (Coleoptera). Annali del Museo civico di Storia naturale di Ferrara 5 (2002): 41–50.
- Goggi G (2004) Indagine faunistica sui Coleotteri delle Isole Pelagie (Sicilia). Giornale italiano di Entomologia 11: 127–143.
- Gridelli E (1930) Risultati zoologici della missione inviata dalla R. Società Geografica Italiana per l'esplorazione dell'Oasi di Giarabub (1926–1927). Coleotteri. Annali del Museo Civico di Storia Naturale Giacomo Doria 54: 1–480 + 1 map.
- Gridelli E (1939) Coleoptera Staphylinidae, Diversicornia, Heteromera, Lamellicornia, Chrysomelidae (Partim). Missione Biologica nel Paese dei Borana. Volume secondo Raccolte Zoologiche Parte prima. Reale Accademia d'Italia, Centro Studi per l'Africa Orientale Italiana, Roma 4: 311–615 + pls. VIII–XVI.
- Gridelli E (1940) Coleotteri dell'Africa Orientale Italiana. 11° Contributo. Materiali per lo studio della fauna eritrea raccolti nel 1901–03 dal Dott. Alfredo Andreini. Memorie della Società Entomologica Italiana 18(2) (1939): 219–258.

- Haack RA (2006) Exotic bark- and wood-boring Coleoptera in the United States: recent establishments and interceptions. Canadian Journal of Forest Research 36(2): 269–288. doi: 10.1139/x05-249
- Haack RA, Cavey JF (2000) Insects intercepted on solid wood packing materials at United States ports-of-entry: 1985–1998. In: Quarantine pests, risks for the forestry sector and their effects on foreign trade. Proceedings on CD-ROM of Silvotecna 14, 27–28 June 2000, Concepcion, Chile. CORMA, Concepcion, Chile, 16 pp.
- Haack RA, Slansky F (1987) Nutritional ecology of wood-feeding Coleoptera, Lepidoptera, and Hymenoptera. In: Slansky F, Rodriguez JG (Eds) Nutritional Ecology of Insects, Mites, Spiders, and Related Invertebrates. A Wiley-Interscience Publication, New York, NY, 449–486.
- Halperin J, Damoiseau R (1980) The bostrychid beetles (Coleoptera) of Israel. Israel Journal of Entomology 14: 47–53.
- Halperin J, Geis K-U (1999) Lyctidae (Coleoptera) of Israel, their damage and its prevention. Phytoparasitica 27(4): 257–262. doi: 10.1007/BF02981481
- Halstead DGH, Mifsud D (2003) Silvanidae and Laemophloeidae (Coleoptera: Cucujoidea) from the Maltese Islands (Central Mediterranean). The Central Mediterranean Naturalist 4(1): 41–46.
- Hamad NE-DF, Aly MZJ (1980) Seasonal fluctuations of certain species of nocturnal Coleoptera as indicated by a light trap at Quena. Qatar University Science Bulletin 5: 287–304.
- Hanna HM (1970) Studies on catches of Coleoptera in a light trap, at Assiut. Bulletin de la Société entomologique d'Egypte 53 (1969): 591–613.
- Haslam SM, Sell PD, Wolseley PA (1977) A flora of the Maltese Islands. Malta University Press, Msida, I–XXI + 560 pp.
- Háva J, Mifsud D (2006) The dermestid beetles (Coleoptera: Dermestidae) of the Maltese archipelago (Central Mediterranean). Studies and Reports of District Museum Prague-East, taxonomical Series 2: 51–63.
- Hellrigl K (2006) Über Auftreten von Holzinsekten in Häusern. Forest Observer 2/3: 333–348.
- Heyden (von) L, Reitter E, Weise J (Eds) (1891) Catalogus Coleopterorum Europae, Caucasi et Armeniae Rossicae. Berlin, Mödling, Paris, 420 pp.
- Hoffman WA (1933) *Rhizopertha dominica* F. as a Library Pest. Journal of Economic Entomology, 26: 293–294.
- Hopkins AD (1911) Appendix. Notes on habits and distribution, with list of described species. In: Kraus EJ (Ed.) Technical Papers on Miscellaneous Forest Insects. III. A revision of the powder-post beetles of the family Lyctidae of the United States and Europe. U.S. Department of Agriculture, Bureau of Entomology, Technical Series, 20(III): 130–138.
- Hoppe T (1981) Pests of stored products in Switzerland: abundance and insecticide resistance. Mitteilungen der Schweizerischen Entomologischen Gesellschaft 54(1/2): 3–13.
- Horion A (1961) Faunistik der Mitteleuropäischen Käfer. Band. VIII. Clavicornia. 2. Teil. (Thorictidae bis Cisidae) Teredilia, Coccinellidae. Überlingen – Bodensee, Kommissionsvergal Buchdruckerei Aug. Feyel, I–XV + 375 pp.
- Horion A (1972) Zwölfter Nachtrag zum Verzeichnis der mitteleuropäischen Käfer. Entomologische Blätter 68(1): 9–42.

- Howarth FG, Ramsay GW (1991) The conservation of island insects and their habitats. In: Collins NM, Thomas JA (Eds) The conservation of insects and their habitats. 15th Symposium of the Royal Entomological Society of London. Academic Press, London, 71–107.
- Humble LM (2010) Pest risk analysis and invasion pathways – insects. New Zealand Journal of Forestry Science (supplement) 40: 57–72.
- Hyde MB, Daubney CG (1960) A study of grain storage fossae in Malta. Tropical Science 2: 115–129.
- Iablokoff-Khnzorian SM (1976) Notes sur l’édéage des Coléoptères. Nouvelle Revue d’Entomologie 6(3): 229–238.
- Iablokoff-Khnzorian SM (1980) Le segment genital male des coléoptères et son importance phylogénétique (Coleoptera). Deutsche entomologische Zeitschrift 27(4): 251–295.
- ICZN (1985) International Code of Zoological Nomenclature. Third edition adopted by the XX general assembly of the International Union of Biological Sciences. International Trust for Zoological Nomenclature, London, I–XX + 338 pp.
- Inglebert H (2004) 1^{er} Supplément au Catalogue des Coléoptères de Paris Intra-Muros. L’Entomologiste 60 (5): 213–228.
- Ishaque M, Ali A, Qayyum HA (1982) Effect of some Fumigants and Contact Insecticides on Insect pests of Stored Wheat. Pakistan Entomologist 4(1/2): 1–6.
- Ivie MA (2002) 69. Bostrichidae Latreille 1802. In: Arnett RH Jr (†), Thomas MC, Skelley PE, Frank JH (Eds) American Beetles Polyphaga: Scarabaeoidea through Curculionoidea. CRC Press, Boca Raton, London, New York, Washington 2, 233–244.
- Ivie MA (2010) Additions and corrections to Borowski & Węgrzynowicz’s world catalogue of Bostrichidae (Coleoptera). Zootaxa 2498: 28–46.
- Ivinskis P, Rimšaitė J, Ostrauskas H, Talunytė L (2009) Alien Insects and Spiders Species and Species Spreading Naturally in Lithuania. In: Grasserbauer M, Sakalauskas L, Zavadskas EK (Eds) 5th International Vilnius Conference EURO Mini Conference “Knowledge-Based Technologies and OR Methodologies for Strategic Decisions of Sustainable Development” (KORSD-2009), September 30–October 3, 2009, Vilnius, Lithuania, 451–455.
- Iwata R, Nakano K (1982) Occurrence of *Sinoxylon conigerum* Gerstäcker (Coleoptera; Bostrichidae: Bostrychinae) on a picture frame manufactured in Java. The Society of House and household Pest Science, Japan 28(1): 81–83. [in Japanese]
- Iwata R, Nishimoto K (1981) Observations on the External Morphology and the Surface Structure of *Lyctus brunneus* (Stephens) (Coleoptera, Lyctidae) by Scanning Electron Microscopy. I. Larvae and Pupae. Kontyū 49(4): 542–557.
- Iwata R, Nishimoto K (1982) Observations on the External Morphology and the Surface Structure of *Lyctus brunneus* (Stephens) (Coleoptera, Lyctidae) by Scanning Electron Microscopy. II. Adult and Eggs. Kontyū 50(1): 10–22.
- Jabłoński T, Hilszczański J, Kolk A, Ślusarski S, Sukovata L (2007) Invasive forest insect species in Poland (Abstract). In: Evans H, Oszako T (Eds) Alien Invasive Species and International Trade. Forest Research Institute, Warsaw, 166.
- Jeannel R (1955) L’édéage. Initiation aux recherches sur la systématique des Coléoptères. Muséum National d’Histoire naturelle (Publications du Muséum N. 16), Paris, 155 pp.

- Jeannel R, Paulian R (1944) Morphologie abdominale des Coléoptères et systématiques de l'ordre. Revue française d'Entomologie 11: 65–110.
- Kahlen M, Hellrigl K (1996) Coleoptera - Käfer (Deck- oder Hartflügler). In: Hellrigl K (Ed) Die Tierwelt Südtirols. Veröffentlichungen des Naturmuseums Südtirol, Bozen 1: 393–511.
- Kalshoven LGE (1963a) Notes on the biology of Indonesian Bostrichidae (Coleoptera). Entomologische Berichten 23: 242–257.
- Kalshoven LGE (1963b) Ecological data on some neotropical Scolytidae, Platypodidae, and Bostrichidae (Coleoptera), mainly of Surinam. Beaufortia 9: 232–240.
- Kaszab Z (1959) Ergebnisse der zoologischen Aufsammlungen des Ungarischen Natur-wissenschaftlichen Museums in Ägypten im Jahre 1957. 5. Coleoptera. Annales Historico-Naturales Musei Nationalis Hungarici 59: 283–291.
- Kenis M (2005) 4 Insects – Insecta. In: Wittenberg R (Ed.) An inventory of alien species and their threat to biodiversity and economy in Switzerland. CABI Bioscience Switzerland Centre report to the Swiss Agency for Environment, Forests and Landscape, 131–212.
- Kenis M, Branco M (2010) Impact of alien terrestrial arthropods in Europe. Chapter 5. In: Roques A, Kenis M, Lees D, Lopez-Vaamonde C, Rabitsch W, Rasplus J-Y, Roy DB (Eds) Alien terrestrial arthropods of Europe. BioRisk 4(1): 51–71. doi: 10.3897/biorisk.4.42
- Khalilov BB (1972) Pests of grape vines in Azerbaijan. Zashchita Rastenii 17(2): 36–40.
- King GA (2009) The Alien Presence: Palaeoentomological Approaches to Trade and Migration. PhD in Archaeology University of York, York, 411 pp.
- Knopf HE (1972) Contributions to the Knowledge of the Insects Fauna of Trees of Iraq. Part I. Coleoptera. Zeitschrift für Angewandte Entomologie 69: 82–87. doi: 10.1111/j.1439-0418.1971.tb03186.x
- Kocher L (1956) Catalogue commenté des Coléoptères du Maroc, fascicule IV. Clavicornia et groupes voisins. Travaux de l'Institut Scientifique Chérifien 11: 1–136.
- Köhler F, Klausnitzer B (Eds) (1998) Verzeichnis der Käfer Deutschlands. Entomologische Nachrichten und Berichte (Dresden) Beiheft 4: 1–185.
- Krajewski A, Mazurek A (2009) Exotic species of wood borers in investigations of Wood Protection Division SGGW in Warsaw in years 1997. Annals of Warsaw University if Life Science – SGGW Forestry and Wood Technology 71: 395–399.
- Kraus EJ (1911) Technical Papers on Miscellaneous Forest Insects. III. A revision of the powder-post beetles of the family Lyctidae of the United States and Europe. U.S. Department of Agriculture, Bureau of Entomology, Technical Series 20(III): I–III + 111–129.
- Kucerová Z, Stesík V (2008) Differences in egg morphology of the stored-grain pests *Rhyzopertha dominica* and *Prostephanus truncatus* (Coleoptera: Bostrichidae). Journal of Stored Products Research 44(1): 103–105. doi: 10.1016/j.jspr.2007.06.005
- Lawrence JF (2010) 6.3. Bostrichidae Latreille, 1802. In: Leschen RAB, Beutel R, Lawrence JF (Eds) Coleoptera, Beetles. Volume 2: Morphology and Systematics (Elateroidea, Bostri-chiformia, Cucujiformia partim). Arthropoda Insecta. Handbook of Zoology. De Gruyter, Berlin & New York, 209–217.
- Lawrence JF, Slipinski A (2013) *Loranthophila*, a new genus of Australian Lyctinae (Coleoptera: Bostrichidae). Zootaxa 3737(3): 295–300. doi: 10.11646/zootaxa.3737.3.8

- Leal I, Allen E, Humble L, Sela S, Uzunovic A (2010) Phytosanitary Risks Associated with the Global Movement of Forest Products: A Commodity-based Approach. Pacific Forestry Centre III. Series: Information report (Pacific Forestry Centre) BC-X-419: I–IV + 42 pp.
- Lepesme P (1944) Encyclopédie entomologique, 22. Les Coléoptères des denrées alimentaires et des produits industriels entreposés. P. Lechevalier, Paris, 336 pp.
- Lesne P (1899) Revision des Coléoptères de la famille des Bostrichidae. 3^e Mémoire. Bostrichinae. Annales de la Société Entomologique de France 67: 438–621.
- Lesne P (1901a) Révision des Coléoptères de la famille des bostrychides. 4^e Mémoire. Bostrychinae sens. strict. – II. Les *Xylopertha*. Annales de la Société Entomologique de France 69 (1900): 473–639.
- Lesne P (1901b) Synopsis des Bostrychides paléartiques. L’Abeille 30 (1900–1906): 73–136.
- Lesne P (1904) Supplément au Synopsis des Bostrychides Paléartiques. L’Abeille 30 (1900–1906): 153–168 + pls. 1–4.
- Lesne P (1905) Notes additionnelles et rectificatives sur les Bostrychides paléartiques. L’Abeille 30 (1900–1906): 249–251.
- Lesne P (1906) Revision des Coléoptères de la famille des Bostrichidae. 5^e Memoire. Sinoxylinae. Annales de la Société Entomologique de France 75(4): 445–561.
- Lesne P (1909) Révision des Coléoptères de la famille des Bostrichidae. 6^e Mémoire : Diapatiinae et Apatinae. Annales de la Société entomologique de France 78: 471–574 + pls. 13–17.
- Lesne P (1924) Encyclopédie entomologique, 3. Les Coléoptères Bostrichides de l’Afrique tropicale française. Presses universitaires de France & P. Lechevalier, Paris, 301 pp.
- Lesne P (1932) Les formes d’adaptation au commensalisme chez les Lyctides. Livre du centenaire / Societe entomologique de France, Paris, 619–627.
- Lesne P (1935) Les Bostrichides de l’Arabie. Revue française d’Entomologie 1: 268–272.
- Lesne P (1938) Coleopterorum Catalogus, 10 pars 161 Bostrichidae. Auspicis et auxilio W. Junk, ed. S. Schenkling, s’Gravenhage, 84 pp.
- Lesne P (1945) Quelques remarques sur le *Rhizopertha dominica* F. [Col. Bostrichidae]. Revue française d’Entomologie 8(4): 145–151.
- Liberto A, Audisio P (2005) Sulla presenza in Sardegna di *Teloclerus compressicornis* (Coleoptera Cleridae). Bollettino della Società entomologica italiana 137(1): 45–48.
- Lillig M, Mifsud D, Grimm R (2012) Faunistic and taxonomic updates on the Tenebrionidae of Malta (Coleoptera). Bulletin of the Entomological Society of Malta 5: 111–119.
- Liu L-Y, Beaver RA, Yang J-T (2006) The Bostrichidae (Coleoptera) of Taiwan: a key to species, new records, and a lectotype designation for *Sinoxylon mangiferae* Chujo. Zootaxa 1307: 1–33.
- Liu L-Y, Schönitzer K (2011) Phylogenetic analysis of the family Bostrichidae auct. at supra-generic levels (Coleoptera: Bostrichidae). Mitteilungen der Münchner entomologischen Gesellschaft 101: 99–132.
- Liu L-Y, Schönitzer K, Yang J-T (2008) A review of the literature on the life history of Bostrichidae (Coleoptera). Mitteilungen der Münchner Entomologischen Gesellschaft 98: 91–97.
- Lodos N (1985) The first record of *Bostrychus capucinus* (L.) as an important pest of furnitures in Turkey (Coleoptera: Bostrichidae) [in Turkish]. Turkiye Bitki Koruma Dergisi 9(3): 183–188.
- López-Colón JI (2000) Los Bostrichidae Latreille, 1802 de la fauna de Marruecos (Coleoptera). Biocosme Mésogén 16(4) (1999): 171–221.

- López-Colón JI, Melic A, González Peña CF, Beltrán Valen JR, Blasco Zumeta J (2001) Insecta: Coleoptera (Familia 40): Bostrichidae Latreille, 1802. Catalogus de la Entomofauna Aragonesa 25: 15–28.
- Lucht WH (1987) Die Käfer Mitteleuropas. Katalog. Goecke & Evers, Krefeld, 342 pp.
- Luciano P (1982) Nuovi danni causati in Sardegna da *Apate monachus* Fabr. (Coleoptera Bostrichidae). Studi sassaresi. Sezione III: Annali della Facoltà di Agraria dell'Università di Sassari 29: 67–71.
- Luigioni P (1929) I Coleotteri d'Italia. Catalogo sinonimico-topografico-bibliografico. Memorie della Pontificia Accademia delle Scienze, I Nuovi Lincei (2) 13: [4 +] 1–1159 [+ 1].
- Lundberg S, Palm T, Trottestam O (1987) Skalbaggssstudier på Siciliens nordkust. II. Gräsmark, flodmynningar, havsstrand m m. Entomologisk Tidskrift 108: 123–129.
- Machado A, Oromí P (2000) Elenco de los Coleópteros de las Islas Canarias – Catalogue of the Coleoptera of the Canary Islands. Instituto de Estudios Canarios Monografía 70: 1–308.
- Maes JM (1995) Los Bostrichidae (Coleoptera) de Nicaragua. Revista Nicaraguense de Entomología 32: 5–17.
- Mantisi C (2001) Storia forestale dell'isola di Lampedusa con brevi note su Linosa e Lampione (Arcipelago delle Pelagie Provincia di Agrigento). Sicilia Foreste, Supplemento, 13: 1–120.
- Mart C, Erkiliç L, Bolu H, Uygun N, Altin M (1995) Species and pest control methods used in pistacchio orchards of Turkey. Acta Horticulturae (ISHS) 419: 379–386.
- Martínez de la Escalera M (1914) Los Coleópteros de Marruecos. Trabajos del Museo National de Ciencias Naturales, Serie Zoológica 11: 1–553.
- Mattson W, Vanhanen H, Veteli T, Sivonen S, Niemela P (2007) Few immigrant phytophagous insects on woody plants in Europe: legacy of the European crucible? Biological Invasions 9: 957–974. doi: 10.1007/s10530-007-9096-y
- Majka CG (2007) The Derodontidae, Dermestidae, Bostrichidae, and Anobiidae of the Maritime Provinces of Canada (Coleoptera: Bostrichiformia). Zootaxa 1573: 1–38.
- McCaffrey S (2011) Powderpost beetle (*Scobicia chevrieri*). Updated on 11/23/2011. Available online: PaDIL - <http://www.padil.gov.au>
- Mifsud D (2002) Longhorn Beetles (Coleoptera, Cerambycidae) of the Maltese Islands (Central Mediterranean). The Central Mediterranean Naturalist 3(4): 161–169.
- Mifsud D, Bílý S (2002) Jewel Beetles (Coleoptera, Buprestidae) from the Maltese Islands (Central Mediterranean). The Central Mediterranean Naturalist 3(4): 181–188.
- Mifsud D, Knížek M (2009) The Bark Beetles (Coleoptera: Scolytidae) of the Maltese Islands (Central Mediterranean). Bulletin of the Entomological Society of Malta 2: 25–52.
- Mifsud D, Falzon A, Malumphy C, De Lillo E, Porcelli F (2012) On some arthropods associated with *Ficus* spp. in the Maltese Islands. Bulletin of the Entomological Society of Malta 5: 5–34.
- Moleas T (1988) Osservazioni epidemiologiche su *Sinoxylon sexdentatum* Oliv. e *Amphicerus* (= *Schistoceros*) *bimaculatus* Oliv. (Coleoptera, Bostrichidae) sulla vite in Puglia. Informatore fitopatologico 38(1): 55–58.
- Mostafa SAS, Dabbour AI, Nassif MA, Aziz MIA (1981) Insects damaging stored products in Saudi Arabia. Anzeiger fur Schadlingskunde Pflanzenschutz Umweltschutz 54(12): 184–187. doi: 10.1007/BF01902237
- Mourikis PA, Tsourgianni A, Chitzanidis A (1998) Pistachio nut insect pests and means of control in Greece. Acta Horticulturae (ISHS) 470: 604–611.

- Müller G (1923) Materiali per una fauna coleotterologica delle isole e degli scogli dell'Adriatico. Liburnia, Rivista della Sezione di Fiume del Club Alpino Italiano 16: 3–30.
- Muscarella M, Sparacio I, Liberto A, Nardi G (2013) The genus *Lichenophanes* Lesne, 1899 in Italy (Coleoptera Bostrichidae) and short considerations on the saproxylophagous beetle-fauna of Nebrodi Mountains (Sicily). *Biodiversity Journal* 4(4): 451–466.
- Myers N, Mittermeier RA, Mittermeier CG, da Fonseca GAB, Kent J (2000) Biodiversity hot-spots for conservation priorities. *Nature* 403: 853–858. doi: 10.1038/35002501
- Nardi G (1997) Coleoptera Dermestidae, Lyctidae, Bostrichidae, Ptinidae. In: Zapparoli M (Ed.) *Gli Insetti di Roma*. Comune di Roma, Dipartimento X Area Risorsa Suolo e Tutela Ambientale, Quaderni dell'Ambiente 6: 177–179.
- Nardi G (2004a) Fauna Europaea: Lyctidae. In: Audisio P (Ed.) *Fauna Europaea: Coleoptera 2, Beetles*. Fauna Europaea version 1.0. <http://www.faunaeur.org> [accessed 12 July 2014 as version 2.6.2 of 29 August 2013]
- Nardi G (2004b) Fauna Europaea: Bostrichidae. In: Audisio P (Ed.) *Fauna Europaea: Coleoptera 2, Beetles*. Fauna Europaea version 1.0. <http://www.faunaeur.org> [accessed 12 July 2014 as version 2.6.2 of 29 August 2013]
- Nardi G, Mifsud D (2003) A review of the Anthicidae of the Maltese Islands (central Mediterranean) (Coleoptera). *Fragmenta entomologica* 35(2): 77–127.
- Nardi G, Ratti E (1995) Coleoptera Lyctidae, Bostrychidae e Anobiidae di Pantelleria e Lampedusa. In: Massa B (Ed.) *Arthropoda di Lampedusa, Linosa e Pantelleria (Canale di Sicilia, Mar Mediterraneo)*. Il Naturalista siciliano 19 (Supplemento): 477–492.
- Nardi G, Zahradník P (2004) Bostrichidae and Anobiidae (Coleoptera). In: Cerretti P, Hadersen S, Mason F, Nardi G, Tisato M, Zapparoli M (Eds) *Invertebrati di una foresta della Pianura Padana, Bosco della Fontana, Secondo contributo. Conservazione Habitat Invertebrati*, 3. Cierre Grafica Editore, Verona, 125–139.
- Nguyen DT (2006) Analysis of the behaviour of *Rhyzopertha dominica* (F.) (Coleoptera: Bostrichidae) towards host volatiles. Natural Resources Institute, University of Greenwich, I–XVI + 106 pp.
- Nicoli Aldini R (2003) Insetti delle derrate e dell'ambiente antropico di recente introduzione in Italia. In: Cravedi P (Ed.) *Atti del 7° Simposio "La difesa antiparassitaria nelle industrie alimentari e la protezione degli alimenti"*, Piacenza, 18–20 settembre 2002. Chiriotti Editori, Pinerolo, 118–143.
- Nicoli Aldini R (2004) Insetti delle derrate e dell'ambiente antropico di recente introduzione in Italia. *Tecnica Molitoria* 55(10): 977–999.
- Nieto A, Alexander KNA (2010) European Red List of Saproxylic Beetles. Publications Office of the European Union, Luxembourg, I–VIII + 45 pp.
- Niloufari P (1985) Studies on the infestation behaviour of the powder-post beetle *Lyctus brunneus* (Steph.) and its physical control in the wood yards of the Caspian forests of Iran. International Research Group on Wood Preservation, IRG/WP 1271.
- Norato G (1957) Notizie su due parassiti del grano di recente introduzione in Sicilia. *Bullettino dell'Istituto di Entomologia agraria e dell'Osservatorio di Fitopatologia di Palermo* 2 (1956–1957): 245–249.
- Normand H (1936) Contribution au catalogue des Coléoptères de la Tunisie (9^e fascicule). *Bulletin de la Société d'Histoire naturelle de l'Afrique du Nord* 27: 144–164.

- Novak P (1952) Kornjaši jadranskog primorja. (Coleoptera). Jugoslavenska Akademija Znanosti i Umjetnosti, Split, 524 pp.
- Olivier [AG] (1790) Encyclopédie méthodique ou par ordre de matières; par une société de gens de lettres, de savans et d'artistes; précédée d'un vocabulaire universel, servant de table pour tout l'ouvrage, ornée des portraits de Mm. Diderot et d'Alembert, premiers éditeurs de l'Encyclopédie. Histoire Naturelle. Insectes. Tome cinquième. C.-J. Panckoucke, Paris, 793 pp.
- O'Mahony E (1949) *Lyctus brunneus* Steph (Col., Lyctidae) and *Heterobostrichus* [sic!] *aqualis* Waterh. (Col., Bostrichidae) in Ireland. Entomologist's Monthly Magazine 85: 48.
- Oromí P, de la Cruz S, Báez M (2009) Coleoptera. In: Arechavaleta M, Rodríguez S, Zurita N, García A (Eds) Lista de especies silvestres de Canarias. Hongos, plantas y animales terrestres. Gobierno de Canarias, Litografía Gráficas Sabater, Tenerife, 254–301.
- Oromí P, Martín E, Zurita N, Cabrera A (2005) Coleoptera [Staphylinidae excepted]. In: Arechavaleta M, Zurita N, Marrero MC, Martín JL (Eds) Lista preliminar de especies silvestres de Cabo Verde (hongos, plantas y animales terrestres) Consejería de Medio Ambiente y Ordenación Territorial. Gobierno de Canarias, Santa Cruz de Tenerife, 78–86.
- Oromí P, Serrano A, Borges PAV (2010) Coleoptera [Coccinellidae, Elateridae, Latridiidae and Staphylinidae excepted]. In: Borges, PAV, Costa A, Cunha R, Gabriel R, Gonçalves V, Martins AF, Melo I, Parente M, Raposeiro P, Rodrigues P, Santos RS, Silva L, Vieira P, Vieira V (Eds) A list of the terrestrial and marine biota from the Azores. Princípia, Cascais, 222–232.
- Ovtchinnikov SV (1996) Bostrichidae. In: Tarbinsky YS (Ed.) Genetical Fund Cadastre of Kyrgyzstan. Volume III. Superclassis Hexapoda (Entognatha and Insecta). Bishkek, 137.
- Parmesan C (2006) Ecological and Evolutionary Responses to Recent Climate Change. Annual Review of Ecology, Evolution, and Systematics 37: 637–669. doi: 10.1146/annurev.ecolsys.37.091305.110100
- Pasta S, La Mantia T (2002) Il paesaggio vegetale e le sue modificazioni in età storica. In: Corti C, Lo Cascio P, Masseti M, Pasta S (Eds) Storia naturale delle Isole Pelagie. L'Epos, Palermo, 129–133.
- Paulian R (1988) Biologie des Coleopteres. Edition Lechevalier, Paris, I–XI + 719 pp.
- Peck SB, Heraty J, Landry B, Sinclair BJ (1998) The introduced insect fauna of an oceanic archipelago: The Galapagos Islands, Ecuador. American Entomologist 44: 218–237. doi: 10.1093/ae/44.4.218
- Peres Filho O, Teixeira ÉP, Becerra MLM, Dorval A, Berti Filho E (2006) First record of *Sinoxylon conigerum* Gerstäcker (Coleoptera: Bostrichidae) in Brazil. Neotropical Entomology 35(5): 712–713. doi: 10.1590/S1519-566X2006000500023
- Peres Filho O, Dorval A, Miranda de Souza Noquelli MJ (2007) Coleópteros em plantio de *Eucalyptus camaldulensis* no estado de Mato Grosso. Floresta e Ambiente 14(1): 45–51.
- Peris D, Delclòs X, Soriano C, Perrichot V (2014) The earliest occurrence and remarkable stasis of the family Bostrichidae (Coleoptera: Polyphaga) in Cretaceous Charentes amber. Palaeontologia Electronica 17(1): 14A, 8 pp. <http://palaeo-electronica.org/content/2014/706-new-cretaceous-auger-beetle>
- Peters BC, Creffield JW, Eldridge RH (2002) Lyctine (Coleoptera: Bostrichidae) pests of timber in Australia: A literature review and susceptibility testing protocol. Australian Forestry 65(2): 107–119. doi: 10.1080/00049158.2002.10674861

- Peretz I, Cohen M (1961) *Apate monachus* in Israel. FAO Plant Protection Bulletin 9: 76–79.
- Peyerimhoff P de (1919) Notes sur la biologie de quelques Coléoptères phytophages du Nord-Africain (troisième série) avec les descriptions de cinq espèces nouvelles et de sept sous-espèces ou variétés. Annales de la Société Entomologique de France 88: 169–258.
- Piras L, Pisano P (1972) Secondo contributo alla conoscenza faunistica della Sardegna: la costa del Sulcis (Sardegna sud-occidentale). Bollettino della Società sarda di Scienze naturali 11: 3–28.
- Pisano P, Viarengo M, Puddu F (2003) Animali di Sardegna. Gli insetti. Carlo Delfino Editore, Sassari, 368 pp.
- Podobivskii SS (1991) *Rhizopertha dominica* in the Ukraine. Zashchita Rastenii 12: 38–39.
- Poggi R, Brussino G, Scarpelli F (1994) Intercettazione in Piemonte di *Sinoxylon conigerum* Gerstäcker (Coleoptera: Bostrichidae). Atti XVII Congresso nazionale italiano di Entomologia, Udine, 13–18 giugno 1994, 217–219.
- Pollini A (1998) Manuale di entomologia applicata. Edagricole, Bologna, I–X + 1426 pp.
- Potter C (1935) The biology and distribution of *Rhizopertha dominica* (Fab.). Transactions of the Royal entomological Society of London 83: 449–482. doi: 10.1111/j.1365-2311.1935.tb02995.x
- Porta A (1929) Fauna Coleopterorum Italica. III. Diversicornia. Stabilimento Tipografico Piacentino, Piacenza, 466 pp.
- Porta A (1934) Fauna Coleopterorum Italica Supplementum. Stabilimento Tipografico Piacentino, Piacenza, I–VII + 208 pp.
- Porta A (1949) Fauna Coleopterorum Italica. Supplementum II. Stabilimento Tipografico Soc. An. G. Gandolfi, Sanremo, 386 pp.
- Porta A (1959) Fauna Coleopterorum Italica. Supplementum III. Stabilimento Tipografico Soc. An. G. Gandolfi, Sanremo, 345 pp.
- Price T, Brownell KA, Raines M, Smith CL, Gandhi KJK (2011) Multiple detections of two exotic auger beetles of the genus *Sinoxylon* (Coleoptera: Bostrichidae) in Georgia, USA. Florida Entomologist 94(2): 354–355. doi: 10.1653/024.094.0235
- Prota A (1963) Osservazioni su alcuni danni causati in Sardegna da *Apate monachus* Fabr. (Coleoptera Bostrichidae). Studi sassaresi. Sezione III: Annali della Facoltà di Agraria dell'Università di Sassari 11(3): 77–88.
- Quddus A, Qayyum HA (1982) Studies on the toxicity of some insecticides against *Rhizopertha dominica* (F.). Pakistan entomologist 4(1–2): 20–26.
- Querner P, Morelli M, Oberthaler E, Strolz M, Schmitz Von Ledebur K, Zatschek I, Femimebarek A, Diehl J, Hözl R, Engelhardt I, Krammer H, Fürnkranz S (2011) Ten years of Integrated Pest Management (IPM) at the Kunsthistorisches Museum in Wien. Journal of Entomological and Acarological Research Ser. II 43(2): 185–190.
- Ragusa E (1896) Catalogo ragionato dei Coleotteri di Sicilia. Il Naturalista siciliano (n.s.) 1(4–7): 69–106.
- Ragusa S, Russo A (1989) Gli artropodi dell'Annona in Calabria. L'informatore agrario 29: 71–74.
- Ratti E (2002) Biodiversità della laguna di Venezia e della costa nordadriatica veneta. Segnalazioni. 72 – *Heterobostrychus aequalis* (Waterhouse, 1884) (Insecta Coleoptera Bostrichidae). Bollettino del Museo civico di Storia naturale di Venezia 53: 284.

- Ratti E (2004) Coleoptera Lyctidae e Bostrichidae intercettati nel porto e negli ambienti urbani di Venezia. Bollettino del Museo civico di Storia naturale di Venezia 55: 121–125.
- Ratti E (2007) Coleotteri alieni in Italia/Alien Coleoptera in Italy. Vers. 2007-05-25. <http://www.visitmuve.it> [accessed 23 October 2013]
- Redolfi De Zan L, Bellotti F, D'Amato D, Carpaneto GM (2014) Saproxylic beetles in three relict beech forests of central Italy: Analysis of environmental parameters and implications for forest management. Forest Ecology and Management 328: 229–244. doi: 10.1016/j.foreco.2014.05.040
- Reichardt H (1964) Bostrichidae (Coleoptera) 5 Sobre a distribuição geográfica de duas espécies de *Apate* introduzidas no Brasil. Papéis avulsos do Departamento de Zoologia. Secretaria da Agricultura 16: 105–108.
- Reitter E (1883) Bostrychidae. In: Heyden (von) L, Reitter E, Weise J (Eds) Catalogus Coleopterorum Europae et Caucasi. Editio Tertia. London, Berlin, Paris, 124–125.
- Robinet C, Roques A (2010) Direct impacts of recent climate warming on insect populations. Integrative Zoology 5(2): 132–142. doi: 10.1111/j.1749-4877.2010.00196.x
- Robinson WH (2013) Urban Insects and Arachnids: A Handbook of Urban Entomology. Cambridge University Press, London, 480 pp.
- Roques A (2010) Alien forest insects in a warmer world and a globalised economy: impacts of changes in trade, tourism and climate on forest biosecurity. New Zealand Journal of Forestry Science, 40 suppl. (2010): S77–S94.
- Rostom ZMF (1993) Survey of some granivorous and non-granivorous insects and mites of stores in Saudi Arabia. Journal of Stored Products Research 29(1): 27–31. doi: 10.1016/0022-474X(93)90017-X
- Rungs C (1946) Sur les hôtes de quelques Coléoptères Bostrychidae observés au Maroc. L'Entomologiste 2(1): 7–11.
- Saccà G (1940) Contributo alla conoscenza dei coleotteri della fauna carticola. Bollettino del R. Istituto di Patologia del Libro 2(1): 1–8.
- Sadok BKM, Gerini V (1988) *Apate monachus*. F: coleoptera bostrychidae. Un insecte qui pourra devenir un fléau aux palmier dattiers dans le Oasis du Governorat de Kebili, en Tunisie. Rivista di Agricoltura subtropicale e tropicale 82: 371–377.
- Saliba LJ (1963) Insect pests of crop plants in the Maltese Islands. Department of Information, Malta, 35 pp.
- Sardar Alam M, Shahid Shaukat S, Ahmed M, Iqbal S, Ahmad A (1999) A Survey of Resistance to Phosphine in Some Coleopterous Pests of Stored Wheat and Rice Grain in Pakistan. Pakistan Journal of Biological Sciences 2(3): 623–626. doi: 10.3923/pjbs.1999.623.626
- Savoldelli S, Regalin R (2009) Infestation of wood pallets by *Sinoxylon unidentatum* (Fabricius) (Coleoptera Bostrichidae) in Italy. Bollettino di Zoologia agraria e di Bachicoltura, Serie II, 41(3): 235–238.
- Schabel HG (2006) Forest Entomology in East Africa. Forest Insects of Tanzania. Springer, I–IX + 328 pp.
- Schatzmayr A, Müller G (1925) Materiali per la fauna entomologica delle isole e degli scogli dell'Adriatico III. L'Isola e gli scogli di Unie. Fiume: rivista semestrale della Società di Studi fumiani in Fiume 31: 63–81.

- Schaufuss LW (1882) Zoologische Ergebnisse von Excursionen auf den Balearen. III. (Addenda und Fortsetzung). *Numquam otiosus* 3 (1879): 527–552.
- Schembri PJ (1993) Physical Geography and Ecology of the Maltese Islands: A Brief Overview. *Options Méditerranéennes*, Série B 7: 27–39.
- Schembri PJ (1997) The Maltese Islands: climate, vegetation and landscape. *GeoJournal* 41(2): 115–125. doi: 10.1023/A:1006828706452
- Schembri PJ (2003) Current state of knowledge of the Maltese non-marine fauna. Malta Environment and Planning Authority. Annual report and accounts 2003: 33–65.
- Schembri PJ, Lanfranco E (1996) Introduced species in the Maltese Islands. In: Baldacchino AE, Pizzuto A (Eds) Introduction of alien species of flora and fauna. Proceedings of a seminar held at Qawra, Malta, 5 March 1996. Environment Protection Department, Floriana, Malta, 29–54.
- Seabra AF de (1943) Contribuições para o Inventário da Fauna Lusitânica. Insecta Coleoptera. Memórias e Estudos do Museu Zoológico da Universidade de Coimbra, Série I 142: I–XX + 1–152.
- Serrano ARM (1981) Dados para a inventariação da fauna Lusitânica: Coleópteros novos para Portugal (Insecta, Coleoptera). *Boletim da Sociedade Portuguesa de Entomologia* 15: 1–7.
- Sharaf N, Akkawi M, Nazer I (1983) Preliminary survey of coleopteran fauna of Jordan. *Dirasat* 10(1): 57–65.
- Sittichaya W, Beaver RA, Liu L-Y, Ngampongsai A (2009) An illustrated key to powder post beetles (Coleoptera, Bostrichidae) associated with rubberwood in Thailand, with new records and a checklist of species found in Southern Thailand. *ZooKeys* 26: 33–51. doi: 10.3897/zookeys.26.88
- Song XH, Wang PP, Zhang HY (2011) Phosphine resistance in *Rhyzopertha dominica* (Fabricius) (Coleoptera: Bostrichidae) from different geographical populations in China. *African Journal of Biotechnology* 10(72): 16367–16373.
- Soro S (1964) Lo *Schistoceros bimaculatus* ed i suoi danni alla vite, in provincia di Sassari. *Informatore fitopatologico* 14: 351–352.
- Sparacio I (1997) Coleotteri di Sicilia. Parte seconda. L'Epos, Palermo, 208 pp.
- Stokland JN, Siitonens J, Jonsson BG (2012) Biodiversity in Dead Wood. Cambridge University Press, Cambridge, I–XII + 509 pp.
- Suma P, Russo A (2005) On the presence of *Prostephanus truncatus* (Horn) (Coleoptera Bostrichidae) in Italy. *Bollettino di Zoologia agraria e di Bachicoltura*, Serie II 37(2): 135–139.
- Surtees G (1961) Spermathecal structure in some Coleoptera associated with stored products. *Proceedings of the Royal entomological Society of London (A)* 36: 144–152. doi: 10.1111/j.1365-3032.1961.tb00260.x
- Tassi F (1967) Reperti 4. *Apate monachus* Fabr. (Col. Bostrichidae). *Bollettino dell'Associazione Romana di Entomologia* 22(3): 46.
- Taylor RWD, Halliday D (1986) The geographical spread of resistance to phosphine by coleopterous pests of stored products. 1986 British Crop Protection Conference. Pests and diseases. Volume 2. Proceedings of a conference held at Brighton Metropole, England, November 17–20, 1986, 607–613.

- Tezcan S (2008) Bati anadolu nar üretim alanlarında önem kazanan bir böcek türü: ikilekeli kubbekibocek (*Schistoceros bimaculatus*) (Coleoptera: Bostrichidae). hasad bitkisel üretim 24(277): 80–84.
- Thuiller W, Lavergne S, Roquet C, Boulangeat I, Lafourcade B, Araujo MB (2011) Consequences of climate change on the tree of life in Europe. *Nature* 470: 531–534. doi: 10.1038/nature09705
- Vigna Taglianti A, Audisio PA, Belfiore C, Biondi M, Bologna MA, Carpaneto GM, De Biase A, De Felici S, Piattella E, Racheli T, Zapparoli M, Zoia S (1993) Riflessioni di gruppo sui corotipi fondamentali della fauna W-paleartica ed in particolare italiana. *Biogeographia, Lavori della Società italiana di Biogeografia* 16 (1992): 159–179.
- Vigna Taglianti A, Audisio PA, Biondi M, Bologna MA, Carpaneto GM, De Biase A, Fattorini S, Piattella E, Sindaco R, Venchi A, Zapparoli M (1999) A proposal for a chorotype classification of the Near East fauna, in the framework of the Western Palearctic region. *Biogeographia, Lavori della Società italiana di Biogeografia* (n.s.) 20: 31–59.
- Viñolas A, Muñoz J, Mencuccini M, Benvenuti F (2013) Nuevos datos sobre *Rhusia parreyssi* (Mulsant, 1856), Melandrydae Leach, 1815 y otros coleópteros interesantes de la sierra de Prades, Tarragona (Coleoptera). *Orsis* 27: 29–51.
- Vrydagh JM (1952) Bostrichidae paléartiques: le genre *Scobicia* Lesne. *Bulletin et Annales de la Société Entomologique de Belgique* 88(III/IV): 58–59.
- Vrydagh JM (1955) Contribution à l'étude des Bostrichidae (Coleoptera Teredilia) 5. – Collection de la « California Academy of Sciences ». *Bullettin. Institut royal des Sciences naturelles de Belgique* 31(53): 1–15.
- Vrydagh JM (1956) Contribution à l'étude des Bostrichidae. 8 – Collection de la « Zoolo- gische Sammlung Bayerischen Staates » à Munich. *Bullettin. Institut royal des Sciences naturelles de Belgique* 32(6): 1–20.
- Vrydagh JM (1958) Contribution à l'étude des Bostrichidae. 16. – Collection du Musée G. Frey a Tutzing. *Entomologische Arbeiten aus dem Museum G. Frey Tutzing bei München* 9: 1068–1077.
- Vrydagh JM (1960a) Contribution à l'étude des Bostrichidae. 21. – Deuxième collection de l'Academie californienne des Sciences. *Bullettin. Institut royal des Sciences naturelles de Belgique* 36(14): 1–20.
- Vrydagh JM (1960b) Contribution à l'étude des Bostrichidae. 23. – Collection de la Section Zoologique du Musée Naturelle Hongrois à Budapest. *Bullettin. Institut royal des Sciences naturelles de Belgique* 36(39): 1–32.
- Vrydagh JM (1961) Contribution à l'étude des Bostrichidae. 27. – Collection du Musée d'Histoire naturelle Senckenberg à Francfort-sur-Main. *Bullettin. Institut royal des Sciences naturelles de Belgique* 37(4): 1–23.
- Vrydagh JM (1962) (†) Contribution à l'étude des Bostrichidae (Coleoptera). 31. Troisième collection du Musée zoologique de l'Université Humboldt à Berlin. *Bullettin. Institut royal des Sciences naturelles de Belgique* 38(4): 1–47.
- Wakil W, Ashfaq M, Shabbir A, Javed A, Sagheer M (2006) Efficacy of diatomaceous earth (protect-it) as a protectant of stored wheat against *Rhyzopertha dominica* (F.) (Coleoptera: Bostrichidae). *Pakistan Entomologist* 28(2): 19–24.

- Wakil W, Riasat T, Lord JC (2013) Effects of combined thiamethoxam and diatomaceous earth on mortality and progeny production of four Pakistani populations of *Rhyzopertha dominica* (Coleoptera: Bostrichidae) on wheat, rice and maize. Journal of Stored Products Research 52: 28–35. doi: 10.1016/j.jspr.2012.09.002
- Walther G-R, Roques A, Hulme Pe, Sykes Mt, Pyšek P, Kühn I, Zobel M, Bacher S, Botta-Dukát Z, Bugmann H, Czúcz B, Dauber J, Hickler T, Jarošík V, Kenis M, Klotz S, Minchin D, Moora M, Nentwig W, Ott J, Panov Ve, Reineking B, Robinet C, Semenchenko V, Solarz W, Thuiller W, Vilà M, Vohland K, Settele J (2009) Alien species in a warmer world: risks and opportunities. Trends in Ecology and Evolution 24(12): 686–693. doi: 10.1016/j.tree.2009.06.008
- Wang H, Varma RV, Xu T (1996) Insect Pests of Bamboos in Asia – An Illustrated Manual. International Network for Bamboo and Rattan (INBAR), Beijing, Eindhoven, New Delhi, 200 pp.
- Westcott RL, Labonte JR, Parsons GL, Johnson PJ (2006) New records and other notes for Oregon Coleoptera. Zootaxa 1142: 1–33.
- Winkler A (1927) Bostrychidae. In: Winkler A (Ed.) Catalogus Coleopterorum regionis palaeoarcticae. Pars 7. Albert Winkler Verlag, Wien, 795–798.
- Yan X, Zhou H, Shen Z, Li W, Guo D, Song Y, Lan S, Zhang J (2010) National investigations of stored grain arthropods in China. 10th International Working Conference on Stored Product Protection, 212–218.
- Yucel A (1988) Investigation on determining flour beetles and their damage in meal factories and mills in south-eastern Anatolia. Bitki Koruma Bulteni 28(1/2): 57–77.
- Zahradník P, Háva J (2014) Catalogue of the world genera and subgenera of the superfamilies Derodontoidae and Bostrichoidea (Coleoptera: Derodontiformia, Bostrichiformia). Zootaxa 3754(4): 301–352. doi: 10.11646/zootaxa.3754.4.1
- Zanardi D, Deligia S, Piras S (1969) Contributo alla conoscenza dell' *Apate monachus* Fabr. var. *rufiventris* Lucas della Sardegna e prime indicazioni di lotta. Giornate Fitopatologiche, Cagliari, 19–20 maggio, Atti, 215–226.
- Zanon V (1922) Contributo alla conoscenza della fauna entomologica di Bengasi. Coleotteri. Memorie della Società entomologica italiana 1(2): 112–139.
- Zapparoli M (2008) La componente alloctona nella entomofauna italiana: aspetti generali. Atti Accademia Nazionale Italiana di Entomologia 55 (2007): 97–101.
- Zavattari E (1934) Prodromo della fauna della Libia. Tipografia già Coperativa, Pavia, I–VIII + 1234 pp.
- Ziae M, Safaralizadeh MH, Shayesteh N (2006) Effects of temperature and exposure interval on the toxicity of Silico Sec® against two stored products insects. Pakistan Entomologist 28(1): 45–50.
- Zidan F, Obretenchev D (2009) Laboratory assessment of presence and level of resistance to the phosphine in populations of the lesser grain borer *Rhyzopertha dominica* F. Coleoptera, Bostrichidae in storages and silos in Bulgaria. Selskostopanska Nauka 42(6): 37–46. [in Bulgarian]
- Zocchi R (1971) Contributo alla conoscenza dell'entomofauna delle Tamerici in Italia. Redia 52 (1970–1971): 31–129.