



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.

Restriction of Measles Virus Gene Expression in Acute and Subacute Encephalitis of Lewis Rats

SIBYLLE SCHNEIDER-SCHAULIES,* UWE G. LIEBERT,* KNUT BACZKO,* ROBERTO CATTANEO,† MARTIN BILLETER,† AND VOLKER TER MEULEN*¹

**Institut für Virologie und Immunbiologie, Universität Würzburg, Versbacher Strasse 7.8700 Würzburg, Federal Republic of Germany, and*
†*Institut für Molekularbiologie I, Universität Zürich, Hänggerberg, 8006 Zürich, Switzerland*

Received February 15, 1989; accepted April 18, 1989

Measles virus (MV) replication in brain tissue of Lewis rats with acute (AE) and subacute (SAME) encephalitis was characterized by biochemical techniques. Messenger RNAs specific for measles virus nucleocapsid (N), phospho (P)-, matrix (M), fusion (F), and haemagglutinin (H) protein were detected in all brain extracts examined. The quantity of the individual MV mRNA species was quite different in comparison to lytically infected Vero cells. A steep gradient of MV transcripts was found in brain tissue which is most likely due to strongly attenuated transcription of mRNAs along the viral genome, representing particularly low transcription of the glycoprotein genes. In addition, *in vitro* translation assays only revealed synthesis of N and P protein in consistent fashion. The mRNAs for the glycoproteins did not direct the synthesis of detectable viral proteins whereas the M mRNA revealed some activity in animals with AE. The data indicate a strong restriction of the MV envelope gene expression in infected brain tissue, which is independent of the incubation time and type of the central nervous system (CNS) disease. This phenomenon which is similar to the findings observed in measles inclusion body encephalitis and subacute sclerosing panencephalitis suggest that host factors may initially be responsible for the initiation of transcriptional and translational alterations. © 1989 Academic Press, Inc.

INTRODUCTION

Measles virus (MV) is responsible for measles inclusion body encephalitis (MIBE) and subacute sclerosing panencephalitis (SSPE) in man (ter Meulen and Carter, 1982). Both diseases develop after long incubation periods on the basis of a persistent viral infection. To date the pathogenetic mechanisms leading to these CNS infections are not completely understood but studies of the viral gene expression have recently led to a better understanding of the state of measles virus persistence in brain cells of these two human CNS diseases. It has been found that MV persistence in brain tissue is characterized by a restriction of envelope viral gene expression implicating MV transcriptional and translational defects (Baczko *et al.*, 1986, 1988; Cattaneo *et al.*, 1986, 1987a, 1988; Liebert *et al.*, 1986). However, these observations do not help to explain which factors are involved in the establishment of persistence since the studies have been carried out on autopsy material. Therefore, it is not possible to determine which of the defects observed in MV replication in brain tissue of patients with MIBE or SSPE is a consequence of long-lasting virus host-cell interaction and which are the initial events determining persistence once MV has entered the brain.

To study these parameters we analyzed MV gene expression in a recently described animal model system

(Liebert and ter Meulen, 1987). Following intracerebral inoculation of a neurotropic rat brain adapted CAM-RBH strain of MV, Lewis rats develop different types of CNS diseases. Newborn animals invariably succumb to an acute encephalitis (AE) within a few days following infection. Weanling animals develop either an AE or a subacute encephalomyelitis (SAME) weeks or months after infection. Previous analyses of MV structural proteins in infected brain tissue by immunohistological staining procedures using monoclonal antibodies have suggested a restriction of MV envelope gene expression in all animals. Already in animals with AE, only one third of the infected brain cells expressed MV envelope proteins, and this was observed to decrease further to almost undetectable levels in brains of animals with SAME (Liebert and ter Meulen, 1987). We show in this study that the restriction of MV gene expression in infected brain tissue of Lewis rats results from transcriptional and translational alterations. The data obtained indicate that the defects observed in measles virus replication in acute or subacute measles encephalomyelitis in Lewis rats parallel many findings described in MIBE and SSPE.

MATERIALS AND METHODS

Infection of animals and tissue culture

Inoculation of the rat brain adapted CAM/R 40 strain of measles virus was performed as described pre-

¹ To whom requests for reprints should be addressed.

viously (Liebert and ter Meulen, 1987). Rats were infected in the left cerebral hemisphere with 25 μ l of CAM/RBH (4×10^5 TCID₅₀/ml) using a dispenser syringe. The animals were killed under ether narcosis. Brain material was obtained immediately after death and RNA extracted. Vero cells were infected with CAM/RBH virus at a m.o.i. of 0.1.

Immunohistology

Immunohistological studies on brain sections were carried out with a library of monoclonal antibodies directed against MV structural proteins as described previously (Liebert and ter Meulen, 1987).

Extraction of RNA

Total RNA from MV-infected Vero cells and from brain tissue was extracted as described previously (Chirgwin *et al.*, 1979; Baczko *et al.*, 1984). Cell homogenate from infected Vero cells was obtained by lysis in guanidinium isothiocyanate buffer when CPE was almost complete. Organ material was homogenized in guanidinium isothiocyanate buffer by using a Potter homogenizer. The homogenate was overlaid on a CsCl cushion (1.7 g/cm³) and centrifuged in an SW 28 rotor (Beckman) at 20° and 24,000 rpm overnight. The RNA pellet was dissolved in SEH buffer (100 mM NaCl, 10 mM Hepes, pH 7.0, 1 mM EDTA) containing 0.1% SDS and ethanol precipitated. The average yield was 600 μ g of total RNA/g tissue. Polyadenylated RNA was selected from total RNA by one cycle of oligo(dT)-cellulose chromatography (Sigma).

S1-Nuclease protection assay

Total RNA extracted from different rat organs (see Results) was screened for the presence of N-specific sequences by S1-analysis as described previously (Favaloro *et al.*, 1980; Billeter *et al.*, 1984). Briefly, single-stranded M13 hybridization probes covering the 3' end of the MV genome (*Pst*I/*Bam*HI fragment, nucleotides 0–230) (Billeter *et al.*, 1984) were labeled with [³²P]dATP by primer extension according to the conditions for sequence determination by chain termination (Davis *et al.*, 1986) and hybridized to 10 μ g of total RNA for 16 hr at 42° in a buffer containing 80% formamide. Nuclease S1 digestion was performed for 60 min at 30° with 700 units S1-nuclease/ml in the appropriate buffer. Following phenol extraction and ethanol precipitation, protected fragments were separated on an 8% polyacrylamide sequencing gel.

Northern blots

MV-specific RNAs were analyzed as described previously (Cattaneo *et al.*, 1987b). Briefly, 1 μ g of +pA-

RNA isolated from infected Vero cells or rat brain tissue was loaded on a 3-cm broad slot of a 1.5% formaldehyde-containing agarose gel together with a mixture of standard RNAs described below (usually 2 fmol each). After electrophoresis, the gel was blotted onto nitrocellulose which was cut in strips for hybridization. The strips were hybridized with [³²P]CTP-labeled RNA probes transcribed from MV-specific cDNA templates specific for all MV structural genes except L cloned in the pGem-1 vector. RNA probes used for all hybridization experiments were the precise complements of the external standard RNAs mentioned above and were transcribed from the following cDNA fragments: pGem N, 851-bp *Eco*RV/*Xba*I fragment; pGem P, 531-bp *Bal*HI/*Hind*III fragment; pGem M, 520-bp *Pst*I/*Sma*I fragment; pGem F, 320-bp *Taq*I/*Dde*I fragment; pGem H, 787-bp *Bgl*II/*Ava*I fragment (Cattaneo *et al.*, 1987b). Hybridization was performed at 54° in buffer containing 50% formamide. Following autoradiography, bands corresponding to the monocistronic MV mRNAs and the external standard RNAs were excised and the amount of radioactivity retained was determined by scintillation counting.

In situ hybridization

MV-specific RNA was localized in brain sections with ³⁵S-labeled DNA and RNA probes using modifications of the procedures described by Haase *et al.* (1984) and Cox *et al.* (1986). Paraffin sections (5–6 μ m thick, deparaffinized through xylene) or cryostat sections were rehydrated through graded ethanols, air-dried, immersed in 0.2 M HCl (20 min, room temperature), 2X SSC (30 min, 70°), 25 μ g/ml proteinase K (Calbiochem) in 2 mM CaCl₂, 10 mM Tris, pH 7.2 (15 min, 37°), distilled water (3 \times 5 min), freshly mixed 0.25% acetic anhydride in 0.1 M triethanolamine (10 min, room temperature), and finally dehydrated through graded ethanol and air-dried. ³⁵S-labeled probes were obtained either by nick-translation of the pGem N clone in the presence of [³⁵S]dATP (Amersham) or by *in vitro* transcription of the MV-specific pGem-1 clones in the presence of [³⁵S]UTP (Amersham) in anti-mRNA sense orientation using SP6- or T7-RNA polymerase, respectively (Boehringer). Specific activity for the strand-specific RNA probes was in the order of 5×10^8 and 5×10^7 dpm/ μ g for the nick-translated probe.

Labeled probes were applied to the sections at a concentration of 0.2 μ g/ml in a mixture containing 50% deionized formamide, 10% dextran sulfate, 0.3 M NaCl, 10 mM Tris, pH 7.2, 1 mM EDTA, 0.2% Ficoll, 0.2% polyvinylpyrrolidone, 0.2% bovine serum albumin, 200 μ g/ml +pA-RNA, 100 μ g/ml yeast tRNA, and 50 mM DTT. Siliconized coverslips were applied to the

sections and sealed with rubber cement. Hybridization was performed at 48° overnight for RNA probes and at room temperature for 48 hr for the DNA probe.

Following hybridization, slides were immersed in 2× SSC for 10 min at room temperature and afterward in hybridization wash mixture (HWM: 50% formamide, 0.6 M NaCl, 10 mM Tris-HCl, pH 7.4, 2 mM EDTA) (for RNA probes, 30 min at 45°, for the DNA probe, 30 min at 30°). Following two rinses in 0.1× SSC for 10 min, sections were dehydrated in graded ethanol containing 0.3 M ammonium acetate. Slides were air-dried and dipped in Kodak NTB2 emulsion diluted 1:1 with 0.6 M ammonium acetate at 43°. They were kept at 4° for 3 days, developed with Kodak D 19 developer, fixed with 30% sodium thiosulfate, and counterstained with hematoxylin. Controls included hybridization of MV-specific probes with brain sections of noninfected Lewis rats and labeled pGem-1 vector transcripts or a murine coronavirus JHM-specific transcript to brain sections from MV-infected Lewis rats. The murine coronavirus JHM probe was kindly provided by Dr. S. Siddell, Würzburg.

In vitro translation and immunoprecipitation

The ability of the different MV mRNAs obtained from the rat brains to direct the synthesis of the respective proteins was investigated using an *in vitro* translation system derived from rabbit reticulocyte lysate (Amersham) following a procedure described previously (Baczko *et al.*, 1984). Proteins made *in vitro* were immunoprecipitated using a polyclonal rabbit-anti-MV serum and monoclonal antibodies against MV structural proteins. Total translates and immunoprecipitated proteins were analyzed on a discontinuous 12% polyacrylamide gel as described (Carter *et al.*, 1982; Baczko *et al.*, 1984).

RESULTS

Detection of measles virus (MV) transcripts in different organs of infected Lewis rats

To determine if MV replication is restricted to brain cells following intracerebral inoculation of MV into weanling Lewis rats, RNA from different organs was extracted at various time intervals following infection (3–12 days postinfection). Total RNA obtained from tissue of two different brain regions and spinal cord as well as lung, liver, kidney, bone marrow, thymus, spleen, and peripheral blood mononuclear cells was screened for MV-specific transcripts by nuclease S1 analysis using ³²P-labeled M 13 probes derived of the 3' end of the N gene in both orientations. As summarized in Table 1, MV-specific transcripts were always detectable in

TABLE 1

EXPRESSION OF MV-SPECIFIC TRANSCRIPTS IN INFECTED LEWIS RATS

dpi ^a	Brain ^b	Other organs ^c
3	0/3 ^d	0/3
4–5	1/3	0/3
6–11	3/3	0/3
12	3/3	1/3 ^e

Note. Ten micrograms total RNA derived from different organs from weanling Lewis rats infected with CAM/RBH was analyzed for N-gene-specific sequences of either genomic or mRNA expression by S1-nuclease protection assay using [³²P]dATP-labeled M13 probes.

^a Samples from infected weanling Lewis rats from 3 to 12 days postinfection (dpi) were examined. The outcome of clinical disease was not considered.

^b Two different regions of the brains were tested.

^c Including lung, liver, kidney, thymus, bone marrow, spleen, peripheral blood mononuclear cells, and spinal cord.

^d Number of positive/number of tested animals.

^e One animal at dpi 12 was positive for bone marrow and thymic.

brain specimens as early as 4–5 days following infection but not in other tissues. In only one animal was a weak signal seen in total RNA derived from thymus and bone marrow.

Expression of MV-specific transcripts in brain material from CAM/RBH-infected Lewis rats

MV-specific transcription in brain tissue was analyzed by Northern blot using +pA-RNA fractions isolated from brain material of CAM-infected Lewis rats with different types of CNS diseases. For comparison, +pA-RNA from Vero cells infected with CAM/RBH virus was included in this study. Figure 1 shows the pattern of MV transcription in Vero cells (Fig. 1A), brain tissue of a newborn and a weanling Lewis rat with AE (Figs. 1B and C), and an animal with SAME (Fig. 1D). All MV structural genes investigated were expressed as monocistronic and bi- or polycistronic polyadenylated transcripts in all RNA specimens. However, the absolute amount and transcriptional efficiency of the individual MV-specific mRNAs in Vero cells were clearly different from those found in brain material. Similar to the results obtained for HeLa cells lytically infected with the Edmonston strain of MV (Cattaneo *et al.*, 1987b), we detected a polar expression gradient for CAM-infected Vero cells reflecting an almost linearly decreasing transcriptional attenuation of the structural genes along the gene order. In contrast, the gradient of mRNAs derived from brain was steeper and did not differ significantly between AE in newborn or weanling rats and in SAME (Figs. 1B–D). As in the lytic infection, the mRNAs for the first three genes, N, P, and M genes,




Fig. 1. Northern blot analysis of +pA-RNA derived from infected Vero cells (Fig. 1A), and from brain tissue of CAM/RBH-infected Lewis rats: a newborn animal 5 days postinfection (Fig. 1B), a weanling animal at Day 10 postinfection (Fig. 1C) both with AE, and an animal with SAME (Fig. 1D) at Day 30 postinfection. Isolated +pA samples were mixed with 2 fmol standard RNA transcripts (except Fig. 1C where there is only 1 fmol of the H standard that is hardly visible) and separated on a 1.5% formaldehyde-containing agarose gel. Filter strips were hybridized to ^{32}P -labeled strand-specific RNA probes and following hybridization were ordered according to the MV gene order. ▲, signals for the monocistronic; Δ, signals for the polycistronic RNAs; and □, signals for the standard RNA transcripts migrating corresponding to their individual lengths.

were efficiently transcribed as visualized by strong hybridization signals. However, transcription of the glycoprotein-specific mRNAs for the F and the H genes was significantly reduced in brain-derived RNAs. This was already evident in the acute form of infection (Figs. 1B and C) and was further enhanced in the subacute type of disease (Fig. 1D).

This typical expression pattern was observed in the large number of the infected animals we studied and was independent of the incubation period and type of CNS disease. In the case of AE, 20 newborn and 28 weanling Lewis rats were examined while for SAME 30 animals were studied.

Quantitation of the signals obtained for the monocistronic mRNAs revealed that the amount of N gene transcription proved to be about 5 to 7-fold lower in brain material as compared to lytically infected tissue culture (Table 2). The expression gradient, generally steeper for the brain-derived MV-specific mRNAs, showed the most obvious restriction occurring at the M–F gene

boundary. The values for the absolute expression/10 pg RNA varied slightly for the individual mRNAs (Table 2). With increasing age and incubation period the expression gradient of MV transcription became steeper, affecting in particular the envelope protein genes (Figs. 1C and D, Table 2). For negative control, MV-specific probes were hybridized to +pA-RNA from the brain of noninfected Lewis rats. No signal except the hybridization of the standard RNAs was observed (data not shown).

Analysis of the biological activity of the isolated +pA-RNAs from brain material

The finding of a transcriptional restriction of MV-genes, together with the observation of a reduced expression of envelope proteins in brain tissue of Lewis rats with AE and SAME (Liebert and ter Meulen, 1987), indicated that further characterization of the brain-derived mRNAs might be instructive. mRNAs from brain material of Lewis rats with AE and SAME were isolated, and their ability to direct the synthesis of the corresponding proteins was analyzed by *in vitro* translation (Fig. 2). Immunoprecipitation of MV-specific proteins synthesized *in vitro* from +pA-RNA derived from CAM-infected Vero cells revealed signals for the P, H, N, and M proteins (Figs. 2A and 2B, lane 2). +pA-RNAs derived from brain material from either newborn Lewis rats with AE (Fig. 2A, lanes 3 and 4) or weanling animals with AE (Fig. 2A, lane 5 and 6) which directed the synthesis of the N and P protein. However, the signals obtained for the P proteins were significantly reduced compared to the lytic control (Fig. 2A, lane 2). In lanes 3–6 of Fig. 2A, a band can be seen at the position of the M protein. +pA-RNAs derived from the brains of Lewis rats with SAME revealed only N and small amounts of P-specific translation products (Fig. 2B, lanes 4–7). In lanes 1 and 2 in Figs. 2A and 2B a faint band with an approximate molecular weight of 70 kDa which represents MV-specific H protein can be seen. This band was not detectable in the other lanes. In addition, in lanes 1 and 2 of Figs. 2A and 2B a protein of approximately 46 kDa is visible which could not be identified by the application of our monoclonal antibodies directed against MV structural proteins. F protein expression could not be examined since our monoclonal and monospecific polyclonal anti-F antibodies did not immunoprecipitate this protein from the *in vitro* synthesized products.

Sensitivity of the *in Vitro* translation assay

In order to exclude the possibility that the failure to detect M protein synthesis in the *in vitro* translation experiments using polyadenylated RNAs from brain of animals with SAME results from low sensitivity of the *in*

TABLE 2

QUANTITATIVE ANALYSIS OF MV-SPECIFIC GENE EXPRESSION IN BRAINS OF CAM-INFECTED LEWIS RATS

MV-specific mRNA	Vero/CAM		AE newborn rat		AE weanling rat		SAME weanling rat	
	Copies ^a	% ^b	Copies	%	Copies	%	Copies	%
N	21,400	100	5500	100	4300	100	4400	100
P	17,800	82	3300	59	2000	47	1900	43
M	14,200	67	2800	51	1400	33	1600	36
F	10,600	49	850	16	650	15	350	8
H	8,400	39	450	8	300	7	200	5

^a Copy numbers of the individual mRNA were determined as described (Cattaneo *et al.*, 1987b) and refer to 10 pg of RNA, the amount estimated per cell. Radioactivity bound to the nitrocellulose filters from Figs. 1A–D at the positions of the standard RNA transcripts and the corresponding monocistronic mRNA was measured by scintillation counting.

^b Relative expression of the MV structural genes was determined in relation to the individual N-gene expression.

in vitro translation system employed, +pA from lytically infected Vero cells was translated at different concentrations. The amount of RNA of infected cells was adjusted to the individual copy number calculated for the N-, M-, and H-specific mRNAs in the brain from animals with SAME. Figure 3 shows the results of the *in vitro* translation experiment using serially diluted +pA-RNA from infected cells (lanes 1–7) and from the brain of an animal with SAME (lane 8). The positions of the L-, P-, H-, N-, and M-specific translation products are indicated as well as the position of the 46-kDa protein (designated X) also seen in Figs. 2A and 2B. As an internal

control for equal amounts of +pA-RNA translated *in vitro* (1 µg each) a cellular protein band marked I is recognizable in each lane (Fig. 3A) with approximately the same intensity. In lane 3 of Fig. 3 the concentration of +pA-RNA represents about 5000 copies of the N-mRNA, in lane 4, about 1400 copies of the M-mRNA, and in lane 6, about 200 copies of the H-mRNA. In the total translate (Fig. 3A) all three viral proteins are detectable at the adjusted concentrations in infected Vero cell lysate as well as in the immune precipitate (Fig. 3B), except for the haemagglutinin which was not clearly precipitable as a distinct band. The intensity of the N-specific bands in lanes 3 and 8 was comparable. The M-specific protein, additionally identified by using a monospecific antiserum (data not shown), could be immunoprecipitated at the appropriate concentration (lane 4), but not visualized in the brain-derived RNA (lane 8).




Fig. 2. *In vitro* translation and immunoprecipitation of isolated +pA-RNA from CAM-infected tissue culture cells and brain material. 1 µg of +pA-RNA from lytically infected Vero cells and brain material from newborn and weanling Lewis rats with AE and animals with SAME was translated *in vitro* using a rabbit reticulocyte system. Total translation products were immunoprecipitated employing a polyclonal rabbit-anti-MV hyperimmune serum. (A) Lane 1, total translate of lytically infected Vero cells. Lanes 2–6, immunoprecipitation of total translate from: lane 2, lytically infected Vero cells; lanes 3 and 4, newborn Lewis rats with AE at 5 and 6 days postinfection (dpi); lanes 5 and 6, weanling Lewis rats with AE at dpi 6 and 8; lane 7, marker proteins. (B) Lanes 1 and 2 according to Fig. 2A, lanes 3–7 immunoprecipitation of total translate from: lane 3, an uninfected Lewis rat; lanes 4–7, animals with SAME at dpi 21, 23, 32, and 40; lane 8, marker proteins.

Expression of MV transcripts at the single cell level

The results of the Northern blot analysis and the *in vitro* translation studies indicated the development of transcriptional and translational alterations in the course of MV infection in brain tissue. In order to determine whether this phenomenon could also be demonstrated at the single cell level *in situ* hybridization on brain sections was carried out. For this purpose, we first determined the distribution and localization of MV-infected cells in the brain of an animal with SAME using a double-stranded N-gene-specific probe labeled by nick-translation with ³⁵S (Fig. 4). MV-specific signals occurred mainly in the gray matter of the brain, but were also found in the white matter.

To analyze the expression of the individual MV-specific mRNAs a brain area of an animal with SAME was chosen in which only N and P proteins but no M, F, and




FIG. 3. *In vitro* translation and immunoprecipitation of isolated +pA-RNA from CAM-infected and uninfected Vero cell culture and brain material from an animal with SAME. +pA-RNA from CAM-infected Vero cells was diluted to decreasing concentrations with +pA-RNA from uninfected Vero cells to a total amount of 1 μ g +pRNA and translated using a rabbit reticulo system. Total translation products were immunoprecipitated employing a polyclonal rabbit-anti-MV hyperimmune serum. (A) Total translation products of 1 μ g (lane 1), 0.5 μ g (lane 2), 250 ng (lane 3, about 5000 copies N-mRNA/10 pg RNA, 100 ng (lane 4 about 1400 copies M-mRNA/10 pg RNA), 50 ng (lane 5), 25 ng (lane 6, about 200 copies H-mRNA/10 pg RNA), and 10 ng (lane 7) +pA-RNA from infected cells, and 1 μ g of +pA from an animal with SAME (lane 8). Lane 9, ¹⁴C-labeled protein marker. (B) Immunoprecipitation of the total translation products from Fig. 3A, lanes 1-8 using a polyclonal rabbit-anti-MV hyperimmune serum. Lane 9, ¹⁴C-labeled protein marker.

H proteins were detectable by immunohistology. From this area, serial frozen sections were alternately stained for immunohistology or hybridized to the same

strand-specific RNA probes used for the Northern blot analyses showing the expression of the N- (Fig. 5A), P- (not shown); M- (Fig. 5B), F- (Fig. 5C), and H- (Fig. 5D)




FIG. 4. *In situ* hybridization of a paraffin-embedded 6- μ m-thick brain section of an animal with SAME using an N-gene specific, double-stranded probe labeled by nick-translation with [³⁵S]dATP. Tissue was counterstained with hematoxylin/eosin, magnification is 200 \times .




FIG. 5. *In situ* hybridization analysis of MV-specific transcripts in the brain of a Lewis rat with SAME. Serial frozen sections from the brain of a Lewis rat with SAME were hybridized to [³⁵S]UPT-labeled RNA probes as described. Following autoradiography, tissue was counterstained with hematoxylin/eosin. (A) N-mRNA, (B) M-mRNA, (C) F-mRNA, (D) H-mRNA. The areas shown in (A)–(D) are serial sections and are derived from the same brain area. Magnification is 400 ×.

mRNAs. About the same number of cells showed hybridization signals for the N-, P-, and M-mRNA with a gradual reduction in intensity from N to M. In contrast, F- and H-specific signals were detectable in a smaller number of cells. The presence of M-, F-, and H-specific mRNA in detectable amounts in the same brain area where no expression of the corresponding proteins were seen could either reflect the higher sensitivity of the RNA detection method or suggest the existence of translational defects for the F- and H-specific mRNAs. A similar hybridization pattern was obtained with material from newborn and weanling Lewis rats with AE (data not shown), in which, in contrast to SAME, 20–30% of infected brain cells expressed MV envelope proteins (Liebert and ter Meulen, 1987).

DISCUSSION

In the present study we analyzed MV gene expression at the transcriptional and translational level in brain material derived from infected Lewis rats showing different types of CNS diseases. The transcription of the glycoprotein-specific mRNAs was generally restricted, the degree of transcriptional restriction being related to the age of the animals and the type of dis-

ease. In addition, the synthesis of the viral M proteins and probably also the H proteins was shown to be inhibited, due to a failure in translation of the corresponding mRNA in Lewis rats with SAME. Thus, data obtained at the molecular level support the *in vivo* observation of a restricted expression of the M, F, and H proteins in the brains of infected rats (Liebert and ter Meulen, 1987).

Transcriptional restriction of the glycoprotein-specific mRNAs could be seen as soon as MV-specific mRNA was detectable in infected brains. This phenomenon was independent both of the incubation time after infection and the humoral immune response of the animals, since measles antibodies in serum and CSF could only be detected in diseased weanling rats (Dörries *et al.*, 1988). The fact that a restriction of MV transcription has already occurred in newborn rats a few days after infection suggests that brain-specific host cell factors may primarily interfere with MV gene expression. At present we cannot determine whether brain cells lack factors necessary for efficient MV transcription or contain factors actively interfering with MV gene expression leading to a premature termination of MV transcripts.

Interestingly, restriction occurred predominantly at the M-F gene boundary located within the only long noncoding region of the MV genome spanning about 1000 nucleotides. This genomic region is indeed peculiar, since three open reading frames have been defined, but no translation products of these reading frames have been detected *in vivo* or *in vitro* (Bellini *et al.*, 1986; Wong *et al.*, 1987; Greer *et al.*, 1987). Moreover, no preferential accumulation of mutations has occurred in this genomic region comparing the sequences of lytic MV strains (Cattaneo *et al.*, 1988), in spite of the high variability of viral RNA genomes (Steinhauer and Holland, 1987; Domingo and Holland, 1987). Furthermore, noncoding sequences at the 3' end of the M mRNA have also been defined for other paramyxoviruses (Bellini *et al.*, 1986; Blumberg *et al.*, 1984; Hidaka *et al.*, 1984). Although there is only chance homology between those noncoding regions within the different viruses, the analogy in location and the conservation of these particular sequences could argue for a regulatory function in the expression of viral genes. Interestingly, the 5' untranslated region of the MV F gene has been shown to contain an unusually high content of cytosine residues (44%) (Buckland *et al.*, 1987) with an overall GC content of 64%. This may not only influence translational efficiency of the corresponding mRNA as has been shown for cellular genes (Takeishi *et al.*, 1985; Ordahl *et al.*, 1984; Sap *et al.*, 1986) but also transcriptional regulation. The viral polymerase complex could slowdown or stall along this region which could possibly favor termination. Alternatively, the restriction of MV transcription could possibly result from a selective instability of the glycoprotein-specific mRNAs in brain tissue. This has been described in other viral and cellular systems in which instability of viral and cellular transcripts occur, probably due to the activity of endogenous RNAses (for review, see Brawerman, 1987; Pontecorvi *et al.*, 1988).

The finding that during AE in newborn Lewis rats the restriction observed is less pronounced than in weanling animals with AE may support the hypothesis that cellular differentiation of the developing brain could influence the restriction of viral gene expression. It is known from tissue culture experiments that viral replication is down-regulated with increased differentiation of the infected cell (Robbins and Rapp, 1980; Miller and Carrigan, 1982; Yoshikawa and Yamanouchi, 1984). The level of endogenous signal transmitters (cAMP and cGMP) and the mitotic activity (Joseph *et al.*, 1975; Lucas *et al.*, 1978; McChesney *et al.*, 1987) have been shown to influence MV replication in different tissue culture systems especially of neuronal and lymphoid origin. Similar phenomena may occur in our animal model and may interfere with the replication of the measles virus.

Besides the transcriptional alterations, translation of the envelope proteins has been shown to be restricted. This could be documented for the M-specific mRNA by dilution experiments with RNA from infected cells (Fig. 3) and at the single cell level for the F and H protein. The basis underlying these restrictions probably resides in the mRNAs themselves because they occur not only *in vivo* but also in *in vitro* translation experiments. Cap-dependent translation alterations have been shown for VSV methylation mutants (Horikami and Moyer, 1982), but mutational events should also be considered. Although the time course for mutations seems to be very short one should keep in mind that RNA-editing mechanisms have been shown recently for the P genes of negative strand RNA viruses (Thomas *et al.*, 1988; Cattaneo *et al.*, 1989).

The observed alterations of MV transcription and translation in infected brain tissue of rats are very similar to the changes seen in MV persistence in human CNS diseases (Baczko *et al.*, 1986; Cattaneo *et al.*, 1987a). In MIBE as well as in SSPE a restriction at a transcriptional level is found, affecting mainly MV M, F, and H genes. This leads to a rather steep gradient of mRNA transcripts according to the location of the individual genes in the MV genome (Cattaneo *et al.*, 1987b). In *in vitro* translation experiments using isolated +pA RNA from human brain tissue only MV N and P proteins were always detectable whereas the viral envelope proteins were either absent or only occasionally immunoprecipitated (Baczko *et al.*, 1986). These *in vitro* findings were supported by immunohistological investigations (Liebert *et al.*, 1986). Alterations in translation of MV genes caused by mutations have been observed in some cases of SSPE and in MIBE (Cattaneo *et al.*, 1988). In one case of MIBE a mutation caused elimination of the initiation codon of the M gene (Cattaneo *et al.*, 1988), and in one SSPE case a mutation introduced a stop codon in the M gene sequence (Cattaneo *et al.*, 1986, 1988). However, mechanisms other than mutations could be implicated in alteration of translation of MV protein (Ogura *et al.*, 1987, 1988).

The observation of restricted MV transcription and translation in infected rat brain a few days after infection demonstrates that this phenomenon is independent of long incubation periods. One can assume that similar events may have taken place in SSPE or MIBE a long time before the onset of disease. By this mechanism, which appears to be host-cell dependent, infected brain cells may not be destroyed by virus replication. In the absence of F protein, no cell fusion occurs and the defect of M protein synthesis blocks the maturation of complete virus particles thus keeping the infection strictly cell-associated. Moreover, the overall restriction of the expression of MV surface glycoproteins saves the infected cells from being recognized

and destroyed by antiviral antibodies. These events probably prevent a lytic infection of MV and allow the establishment of persistence. Since the ribonucleo-capsid-complex is infectious (Rozenblatt *et al.*, 1979), the infection is probably maintained by spread via cell processes. The pathogenetic role played by the observed mutations in the two human diseases is not understood. It is conceivable that mutations amplify the restriction of gene expression and support persistence rather than initiate MV gene restriction.

ACKNOWLEDGMENTS

We thank the Deutsche Forschungsgemeinschaft and Schweizerischer Nationalfonds for financial support, Dr. B. K. Rima for helpful discussions, J. Roller for technical assistance, Fritz Ochsenbein for preparing the photographs, and H. Kriesinger for typing the manuscript.

REFERENCES

- BACZKO, K., CARTER, M. J., BILLETER, M., and TER MEULEN, V. (1984). Measles virus gene expression in subacute sclerosing panencephalitis. *Virus Res.* **1**, 585–595.
- BACZKO, K., LIEBERT, U. G., BILLETER, M., CATTANEO, R., BUDKA, H., and TER MEULEN, V. (1986). Expression of defective measles virus genes in brain tissues of patients with subacute sclerosing panencephalitis. *J. Virol.* **59**, 472–478.
- BACZKO, K., LIEBERT, U. G., CATTANEO, R., BILLETER, M. A., ROOS, R. P., and TER MEULEN, V. (1988). Restriction of measles virus gene expression in measles inclusion body encephalitis. *J. Infect. Dis.* **158**, 144–150.
- BELLINI, W. J., ENGLUND, G., RICHARDSON, C. D., ROZENBLATT, S., and LAZZARINI, R. A. (1986). Matrix genes of measles virus and canine distemper virus: cloning, nucleotide sequences, and deduced amino acid sequences. *J. Virol.* **58**, 408–416.
- BILLETER, M., BACZKO, K., SCHMID, A., and TER MEULEN, V. (1984). Cloning of DNA Corresponding to four different measles virus genomic regions. *Virology* **132**, 149–159.
- BLUMBERG, B. M., ROSE, K., SIMONA, M. G., ROUX, L., GIORGI, C., and KOLAKOFSKY, D. (1984). Analysis of the Sendai virus M gene and protein. *J. Virol.* **52**, 656–663.
- BRAWERMAN, G. (1987). Determinants of Messenger RNA Stability. *Cell* **48**, 5–6.
- BUCKLAND, R., GERALD, C., BARKER, R., and WILD, T. F. (1987). Fusion glycoprotein of measles virus: Nucleotide sequence of the gene and comparison with other paramyxoviruses. *J. Gen. Virol.* **68**, 1695–1703.
- CARTER, M. J., WILLCOCKS, M. M., LÖFFLER, S., and TER MEULEN, V. (1982). Relationships between monoclonal antibody binding sites on the measles virus haemagglutinin. *J. Gen. Virol.* **63**, 113–120.
- CATTANEO, R., KAELIN, K., BACZKO, K., and BILLETER, M. A. (1989). Measles virus editing provides an additional cystein-rich protein. *Cell* **56**, 759–764.
- CATTANEO, R., REBMANN, G., BACZKO, K., TER MEULEN, V., and BILLETER, M. A. (1987). Altered ratios of measles virus transcripts in diseased human brains. *Virology* **160**, 523–526.
- CATTANEO, R., REBMANN, G., SCHMID, A., BACZKO, K., TER MEULEN, V., and BILLETER, M. A. (1987). Altered transcription of a defective measles virus genome derived from diseased human brain. *EMBO J.* **6**, 681–687.
- CATTANEO, R., SCHMID, A., ESCHLE, D., BACZKO, K., TER MEULEN, V., and BILLETER, M. A. (1988). Biased hypermutation and other genetic changes in defective measles viruses in human brain infections. *Cell* **55**, 255–265.
- CATTANEO, R., SCHMID, A., REBMANN, G., BACZKO, K., TER MEULEN, V., BELLINI, W. J., ROZENBLATT, S., and BILLETER, M. A. (1986). Accumulated measles virus mutations in a case of subacute sclerosing panencephalitis: Interrupted matrix proteins reading frame and transcription alteration. *Virology* **154**, 97–107.
- CHIRGWIN, J. M., PRZYBYLA, A. E., MACDONALD, R. J., and RUTTER, W. J. (1979). Isolation of biologically active ribonucleic acids from sources enriched in ribonuclease. *Biochemistry* **18**, 5294–5299.
- COX, K. H., DELEON, D. V., ANGERER, L. M., and ANGERER, R. C. (1984). Detection of mRNAs in sea urchin embryos by in situ hybridization using asymmetric RNA probes. *Dev. Biol.* **101**, 485–502.
- DAVIS, L. G., DIBNER, M. D., and BATTEY, J. R. (1986). "Basic Methods in Molecular Biology." Elsevier, New York/Amsterdam/London.
- DOMINGO, E., and HOLLAND, J. J. (1987). High error rates, population equilibrium and evolution of RNA replication systems. In "RNA Genetics" (E. Domingo, P. Ahlquist, and J. J. Holland, Eds.). CRC Press, Boca Raton, FL.
- DÖRRIES, R., LIEBERT, U. G., and TER MEULEN, V. (1988). Comparative analysis of virus-specific antibodies and immunoglobulins in serum and cerebrospinal fluid of subacute measles virus induced encephalomyelitis (SAME) in rats and subacute sclerosing panencephalitis (SSPE). *J. Neuroimmunol.* **19**, 339–352.
- FAVALORO, J. R., TREISMAN, R., and KAMEN, R. (1980). Transcription maps of polyoma virus-specific RNA: Analysis by two-dimensional nuclease S1 gel mapping. In "Methods in Enzymology" (L. Grossman and K. Maldave, Eds.), Vol. 65, pp. 718–749. Academic Press, New York.
- GREER, P. A., HASEL, K. W., and MILLWARD, S. (1986). Cloning and in vitro expression of the measles virus matrix gene. *Biochem. Cell Biol.* **64**, 1038–1043.
- HAASE, A. T., BRAHIC, M., STOWING, L., and BEUTH, H. (1984). Detection of viral nucleic acids by in situ hybridization. "Methods in Virology." Vol. 7, pp. 189–226. Academic Press, San Diego.
- HIDAKA, Y., KANDA, Y., IWASAKI, K., NOMOTO, A., SHIODA, T., and SHIBUTA, H. (1984). Nucleotide sequence of a Sendai virus genome region covering the entire M gene and the 3' proximal 1013 nucleotides of the F gene. *Nucleic Acids Res.* **12**, 7965–7973.
- HORIKAMI, S. M., and MOYER, S. A. (1982). Host range mutants of VSV defective in in vitro RNA methylation. *Proc. Natl. Acad. Sci. USA* **79**, 7694–7698.
- JOSEPH, B. S., LAMPERT, P. W., and OLDSTONE, M. B. A. (1975). Replication and persistence of measles virus in defined subpopulations of human leukocytes. *J. Virol.* **16**, 1638–1649.
- LIEBERT, U. G., BACZKO, K., BUDKA, H., and TER MEULEN, V. (1986). Restricted expression of measles virus proteins in brains from cases of subacute sclerosing panencephalitis. *J. Gen. Virol.* **67**, 2435–2444.
- LIEBERT, U. G., and TER MEULEN, V. (1987). Virological aspects of measles virus induced encephalomyelitis in Lewis and BN rats. *J. Gen. Virol.* **68**, 1715–1722.
- LUCAS, C. J., UBELS-POSTMA, J. C., REZEE, A., and GALAMA, J. M. D. (1978). Activation of measles virus from silently infected human lymphocytes. *J. Exp. Med.* **148**, 940–952.
- MCCHESNEY, M. B., KEHRL, J. H., VALSAMAKIS, A., FAUCI, A. S., and OLDSTONE, M. B. A. (1987). Measles virus infection of B lymphocytes permits cellular activation but blocks progression through the cell cycle. *J. Virol.* **61**, 3441–3447.
- TER MEULEN, V., and CARTER, M. J. (1982). Morbillivirus persistent infections in animals and man. In "Virus Persistence" (B. W. J. Mahy, A. C. Minson, and G. K. Darby, Eds.), pp. 97–132. Cambridge Univ. Press, London/New York.

- MILLER, C. A., and CARRIGAN, D. R. (1982). Reversible repression and activation of measles virus infection in neural cells. *Proc. Natl. Acad. Sci. USA* **79**, 1629–1633.
- OGURA, H., BACZKO, K., RIMA, B. K., and TER MEULEN, V. (1987). Selective inhibition of translation of the mRNA coding for the measles virus membrane protein at elevated temperatures. *J. Virol.* **61**, 472–479.
- OGURA, H., RIMA, B. K., and TER MEULEN, V. (1988). Restricted synthesis of the fusion protein of measles virus at elevated temperatures. *J. Gen. Virol.* **69**, 925–929.
- ORDAHL, C. P., EVANS, G. L., COOPER, T. A., KUNZ, G., and PERRIARD, J. C. (1984). Complete cDNA-derived amino acid sequence of chick muscle creatine kinase. *J. Biol. Chem.* **259**, 15,224–15,227.
- PONTECORVI, A., TATA, J. R., PHYLLAIER, M., and ROBBINS, J. (1988). Selective degradation of mRNA: The role of short-lived proteins in differential destabilization of insulin-induced creatine phosphokinase and myosin heavy chain mRNAs during rat skeletal muscle L6 cell differentiation. *EMBO J.* **7**(5), 1489–1495.
- ROBBINS, S. J., and RAPP, F. (1980). Inhibition of measles virus replication by cAMP. *Virology.* **106**, 317–326.
- ROZENBLATT, S., KOCH, T., PINHASI, O., and BRATASIN, S. (1979). Infective substructures of measles virus from acutely and persistently infected cells. *J. Virol.* **32**, 329–333.
- SAP, J., MUNOZ, A., DAMM, K., GOLDBERG, Y., GHYSDAEL, J., LEUTZ, A., BENG, H., and VENNSTROM, B. (1986). The c-erb-A protein is a high-affinity receptor for thyroid hormone. *Nature (London)* **324**, 635–640.
- STEINHAUER, D. A., and HOLLAND, J. J. (1987). Rapid evolution of RNA viruses. *Annu. Rev. Microbiol.* **41**, 409–433.
- TAKEISHI, K., KANEDA, S., AYSUSAWA, D., SHIMIZU, K., GOTOH, O., and SENO, T. (1985). Nucleotide sequence of a functional cDNA for human thymidylate synthase. *Nucleic Acids Res.* **13**, 2035–2043.
- THOMAS, S. M., LAMB, R. A., and PATERSON, R. G. (1988). Two mRNAs that differ by two nontemplated nucleotides encode the amino co-terminal proteins P and V of the paramyxovirus SV5. *Cell* **54**, 891–902.
- WONG, T. C., WIPF, G., and HIRANO, A. (1987). The measles virus matrix gene and gene product defined by *in vitro* and *in vivo* expression. *Virology* **157**, 497–508.
- YOSHIKAWA, Y., and YAMANOUCHI, K. (1984). Effects of papaverine treatment on replication of measles virus in human neural and non-neural cells. *J. Virol.* **50**, 489–496.