




Ca²⁺/Calmodulin-Dependent Protein Kinase II (CaMKII) β -Dependent Phosphorylation of GABA_{B1} Triggers Lysosomal Degradation of GABA_B Receptors via Mind Bomb-2 (MIB2)-Mediated Lys-63-Linked Ubiquitination

Khaled Zemoura^{1,2} · Karthik Balakrishnan^{1,3} · Thomas Grampp¹ · Dietmar Benke^{1,3,4} 

Received: 18 January 2018 / Accepted: 22 May 2018 / Published online: 7 June 2018
© The Author(s) 2018

Abstract

The G protein-coupled GABA_B receptors, constituted from GABA_{B1} and GABA_{B2} subunits, are important regulators of neuronal excitability by mediating long-lasting inhibition. One factor that determines receptor availability and thereby the strength of inhibition is regulated protein degradation. GABA_B receptors are constitutively internalized from the plasma membrane and are either recycled to the cell surface or degraded in lysosomes. Lys-63-linked ubiquitination mediated by the E3 ligase Mind bomb-2 (MIB2) is the signal that sorts GABA_B receptors to lysosomes. However, it is unknown how Lys-63-linked ubiquitination and thereby lysosomal degradation of the receptors is regulated. Here, we show that Ca²⁺/calmodulin-dependent protein kinase II (CaMKII) promotes MIB2-mediated Lys-63-linked ubiquitination of GABA_B receptors. We found that inhibition of CaMKII in cultured rat cortical neurons increased cell surface GABA_B receptors, whereas overexpression of CaMKII β , but not CaMKII α , decreased receptor levels. This effect was conveyed by Lys-63-linked ubiquitination of GABA_{B1} at multiple sites mediated by the E3 ligase MIB2. Inactivation of the CaMKII phosphorylation site on GABA_{B1}(Ser-867) strongly reduced Lys-63-linked ubiquitination of GABA_B receptors and increased their cell surface expression, whereas the phosphomimetic mutant GABA_{B1}(S867D) exhibited strongly increased Lys-63-linked ubiquitination and reduced cell surface expression. Finally, triggering lysosomal degradation of GABA_B receptors by sustained activation of glutamate receptors, a condition occurring in brain ischemia, was accompanied with a massive increase of GABA_{B1}(Ser-867) phosphorylation-dependent Lys-63-linked ubiquitination of GABA_B receptors. These findings indicate that CaMKII β -dependent Lys-63-linked ubiquitination of GABA_{B1} at multiple sites controls sorting of GABA_B receptors to lysosomes for degradation under physiological and pathological condition.

Keywords GABA receptor · CaMKII · Lysosome · Protein degradation · Ubiquitination · E3 ubiquitin ligase

Khaled Zemoura and Karthik Balakrishnan contributed equally to this work.

✉ Dietmar Benke
benke@pharma.uzh.ch

- ¹ Institute of Pharmacology and Toxicology, University of Zurich, Winterthurerstrasse 190, 8057 Zurich, Switzerland
- ² Present address: Picower Institute for Learning and Memory, Massachusetts Institute of Technology, 77 Massachusetts Avenue, Cambridge, MA 02139-4307, USA
- ³ Neuroscience Center Zurich, University of Zurich and ETH Zurich, Winterthurerstrasse 190, CH-8057 Zurich, Switzerland
- ⁴ Drug Discovery Network Zurich (DDNZ), Winterthurerstrasse 190, CH-8057 Zurich, Switzerland

Introduction

γ -Aminobutyric acid (GABA), the main inhibitory neurotransmitter in the brain, activates the heterodimeric G protein-coupled GABA_B receptors, which are assembled from GABA_{B1} and GABA_{B2} subunits. GABA_B receptors are abundantly expressed in pre- and postsynaptic compartments of inhibitory as well as excitatory neurons to regulate their excitability [1]. GABA_B receptors are expressed by the majority of neurons and are involved in the regulation of virtually all important brain functions, such as neuronal network activity, synaptic plasticity, and neuronal development [2–5]. Accordingly, dysfunction of GABA_B receptor signaling has been implicated in a variety of neurological disorders [6–9].

For understanding the contribution of GABA_B receptors to physiological and pathological mechanisms, the elucidation of its regulation is essential. A main factor regulating GABA_B receptor signaling is the dynamic control of their cell surface expression. In this respect, protein degradation is one important mechanism that regulates receptor availability. The two main cellular protein degradation systems, which are proteasomes and lysosomes, contribute to the regulation of cell surface GABA_B receptors in different cellular compartments. In the endoplasmic reticulum (ER), the amount of newly synthesized GABA_B receptors that are trafficked to the cell surface is determined by proteasomal degradation via the ER-associated degradation (ERAD) machinery [10]. The activity state of the neuron controls the rate of Lys-48-linked ubiquitination of the GABA_{B2} subunit required for proteasomal receptor degradation [11]. In contrast, GABA_B receptors internalized from the cell surface are degraded in lysosomes [12–16]. Sorting the receptors to lysosomes is mediated most likely via the endosomal sorting complex required for transport (ESCRT) machinery [16], which targets ubiquitinated membrane proteins to lysosomes. Accordingly, lysosomal degradation of GABA_B receptors depends on Lys-63-linked ubiquitination of the GABA_{B1} subunit at multiple sites mediated by the E3 ubiquitin ligase MIB2 [17]. However, the mechanism that regulates Lys-63-linked ubiquitination of GABA_{B1}, and thereby lysosomal degradation of the receptors, was unknown. Here, we show that phosphorylation of Ser-867 of GABA_{B1} by CaMKII β regulates the extent of MIB2-mediated K63-linked ubiquitination of GABA_{B1} and thereby the amount of lysosomal degradation.

Materials and Methods

Antibodies

Mouse anti-HA (clone HA-7, 1:1000 for immunofluorescence, 1:500 for in situ PLA, Sigma-Aldrich catalog no. H9658, lot no. 024M4773), rabbit CaMKII (1:500 for immunofluorescence, 1:100 for in situ PLA; Abcam catalog no. ab52476, lot no. GR181543-20), mouse EEA1 (1:100 for in situ PLA; BD Biosciences catalog no. 61047, lot no. 76250), rabbit GABA_{B1b} directed against the N-terminus of GABA_{B1b} (affinity-purified, 1:200 for immunofluorescence, custom made by GenScript) [18], rabbit GABA_{B2} directed against the N-terminus of GABA_{B2} (affinity-purified, 1:500 for immunofluorescence; custom made by GenScript) [19], guinea pig GABA_{B2} (1:500 for immunofluorescence; Millipore catalog no. AB2255, lot no. 2484228), mouse GABA_{B1} (1:50 for in situ PLA; NeuroMab, clone N93A/49, catalog no. 75-183), rabbit ubiquitin K48-specific (clone Apu2, 1:50 for in situ PLA; Millipore, catalog no. 05-1307, lot no. 2385989), rabbit ubiquitin K63-specific (clone Apu3, 1:50 for in situ PLA; Millipore, catalog no. 05-1308, lot no. 2575910), rabbit MIB2 (1:1000 for immunofluorescence,

1:250 for in situ PLA; MyBioSource, catalog no. MBS2014413, lot no. A20160407515), mouse phosphoserine (PSR-45, 1:150 for in situ PLA; Sigma-Aldrich, catalog no. P5747, lot no. 014M4791V), goat Rab7 (1:250 for in situ PLA; Santa Cruz Biotechnology, catalog no. sc-11303, lot no. BO911), and mouse Rab11 (clone 47, 1:25 for in situ PLA; Millipore, catalog no. 05-853, lot no. JBC1868959). Secondary antibodies were labeled with either Alexa Fluor 488 (1:1000, [Invitrogen](#)), Cy-3 (1:500, Jackson ImmunoResearch Laboratories), or Cy-5 (1:300, Jackson ImmunoResearch Laboratories).

Drugs

The following chemicals were used for this study: glutamate (50 μ M, Sigma-Aldrich) and KN93 (10 μ M, Tocris Bioscience). [³H]CGP 54626 (30 Ci/mmol) was purchased from ANAWA Trading SA and CGP 56999A was kindly provided by Novartis.

Plasmids

The following plasmids were used for this study: HA-tagged GABA_{B1a} [20]; GABA_{B2} [21]; HA-tagged GABA_{B1a}(K689/699R), GABA_{B1a}(K893R), and GABA_{B1a}(K961R) [17]; GFP-tagged CaMKII α (Addgene plasmid 21226) [22] and its functionally inactive mutant GFP-tagged CaMKII α (K42M) (here designated CaMKII α (DN)) (kindly provided by U. Bayer, University of Colorado Denver-AMC); GFP-tagged CaMKII β (Addgene plasmid 21227) [22] and its functionally inactive mutant EGFP-tagged CaMKII β (K43R) (here designated CaMKII β (DN)) (Addgene plasmid 21225) [23]; HA-tagged ubiquitin (Addgene plasmid 17608), HA-tagged ubiquitin (KO) (Addgene plasmid 17603), HA-tagged ubiquitin (K63) (Addgene plasmid 17606), and HA-tagged ubiquitin (K48R) (Addgene plasmid 17604) [24]; and HA-tagged ubiquitin (K63R) (kindly provided by L.-Y. Liu-Chen, Temple University, Philadelphia, USA).

Mutation of GABA_{B1a}

Serine 867 in GABA_{B1a} was mutated either to alanine or to aspartate by GenScript.

Culture and Transfection of Cortical Neurons

Primary cultures (co-cultures of neurons and glia cells) of the cerebral cortex were prepared from 18-day-old embryos of Wistar rats as described previously [25]. Neurons were used after 11 to 15 days in culture. Plasmid DNA was transfected into neurons by magnetofection using Lipofectamine 2000 ([Invitrogen](#)) and CombiMag (OZ Biosciences) exactly as specified in Buerli et al. [25]. In our hands, this method

reliably yields 50–100 transfected neurons, when transfection was performed with 11–13-day-old cultures.

Culture and Transfection of HEK 293 Cells

HEK (human embryonic kidney) 293 cells were cultured in Dulbecco's modified Eagle's medium (DMEM, Gibco Life Technologies) containing 10% fetal bovine serum (Gibco Life Technologies) and penicillin/streptomycin (Gibco Life Technologies). Plasmids were introduced into HEK 293 cells using the polyethylenimine method according to the jet-PEI protocol (Polyplus Transfection).

Immunocytochemistry and Confocal Laser Scanning Microscopy

Immunofluorescence staining was done as described previously [13, 26]. For detection of total GABA_B receptors, neurons were fixed for 15–20 min at room temperature with 4% paraformaldehyde followed by permeabilization with 0.2% Triton X-100 and immunostaining. For analysis of cell surface GABA_B receptors, living neurons were incubated with antibodies recognizing the extracellularly located N-terminal domain of GABA_{B1} or GABA_{B2} for 1 h at 4 °C. After washing, the neurons were incubated with fluorophore-labeled secondary antibodies for 1 h at 4 °C and subsequently fixed with 4% paraformaldehyde.

Images of stained neurons were recorded by laser scanning confocal microscopy (LSM 510 Meta, LSM 700 or LSM 710; Zeiss). Five to eight optical sections spaced by 0.3 μm were taken with a ×40, ×63, or ×100 plan-fluar oil differential interference contrast objective (Zeiss) at a resolution of 1024 × 1024 pixels. Total and cell surface fluorescence signals were quantified using the ImageJ software as described previously [26]. To determine cell surface expression of GABA_B receptors, all optical sections were merged into one image. Then, the inner as well outer border of somatic staining was carefully outlined and the mean intensity values were measured. Values for cell surface signals were obtained by subtracting the values of the inner border from those of the outer border. For analysis of total GABA_B receptor expression, the somata of neurons were carefully outlined and the mean intensity values were measured. For each image, background staining was determined in an area containing no specific signal and subtracted from the mean intensity values. All values were normalized to the analyzed cell area.

Proteins overexpressed in neurons were either tagged with GFP or the HA epitope, which were used to identify transfected neurons and to judge the protein expression level. In case of overexpression studies, transfected neurons with comparable expression of the protein of interest were included into the analysis.

In Situ Proximity Ligation Assay

In situ proximity ligation assay (PLA) is an antibody-based technology for the detection of protein-protein interactions and posttranslational modifications of proteins [27, 28]. We applied in situ PLA for the analysis of GABA_B receptor K48-linked or K63-linked ubiquitination, serine phosphorylation, and CaMKII/GABA_B receptor interaction as well as for the co-localization with the endosomal marker proteins EEA1, Rab7, and Rab11. In situ PLA was performed using Duolink PLA probes and detection reagents (Sigma-Aldrich) according to the manufacturer's instructions as described previously [26]. Quantification was done by counting individual in situ PLA spots using the Image J software. The optical sections of each image stack were merged into one image and the number of maxima was determined after setting an appropriate noise tolerance (noise tolerance was kept constant for all images of an experiment). The number of spots was normalized to the area analyzed and to the expression level of GABA_B receptors. In some experiments, signals were too abundant to resolve individual spots precisely. In those cases, the fluorescence intensity of signals was determined.

Internalization Assay

Neurons were placed on ice and cell surface GABA_B receptors were labeled with GABA_{B2} antibodies for 1 h at 4 °C. After extensive washes, neurons were incubated in 37 °C warm medium for exactly 1, 2, 3, 5, or 10 min to permit receptor endocytosis. Endocytosis was terminated by replacing the medium with ice-cold buffer. After incubation with secondary antibody for 1 h at 4 °C, the neurons were fixed with 4% PFA and remaining cell surface GABA_B receptors were determined by laser scanning confocal microscopy.

Western Blot Analysis

For Western blot analysis, neuron/glia co-cultures were grown for 13–14 days on 6-cm culture dishes plated with 450,000 cells obtained from the cerebral cortex of E18 rat embryos. After incubation with KN93, cultures were immediately washed twice with ice-cold PBS, harvested, and homogenized by sonication. After protein determination using the Bradford protein assay (BioRad), the samples were incubated with Laemmli sample buffer (BioRad) for 2 h at 37 °C and aliquots containing 30 μg protein were subjected to sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE) using 7.5% mini-gels (Mini Protean 3, BioRad). Proteins were transferred onto nitrocellulose membranes in a Mini Trans-Blot Module (BioRad) at 365 mA for 120 min using 192 mM glycine, 25 mM Tris, 0.1% SDS, 20% methanol as transfer buffer. After blotting, the transferred proteins were stained with AmidoBlack and immediately imaged using the

E-box VX2 gel imager (Vilber). For immunodetection, the blots were blocked for 1–2 h in PBST (PBS pH 7.4, 0.05% Tween 20) containing 5% nonfat dry milk at room temperature, followed by incubation with antisera overnight at 4 °C in PBST containing 5% non-fat dry milk. The blots were then washed five times for 10 min with TBST and incubated with secondary antibodies conjugated to horseradish peroxidase for 1 h at room temperature. Following extensive washing (see above), immunoreactivity was detected by the chemoluminescence method (SuperSignal West Dura, Thermo Scientific) using a Fujifilm LAS-1000 imager. Immunoreactivity was quantified with the Image Studio software (LI-CORE Biosciences) and normalized to total protein in the corresponding lanes (determined by Amidoblack staining, see above).

Radioligand Binding

For radioligand binding, 450,000 cells derived from the cerebral cortex of E18 rat embryos were plated onto 6-cm polylysine-coated culture dishes and kept in culture for 14 days. After drug incubation, cells were immediately washed two times with ice-cold PBS and harvested. Cells were resuspended in 500 μ l binding buffer containing protease inhibitors (50 mM Tris pH 7.4, 2.5 mM CaCl₂, complete mini, Sigma-Aldrich) and homogenized using a Potter S homogenizer (B. Braun Biotech International). Aliquots of the homogenate were incubated with 6.7 nM [³H]CGP 54626 in binding buffer for 90 min at room temperature. The incubation was terminated by rapid filtration onto glass fiber filters using a 12-channel semi-automated cell harvester (Scatron) and washed with ice-cold binding buffer. Non-specific [³H]CGP 54626 binding was determined in the presence of 10 μ M CGP 56999A. The radioactivity retained by the filters was determined by liquid scintillation counting using a Tricarb 2500 liquid scintillation analyzer.

Statistics

The statistical analyses were done with GraphPad Prism 5. The tests used and *p* values are given in the figure legends. Differences were considered statistically significant when *p* < 0.05.

Results

CaMKII β Controls Expression of GABA_B Receptors


Sustained activation of glutamate receptors, which occurs in cerebral ischemia, downregulates GABA_B receptors by preferentially sorting constitutively internalized receptors to lysosomes for degradation instead of recycling them to the cell

surface [29–32]. Phosphorylation of GABA_{B1} at Ser-867 by CaMKII plays a key role in this mechanism [29]. However, it was unknown whether CaMKII is also involved in the regulation of GABA_B receptor expression under normal physiological conditions. Therefore, we tested whether inhibition of CaMKII by KN93 affects total expression of GABA_B receptors in cultured cortical neurons. Incubation of neurons with KN93 transiently increased the expression of GABA_{B1} and GABA_{B2}, which peaked at about 7.5 min as tested by Western blotting (Fig. 1a). This was confirmed by radioligand binding experiments on homogenates prepared from the cultures using the GABA_B receptor antagonist [³H]CGP 54626 (Fig. 1b). Finally, we monitored cell surface expression of GABA_B receptors by staining living neurons at 4 °C with antibodies directed against the extracellular located N-terminal domain of the receptors. In line with the experiments on total receptor expression, incubation of neurons with KN93 for 7.5 min considerably increased the level of GABA_B receptors at the cell surface ($148 \pm 64\%$ of control; Fig. 1c). The cause for the transient nature of blocking CaMKII on cell surface expression of the receptors is currently unclear. It might well be that a so far unknown homeostatic response of the neurons downregulates the increased cell surface receptors to normal levels. However, our results clearly indicate that the expression of the receptors in the plasma membrane is indeed controlled by CaMKII under basal conditions.

CaMKII α and CaMKII β are the main CaMKII isoforms expressed in neurons [33]. We therefore tested which isoform is involved in the regulation of GABA_B receptors by transfecting neurons with either CaMKII α , CaMKII β , or their functionally inactive mutants (CaMKII α (DN), CaMKII β (DN)) and determined cell surface expression of GABA_B receptors using GABA_{B2} antibodies. While transfecting CaMKII α or CaMKII α (DN) did not affect cell surface expression of GABA_B receptors, overexpression of CaMKII β decreased ($43 \pm 17\%$ of EGFP-transfected control neurons, Fig. 2a) and its nonfunctional mutant CaMKII β (DN) increased cell surface levels of receptors ($186 \pm 29\%$ of EGFP-transfected control neurons, Fig. 2b).

It was previously shown that CaMKII interacts with GABA_{B1} [29]. Next, we tested whether CaMKII β or CaMKII α interacts with GABA_B receptors under basal conditions by overexpressing either CaMKII α or CaMKII β in neurons and testing for changes in interaction levels with GABA_B receptors using in situ PLA with antibodies against GABA_{B1} and CaMKII (Fig. 2b). In line with our experiments described above, overexpression of CaMKII α did not affect the level of GABA_B receptor/CaMKII interaction (PLA signal EGFP-transfected control, $100 \pm 32\%$; CaMKII α , $80 \pm 30\%$), but transfection of CaMKII β significantly increased the extent of interaction ($216 \pm 58\%$ of control). The interaction of CaMKII β with GABA_B receptors was activity-dependent as blocking CaMKII with KN93 reduced the interaction level in

Fig. 1 CaMKII regulates expression of GABA_B receptors. Neuronal cultures were incubated for the indicated times with the CaMKII inhibitor KN93 (10 μM) and tested for GABA_B receptor expression by Western blotting using antibodies against GABA_{B1} and GABA_{B2} (a) or radioligand binding with the GABA_B receptor antagonist [³H]CGP 54626 (b). In addition, cell surface expression of GABA_B receptors was analyzed by immunocytochemistry using GABA_{B1} antibodies (c). Untreated cultures served as controls. **a** Mean ± SD of 8–10 independent cultures for GABA_{B1} and 7–9 cultures for GABA_{B2}; **b** Mean ± SD of 4–6 independent cultures; **c** Mean ± SD of 39–43 neurons from three independent experiments. Scale bar in representative immunofluorescence images depicted in c represents 5 μm. **p* < 0.05, ***p* < 0.01, ****p* < 0.001; one-way ANOVA, Dunnett's multiple comparison test



neurons transfected with CaMKIIβ (PLA signal. $110 \pm 40\%$ of control).

These results indicate that CaMKIIβ interacts with GABA_B receptors in an activity-dependent manner to regulate the cell surface expression of the receptors.

CaMKII Triggers Sorting of GABA_B Receptors to Lysosomal Degradation

It had been suggested that phosphorylation of GABA_{B1} Ser-867 by CaMKII may mediate internalization of the receptors [29]. We therefore tested whether inhibition of CaMKII by KN93 affects the internalization of GABA_B receptors. However, blocking CaMKII with KN93 did affect neither the kinetics (half live; control, 1.2 min; KN93, 1.6 min) nor the extent (control, $64 \pm 19\%$; KN93, $65 \pm 20\%$) of receptor internalization (Fig. 3a).

To gain insight into the mechanism that is affected by CaMKII inhibition, we tested the co-localization of GABA_B receptors with the endosomal marker proteins EEA1 (marker for early endosomes), Rab7 (marker for late endosomes and lysosomes), and Rab11 (marker for recycling endosomes) [34] using in situ PLA. Inhibition of CaMKII activity in neurons with KN93 did not affect the co-localization of GABA_B

receptors with EEA1 ($94 \pm 26\%$ of control; Fig. 3b), supporting our observation of an unchanged internalization rate of the receptors. Instead, blocking CaMKII decreased the co-localization of GABA_B receptors with Rab7 ($39 \pm 28\%$ of control; Fig. 3b) and increased the co-localization of GABA_B receptors with Rab11 ($145 \pm 32\%$ of control; Fig. 3b). Thus, inhibition of CaMKII appears to prevent recruitment of GABA_B receptors to the lysosomal degradation pathway (Rab7) and increase recycling of the receptors (Rab11). This observation indicates that basal CaMKII activity is involved in targeting GABA_B receptors to lysosomes for degradation.

CaMKIIβ Promotes Lys-63-Linked Ubiquitination of GABA_B Receptors

The results so far support the hypothesis that CaMKIIβ regulates lysosomal degradation of GABA_B receptors. We had previously shown that lysosomal degradation of GABA_B receptors depends on Lys-63-linked ubiquitination of GABA_{B1} at multiple sites via the E3 ligase MIB2 [17]. It was therefore likely that CaMKIIβ controls Lys-63-linked ubiquitination of GABA_B receptors.





Fig. 2 CaMKII β interacts in an activity-dependent manner with GABA_B receptors to regulate their cell surface expression. **a** CaMKII β but not CaMKII α regulates cell surface expression of GABA_B receptors. Cortical neurons were transfected with either EGFP (control), CaMKII α , CaMKII β , or their functionally inactive mutants CaMKII β (DN) or CaMKII α (DN). Two days after transfection, cell surface expression of GABA_B receptors was assessed using GABA_{B2} antibodies. Transfected neurons were identified via the GFP-tag of transfected CaMKII constructs (not shown). Left, representative images of the soma of stained neurons (scale bar, 5 μ m). Right, the graphs show the quantification of fluorescence intensities. Fluorescence intensities for GABA_{B2} in EGFP-transfected neurons were set to 100%. The data represent the mean \pm

SD of 26–30 neurons from three independent experiments. *** $p < 0.001$, ^{ns} $p > 0.05$; one-way ANOVA, Dunnett's multiple comparison test. **b** CaMKII β interacts with GABA_B receptors in an activity-dependent manner. Neurons were transfected with either EGFP (control), CaMKII α , or CaMKII β and tested for interaction of CaMKII with GABA_B receptors in the presence or absence of KN93 by in situ PLA using antibodies directed against CaMKII and GABA_{B1}. Left, representative images of in situ PLA signals (white dots) in the soma of neurons (scale bar, 5 μ m). Right, the graph shows the quantification PLA signals. The data represent the mean \pm SD of 30 neurons per condition from three independent experiments. *** $p < 0.001$, ^{ns} $p > 0.05$; one-way ANOVA, Dunnett's multiple comparison test

To test this hypothesis, we first analyzed the effect of various ubiquitin mutants on the downregulation of cell surface GABA_B receptors induced by overexpression of CaMKII β . Overexpression of CaMKII β in neurons reduced cell surface GABA_B receptors to $43 \pm 18\%$ of control neurons transfected with EGFP (Fig. 4). Additional overexpression of wild-type ubiquitin (Ub, $36 \pm 25\%$ of control) and a ubiquitin mutant that permits only Lys-63-linked ubiquitination (Ub(K63), $35 \pm 18\%$ of control) as well as a mutant that prevents Lys-48-linked ubiquitination (Ub(K48R), $39 \pm 24\%$ of control) did not affect CaMKII β -

induced downregulation of the receptors ($43 \pm 18\%$ of control). However, overexpressing a ubiquitin mutant in which all lysine residues were mutated to arginine to entirely block polyubiquitination (Ub(KO), $109 \pm 22\%$ of control) as well as a ubiquitin mutant that specifically prevents Lys-63-linked ubiquitination (Ub(K63R), $124 \pm 40\%$ of control) completely prevented downregulation of the receptors or even increased cell surface expression of the receptors (Fig. 4). These results indicate that downregulation of GABA_B receptors induced by overexpression of CaMKII β is mediated by Lys-63-linked ubiquitination.


Fig. 3 CaMKII triggers sorting of GABA_B receptors to lysosomes. **a** Blocking CaMKII activity does not affect internalization of GABA_B receptors. GABA_{B2} antibody-labeled cell surface receptors of living neurons were incubated for the indicated time intervals at 37 °C to permit endocytosis. Subsequently, remaining cell surface GABA_{B2}-labeled receptors were quantified by immunofluorescence microscopy. The immunofluorescence signals of neurons at time 0 (not exposed to 37 °C) were set to 100%. Data were fitted by nonlinear regression to one phase decay. The data represent the mean ± SD of 35–46 neurons per time point from two independent experiments. **b** Blocking CaMKII activity decreased the colocalization of GABA_B receptors with Rab7 and increased the colocalization with Rab11. Neurons were either treated or not with KN93 and subjected to in situ PLA using antibodies directed against GABA_{B1} and Rab11 (marker for recycling endosomes), Rab7 (marker for the lysosomal pathway), or EEA1 (marker for early endosomes), respectively. Left, representative images (scale bar 5 μm). Right, quantification of PLA signals (white dots in images). The PLA signals were normalized to the cell area. The control condition was set to 100%. The data represent the mean ± SD of 30 neurons (Rab11) and 40 neurons (Rab7, EEA1) from two independent experiments. *****p* < 0.0001, ^{ns}*p* > 0.05; two-tailed unpaired *t* test



Next, we tested for direct CaMKII β -induced Lys-63-linked ubiquitination of GABA_B receptors using in situ PLA and antibodies directed against GABA_{B1} and Lys-63-linked ubiquitin. Inhibition of CaMKII with KN93 significantly reduced basal Lys-63-linked ubiquitination of the receptor ($69 \pm 43\%$ of control, Fig. 5a). In addition, overexpression of the functionally inactive CaMKII β mutant CaMKII β (DN) in neurons considerably reduced Lys-63-linked ubiquitination of the

receptors ($46 \pm 24\%$, Fig. 5b) as compared to neurons transfected with wild-type CaMKII β . In contrast, overexpression of the functionally inactive mutant of CaMKII α (CaMKII α (DN)) had no effect on the level of Lys-63-linked ubiquitinated GABA_B receptors as compared to wild-type CaMKII α . Finally, overexpression of neither CaMKII α (DN) nor CaMKII β (DN) affected Lys-48-linked ubiquitination of the receptors, which is required for their proteasomal

Fig. 4 CaMKII β -induced downregulation of cell surface GABA_B receptors depends on Lys-63-linked ubiquitination. Neurons were transfected with either EGFP (control), CaMKII β , or CaMKII β plus ubiquitin (Ub) or one of its mutants (Ub(K63); only Lys-63 linkages, Ub(K63R); no Lys-63 linkages, Ub(K48R); no Lys-48 linkages, Ub(KO); no polyubiquitin linkages (only mono-ubiquitination). Two days after transfection, neurons were stained for GABA_B receptors using GABA_{B2} antibodies. Only interference with K63-linked ubiquitination prevented CaMKII β -mediated downregulation of the receptors. Representative images of stained neuronal somata (scale bar, 5 μ m). The scatter plot shows quantification of fluorescence intensities. The fluorescence signal of neurons transfected with EGFP was set to 100%. The data represent the mean \pm SD of 28–30 neurons from two independent experiments. *** p < 0.001, ^{ns} p > 0.05; one-way ANOVA, Dunnett's multiple comparison test



degradation (Fig. 5c). These results indicate that CaMKII β is involved in the regulation of Lys-63-linked ubiquitination of GABA_B receptors.


We recently identified in GABA_{B1} several lysine residues (Lys-697/698, Lys-892, and Lys-960) that are substrates for Lys-63-linked ubiquitination and important for targeting the receptors to lysosomal degradation [17]. To further substantiate the finding that CaMKII regulates lysosomal degradation of GABA_B receptors by promoting Lys-63-linked ubiquitination of GABA_{B1}, we tested whether blocking CaMKII affects cell surface expression of three GABA_{B1a}(K->R) mutants in which Lys-697/698, Lys-892, or Lys-960 was mutated to arginine (GABA_{B1a}(K697/698R), GABA_{B1a}(K892R), and GABA_{B1a}(K960R)). In contrast to receptors containing wild-type GABA_{B1}, which displayed increased cell surface expression upon treatment with KN93 (GABA_{B1}: 164 \pm 149%, GABA_{B2}: 142 \pm 91% of control; Fig. 6), the expression level of receptors containing any of the three GABA_{B1a}(K->R) mutants remained unaffected by inhibition of CaMKII (GABA_{B1a}(K697/698R), control: 552 \pm 426%, KN93: 490 \pm 456%; GABA_{B1a}(K892R), control: 421 \pm 284%, KN93: 370 \pm 272%; GABA_{B1a}(K960R),

control: 273 \pm 256%, KN93: 281 \pm 270% of wild-type control; GABA_{B2} co-expressed with GABA_{B1a}(K697/698R), control: 160 \pm 115%, KN93: 140 \pm 117%; GABA_{B2} co-expressed with GABA_{B1a}(K892R), control: 184 \pm 106%, KN93: 163 \pm 88%; GABA_{B2} co-expressed with GABA_{B1a}(K960R), control: 147 \pm 84%, KN93: 145 \pm 78% of wild-type control; Fig. 6). In conclusion, these findings suggest that CaMKII is involved in K63-linked ubiquitination of GABA_{B1} at K697/698 and K892 as well as K960.

CaMKII Regulates Cell Surface Receptors by Promoting MIB2-Mediated Lys-63-Linked Ubiquitination via Phosphorylation of GABA_{B1} at Ser-867

CaMKII phosphorylates GABA_{B1} at Ser-867 [29]. To prove that direct phosphorylation of Ser-867 in GABA_{B1} is involved in the regulation of GABA_B receptor cell surface expression, we silenced this phosphorylation site by mutating it to alanine (GABA_{B1}(S867A)) or rendered the site permanently active by mutating it to aspartate (GABA_{B1}(S867D)). First, we verified that CaMKII β indeed phosphorylates GABA_{B1} at Ser-867 by

Fig. 5 CaMKII regulates K63-linked ubiquitination of GABA_B receptors. **a** Inhibition of CaMKII activity reduced K63-linked ubiquitination of GABA_B receptors. Neurons were incubated for 7.5 min in the absence (control) or presence of KN93 and analyzed for K63-linked ubiquitination of GABA_B receptor by in situ PLA using antibodies directed against GABA_{B1}- and K63-linked ubiquitin (white dots in representative images, scale bar 5 μm). Right, quantification of in situ PLA signals. The data represent the mean ± SD of 33 (control) and 30 (KN93) neurons from two independent experiments. ***p* < 0.01, two-tailed unpaired *t* test. **b, c** CaMKIIβ promotes Lys-63-linked ubiquitination of GABA_B receptors. Neurons were transfected with either CaMKIIα or CaMKIIβ or their functionally inactive mutants CaMKIIα(DN) or CaMKIIβ(DN). After 2 days, GABA_B receptors were tested for Lys-63-linked (b) or Lys-48-linked (c) ubiquitination by in situ PLA using antibodies directed against GABA_{B1} and Lys-63 or Lys-48 ubiquitin. Left, representative images depicting PLA signals (white dots, scale bars 5 μm). Right, quantification of PLA signals. The data represent the mean ± SD of 17–21 neurons from two independent experiments. ***p* < 0.01, ****p* < 0.0001, ^{ns}*p* > 0.05; two-tailed unpaired *t* test



in situ PLA using antibodies directed against GABA_{B1} and phosphoserine. Upon expression in HEK 293 cells, wild-type GABA_B receptors displayed a basal level of serine phosphorylation, which was more than threefold enhanced by co-expression with CaMKIIβ (PLA signal wt-GABA_{B1}: 98 ± 5, GABA_{B1}+CaMKIIβ: 358 ± 14; Fig. 7a). In contrast, GABA_B receptors in which Ser-867 of GABA_{B1} was inactivated by mutation to alanine, only basal level of serine phosphorylation was detected in the presence of CaMKIIβ (PLA signal GABA_{B1}(S867A)+CaMKIIβ: 108 ± 6 compared to wt-GABA_{B1}: 98 ± 5, Fig. 7a). These results confirm that CaMKIIβ specifically phosphorylates GABA_{B1} at Ser-867.

Next, we tested the effect of inactivating or permanently mimicking phosphorylation of GABA_{B1} Ser-867 on cell

surface expression of the receptors by expressing GABA_{B1}(S867A) or GABA_{B1}(S867D) in cortical neurons. Inactivation of GABA_{B1} Ser-867 considerably increased cell surface expression of receptors containing GABA_{B1}(S867A) (271 ± 119% of control, Fig. 7b) upon transfection in neurons, whereas receptors containing the phosphomimetic GABA_{B1}(S867D) mutant were expressed at reduced levels (50 ± 22% of control, Fig. 7b). In line with this finding, receptors containing the CaMKII phosphorylation-deficient GABA_{B1}(S867A) mutant exhibited reduced K63-linked ubiquitination (47 ± 32% of control, Fig. 7c), while receptors containing the phosphomimetic GABA_{B1}(S867D) mutant showed a considerably higher level of Lys-63-linked ubiquitination (226 ± 76% of control, Fig. 7c). These findings




Fig. 6 Blocking CaMKII did not affect the expression levels of GABA_{B1a}(K->R) mutants. Neurons were transfected with HA-tagged wild-type GABA_{B1a} or HA-tagged GABA_{B1a}(K->R) mutants along with GABA_{B2} and tested for cell surface expression of GABA_{B1} and GABA_{B2} after treating the neurons with 10 μM KN93 for 7.5 min. Left, representative images of untreated neurons (control, left panels) and of neurons treated with KN93 (right panels, scale bar 5 μm). The corresponding graphs show the quantification of fluorescence signals. The fluorescence intensity of neurons transfected with wild-type GABA_{B1} from untreated neurons (control) was set to 100%. The data represent the mean ± SD of 45–60 neurons per experimental condition derived from three independent experiments. Please note that the considerably lower increase in GABA_{B2} cell surface expression as compared with mutant GABA_{B1} was due to the fact that in the case of GABA_{B1}, only transfected subunits were detected (HA-tagged) but in the case of GABA_{B2}, transfected as well as endogenously expressed subunits were detected. *** $p < 0.001$, ^{ns} $p > 0.05$; two-tailed unpaired t test

suggest that CaMKII promotes Lys-63-linked ubiquitination of GABA_B receptors via phosphorylation of serine 867 in GABA_{B1}.

We previously showed that Lys-63-linked ubiquitination of GABA_{B1} is mediated by the ubiquitin E3 ligase MIB2 [17]. We therefore tested for the involvement of MIB2 in CaMKIIβ-induced downregulation of GABA_B receptors. Overexpression of MIB2 or CaMKIIβ reduced cell surface expression of GABA_B receptors in neurons to a similar extent (MIB2: 40 ± 16%, CaMKIIβ: 45 ± 17% of EGFP-transfected control neurons; Fig. 8a). Co-transfection of MIB2 and CaMKIIβ in neurons did not further increase downregulation of the receptors (43 ± 14% of EGFP-transfected control neurons; Fig. 8a), indicating that both enzymes affect GABA_B receptors via the same pathway. In line with this observation,





Fig. 7 Phosphorylation of GABA_{B1} Ser-867 controls cell surface expression and Lys-63-linked ubiquitination of GABA_B receptors. **a** CaMKIIβ phosphorylates GABA_{B1} at Ser-867. HEK293 cells were transfected either with GABA_{B1}/GABA_{B2} with or without CaMKIIβ or with GABA_{B1}(S867A)/GABA_{B2} and CaMKIIβ. Two days after transfection, the cells were tested for serine phosphorylation by in situ PLA using antibodies directed against GABA_{B1} and phosphoserine. Left, representative images (scale bar, 10 μm; wt, wild type). Right, quantification of PLA signals (white dots in images). The PLA signals were normalized to the cell area as well as to the expression level of GABA_{B1}. The data represent the mean ± SD of 50 cells from two independent experiments. *** $p < 0.001$, ^{ns} $p > 0.05$; one-way ANOVA, Dunnett's multiple comparison test. **b, c** Inactivation or constitutively mimicking phosphorylation of

GABA_{B1} Ser-867 affects cell surface expression (**b**) as well as Lys-63-linked ubiquitination (**c**) of the receptors. Neurons were transfected with wild-type HA-tagged GABA_{B1a} (control), with the phosphorylation-deficient HA-tagged GABA_{B1a}(S867A) mutant or with the phosphomimetic HA-tagged GABA_{B1a}(S867D) mutant along with GABA_{B2} and tested for cell surface expression using HA antibodies (**b**) as well as for Lys63-linked ubiquitination by in situ PLA with HA and Lys-63 antibodies (**c**). Left, representative images (scale bars, 5 μm). Right, quantification of fluorescence intensities (**b**) and PLA signals (**c**). The data represent the mean ± SD of 23–27 (**b**) and 18–20 (**c**) neurons from two independent experiments. * $p < 0.05$, *** $p < 0.001$; one-way ANOVA, Dunnett's multiple comparison test



MIB2 was unable to downregulate cell surface GABA_B receptors when the nonfunctional CaMKII β mutant CaMKII β (DN) was simultaneously overexpressed with MIB2 (CaMKII β (DN): $195 \pm 42\%$, CaMKII β (DN)+MIB2: $200 \pm 43\%$ of EGFP-transfected control neurons, Fig. 8a).

Next, we tested whether MIB2-mediated downregulation of GABA_B receptors depends on phosphorylation of GABA_{B1} Ser-867. Co-transfection of GABA_{B1a,2} and MIB2 in neurons strongly reduced cell surface expression of

receptors containing transfected GABA_{B1a} ($28 \pm 11\%$ of control, Fig. 8b). However, MIB2 failed to affect expression of the elevated or reduced cell surface expression levels of receptors containing GABA_{B1a}(S867A) or GABA_{B1a}(S867D) (S867A: $204 \pm 39\%$, S867A+MIB2: $211 \pm 38\%$, S867D: $23 \pm 8\%$, S867D+MIB2: $24 \pm 9\%$ of control, Fig. 8b). Together, these results indicate that phosphorylation of GABA_{B1} at Ser-867 by CaMKII β is required for MIB2-mediated downregulation of GABA_B receptors.

Fig. 8 The ubiquitin E3 ligase MIB2 is involved in CaMKII β -mediated regulation of cell surface GABA_B receptors. **a** Overexpression of MIB2 in addition to CaMKII does not enhance downregulation of cell surface GABA_B receptors. Neurons were transfected with the indicated plasmid and tested for cell surface expression of GABA_B receptors using GABA_{B2} antibodies 2 days after transfection. Co-expression of MIB2 did not further enhance downregulation of cell surface receptors, indicating the involvement in the same pathway. Upper panels show representative images (scale bars, 5 μ m). The graphs depict quantification of the fluorescence signals. The data represent the mean \pm SD of 30 neurons per condition derived from two independent experiments. ^{ns} $p > 0.05$, ^{****} $p < 0.0001$; two-way ANOVA with Bonferroni's multiple comparison test (interaction: $F(2,234) = 32.78$, $p < 0.0001$). **b** MIB2-induced downregulation of cell surface receptors depends on phosphorylation of GABA_{B1} Ser-867. Neurons were transfected with wild-type HA-tagged GABA_{B1a} (control), with the phosphorylation-deficient HA-tagged GABA_{B1a}(S867A) mutant or with the phosphomimetic HA-tagged GABA_{B1a}(S867D) mutant along with GABA_{B2} and with or without MIB2. Two days after transfection, neurons were tested for cell surface expression of tagged GABA_{B1} using HA antibodies. Upper panels show representative images (scale bars, 5 μ m). The graph depicts quantification of the fluorescence signals. The data represent the mean \pm SD of 30 neurons per condition derived from two independent experiments. ^{ns} $p > 0.05$, ^{****} $p < 0.0001$; two-way ANOVA, Bonferroni's multiple comparison test (interaction: $F(2,174) = 49.09$, $p < 0.0001$). **c** GABA_{B1} phosphomutants display altered MIB2 interaction. HEK 293 cells were transfected with wild-type GABA_{B1}, GABA_{B1}(S867A), or GABA_{B1}(S867D) together with GABA_{B2} and MIB2. Two days after transfection, cells were analyzed for interaction of GABA_B receptors with MIB2 by in situ PLA using antibodies directed against GABA_{B1} and MIB2. Upper panels show representative images (scale bar, 10 μ m). The graph depicts quantification of the PLA signals. The data represent the mean \pm SD of 35 neurons per condition derived from two independent experiments. ^{**} $p < 0.01$, ^{***} $p < 0.001$; one-way ANOVA, Dunnett's multiple comparison test

Finally, we tested the hypothesis that CaMKII-mediated phosphorylation of GABA_{B1} might affect the interaction of MIB2 with GABA_B receptors. Upon transfection in HEK 293 cells, MIB2 displayed considerably lower interaction with receptors containing the phospho-deficient mutant GABA_{B1a}(S867A) ($25 \pm 15\%$ of control, Fig. 8c) and a higher level of interaction with receptors containing the phosphomimetic mutant GABA_{B1a}(S867D) ($126 \pm 44\%$ of control, Fig. 8c) than wild-type receptors. This result indicates that phosphorylation of GABA_{B1} at Ser-867 promotes the interaction of GABA_B receptors with MIB2.

Sustained Activation of Glutamate Receptors Increases Lys-63-Linked Ubiquitination of GABA_B Receptors via Phosphorylation of GABA_{B1} Ser-867

Glutamate-induced downregulation of GABA_B receptors via lysosomal degradation depends on CaMKII-mediated phosphorylation of GABA_{B1} Ser-867 [29] as well as on Lys-63-linked ubiquitination of GABA_{B1} at Lys-697/698, Lys-892, and Lys-960 [17]. To investigate the role of CaMKII in this process, we first analyzed the effect of sustained glutamate


treatment on the expression level of CaMKII as well as on the degree of CaMKII-GABA_B receptor interaction. Treatment of neurons for 60 min with glutamate significantly upregulated CaMKII ($233 \pm 61\%$ of control, Fig. 9a) and considerably increased the interaction of CaMKII with GABA_B receptors ($165 \pm 74\%$ of control, Fig. 9b).

We then analyzed whether glutamate-induced K63-linked ubiquitination of GABA_B receptors depends on phosphorylation of Ser-867. Wild-type GABA_{B1} or the two mutants GABA_{B1}(S867A) and GABA_{B1}(S867D) were expressed in cortical neurons and exposed for 60 min to glutamate. Under this condition, cell surface wild-type GABA_B receptors were downregulated to $30 \pm 20\%$ as compared to untreated controls (Fig. 9c). However, receptors containing the phospho-deficient mutant GABA_{B1}(S867A) showed enhanced cell surface expression under control conditions ($240 \pm 99\%$ of wild-type controls) and were not affected by glutamate ($240 \pm 94\%$ of wild-type controls, Fig. 9c). In contrast, receptors containing the phosphomimetic mutant GABA_{B1}(S867D) displayed strongly reduced expression under control conditions ($45 \pm 24\%$ of wild-type controls, Fig. 9c) and remained unaffected after glutamate treatment ($44 \pm 23\%$ of wild-type controls, Fig. 9c). A complementary pattern was observed for Lys-63-linked ubiquitination of the respective receptors as tested by in situ PLA. Treatment of cultured neurons with glutamate strongly increased K63-linked ubiquitination of wild-type GABA_B receptors (control: $100 \pm 60\%$, glutamate: $247 \pm 84\%$, Fig. 9d), whereas receptors containing the phospho-deficient mutant GABA_{B1}(S867A) displayed only marginal Lys-63-linked ubiquitination under both conditions (control: $10 \pm 13\%$, glutamate: 11 ± 13 , Fig. 9d) and the phosphomimetic mutant GABA_{B1}(S867D) exhibited strongly enhanced Lys-63-linked ubiquitination (control: 175 ± 126 , glutamate: 181 ± 121 , Fig. 9d).

These findings suggest that sustained activation of glutamate receptors induces GABA_{B1}-Ser-867 phosphorylation-mediated K63-linked ubiquitination of GABA_B receptors, promoting their lysosomal degradation.

Discussion

The expression of GABA_B receptors available for signal transduction at the cell surface critically depends on their rate of degradation. The two main cellular degradation systems—proteasomes and lysosomes—control the abundance of GABA_B receptors in distinct cellular compartments in response to the activity state of the neuron [10, 11, 17]. In the ER, the amount of newly synthesized GABA_B receptors destined for trafficking to the plasma membrane is regulated by proteasomal degradation via the ERAD machinery [10], whereas cell surface GABA_B receptors are degraded in lysosomes [12–16]. Both degradation pathways require



ubiquitination of the receptor as targeting signals. Lys-48-linked ubiquitination of GABA_{B2} at Lys-767/771 tags GABA_B receptors for proteasomal degradation [10] and Lys-

63-linked ubiquitination of GABA_{B1} at several sites sorts the receptors to lysosomes [17]. Interestingly, Lys-48- and Lys-63-linked ubiquitination appears to be largely segregated to

Fig. 9 Glutamate-induced downregulation of GABA_B receptors. **a** Prolonged glutamate treatment increased CaMKII expression. Neurons were treated with 50 μM glutamate for 60 min and analyzed after additional 16 h for CaMKII expression. Upper panel shows representative images (scale bar, 5 μm). The data represent the mean ± SD of 45 neurons per experimental condition derived from two independent experiments. *****p* < 0.0001, two-tailed unpaired *t* test. **b** Prolonged glutamate treatment increased the interaction of CaMKII with GABA_B receptors. Neurons were treated with 50 μM glutamate for 60 min and analyzed after additional 16 h for the interaction of CaMKII with GABA_B receptors via in situ PLA using CaMKII and GABA_{B1} antibodies. Upper panel shows representative images (scale bar, 5 μm). The data represent the mean ± SD of 39 neurons per experimental condition derived from two independent experiments. *****p* < 0.0001, two-tailed unpaired *t* test. **c, d** Glutamate-induced downregulation of GABA_B receptors is mediated by GABA_{B1} Ser-867 phosphorylation-induced Lys-63-linked ubiquitination. Neurons transfected with wild-type HA-tagged GABA_{B1a}, with the phosphorylation-deficient HA-tagged GABA_{B1a}(S867A) mutant or with the phosphomimetic HA-tagged GABA_{B1a}(S867D) mutant along with GABA_{B2} were incubated for 60 min in the absence (control) or presence of 50 μM glutamate. Neurons were analyzed for cell surface expression of transfected GABA_{B1} using HA antibodies (**c**) or analyzed for Lys-63-linked ubiquitination by in situ PLA using HA and Lys-63 antibodies (**d**). Upper panels show representative images (scale bars, 5 μm). The graphs depict quantification of the fluorescence signals (**c**) and in situ PLA signals (**d**). The data represent the mean ± SD of 28–30 (**c**) and 28–36 (**d**) neurons per condition derived from three independent experiments. ^{ns}*p* > 0.05, *****p* < 0.0001; two-way ANOVA, Bonferroni's multiple comparison test (**c** interaction: *F*(2,170) = 6.27, *p* < 0.005; **d** interaction: *F*(2, 205) = 17.22, *p* < 0.0001)

GABA_{B2} and GABA_{B1} [10, 17], respectively, which might be explained by a selective interaction of the subunits with the respective E3 ligases. However, the mechanisms regulating lysosomal degradation of GABA_B receptors in response to changes in the physiological state of the neuron were unclear.

Under normal physiological conditions, cell surface GABA_B receptors constitutively internalize to early endosomes and recycle to the plasma membrane or are sorted to lysosomes for degradation [12–14, 35, 36]. Sorting the receptors to recycling or degradation must be precisely regulated in order to provide the required number of cell surface receptors for signal transduction under a given physiological condition. However, after prolonged activation of glutamate receptors, GABA_B receptors are rapidly downregulated by shifting the recycling/degradation balance towards lysosomal degradation [30–32]. This downregulation of GABA_B receptors critically depends on CaMKII-mediated phosphorylation of GABA_{B1} on Ser-867 [29] as well as on Lys-63-linked ubiquitination of GABA_{B1} mediated by the E3 ligase MIB2 [17]. As the level of CaMKII activity is regulated by the intracellular Ca²⁺ concentration, it is in an ideal position to link the increased neuronal activity after prolonged activation of glutamate receptors to lysosomal degradation of GABA_B receptors at least under pathological conditions. We, therefore, hypothesized that CaMKII regulates lysosomal degradation also under physiological conditions. Indeed, we found that

pharmacologically blocking CaMKII significantly increased cell surface expression of GABA_B receptors. This increase was accompanied by a reduced co-localization of the receptors with the late endosomal marker Rab7, indicating that inhibition of CaMKII prevented sorting of GABA_B receptors to lysosomes. Our data also imply that a considerable fraction of GABA_B receptors is constitutively degraded under basal conditions since blocking CaMKII activity for only 7.5 min was sufficient to induce a significant increase (~150%) of cell surface GABA_B receptors.

CaMKII is a large protein complex constituted from 12 catalytically active subunits of different isoforms (CaMKIIα-δ) [33]. In brain, CaMKIIα and CaMKIIβ are the predominant subunits and in forebrain neurons the CaMKII holoenzyme largely consists of nine α subunits and three β subunits [37, 38]. Interestingly, we found that overexpression of CaMKIIβ, but not CaMKIIα, increased the level of GABA_B receptor/CaMKII interaction in an activity-dependent manner (inhibition of CaMKII activity prevented the increase in interaction), resulting in decreased cell surface receptor expression. Consistent with this observation, overexpressing its functionally inactive mutant CaMKIIβ(DN), which was expected to inhibit CaMKIIβ activity and thus lysosomal degradation of the receptors, increased cell surface expression of GABA_B receptors. As intracellular Ca²⁺ concentrations under basal conditions are low, this finding fits well to the observation that CaMKIIβ exhibits a ninefold higher affinity to Ca²⁺/calmodulin than CaMKIIα [38]. Thus, our data suggest that under normal physiological conditions basal CaMKIIβ activity determines the level of lysosomal degradation of GABA_B receptors.

Phosphorylation often regulates ubiquitination of proteins, thereby promoting their degradation [39–41]. Several lines of evidence indicate that regulation of cell surface GABA_B receptors by CaMKIIβ is conveyed by MIB2-mediated Lys-63-linked ubiquitination of GABA_{B1}. First, overexpression in neurons of ubiquitin mutants unable to form Lys-63-linkages prevented downregulation of GABA_B receptors induced by overexpression of CaMKIIβ, while a mutant that only can form Lys-63-linkages, but not any other kind of linkages, did not inhibit downregulation of the receptors. Second, blocking CaMKII activity pharmacologically or by overexpression of the functionally inactive mutant CaMKIIβ(DN) reduced Lys-63-linked ubiquitination of GABA_B receptors. Third, the cell surface expression of three GABA_{B1a}(K->R) mutants with inactivated Lys-63-linked ubiquitination sites, which are indispensable for lysosomal targeting of the receptors [17], remained unaffected by CaMKII inhibition. Fourth, overexpression of the E3 ligase MIB2, which downregulates GABA_B receptors via lysosomal degradation [17], neither enhanced CaMKIIβ-mediated downregulation of cell surface receptors nor affected the expression of a GABA_{B1} mutant with inactivated CaMKII phosphorylation site (GABA_{B1}(S867A)). Finally, our experiments with the CaMKII

phosphorylation-deficient and phosphomimetic GABA_{B1}(S867) mutants indicate that direct phosphorylation of GABA_{B1} Ser-867 promotes Lys-63-linked ubiquitination and thereby degradation of GABA_B receptors. Under normal physiological as well as conditions of over-excitation via prolonged activation of glutamate receptors, the phospho-deficient mutant GABA_B(S867A) displayed only marginal Lys-63-linked ubiquitination and increased cell surface expression whereas the phosphomimetic mutant GABA_{B1}(S867D) exhibited strongly increased Lys-63-linked ubiquitination and reduced cell surface expression.

The precise mechanism by which CaMKII β promotes Lys63-linked ubiquitination of GABA_{B1} remains unresolved. Phosphorylation of GABA_{B1}(S867) might be the signal that targets the receptors to an endosomal compartment where they are Lys-63-linked ubiquitinated by MIB2. Alternatively, phosphorylation of GABA_{B1}(S867) might induce a conformational change in the receptor that uncovers the MIB2 interaction site or exposes the lysine residues for ubiquitination. We favor the latter hypothesis, which is supported by our finding that MIB2 displayed increased interaction with the phosphomimetic mutant GABA_{B1}(S867D) and reduced interaction with the phospho-deficient mutant GABA_{B1}(S867A). However, further experimentation is required to reveal the mechanism as to how CaMKII β promotes Lys-63-linked ubiquitination of GABA_{B1}.

In addition to CaMKII β controlling cell surface GABA_B receptor expression under normal physiological conditions, it rapidly downregulates the receptors upon sustained activation of glutamate receptors [29–32], a pathological condition that occurs in cerebral ischemia and leads to excitotoxic cell death [42, 43]. Downregulation of GABA_B receptors is triggered by increased Ca²⁺ influx through NMDA receptors and voltage-gated Ca²⁺ channels [30], leading to enhanced CaMKII-mediated phosphorylation of GABA_{B1} on Ser867 [29]. This causes the re-routing of GABA_B receptors from the normal constitutive recycling pathway to lysosomal degradation. Our present results show that this pathological downregulation of GABA_B receptors is maintained after removal of glutamate due to the upregulation of CaMKII expression and the increased interaction of CaMKII with GABA_B receptors. Therefore, interfering with this pathway to normalize GABA_B receptor-mediated inhibition might be a potential approach to counteract over-excitation and to limit neuronal death.

In conclusion, our data indicate that phosphorylation of GABA_{B1} at Ser-867 by CaMKII β induces MIB2-mediated K63-linked ubiquitination of GABA_{B1} at multiple sites, which sorts GABA_B receptors to lysosomes for degradation. This mechanism is expected to fine-tune cell surface expression of GABA_B receptors under physiological conditions and to considerably affect receptor expression in diseases associated with disturbed Ca²⁺ homeostasis.

Acknowledgements We are grateful to Giovanna Bosshard for dissecting the E18 rat cortex.

Funding Information This study was supported by the Swiss National Science Foundation (grant 31003A_138382 to D.B.).

Compliance with Ethical Standards

Conflict of Interest The authors declare that they have no conflicts of interest.

Open Access This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.

References

1. Benke D, Zemoura K, Maier PJ (2012) Modulation of cell surface GABA_B receptors by desensitization, trafficking and regulated degradation. *World J Biol Chem* 3(4):61–72. <https://doi.org/10.4331/wjbc.v3.i4.61> [doi]
2. Pinard A, Seddik R, Bettler B (2010) GABA_B receptors: physiological functions and mechanisms of diversity. *Adv Pharmacol* 58:231–255. [https://doi.org/10.1016/s1054-3589\(10\)58010-4](https://doi.org/10.1016/s1054-3589(10)58010-4)
3. Kohl MM, Paulsen O (2010) The roles of GABA_B receptors in cortical network activity. *Adv Pharmacol* 58:205–229. [https://doi.org/10.1016/s1054-3589\(10\)58009-8](https://doi.org/10.1016/s1054-3589(10)58009-8)
4. Craig MT, McBain CJ (2014) The emerging role of GABA_B receptors as regulators of network dynamics: fast actions from a ‘slow’ receptor? *Curr Opin Neurobiol* 26:15–21. <https://doi.org/10.1016/j.conb.2013.10.002>
5. Gaiarsa JL, Porcher C (2013) Emerging neurotrophic role of GABA_B receptors in neuronal circuit development. *Front Cell Neurosci* 7:206. <https://doi.org/10.3389/fncel.2013.00206>
6. Benarroch EE (2012) GABA_B receptors: structure, functions, and clinical implications. *Neurology* 78(8):578–584. <https://doi.org/10.1212/WNL.0b013e318247cd03>
7. Bowery NG (2006) GABA_B receptor: a site of therapeutic benefit. *Curr Opin Pharmacol* 6(1):37–43. <https://doi.org/10.1016/j.coph.2005.10.002>
8. Cryan JF, Kaupmann K (2005) Don’t worry ‘B’ happy!: a role for GABA_B receptors in anxiety and depression. *Trends Pharmacol Sci* 26(1):36–43
9. Kumar K, Sharma S, Kumar P, Deshmukh R (2013) Therapeutic potential of GABA_B receptor ligands in drug addiction, anxiety, depression and other CNS disorders. *Pharmacol Biochem Behav* 110:174–184. <https://doi.org/10.1016/j.pbb.2013.07.003>
10. Zemoura K, Schenkel M, Acuna MA, Yevenes GE, Zeilhofer HU, Benke D (2013) Endoplasmic reticulum-associated degradation (ERAD) controls cell surface expression of γ -aminobutyric acid, type B receptors. *J Biol Chem* 288(48):34897–34905. <https://doi.org/10.1074/jbc.M113.514745>
11. Zemoura K, Benke D (2014) Proteasomal degradation of γ -aminobutyric acid_B receptors is mediated by the interaction of the GABA_{B2} C terminus with the proteasomal ATPase Rtp6 and regulated by neuronal activity. *J Biol Chem* 289:7738–7746. <https://doi.org/10.1074/jbc.M113.541987>
12. Grampp T, Sauter K, Markovic B, Benke D (2007) γ -Aminobutyric acid type B receptors are constitutively internalized via the clathrin-dependent pathway and targeted to lysosomes for degradation. *J Biol Chem* 282:24157–24165. <https://doi.org/10.1074/jbc.M702626200>

13. Grampp T, Notz V, Broll I, Fischer N, Benke D (2008) Constitutive, agonist-accelerated, recycling and lysosomal degradation of GABA_B receptors in cortical neurons. *Mol Cell Neurosci* 39(4): 628–637. <https://doi.org/10.1016/j.mcn.2008.09.004>
14. Vargas KJ, Terunuma M, Tello JA, Pangalos MN, Moss SJ, Couve A (2008) The availability of surface GABA_B receptors is independent of g-aminobutyric acid but controlled by glutamate in central neurons. *J Biol Chem* 283(36):24641–24648. <https://doi.org/10.1074/jbc.M802419200>
15. Hannan S, Wilkins ME, Dehghani-Tafti E, Thomas P, Baddeley SM, Smart TG (2011) GABA_B receptor internalisation is regulated by the R2 subunit. *J Biol Chem* 286(27):24324–24335. <https://doi.org/10.1074/jbc.M111.220814>
16. Kantamneni S, Holman D, Wilkinson KA, Correa SA, Feligioni M, Ogden S, Fraser W, Nishimune A et al (2008) GISP binding to TSG101 increases GABA_B receptor stability by down-regulating ESCRT-mediated lysosomal degradation. *J Neurochem* 107(1):86–95. <https://doi.org/10.1111/j.1471-4159.2008.05580.x>
17. Zemoura K, Trumpler C, Benke D (2016) Lys-63-linked ubiquitination of γ -aminobutyric acid (GABA), Type B1, at multiple sites by the E3 ligase Mind Bomb-2 targets GABA_B receptors to lysosomal degradation. *J Biol Chem* 291:21682–21693. <https://doi.org/10.1074/jbc.M116.750968>
18. Benke D, Honer M, Michel C, Bettler B, Mohler H (1999) γ -Aminobutyric acid type B receptor splice variant proteins GBR1a and GBR1b are both associated with GBR2 *in situ* and display differential regional and subcellular distribution. *J Biol Chem* 274:27323–27330
19. Benke D, Michel C, Mohler H (2002) Structure of GABA_B receptors in rat retina. *J Recept Signal Transduc* 22:253–266. <https://doi.org/10.1081/RRS-120014600>
20. Kaupmann K, Huggel K, Heid J, Flor PJ, Bischoff S, Mickel SJ, McMaster G, Angst C et al (1997) Expression cloning of GABA_B receptors uncovers similarity to metabotropic glutamate receptors. *Nature* 386(6622):239–246. <https://doi.org/10.1038/386239a0>
21. Kaupmann K, Malitschek B, Schuler V, Heid J, Froestl W, Beck P, Mosbacher J, Bischoff S et al (1998) GABA_B-receptor subtypes assemble into functional heteromeric complexes. *Nature* 396(6712):683–687. <https://doi.org/10.1038/25360>
22. Shen K, Teruel MN, Subramanian K, Meyer T (1998) CaMKII β functions as an F-actin targeting module that localizes CaMKII α / β heterooligomers to dendritic spines. *Neuron* 21(3):593–606
23. Shen K, Meyer T (1999) Dynamic control of CaMKII translocation and localization in hippocampal neurons by NMDA receptor stimulation. *Science* 284(5411):162–166
24. Lim KL, Chew KC, Tan JM, Wang C, Chung KK, Zhang Y, Tanaka Y, Smith W et al (2005) Parkin mediates nonclassical, proteasomal-independent ubiquitination of synphilin-1: implications for Lewy body formation. *J Neurosci* 25(8):2002–2009. <https://doi.org/10.1523/JNEUROSCI.4474-04.2005>
25. Buerli T, Pellegrino C, Baer K, Lardi-Studler B, Chudotvorova I, Fritschy JM, Medina I, Fuhrer C (2007) Efficient transfection of DNA or shRNA vectors into neurons using magnetofection. *Nat Protoc* 2(12):3090–3101. <https://doi.org/10.1038/nprot.2007.445>
26. Maier PJ, Zemoura K, Acuna MA, Yevenes GE, Zeilhofer HU, Benke D (2014) Ischemia-like oxygen and glucose deprivation mediates down-regulation of cell surface γ -aminobutyric acid_B receptors via the endoplasmic reticulum (ER) stress-induced transcription factor CCAAT/enhancer-binding protein (C/EBP)-homologous protein (CHOP). *J Biol Chem* 289(18):12896–12907. <https://doi.org/10.1074/jbc.M114.550517>
27. Leuchowius KJ, Jarvius M, Wickstrom M, Rickardson L, Landegren U, Larsson R, Soderberg O, Fryknas M et al (2010) High content screening for inhibitors of protein interactions and post-translational modifications in primary cells by proximity ligation. *Mol Cell Proteomics* 9(1):178–183. <https://doi.org/10.1074/mcp.M900331-MCP200>
28. Soderberg O, Gullberg M, Jarvius M, Ridderstrale K, Leuchowius KJ, Jarvius J, Wester K, Hydbring P et al (2006) Direct observation of individual endogenous protein complexes *in situ* by proximity ligation. *Nat Methods* 3(12):995–1000. <https://doi.org/10.1038/nmeth947>
29. Guetg N, Aziz SA, Holbro N, Turecek R, Rose T, Seddik R, Gassmann M, Moes S et al (2010) NMDA receptor-dependent GABA_B receptor internalization via CaMKII phosphorylation of serine 867 in GABA_{B1}. *Proc Natl Acad Sci U S A* 107(31): 13924–13929. <https://doi.org/10.1073/pnas.1000909107>
30. Maier PJ, Marin I, Grampp T, Sommer A, Benke D (2010) Sustained glutamate receptor activation down-regulates GABA_B receptors by shifting the balance from recycling to lysosomal degradation. *J Biol Chem* 285(46):35606–35614. <https://doi.org/10.1074/jbc.M110.142406>
31. Terunuma M, Vargas KJ, Wilkins ME, Ramirez OA, Jaureguiberry-Bravo M, Pangalos MN, Smart TG, Moss SJ et al (2010) Prolonged activation of NMDA receptors promotes dephosphorylation and alters postendocytic sorting of GABA_B receptors. *Proc Natl Acad Sci U S A* 107(31):13918–13923. <https://doi.org/10.1073/pnas.1000853107>
32. Kantamneni S, Gonzalez-Gonzalez IM, Luo J, Cimarosti H, Jacobs SC, Jaafari N, Henley JM (2014) Differential regulation of GABA_B receptor trafficking by different modes of N-methyl-D-aspartate (NMDA) receptor signaling. *J Biol Chem* 289(10):6681–6694. <https://doi.org/10.1074/jbc.M113.487348>
33. Hell JW (2014) CaMKII: claiming center stage in postsynaptic function and organization. *Neuron* 81(2):249–265. <https://doi.org/10.1016/j.neuron.2013.12.024>
34. Ng EL, Gan BQ, Ng F, Tang BL (2012) Rab GTPases regulating receptor trafficking at the late endosome-lysosome membranes. *Cell Biochem Funct* 30(6):515–523. <https://doi.org/10.1002/cbf.2827>
35. Pooler AM, Gray AG, McIlhinney RA (2009) Identification of a novel region of the GABA_{B2} C-terminus that regulates surface expression and neuronal targeting of the GABA_B receptor. *Eur J Neurosci* 29(5): 869–878. <https://doi.org/10.1111/j.1460-9568.2009.06636.x>
36. Wilkins ME, Li X, Smart TG (2008) Tracking cell surface GABA_B receptors using an α -bungarotoxin tag. *J Biol Chem* 283(50): 34745–34752. <https://doi.org/10.1074/jbc.M803197200>
37. Miller SG, Kennedy MB (1985) Distinct forebrain and cerebellar isoforms of type II Ca²⁺/calmodulin-dependent protein kinase associate differently with the postsynaptic density fraction. *J Biol Chem* 260(15):9039–9046
38. Brocke L, Chiang LW, Wagner PD, Schulman H (1999) Functional implications of the subunit composition of neuronal CaM kinase II. *J Biol Chem* 274(32):22713–22722
39. Lin HK, Wang L, Hu YC, Altuwaijri S, Chang C (2002) Phosphorylation-dependent ubiquitylation and degradation of androgen receptor by Akt require Mdm2 E3 ligase. *EMBO J* 21(15): 4037–4048
40. Schroeder B, Srivatsan S, Shaw A, Billadeau D, McNiven MA (2012) CIN85 phosphorylation is essential for EGFR ubiquitination and sorting into multivesicular bodies. *Mol Biol Cell* 23(18):3602–3611. <https://doi.org/10.1091/mbc.E11-08-0666>
41. Su H, Chen M, Sands JM, Chen G (2013) Activation of the cAMP/PKA pathway induces UT-A1 urea transporter monoubiquitination and targets it for lysosomal degradation. *Am J Physiol Renal Physiol* 305(12):F1775–F1782. <https://doi.org/10.1152/ajprenal.00393.2013>
42. Kostandy BB (2012) The role of glutamate in neuronal ischemic injury: the role of spark in fire. *Neurol Sci* 33(2):223–237. <https://doi.org/10.1007/s10072-011-0828-5>
43. Lipton P (1999) Ischemic cell death in brain neurons. *Physiol Rev* 79(4):1431–1568. <https://doi.org/10.1152/physrev.1999.79.4.1431>