

Comparison of pharmacist and public views and experiences of community pharmacy medicines-related services in England

Ruth M Rodgers
Shivaun M Gammie
Ruey Leng Loo
Sarah A Corlett
Janet Kraska

Medway School of Pharmacy,
The Universities of Greenwich and
Kent, Chatham Maritime, UK

Background: Services provided by community pharmacists designed to support people using medicines are increasing. In England, two national services exist: Medicine Use Reviews (MUR) and New Medicines Service (NMS). Very few studies have been conducted seeking views of the public, rather than service users, on willingness to use these services or expectations of these services, or determined whether views align with pharmacist perceptions.

Objective: To compare the perceptions of pharmacists and the general public on medicines-related services, particularly MUR and NMS services.

Methods: Two parallel surveys were conducted in one area of England: one involved the general public and was administered using a street survey, and the other was a postal survey of community pharmacists. Similar questionnaires were used, seeking views of services, awareness, reasons for using services, and perceived benefits.

Results: Response rates were 47.2% (1,000/2,012 approached) for the public and 40.8% (341/836) for pharmacists. Few people had experienced a discussion in a private consultation room or were aware of the two formal services, although their willingness to use them was high. Pharmacists estimated time spent on service provision as 10 minutes for MUR and 12 minutes for NMS, which aligned with acceptability to both pharmacists and the public. Pharmacists underestimated the willingness of the public to wait for an informal discussion or to make appointments for formal services. Both pharmacists and the public had high expectations that services would be beneficial in terms of increasing knowledge and understanding, but public expectations and experiences of services helping to sort out problems fell well below pharmacists' perceptions. People who had experienced a pharmacy service had different perceptions of pharmacists.

Conclusion: Views differed regarding why people use services and key aspects of service delivery. For services to improve, the pharmacy profession needs a better awareness of what the public, especially those with potential to benefit from services, view as acceptable and desirable.

Keywords: community pharmacy, public opinion, pharmacist perceptions, medicines-related services

Introduction

Community pharmacists are increasingly being commissioned to provide cognitive services to support and improve medicines use, such as the Home Medicine Review (HMR) service in Australia and Medication Therapy Management (MTM) services in the US. In England, the Medicines Use Review (MUR) service was introduced in 2005 and by 2014/15, 3.2 million MURs were provided by 93.5% (10,916) of community pharmacies in England.¹ This service aims to improve patients' knowledge and use of medicines and help reduce

Correspondence: Janet Kraska
Medway School of Pharmacy,
The Universities of Greenwich and Kent,
Chatham Maritime, Kent ME4 4TB, UK
Tel +44 1634 202950
Fax +44 01634 883827
Email j.kraska@kent.ac.uk

medicines wastage. Changes in 2011 require that at least half the MURs provided must be targeted toward patients with respiratory disease, those taking high-risk medicines (nonsteroidal anti-inflammatory drug, anticoagulant, antiplatelet, or diuretic) and those recently discharged from a hospital.

An early evaluation found that 93% of pharmacists viewed the MUR as potentially improving patients' use of medicines, including adherence and cost-effectiveness,² believing that patients would want pharmacists to review their medicines. Later in-depth work suggested that MURs were poorly integrated into pharmacy workflow, most being provided on an ad hoc basis, conducted with people using few medicines, in part due to dispensing pressures and target numbers imposed by employers.³

Most studies seeking the views of patients or pharmacists on the MUR service have been small scale,^{2,4-8} showing that patients seldom request an MUR, most are initiated by pharmacists and patients feel obliged to undergo one to help the pharmacist. An in-depth study of MUR consultations found that the experience did not increase patients' knowledge and it rarely affected medicine use.³ These findings echo earlier work showing that patients were ambivalent about receiving a medicines management consultation with a pharmacist, viewing it as providing reassurance.⁹

A second formal service, the New Medicines Service (NMS), was introduced in England in 2011, targeted toward patients with asthma, COPD, diabetes, hypertension, or taking an antiplatelet/anticoagulant, aiming to help reduce symptoms and complications of these conditions and identify any related issues, including the need for further information and support. The service involves two or three consultations, only the first of which should be face-to-face interaction. A total of 775,998 NMS were provided by 9,308 (79.7%) English community pharmacies in 2014/15.¹ An evaluation of this service found that pharmacists considered it as an opportunity to educate and support patients, while patients were unaware of it, but viewed it positively once experienced.¹⁰

Studies in several countries suggest that awareness of medicines-related cognitive services among the general public is limited. In Australia and the US, both awareness and expectations of these services are low.¹¹ Being worried about medicines was a potential motivating factor for using the Australian HMR service,¹² whereas in Sweden people taking-up offers of pharmaceutical care services were "worried, vulnerable, and information-seeking".¹³ Australians considered improving ability to manage medicines, reducing concerns, and increasing knowledge were further motivating factors for having a HMR, but their expectations that the

review would achieve these was low.¹⁴ The Royal Pharmaceutical Society in the UK has highlighted the need for greater public awareness of pharmacy services.¹⁵

Most studies of medicines-related services provided by community pharmacists have obtained views of service users rather than the public.¹⁶⁻¹⁸ No work has determined views on people's willingness to make appointments, how much time they are willing to spend discussing their medicines with a pharmacist, and the use of consultation rooms for medicines-related services, although leaflets promoting services mention the need to book a consultation with a pharmacist.¹⁹ Moreover, no studies have compared the views of the public with pharmacists' perceptions of these English medicine services.

This study therefore aimed to obtain the perceptions and experiences of both pharmacists and the general public on medicines-related services in general and specifically on MUR and NMS services, and compare pharmacist and public perceptions.

Methods

Two surveys were conducted in parallel, one involving the general public and the other community pharmacists, between September and December 2012, following approval from Medway School of Pharmacy Research Ethics committee. Members of the public provided verbal consent and the pharmacist questionnaire contained a statement that consent was implied by its return.

Questionnaire development and piloting

Public questionnaire

A previously validated questionnaire seeking public views on pharmacy public health services²⁰ was adapted using findings from focus groups seeking public views on medicines-related services, together with relevant literature and the team's experience.¹⁷ It included mostly closed questions covering: use of pharmacies, medicines and of medicines-related services, awareness of services, expectations and willingness to use services, reasons for using services and perceived benefits, as well as questions covering specific aspects, such as data sharing, appointments, and waiting times ([Supplementary materials](#)). Open questions sought additional reasons for using services and not using pharmacies to seek advice about medicines. Demographic data included gender, age, ethnicity, educational level, and postcode for assessment of deprivation status.

Pharmacist questionnaire

This was developed using findings from focus groups that included both the public and community pharmacists,

ensuring overlap with the public questionnaire to enable comparisons. The questionnaire covered pharmacists' views and experiences of providing medicine advisory services and their perceptions of the aspects included in the public questionnaire ([Supplementary materials](#)). Open-ended questions sought views on reasons people may use services and whether different services are needed. Demographic data included gender, years qualified, role in pharmacy, pharmacy type and location, and estimated MUR and NMS service delivery.

Piloting

For the public survey, 25 members of the public known to researchers completed the questionnaire using an interviewer-assisted method, recording time for completion, and questioning about ease of use and understanding. Five pharmacists known to the team, working outside the study area, were asked to complete and return the pharmacist questionnaire by post, with additional questions to assess the ease of use. Minor amendments were made to both questionnaires in the light of comments received.

Recruitment and data collection

Public survey

This survey was conducted using interviewer-assisted completion by ten students who had received training to ensure a consistent approach. The questionnaires were completed face-to-face with members of the public recruited at high street locations in ten towns in one county in South East England (Kent). The towns were selected to include subjects from differing socioeconomic and deprivation status. A quota sampling method, based on 2011 census data, was used to ensure that the sample was representative of the county in terms of gender and age, with a target of 100 participants per town. Passers-by were approached by a researcher and invited to participate. Initial screening questions excluded people <18 years of age and qualified or trainee health care professionals. The number who declined to complete the survey was recorded.

Pharmacist survey

This was administered as a postal questionnaire sent to all 836 community pharmacies in three counties in South East England (Kent, Surrey, and Sussex). Nonresponders received a second mail and a telephone call with a further copy of the questionnaire, if requested.

Data analysis

Data were analyzed using SPSS v22 (IBM Corporation, Armonk, NY, USA). Use of medicines by the public was

dichotomized into any or none, and frequency of pharmacy use was dichotomized into frequent (at least once per month) or infrequent (less than once a month/never). Chi-squared tests were used to assess differences in responses between pharmacists and public to similar questions and to evaluate the effect of medicines and pharmacy use on experiences of and willingness to use services. Missing data were excluded from analysis. Due to the large number of comparisons made, a P -value of <0.001 was regarded as statistically significant. Free-text responses to open-ended questions were categorized and quantified.

Results

Response rates and demographic characteristics

The response rate for the public questionnaire was 47.2% (1,000 from 2,012 people approached) and 40.8% (341) for the pharmacist questionnaire after second mailing and telephone reminder. The public were reasonably representative of the population of Kent in terms of gender, age, ethnicity, and deprivation status: 52.7% (526) were females, 21.8% (218) were aged ≥ 65 years, 712 (72.3%) were white, and there was an adequate distribution across five deprivation quintiles. Demographic details are shown in Table 1, along with use of pharmacies, medicines, and general health status. Regular pharmacy use was associated with both older age and regular medicines use ($P < 0.001$).

Pharmacist respondent characteristics are shown in Table 2, plus self-reported estimated frequency of MUR and NMS provision in the previous month.

Experiences of medicines-related services

The most common reasons selected by the public for using a pharmacy were to have a prescription dispensed (850; 85.1%) or to buy medicines (791; 79.2%), but getting advice on medicines or minor health problems was also selected by over two thirds (664; 66.5% and 660; 66.1%, respectively).

Experiences of different ways of receiving medicines-related advice and sharing of information are shown in Table 3, together with willingness to accept these aspects of services in future. The most frequent way in which advice about medicines received was across the counter, with only 28.8% (288) indicating they had received advice about medicines collected in a private consultation room and only 19.4% (194) for a new medicine (indicating experience of the NMS). Only one-quarter (248; 24.9%) had experienced a review of all their medicines in a private room (indicating experience

Table 1 Demographic details of public respondents

Characteristic	Number (%)	Kent population (%)
Sex (n=999)		
Female	526 (52.7)	51.6*
Male	473 (47.3)	48.4*
Age group (n=1,000)		
<25	140 (14.0)	26.9*
25–34	140 (14.0)	
35–44	170 (17.0)	51.0*
45–54	175 (17.5)	
55–64	157 (15.7)	
65 and over	218 (21.8)	22.1*
Ethnicity (n=985)		
White	712 (72.3)	93.0**
Asian	112 (11.4)	3.6**
Black	90 (9.1)	1.3**
Mixed	62 (6.3)	1.6**
Other	9 (0.9)	0.5**
Deprivation status (n=920)		
1 (highest)	157 (17.1)	20.5
2	166 (18.0)	20.2
3	172 (18.7)	19.9
4	223 (24.2)	19.8
5 (lowest)	202 (22.0)	19.7
Educational level (n=992)		n/a
None/primary/secondary	314 (31.7)	
Further education	315 (31.8)	
Bachelor/higher degree	319 (32.2)	
Current student	44 (4.4)	
Employment status (n=985)		n/a
Full-time employed	411 (41.7)	
Part-time employed	196 (19.9)	
Retired	212 (21.5)	
Not working	166 (16.9)	
Use of prescribed medicines (n=1,000)		n/a
None	395 (39.5)	
≤4	376 (37.6)	
5–8	172 (17.2)	
>8	57 (5.7)	
Use of pharmacies (n=999)		n/a
More than once a month	136 (13.6)	
Once a month	373 (37.3)	
Once every 2–3 months	258 (25.8)	
Less than every 3 months	91 (9.1)	
Never use/do not know	141 (14.1)	

Notes: *Derived from Kent and Medway 2011 Census data of population aged ≥18 years. **Derived from Kent and Medway 2011 Census data of all ages.

Abbreviation: n/a, not available.

of the MUR), with few having had reviews following hospital discharge or because they used a specific medicine. A total of 136 respondents (13.7%) indicated that they had experienced receiving both advice about new medicines and review of all medicines in a private room.

Most pharmacists provided information on the frequency of providing MUR and NMS, with the former being provided

Table 2 Demographic details of pharmacist responders and pharmacies

Characteristic	Number (%)	National data derived from registers (%)
Sex (n=338)		
Female	179 (53.0)	Female 40.6
Male	159 (47.0)	Male 59.4
Years qualified (n=325)		
≤3	78 (24.0)	Registrants in last 3 years/ total register 18.8
4–10	116 (35.7)	Registrants in last 9 years/ total register 29.3
>10	131 (40.3)	Remaining registrants 51.9
Role in pharmacy (n=340)		
Manager/sole pharmacist	269 (79.1)	Owner/employee 76
Second pharmacist	22 (6.5)	
Superintendent	13 (3.8)	
Locum	36 (10.6)	Locum 24
Type of pharmacy (n=340)		
Large chain (≥31 pharmacies)	223 (65.6)	National multiple 55
Medium chain (11–30)	19 (5.6)	Other multiple 15
Small chain (2–10)	33 (9.7)	Chain of ≤4 pharmacies 29
Single pharmacy	65 (19.1)	
Location of pharmacy (n=338)		n/a
High street/ suburban	116 (34.3)	
Shopping precinct/ out of town center	37 (10.9)	
Rural/village	80 (23.7)	
Supermarket	31 (9.2)	
Attached to GP surgery	49 (14.5)	
Other	25 (7.4)	
Number of MURs in the previous month (n=299)		
0	15 (5.0)	95% of CPs provide MURs
1–10	36 (12.0)	
11–20	44 (14.7)	
21–35	95 (31.8)	
36–50	75 (25.1)	
≥51	34 (11.4)	
Number of NMS in the previous month (n=303)		
0	39 (12.9)	80% of CPs provide NMS
1–10	179 (59.1)	
11–20	56 (18.5)	
21–35	24 (7.9)	
≥36	5 (1.7)	

Abbreviations: CP, community pharmacy; GP, general practitioner; n/a, not available; NMS, New Medicines Service; MURs, Medicine Use Reviews.

more frequently (Table 2). The majority of the pharmacists providing data (275/294; 93.5%) indicated that they had conducted some target MURs in the past month: 258 (87.8%) to people taking high-risk medicines and 251 (85.4%) to those using respiratory medicines, but only 75 (25.5%)

Table 3 Experiences of and willingness to use medicines-related services among the public

Service/aspect of service	Had experience of MRS, n (%)	Would be prepared to use MRS in the future, n (%)	
		Yes	Maybe
Received advice concerning medicines just collected			
In private consultation room	288 (28.8)	695 (69.6)	216 (21.6)
In quiet area of pharmacy	325 (32.5)	732 (73.3)	203 (20.3)
Across pharmacy counter	708 (70.9)	820 (82.1)	132 (13.2)
Received advice concerning new medicine just collected			
In private consultation room	194 (19.4)	700 (70.1)	215 (21.5)
In quiet area of pharmacy	213 (21.3)	732 (73.3)	201 (20.1)
Across pharmacy counter	510 (51.0)	801 (80.2)	146 (14.6)
Had a discussion with a pharmacist:			
Because recently discharged from hospital	102 (10.2)	655 (65.6)	219 (21.9)
Because were taking particular medicine	230 (23.0)	710 (71.1)	226 (22.6)
A general review in a private consultation room	248 (24.9)	685 (68.9)	233 (23.4)
Given pharmacist permission to:			
Telephone you to follow-up about advice already provided	111 (11.1)	544 (54.5)	235 (23.5)
Share your information with local NHS	116 (11.6)	553 (55.4)	232 (23.2)
Share your information with your doctor	345 (34.6)	808 (81.0)	132 (13.2)

Abbreviations: MRS, medicines-related service; NHS, National Health Service.

targeted people recently discharged from a hospital. The median time taken to provide a MUR was estimated at 10 minutes (range 2–40; n=306), median time to recruit to the NMS was 5 minutes (range 1–20; n=298), first consultation 8 minutes (1–30; n=264), and follow-up consultation 5 minutes (1–20; n=262).

Willingness to use services

Over 90% of the public showed willingness to use services in future (Table 3), but the proportion favoring receiving advice across the counter was higher than for a private room. Willingness to have a discussion with a pharmacist because of recent hospital discharge was lower than other reasons for doing so, and while 94.2% (940) were willing to allow sharing of information with their doctor, only 78.6% (785) indicated their willingness to allow data sharing with the local National Health Service and 78.0% (779) to allow a pharmacist to provide telephone follow-up. For all the aspects of services, respondents who were frequent pharmacy users and those using regular medicines were significantly more willing to agree than those not using pharmacies or medicines regularly (all $P < 0.001$).

A majority of the public (717; 71.8%) indicated their willingness to make an appointment for a medicines-related service and 72.0% (718) to wait for 10 to 15 minutes to see a pharmacist (Table 4). In contrast, significantly fewer pharmacists considered that people would be willing to make appointments (161; 47.2%; $P < 0.001$) and the majority (191; 56.0%) felt that people would wait no more than

5 minutes. However, perceptions concerning discussion time with a pharmacist were more closely aligned, with 59.0% of pharmacists (193) and 47.8% (477) of the public viewing no more than 15 minutes as acceptable.

Reasons for using pharmacy medicines-related services

There were 167 members of the public who indicated that they were worried about medicines-related problems for

Table 4 Pharmacist and public perceptions concerning aspects of service provision

Perceived willingness concerning:	Pharmacists, n (%)	Public, n (%)	P-value
Time waiting to talk to the pharmacist			
No more than 5 minutes	191 (56.7)	196 (19.6)	<0.001
No more than 10 minutes	111 (32.9)	370 (37.1)	
No more than 15 minutes	32 (9.5)	348 (34.9)	
More than 15 minutes	3 (0.9)	83 (8.3)	
Making an appointment to talk to the pharmacist (% indicating yes)	161 (48.1)	717 (71.8)	<0.001
Time waiting for an appointment			
No more than 4 hours	47 (15.1)	137 (13.8)	<0.001
No more than 1 day	86 (27.6)	379 (37.9)	
No more than 4 days	45 (14.4)	168 (16.9)	
No more than 1 week	104 (33.3)	259 (26.0)	
More than 1 week	30 (9.6)	52 (5.2)	
Time spent talking to pharmacist			
No more than 5 minutes	44 (13.5)	168 (16.7)	0.01
No more than 15 minutes	193 (59.0)	477 (47.8)	
No more than 30 minutes	64 (19.6)	216 (21.6)	
More than 30 minutes	26 (8.0)	137 (13.7)	

themselves or people they cared for. The most frequent cause for worry was side effects, cited by 78, followed by difficulties in remembering dosage regimens (23), concerns over lack of efficacy (13), and the overall number of medicines and the need for these (14). Over two-thirds of respondents (690) indicated that they would consider going to a pharmacy for advice if they did experience problems with a medicine. The majority of the remainder (251) indicated that they would go to their general practitioner (GP) instead. Reasons for choosing to see a GP were given as: greater GP knowledge of them personally, and health or illness in general (13), pharmacist not being able to help/change medicines (11), and expectation of being referred to GP by pharmacist (8).

Potential reasons for choosing to use a formal service (MUR or NMS) were offered to both groups (Figure 1). Almost all pharmacists thought the main reason for using one of these services would be “because the pharmacist asked” (94.9%), which was selected by proportionately fewer members of the public (68.7%). Significantly fewer pharmacists (57.5%) than members of the public (76.0%) thought people would use a service “because they had problems with their medicines.” “Wanting to help the pharmacist out” was selected as a reason by 53.2% of the public, compared with 22.6% of pharmacists.

While most (274; 80.4%) pharmacist respondents considered that patients needed medicines advisory services, only 186 (55.2%) felt they were wanted. Sixty pharmacists (17.6%) thought that different or additional services or changes to the existing formal services were needed.

Among these, eight suggested services needed were monitoring of patients for blood pressure, INR, and other parameters, and four public health services. Eight pharmacists suggested increasing the scope of the NMS to all medicines and 18 suggested changes to the MUR service, which included increased GP involvement, facilitating domiciliary MURs, and ensuring that services are based on need. Several pharmacists also felt that patients may not perceive benefits from the current services. Examples of comments are given in Table 5.

Awareness of MUR and NMS

Pharmacists considered that people would be more aware of the MUR service than the NMS: 132 (38.7%) pharmacists thought that at least one-half of their customers would have heard of the MUR, but only 16 (4.7%) felt this about the NMS. Actual awareness among the public was indeed low, with only 18.2% (182) having heard of the MUR when it was described to them and 8.6% (86) of the NMS. Awareness of both services was significantly higher among people who indicated that they had experienced an MUR (52.0%; 129 experiencing review of all medicines vs 7.1%; 53 not experiencing) or an NMS (29.9%; 58 in those experiencing vs 3.5%; 28 not experiencing). Respondents with experience of an MUR were also more aware of the NMS than those without it (18.5% 46 vs 5.1% 38). Regular medicine users were more aware of both the MUR (147; 24.3%) and NMS (62; 10.2%), compared with 9.1% (36) and 6.1% (24) in nonmedicine users, respectively.




Figure 1 Potential reasons for using medicines-related service identified by pharmacist and the public.
Notes: *Difference between pharmacist and public views; $P < 0.001$ χ^2 test.

Table 5 Examples of views expressed by pharmacist and public respondents on medicine advisory services

Respondent	Comment	Characteristics
Pharmacist views	There is little connection between prescriber/pharmacist with both MUR and NMS. It would be better with MURs if patients were required to have them prior to a clinical review by GP and then prescriber could undertake their review with feedback information to hand. With the NMS a formal direction to enter the service should be made by prescribers and pharmacists should feed back to the prescribers postintervention	Male manager/sole pharmacist in large multiple pharmacy
	MURs and NMS are now a means for increasing pharmacy revenue, we are hounded daily to do MURs and NMS by head office, we are told to do easy ones that don't take long, they are no longer about patient's needs but how much we get for them!	Female locum in small chain pharmacy
	At the moment MUR and NMS very rarely lead to optimization of medicine management and patients often question what was the point of it	Female manager/sole pharmacist in medium chain pharmacy
Public views	Can trust pharmacists more after these experiences. More knowledge – easier access for help than GP	Male, 34 or younger, on no regular medicines
	A pharmacist has more knowledge than I thought and seems more willing to help and is not as patronizing as some GPs	Male, 65 or over, using more than eight medicines
	Pharmacists do a lot more than before, they do not just dispense but they also make sure that I take my medication correctly and ask how I feel.	Female, 65 or over, using up to four medicines
	The pharmacist also makes sure that I visit the doctor when I need to	

Abbreviations: GP, general practitioner; MUR, Medicine Use Review; NMS, New Medicines Service.

Fewer than half the pharmacists (46.3%; 155) thought that patients agreeing to undergo an MUR knew its purpose, whereas more than one-half thought that this was the case for the NMS (57.9%; 194).

Expectations and experiences of benefits from medicines-related services

Expectations of benefit from receiving medicines services among the public were relatively high, with overall >70% indicating that they would expect general benefit, that the pharmacist would help them know more about their medicines, understand how to use their medicines better, and encourage them to use medicines as prescribed (Table 6). Willingness to use both the MUR and NMS was positively associated with expectation of general benefit, as well as willingness to make an appointment and to wait longer to speak to the pharmacist ($P<0.001$). A higher proportion of

pharmacists than the public were positive about all expected benefits, except for increasing adherence, which was perceived as an expected benefit by proportionately more of the public. Expectations related to sorting out problems and reducing the number of medicines/waste were relatively low in both groups.

Both groups perceived actual benefits attained to be lower than potential benefits, with pharmacists again being more likely to perceive benefits than the public, even among regular medicine users (Table 6).

There were 689 respondents who had experienced a benefit from a pharmacy medicine service, 28.0% (193) of whom considered their relationship with the pharmacist changed as a result, 34.6% (218) felt that their view of pharmacists as health care professionals changed, and 35.4% (244) felt that their awareness of pharmacists' knowledge about medicines changed, with the remainder indicating no change.

Table 6 Public and pharmacist expectations and experiences of medicines-related services

Medicines reviews with a pharmacist help people to:	Expectations*		Experiences**		
	Public (%)	Pharmacists (%)	All public (%)	Regular medicine users (%)	Pharmacists (%)
Benefit in general	781 (78.2)	293 (89.3)	550 (55.3)	428 (71.2)	265 (83.9)
Know more about medicines	729 (72.9)	275 (83.1)	441 (44.2)	343 (56.8)	236 (73.8)
Understand better how to use medicines	757 (75.7)	282 (86.0)	480 (48.2)	369 (61.3)	270 (84.1)
Encourage medicines use as doctor expects	774 (77.5)	230 (70.1)	515 (51.7)	396 (65.8)	185 (57.5)
Sort out problems	612 (61.2)	261 (79.3)	298 (29.9)	243 (40.4)	223 (70.1)
Order fewer medicines/reduce waste	441 (44.1)	175 (53.2)	158 (15.8)	138 (22.9)	135 (42.2)

Notes: *Differences between pharmacist and public expectations were all significant ($P<0.001$). **Differences between pharmacist and public experiences were significant ($P<0.001$) with the exception of encouraging medicine use as per doctor expectations.

In total 260 respondents provided additional comments on their experiences of services, 75 of whom indicated that their appreciation of what a pharmacist could offer had changed as a result (Table 5). Many respondents also indicated that they would use a pharmacist more readily, offering reasons such as ease of access and greater knowledge about medicines than GPs. Users of services showed increased awareness of pharmacists' knowledge and potential to benefit them personally, but there was also greater awareness that pharmacists can help with using medicines more than in the past.

Discussion

The majority of the public have received advice about medicines in community pharmacies across the counter and are happy with this, whereas few have received advice in a private consultation room. Both pharmacist and public surveys indicated that MURs are provided more often than NMS, in line with national statistics. The public were willing to use these services, although there was less willingness to allow telephone follow-up, an option for NMS follow-up consultations. Both pharmacists and the public reported that MURs after hospital discharge were infrequent, and willingness to use this service among the public was also low. The time taken to provide an MUR was estimated by pharmacists as ~10 minutes and NMS as 12 minutes, which align with both pharmacist and public perceptions of how much time people are willing to spend talking to a pharmacist about their medicines.

The results confirm pharmacist-driven service provision, almost all pharmacists, and nearly 70% of the public indicating pharmacist invitation as a reason for using formal services. It is concerning that over half the public saw wanting to help the pharmacist as another reason for accepting an invitation and that over half the pharmacists considered patients did not know the purpose of the MUR when they agreed to it. Awareness of both MUR and NMS was low among the public, even in regular medicine users, confirming previous qualitative work,^{16,17} hence the need for pharmacists to invite people to use services. The low uptake of postdischarge MURs and willingness to use this service has been shown previously,²¹ and efforts are being made to increase this, involving hospital pharmacists.²²

Although around two-thirds of people would consider going to a pharmacy for advice about medicines-related problems, a substantial proportion prefer to talk to their GP, primarily due to low expectations that the pharmacist would be able to resolve problems. This was supported by infrequent experiences among the public of problems being resolved, in contrast to pharmacist views, most of whom

perceived services to benefit recipients. Our data suggest that formal cognitive services can change peoples' perceptions of pharmacists and also that using one service may increase awareness, thus potential use, of other services. Consultations with a pharmacist increase the likelihood of seeking advice again, although, as here, most people prefer to discuss medicines with their doctor.²³

The formal services in England were developed with little involvement of patients or public and are aimed primarily at enhancing patient knowledge, understanding, and adherence and, for the MUR, reducing waste. Improving adherence and reducing waste/reordering were, however, considered by both groups as least likely to be achieved. Our data, which concur with the Australian experience,¹² suggest that worries about side effects are among the most common concerns people have about medicines, thus perhaps greater emphasis should be placed on this aspect of medicines during consultations. Relatively little research has explored what people really want from pharmacist-provided medicines-related services, most studies determine views on existing services, and few compare consumer and pharmacist views.²⁴⁻²⁷ As found here, Australian work shows low public awareness of medicines-related services²⁴ and differences in views.²⁵ Australian pharmacists overestimated the importance of being available for consultations and providing advice on minor ailments. Our work showed that pharmacists underestimated the willingness of the public to both wait to speak to them and make appointments for services. Misunderstanding of pharmaceutical care services between pharmacists and patients and divergence in perceived benefits, as well as in frequency of information provision, have also been found.^{26,27}

More work is needed to determine what services are desired by patients and the public and how they should be delivered. Our study suggests that pharmacists providing formal services should determine the willingness of potential users to make appointments and make use of this to engage them more effectively in preparing for their consultation, perhaps by providing leaflets in advance. This could ensure that those who take up the services are those who really perceive a potential for benefit or have problems they wish to discuss with a pharmacist and thus may lead to increased user experiences of benefits.

Strengths and limitations

Both surveys were developed from the findings of qualitative work, however no psychometric testing was carried out. Both achieved relatively good response rates in comparison with other studies involving these groups, although the

pharmacist survey required two reminders. The surveys were administered in the same area of England at the same time and the respondents were reasonably representative of their respective populations. Equivalent questions were used to enable comparisons between pharmacists and public. Data on service provision, obtained to assess sample representativeness, were self-reported. Both surveys may be subjected to recall bias and in addition, the public survey was interviewer-assisted, so may include an element of obsequiousness bias.

Conclusion

The public is willing to use community pharmacy medicines-related services, but many people prefer to discuss problems with a doctor. Perceptions of pharmacists and the public differed with respect to reasons for using services, willingness to wait to see a pharmacist and to make appointments, as well as expected and perceived benefits of services. For services to improve, the pharmacy profession needs a better awareness of what the public, especially those with potential to benefit from services, view as acceptable and desirable. Professional pharmacy organizations and pharmacists themselves also need to make use of these findings to encourage greater uptake of services through promotion and explanation of their potential benefits.

Disclosure

The authors report no conflicts of interest in this work.

References

1. Health and Social care Information Centre. Available from: <http://www/hscic.gov.uk>. Accessed April 21, 2015.
2. Latif A, Boardman H. Community pharmacists' attitudes towards medicines use reviews and factors affecting the numbers performed. *Pharm World Sci*. 2008;30(5):536–543.
3. Latif A, Pollock K, Boardman H. The contribution of the Medicines Use Review (MUR) consultation to counselling practice in community pharmacies. *Patient Educ Couns*. 2011;83(3):336–344.
4. Krska J, Nesbit J, Baylie K, O'Kane A. Patient views on the MUR service. *Int J Pharm Pract*. 2009;17(Suppl 1):A41–A42.
5. Iqbal S, Wood K. Exploring patient opinions of MURs. *Int J Pharm Pract*. 2011;18(Suppl 2):20.
6. Youssef S. Do patients perceive benefit from a structured warfarin MUR? *Int J Pharm Pract*. 2009;17(Suppl 2):B62–B63.
7. Latif A, Boardman H, Pollock J. Reasons involved in selecting patients for a MUR: exploring pharmacist and staff choices. *Int J Pharm Pract*. 2011;19(Suppl 1):31–32.
8. Wilcock M, Harding G. What do pharmacists think of MURs and do they change prescribed medication? *Pharm J*. 2008;281:163–167.
9. Bissell P, Blenkinsopp A, Short D, Mason L. Patients' experiences of a community pharmacy-led medicines management service. *Health Soc Care Community*. 2008;16(4):363–369.
10. Elliott R, Boyd M, Waring J, et al. Understanding and appraising the New Medicines Service in the NHS in England (029/0124). Available from: <http://www.nottingham.ac.uk/~pazmjb/nms/downloads/report/index.html#2/z>. Accessed October 2, 2015. (Archived by WebCite® at <http://www.webcitation.org/6bydND1cL>.)
11. McMillan S, Kelly F, Sav A, King MA, Whitty JA, Wheeler AJ. Consumer and carer views of Australian community pharmacy practice: awareness, experiences and expectations. *J Pharm Health Serv Res*. 2014;5:29–36.
12. Carter SR, Moles R, White L, Chen TF. The effect of experiencing medication-related worry and the social influence of the general practitioner on consumer willingness to use a pharmacist-provided medication management service. *Res Social Admin Pharm*. 2012;8(6):e21.
13. Montgomery AT, Kälve mark Sporrang S, Manap N, Tully MP, Lindblad AK. Receiving a pharmaceutical care service compared to receiving standard pharmacy service in Sweden how do patients differ with regard to perceptions of medicine use and the pharmacy encounter? *Res Social Admin Pharm*. 2010;6(3):185–195.
14. Carter SR, Moles RJ, White L, Chen TF. Patients' willingness to use a pharmacist-provided medication management service: The influence of outcome expectancies and communication efficacy. *Res Social Admin Pharm*. 2012;8(6):487–498.
15. Smith J, Picton C, Dayan M. *Now More than Ever: Why Pharmacy Needs to Act*. London: Royal Pharmaceutical Society; 2014.
16. Kember J, Hodson K, James DH. Public perceptions of community pharmacy: a qualitative study. *Int J Pharm Pract*. 2013;21(Suppl 2):50.
17. Corlett SA, Dodds LJ, Rodgers RM. As pharmacists expand their clinical roles do the general public recognise what they can offer? *Pharmacoepidemiol Drug Saf*. 2013;22(6):672–673.
18. Mackridge AM, Marriott JF. Information-related roles and pharmacists: public opinion. Abstract presented at: Health Services Research and Pharmacy Practice conference; April 3–4, 2006; Bath, UK.
19. Van den Berg M, Donyai P. How was patient empowerment portrayed in information leaflets describing the community pharmacy Medicines Use Review service in the UK? *Patient Educ Couns*. 2010;80(2):274–276.
20. Saramunee K, Krska J, Mackridge AJ, Richards J, Suttajit S, Philips-Howard P. General public views on pharmacy public health services: current situation and opportunities in the future. *Pub Health*. 2015;129(6):705–715.
21. Lam MYY, Dodds L, Corlett SA. Poor uptake of dMURs is not the fault of hospital pharmacy services. *Pharmacoepidemiol Drug Saf*. 2016;25:12–13.
22. Robinson S. Hospital e-referral initiative boosts post-discharge MURs in community pharmacies. *Pharm J*. 2015;295.
23. Tinelli M, Bond C, Blenkinsopp A, Jaffray M, Watson M, Hannaford P. On the behalf of the community pharmacy medicines management evaluation team. Patient evaluation of a community pharmacy medications management service. *Ann Pharmacother*. 2007;41(12):1962–1970.
24. McMillan SS, Wheeler AJ, Sav A, et al. Community pharmacy in Australia: a health hub destination of the future. *Res Social Admin Pharm*. 2013;9(6):863–875.
25. McMillan S, Kelly F, Sav A, King MA, Whitty JA, Wheeler AJ. Australian community pharmacy services: a survey of what people with chronic conditions and their carers use versus what they consider important. *BMJ Open*. 2014;4(12):e006587.
26. Assa-Eley M, Kimberlin CL. Using interpersonal perception to characterize pharmacists' and patients' perceptions of the benefits of pharmaceutical care. *Health Commun*. 2005;17(1):41–56.
27. Puspasari HP, Aslani P, Krass I. Pharmacists' and consumers' viewpoints on counselling on prescription medicines in Australian community pharmacies. *Int J Pharm Pract*. 2010;18(4):202–208.
28. Saramunee K, Krska J, Mackridge AJ, Richards J, Suttajit S, Philips-Howard P. Methodological and economic evaluations of eight survey modes applied to health service research. *J Pharm Health Serv Res*. 2016;7(1):43–52. Available from: <http://onlinelibrary.wiley.com/doi/10.1111/jphs.12121/abstract>. Accessed August 16, 2016.

Patient Preference and Adherence

Dovepress

Publish your work in this journal

Patient Preference and Adherence is an international, peer-reviewed, open access journal that focuses on the growing importance of patient preference and adherence throughout the therapeutic continuum. Patient satisfaction, acceptability, quality of life, compliance, persistence and their role in developing new therapeutic modalities and compounds to optimize

clinical outcomes for existing disease states are major areas of interest for the journal. This journal has been accepted for indexing on PubMed Central. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <http://www.dovepress.com/patient-preference-and-adherence-journal>