













Submitted: 09/05/2022

Accepted: 25/07/2022

Published: 25/08/2022

***In vitro* antimicrobial activity, antibioresistance reversal properties, and toxicity screen of ethanolic extracts of *Heracleum mantegazzianum* Sommier and Levier (giant hogweed), *Centaurea jacea* L. (brown knapweed), and *Chenopodium album* L. (Pigweed): Three invasive plants**

Mbarga Manga Joseph Arsene^{1*} , Podoprigrora Irina Viktorovna¹ , Mefed Kirill Mikhaïlovitch¹ , Anyutoulou Kitio Linda Davares¹ , Kezimana Parfait² , Manar Rehailia² , Senyagin Alexander Nikolayevich¹ , Girich Valentina Stefanovna¹ , Souadkia Sarra¹ , Khabadze Zurab Sulikoevich³ , Chernaia Zoya Anatolyevna⁴  and Das Shommiya⁵ 

¹Department of Microbiology named after V.S. Kiktenko, Institute of Medicine, RUDN University, Moscow, Russia

²Department of Agrobiotechnology, Agrarian Institute, RUDN University, Moscow, Russia

³Department of Therapeutic Dentistry, RUDN University, Moscow, Russia

⁴Department of Microbiology and Virology, Pirogov Russian National Research Medical University, Moscow, Russia

⁵Department of Hospital Therapy with Endocrinology, Hematology and Clinical Laboratory Diagnostics courses, Institute of Medicine, RUDN University, Moscow, Russia

Abstract

Background: Plants, including invasive ones, can play a significant role in the fight against antibiotic resistance and the search for new antimicrobials.

Aims: The present study aimed at assessing the antimicrobial activity, antibioresistance reversal properties, and toxicity of four samples from invasive plants, namely, *Heracleum mantegazzianum* (leaves and flowers), *Chenopodium album* (leaves), and *Centaurea jacea* (flowers).

Methods: The extraction of active compounds was done with ethanol (80%, v/v) and the extraction yields were calculated. Antimicrobial activity was studied using the agar-well diffusion method against *Escherichia coli* ATCC 25922, *Staphylococcus aureus* ATCC 6538, and *Candida albicans* ATCC 10231. Minimum inhibitory concentrations (MIC) and minimum bactericidal concentrations (MBC) were determined using the microdilution method. The antibioresistance reversal properties were assessed using the checkerboard method and the toxicity of the extracts was studied using the larval form of the Greater Wax Moth (*Galleria mellonella*).

Results: The mass yields were 11.9, 15.0, 18.2, and 21.5, respectively, for *C. jacea* flower (CJF), *H. mantegazzianum* flower (HMF), *H. mantegazzianum* leaf (HML), and *C. album* leaf (CAL). The highest inhibition diameters (ID) were found with HMF, CAL, CJF, and HML against *S. aureus* with 26.6, 21.6, 21.0, and 20.0 mm, respectively. Only CJF and HMF were active against *E. coli* with respective ID of 15.3 and 19.0 mm. Except HMF (ID = 13.6 ± 2.0 mm), no other extract was active against *C. albicans*. Moreover, HMF exhibited the lowest MIC (0.5 mg/ml) and the lowest MBC (1 and 4 mg/ml) against both *S. aureus* and *E. coli*. Regarding the synergy test, an additional effect [$0.5 \leq$ fractional inhibitory concentration (FIC) ≤ 1] was found in almost all the combinations antibiotics + extracts excepted for HMF + (Kanamycin or Ampicillin) against *S. aureus* and CJF + Ampicillin against *E. coli* where we found synergy effect (FIC ≤ 0.5). The median lethal doses (LD50s) of HMF, HML, CAL, and CJF were 20.2, 0.58, 13.2, and 4.0 mg/ml, respectively.

Conclusion: Only the ethanolic extract of HMFs showed noteworthy broad spectrum antimicrobial activity.

Keywords: Invasive plants, Antimicrobial activity, Synergy, Toxicity.

Introduction

As the world gradually recovers from the health crisis caused by COVID-19, countries should pay more attention to other silent epidemics, such as resistance to antibiotics, which is constantly growing in medicine, agriculture, and animal breeding (Souadkia *et al.*, 2021; Arsene *et al.*, 2022). Recent estimates have shown that antibiotic resistance is responsible for 700,000 annual

deaths worldwide, 230,000 of which have resulted from multidrug-resistant tuberculosis (WHO, 2019). The WHO estimates that if nothing is done to address this problem, drug-resistant diseases may cause 10 million deaths each year by 2050 and damage to the economy as catastrophic as the 2008–2009 global financial crisis (WHO, 2019).

Antibiotic resistance is defined as the ability of bacteria to resist the inhibitory or destructive activity of an

*Corresponding Author: Mbarga Manga Joseph Arsene. Department of Microbiology named after V.S. Kiktenko, Institute of Medicine, RUDN University, Moscow, Russia. Email: josepharsenembarga@yahoo.fr

antibiotic to which it was initially sensitive (Su *et al.*, 2020). It primarily results from the uncontrolled use of antibacterial drugs both in medicine and in agriculture, which leads to the recurrent exposure of bacteria to sub-lethal doses of antibiotics and results in their adaptation. This adaptation phenomenon results mainly from the enzymatic degradation of antibiotics by bacteria (Elekhawy *et al.*, 2020), the modification of the antibiotic target (Schaenzer and Wright, 2020), change in membrane permeability (Singh *et al.*, 2020), and alternative metabolic pathways (Singh *et al.*, 2020). Interbacterial transmission of antibiotic resistance through horizontal gene transfer (conjugation, transduction, and transformation) has made the situation critical worldwide (Su *et al.*, 2020). To respond to this dangerously worrying situation, research teams from all over the world are permanently evaluating the use of potential alternatives to antibiotics such as nanoparticles (Amer and Awwad, 2021), probiotics (Arsene *et al.*, 2021, 2021b), phage therapy (Ansald *et al.*, 2020), antimicrobial peptides (Andersson *et al.*, 2016), and medicinal plants (Arsene *et al.*, 2021c, 2021d; Mouafo *et al.*, 2021). Among these alternatives, herbal remedies have various advantages because of their availability, fewer reported side effects, cost, high tolerance toward patients, and lack of bacterial resistance (Wojnicz *et al.*, 2012).

In this context, all plants, including invasive ones like *Heracleum mantegazzianum* Sommier and Levier (giant hogweed), *Centaurea jacea* L. (brown knapweed) and *Chenopodium album* L. (Pigweed) deserve to be investigated for their antimicrobial properties. Indeed, among terrestrial plants, species that are invasive tend to be larger and have higher growth rate and shoot allocation than do species that are not invasive (Havel *et al.*, 2015). These characteristics could be an asset if the extracts of such plants prove to be antimicrobial since they are widespread and can be easily cultivable for an eventual large-scale exploitation.

Therefore, the present study aimed at assessing the antimicrobial activity, antibioresistance reversal properties, and the toxicity of the pigweed's and the brown knapweed's leaves, and the giant hogweed's leaves and flowers.

Materials and Methods

Vegetal material

The vegetal materials used in this study were flowers of *H. mantegazzianum* Sommier and Levier and *C. jacea* L., and leaves of *C. album* L and *H. mantegazzianum*. The three plants were collected in June 2021 in the city of Mozhaysk (55°35'02.6"N 35°51'55.5"E, Mozhaysky District, Moscow Oblast, Russia) after identifying the plants using the mobile professional version of PictureThis-Plant Identifier (Glority LLC, 2021). After the harvest, the plant materials were conducted to the laboratory, dried at 37°C until constant mass and dry matter content was calculated. Thereafter, the plants

were grinded and powders with a particle size of less than 1 mm were stored in a sterile airtight container until further use.

Bacterial strains

The microorganisms used for the screening of antimicrobial activity consisted of three standard strains. *Staphylococcus aureus* ATCC 6538 was used as Gram positive model, *Escherichia coli* ATCC 25922 as Gram negative model, and *Candida albicans* ATCC 10231 as fungi model. All the microorganisms were provided by the laboratory of microbiology of RUDN university.

Galleria melonella

Galleria mellonella larvae were obtained commercially from <https://ecobaits.ru/> (ECO BAITs, Moscow, Russia) and stored at 15°C prior to use. Dead larvae and those with dark spots or showing signs of melanisation were discarded.

Chemicals and media

Dimethyl sulfoxide (DMSO) was purchased from BDH Laboratories, VWR International Ltd., USA. We also used BHIB (Brain Heart Infusion Broth, HiMedia™ Laboratories Pvt. Ltd., India), Muller Hinton Agar (MHA HiMedia™ Laboratories Pvt. Ltd., India), Sabouraud Dextrose Broth (SDB, HiMedia™ Laboratories Pvt. Ltd., India), and all other reagents and chemicals used were of analytical grade.

Phytochemical extraction

As we described in our previous investigation ((Arsene *et al.*, 2021b), 30 grams of each vegetal material was weighed and added to 270 ml of the ethanolic solution (80%, v/v) in a conical flask. The flasks were covered tightly and were shaken at 300 rpm for 24 hours and 25°C in a shaker incubator (Heidolph Inkubator 1,000 coupled with Heidolph Unimax 1,010, Germany). The mixtures were then filtered using Whatman filter paper No 1 then concentrated at 40°C in rotary evaporator (IKA RV8) equipped with a water bath IKA HB10 (IKA Werke, Staufen, Germany) and a vacuum pumping unit IKA MVP10 (IKA Werke, Staufen, Germany). The final dried crude extracts were weighed. Extraction volume and mass yield were determined using the following formulas:

$$\text{Volume yield (\%)} = \frac{\text{Volume of the extract after filtration (ml)}}{\text{Initial solvent volume (ml)}} \times 100$$

$$\text{Mass yield (\%)} = \frac{\text{Mass of extracted plant residues (g)}}{\text{Mass of plant raw sample (g)}} \times 100$$

Preparation of antimicrobial solution

For each plant extract, the crude extract was dissolved in the required volume of DMSO (5%, v/v) to achieve a concentration of 64 mg/ml. The extracts were sterilized by microfiltration (0.22 µm; Merck Millipore, Tullagreen, Cork, Ireland) and the solutions obtained

was used to prepare the different concentrations used in the analytical process.

Inoculum preparation

Bacteria were cultured for 24 hours, 0 rpm, and at 37°C in 10 ml of BHIB while the yeast (*C. albicans* ATCC 10231) was cultured in the same volume of SDB and the same conditions (0 rpm, 24 hours at 37°C in 10 ml). After incubation, the cells were collected by centrifugation (7,000 g, 4°C, 10 minutes), washed twice with sterile saline (0.9%), resuspended in 5 ml of sterile saline to achieve a concentration equivalent to McFarland 0.5 using DEN-1 McFarland Densitometer (Grant-bio, Grant instruments Ltd., Cambridge, UK).

Screening of antibacterial activity

Assessment of antimicrobial activity using well diffusion method

The well-agar diffusion method described in our previous investigation (Arsene *et al.*, 2021b) was used to assess the antimicrobial activity of the extracts. Briefly, 15 ml of sterile Muller-Hinton Agar (for bacteria) or Sabouraud Dextrose Agar (SDA) (For *C. albicans*) were poured into petri dishes and 100 µl of each microorganism were spread. Wells with a capacity of 20 µl were drilled on the culture medium and 20 µl (at 100 mg/ml) of each plant extracts were added. The sterile DMSO (5%, v/v) used to prepare the extracts was used as negative control and all the trials was done in triplicate. After incubation at 37°C for 24 hours, the inhibition diameters (ID) were measured.

Determination of minimum inhibitory concentrations (MIC)

MIC is the lowest concentration of antibacterial agent that completely inhibits the visible bacterial growth. The MIC of the extracts was determined using the microbroth dilution method as described in our previous published work (Arsene *et al.*, 2021b). Briefly, 100 µl of broth (BHIB or SDB) was added to all the wells of sterile U-bottom 96-well microplates and extracts preparations (64 mg/ml) were subjected to serial twofold dilution. Each column represented one type of extract and a single strain. For each test well, 10 µl of the respective inoculum (with turbidity equivalent to a 0.5 McFarland scale) was added. Finally, the plates were covered and incubated at 37°C for 24 hours and after incubation, MIC was considered the lowest concentration of the tested material that inhibited the visible growth of the bacteria.

Determination of minimum bactericidal concentration (MBC)

MBCs were determined by subculturing the wells without visible growth (with concentrations \geq MIC) on MHA or SDA plates. Inoculated agar plates were incubated at 37°C for 48 hours and MBC was considered the lowest concentration that did not yield any microbial growth on agar.

Tolerance level

Tolerance level of tested bacterial strains against aqueous and ethanolic extract was determined using the following formula (Mondal *et al.*, 2020):

$$\text{Tolerance} = \frac{\text{MBC}}{\text{MIC}}$$

The characteristic of the antibacterial activity of extracts was determined by the tolerance level indicating the bactericidal or bacteriostatic action against the tested strains. When the ratio of MBC/MIC is ≥ 16 , the antibacterial efficacy of the test agent is considered as bacteriostatic, whereas MBC/MIC ≤ 4 indicates bactericidal activity (Mondal *et al.*, 2020).

Modulation of common antibiotic using the checkerboard

The checkerboard method, commonly used for the determination of synergy between the antibiotics and natural antibacterial compounds, was used for the antibiotic modulation assay (Arsene *et al.*, 2021b). Modulations of ampicillin (AMP), nitrofurantoin (NIT), and kanamycin (KA) were performed with extracts whose MIC was successfully determined. The fractional inhibitory concentration (FIC) index was calculated, as previously described (Trabelsi *et al.*, 2020; Arsene *et al.*, 2021b). Briefly, the individual MICs of the antibiotics (MIC-ATB) and the extract (MIC-extr) on the two targeted strains (*S. aureus* and *E. coli*) were first determined using the microdilution method as described above. Then, the new MIC values (MIC'-ATB and MIC'-extr) were determined after combining the two substances. Combinations of antibiotics + extracts were prepared by mixing the two antimicrobial solutions in 50:50 (v:v) proportions with initial concentrations depending on each stock solutions. To assess the interaction between the antibiotic and the natural extract, the FIC was determined using the using the formula:

$$\text{FIC} = \text{FICA} + \text{FICB},$$

$$\text{With: FICA} = \frac{\text{MIC}'\text{ATB}}{\text{MIC ATB}} \text{ and FICB} = \frac{\text{MIC}'\text{extr}}{\text{MIC extr}}$$

The FIC index was interpreted as follows FIC ≤ 0.5 , synergy; $0.5 \leq \text{FIC} \leq 1$, addition of effects; $1 \leq \text{FIC} \leq 4$, indifference and for FIC > 4 , Antagonism (Trabelsi *et al.*, 2020).

Toxicity assay

The toxicity was evaluated as previously described (Mbarga *et al.*, 2021a). First, each extract was dissolved in sterile phosphate-buffered saline (PBS) to obtain a concentration of 50 mg/ml which was then diluted to obtain concentrations of 40, 20, 10, 5, 2, 1, and 0.5 mg/ml. Furthermore, the larvae were

weighed and only those from 0.2 to 0.5grams were retained. For each concentration of extract, 3 groups of 20 randomly selected *G. mellonella* larvae were used (Fig. 1-A). 20 µl of each dilution were injected using a 0.3 ml Terumo® Myjector® U-100 insulin syringe (VWR, Russia) through the base of the last left proleg as we described in our previous study (Mbarga et al., 2021a), and shown in Figure 1-B. Control groups of 10 larvae injected with 20 µl sterile PBS were also included. After 24 hours of incubation in the dark at 37°C, larvae were examined for mortality, and were considered dead if they were unmoving, failed to reorient themselves when placed on their backs, and failed to respond to stimuli. Percentage survival was plotted as a function of concentration for each of the plants extract using Spline cubic model in the statistical software XLSTAT 2020 (Addinsoft Inc., New York, NY) and median lethal dose (LD50), 90% lethal dose (LD90) and the 100% lethal dose (LD100) values expressed in mg/ml were calculated using each specific spline cubic equation or spline cubic curves obtained. Spline curves were not used when the fit accuracy of the spline equation was less than 90%. The mean weight of each group of 20 larvae was used to extrapolate the LD50, LD90 and LD100 values for each plant extract in g/kg using the formulas:

$$\text{LD50(g/kg body weight)} = \frac{\text{Volume administered (ml)} \times \text{LD50(mg/ml)} \times 10^{-3}}{\text{Body weight (kg)}}$$

$$\text{LD90(g/kg body weight)} = \frac{\text{Volume administered (ml)} \times \text{LD90(mg/ml)} \times 10^{-3}}{\text{Body weight (kg)}}$$

$$\text{LD100(g/kg body weight)} = \frac{\text{Volume administered (ml)} \times \text{LD100(mg/ml)} \times 10^{-3}}{\text{Body weight (kg)}}$$

Ethical approval
Not applicable.

Results

Dry matter and extraction yield

As shown in Table 1, the highest dry matter content was observed on leaf of *C. album* (53.0%) and leaf of *H. mantegazzianum* (35.3%). Flowers of *C. jacea* and *H. mantegazzianum* consisted mainly of water with water contents of 79.3% and 72.5%, respectively. In addition, the volume yield was greater in leaves (89.7% and 81.8%) than in flowers (73.1% and 64.5%). The mass yields were 11.9%, 15.0%, 18.2%, and 21.5%, respectively, for *C. jacea* flower (CJF), *H. mantegazzianum* flower (HMF), *H. mantegazzianum* leaf (HML), and *C. album* leaf.

Antimicrobial activity

Antimicrobial activity was assessed both through ID using the agar well diffusion method and through determination of minimum MIC and MBC. The results of the antimicrobial activity of the extracts against *S. aureus* ATCC 6538, *E. coli* ATCC 25922, and *C. albicans* ATCC 10231 have been grouped in Table 2. All extracts were not active against all the microorganisms tested. Regarding the ID, we found that they ranged from 0.0 ± 0.0 to 26.6 ± 1.5 mm. The highest antimicrobial activity materialized with the highest IDs was observed against *S. aureus* with 26.6 ± 1.5, 21.6 ± 2.8, 21.0 ± 0.9 and 20.0 ± 1.4 mm, respectively, with HMF, *C. album* leaf (CAL), CJF and HML. Only CJF




Fig. 1. (A) Group of 20 larvae and (B) Injection of *G. mellonella* larvae with plant extract through the last left proleg.

Table 1. Dry matter and extraction yield of the plant materials used.

Plant	Dry matter (%)	Water content (%)	Volume yield (%)	Mass yield (%)
<i>H. mantegazzianum</i>	Leaf	35.3	64.7	81.8
	Flower	27.5	72.5	73.1
CJF	20.7	79.3	64.5	11.9
CAL	53.0	47.0	89.7	21.5

Table 2. MIC, MBC, and ID (at 50 mg/ml) of plant extracts.

	Microorganisms	MIC (mg/ml)	MBC (mg/ml)	MBC/MIC	ID (mm)
HML	<i>S. aureus</i> ATCC 6538	4	8	2	20.0 ± 1.4
	<i>E. coli</i> ATCC 25922	64	>64	-	0.0 ± 0.0
	<i>C. albicans</i> ATCC 10231	>64	>64	-	0.0 ± 0.0
<i>C. jacea</i> L. flower	<i>S. aureus</i> ATCC 6538	2	4	2	21.0 ± 0.9
	<i>E. coli</i> ATCC 25922	8	64	8	15.3 ± 1.1
	<i>C. albicans</i> ATCC 10231	>64	>64	-	0.0 ± 0.0
HMF	<i>S. aureus</i> ATCC 6538	0.5	1	2	26.6 ± 1.2
	<i>E. coli</i> ATCC 25922	0.5	4	8	19.0 ± 1.7
	<i>C. albicans</i> ATCC 10231	8	32	4	13.6 ± 2.0
CAL	<i>S. aureus</i> ATCC 6538	1	4	4	21.6 ± 2.8
	<i>E. coli</i> ATCC 25922	>64	>64	-	0.0 ± 0.0
	<i>C. albicans</i> ATCC 10231	>64	>64	-	0.0 ± 0.0

and HMFs were active against *E. coli* with respective ID of 15.3 ± 1.1 and 19.0 ± 1.7 mm. Except HMF (ID = 13.6 ± 2.0 mm), no other extract was active against *C. albicans*. Moreover, MIC and MBC values varied from 0.5 to >64 mg/ml and 1 to >64 mg/ml, respectively. All the extracts were more active on *S. aureus* (0.5 < MIC < 4 mg/ml) compared to their activity on other *E. coli*. Flowers of *H. mantegazzianum* exhibited the lowest MIC (0.5 mg/ml) and the lowest MBC (1 and 4 mg/ml) against both *S. aureus* and *E. coli*. Similarly, unlike other extracts which had indeterminate MIC (>64 mg/ml), only the MIC of HMF was successfully determined (8 mg/ml) against *C. albicans*. We were able to calculate the MBC/MIC ratio in 7/12 of the tests and this ratio varied from two to eight. The lowest MBC/MIC ratio (2) associated with the lowest MIC and MBC was observed with HMFs against *S. aureus*

Synergistic effect between common antibiotics and plant extracts using checkboard method

Table 3 shows the results of modulation of AMP, KA, and NIT with *H. mantegazzianum* leaf (HML), CJF, HMF and CAL against *S. aureus* and *E. coli*. We noted that the FIC ranged from 0.250 to 1.000. The lowest FIC (0.250) associated with the highest fold decrease (8) of the concentration of antibiotic was found with the combinations AMP-CJF and AMP-HMF against *S. aureus*. FICs ≥ 1 were observed in KA-CAL and NIT-HML combinations against *S. aureus* and with KA-CJF and AMP-HML against *E. coli*.

Toxicity assay

The median lethal doses (LD50), 90% (LD90) and 100% (LD100) were determined using the spline cubic survival curves (Fig. 2) from the data obtained on the survival rate of the Greater Wax Moth (*G. mellonella*) 24 hours after the injection with the extracts. As shown in Table 4, the LD50 values of HMF and CAL were 20.2 and 13.2 mg/ml, respectively. In addition, we found that leaves of *H. mantegazzianum* and flowers of *C. jacea* had the lowest LD. The LD50 (mg/ml) of HML was 0.58 (0.27 g/kg bw (body weight)) while that of CJF was 4.0 (2.28 g/kg bw). The concentrations of HMF and CAL capable of killing 100% of the larvae (LD100) were higher than the highest concentration tested (>50 mg/ml) while the LD100 (mg/ml) of the leaves of *H. mantegazzianum* and flowers of *C. jacea* were, respectively, 3.91 (1.73 g/kg bw) and 30.39 (19.21 g/kg bw).

Discussion

Invasive plants are mainly known to cause economic losses in agriculture (Pimentel *et al.*, 2005). This characteristic comes from the fact that they grow very quickly and sometimes with limited resources, even in hostile environments (Pintó-Marijuan and Munne-Bosch, 2013; Havel *et al.*, 2015). Thus, research devoted to these plants is very often aimed at finding effective means to combat them, while very few studies investigate their biological properties (Máximo *et*

Table 3. FIC of the combinations of extracts and antibiotics against *S. aureus* ATCC 6538 and *E. coli* ATCC 25922.

	<i>S. aureus</i> ATCC 6538						<i>E. coli</i> ATCC 25922								
	MIC extr (mg/ml)	MIC ATB (µg/ml)	MIC' extr (mg/ml)	MIC' ATB (µg/ml)	FIC	MIC extr (mg/ml)	MIC ATB (µg/ml)	MIC' extr (mg/ml)	MIC' ATB (µg/ml)	FIC	MIC extr (mg/ml)	MIC ATB (µg/ml)	MIC' extr (mg/ml)	MIC' ATB (µg/ml)	FIC
<i>H. mantegazzianum</i> leaf	AMP	4	4	1	1	0.500	64	32	32	16	1.000	64	32	32	1.000
	Ka		32	1	16	0.750		64	64	16	0.750		16	32	0.750
	NIT		16	2	8	1.000		32	32	8	0.750		8	16	0.750
CJF	AMP	2	4	0.25	0.5	0.250	8	32	32	2	0.375	64	4	4	0.375
	Ka		32	0.5	8	0.500		64	64	4	1.000		4	32	1.000
	NIT		16	0.5	4	0.500		32	32	2	0.500		2	8	0.500
<i>H. mantegazzianum</i> flower	AMP	0.5	4	0.0625	0.5	0.250	0,5	32	32	0.125	0.500	32	0.125	8	0.500
	Ka		32	0.125	4	0.375		64	64	0.125	0.500		0.125	16	0.500
	NIT		16	0.25	2	0.625		32	32	0.125	0.500		0.125	8	0.500
CAL	AMP	1	4	0.25	1	0.500	>64	32	32	-	-	64	-	-	-
	Ka		32	0.5	16	1.000		64	64	-	-		-	-	-
	NIT		16	0.5	4	0.750		32	32	-	-		-	-	-




Fig. 2. Survival curves for *G. mellonella* larvae against ethanolic extract of (A) HML, (B) CJF, (C) HMF, and (D) CAL. Each data point represents the mean percentage survival of three groups of 20 larvae, following injection with 20 μ l of specific concentrations of the selected plant extract and incubation for 24 hours at 37°C.

Table 4. LD50, LD90, and LD100 values for the tested plants using *G. mellonella* model.

Plant materials	LD50		LD90		LD100	
	mg/ml	g/kg of body weight	mg/ml	g/kg of body weight	mg/ml	g/kg of body weight
HML	0.58	0.27	2.31	1.34	3.91	1.73
CJF	4.0	2.28	18.10	10.58	30.39	19.21
HMF	20.2	11.45	>50.00	>40.34	>50.00	>37.93
CAL	13.2	6.94	>50.00	>38.19	>50.00	>39.72

al., 2020). Against the current backdrop of alarming growth in antibiotic resistance, invasive plants that demonstrate worthy antimicrobial activity can be of great use and their rapid growth could be an asset for their potential large-scale exploitation.

In the present study, we investigated the antimicrobial properties of four extracts from three plants recognized as invasive, namely, *H. mantegazzianum* (giant hogweed), *C. jacea* (brown knapweed), and *C. album* (Pigweed). First, as shown in Table 1, we found that

the dry matter content and the extraction mass yields were positively and significantly correlated (Pearson correlation coefficient = 0.974 and $p = 0.026$). Indeed, plants with a high percentage of dry matter (e.g., 53.0% for CAL) had the highest mass yield (21.5% for CAL) while, plants with the lowest dry matter percentage (20.7%) like CJF had the lowest mass yield (11.9%). The same dynamic was observed with CJF and HMFs. This observation is quite strange because we used the same mass of each plant (30grams) after drying them all. However, we can explain it by the tendency of plants to try to reabsorb (during extraction) the amount of water that they lost during drying (Bewley, 1973). This hypothesis was confirmed by observing the volume yields (Table 1), where we found that the plants with the highest dry matter content were associated with the highest volume yields while the plants with the lowest dry matter content were associated with lower volume yields.

Moreover, regarding the antimicrobial activity using well-agar diffusion method, HMF, CAL, CJF, and HML have demonstrated very good activity against *S. aureus* (with ID >20 mm), while only CJF and HMFs were active against *E. coli* with respective ID of 15.3 ± 1.1 and 19.0 ± 1.7 mm. On the one hand, the ID observed in this study showed that the active compounds involved in the antimicrobial activity of these plants can diffuse in aqueous media. Indeed, most of the time, one of the reasons why some plants do not show antibacterial activity is the relatively non-polar nature of the active compounds, which makes them very little diffuse in the aqueous agar matrix used in agar diffusion studies (Eloff, 2019). Moreover, this difference in the antimicrobial activity could be explained by the variability of phytochemical profile of the plants. In addition, the higher sensitivity of Gram-positive bacteria compared to Gram-negative bacteria could be explained by the differences in the composition of their cell wall (Acosta-Gutiérrez et al., 2020). Indeed, Gram-positive bacteria have an outer wall made essentially of peptidoglycan which is an ineffective permeability barrier (Acosta-Gutiérrez et al., 2020), while the outer phospholipid membrane of Gram-negative bacteria makes the outer layer impermeable to lipophilic solutes and constitutes a selective barrier to hydrophilic solutes. These observations were confirmed when determining MIC and MBC where the lowest MIC and MBC were observed on *S. aureus* compared to *E. coli* (Table 2).

These findings are different to that stated by Etame et al. (2019) who found no significant difference in the MIC values of plant extracts against Gram positive and Gram-negative bacteria. Furthermore, Except HMF (ID = 13.6 ± 2.0 mm), no other extract was active against *C. albicans*. This resistance of *C. albicans* compared to bacteria could be ascribed to their membrane composition which is different to those of bacteria. In fact, the higher number of anionic phospholipids

in the membrane of bacteria ease their interaction with antimicrobial compounds and thus increase their sensitivity (Oren et al., 1997; Papo and Shai, 2003). Moreover, flowers of *H. mantegazzianum* exhibited the lowest MIC (0.5 mg/ml) and the lowest MBC (1 and 4 mg/ml) against both *S. aureus* and *E. coli*. The antimicrobial activity of *H. mantegazzianum* can be ascribed to compounds, such as octyl acetate hexyl 2-methylbutyrate, hexyl isobutyrate, and hexyl butyrate, which has been identified as its dominant constituent (Matoušková et al., 2019). Similarly, the activity of *C. album* leaves against *S. aureus* can be ascribed to flavonoid, saponin, cinnamic acid amide, alkaloid chenoalbicin, apocarotenoids, and phenols (Cutillo et al., 2004).

The antimicrobial activity varies widely from plant to plant, and it is necessary to point out that, the antimicrobial activity observed in the present study may be higher (Akinsulire et al., 2018), lower (Romulo et al., 2018), or even equal to that of extracts from other plants, medicinal or not. However, the only advantage that these plants have over some medicinal plants is their invasive aspect which makes them easily available (Máximo et al., 2020) compared to certain seasonal medicinal plants such as *Enantia chlorantha* (Etame et al., 2020), *Aesculus hippocastanum* (Konstantinovitch et al., 2022), *Anthonotha macrophylla* (Essiet et al., 2019), and many others. Notwithstanding this, according to the classification established by Kuete (2010) and Kuete and Efferth (2010), except flowers of *H. mantegazzianum*, the other extracts could be considered as deserving a weak antimicrobial activity independently of the tested strain as they scored MIC value higher than 0.625 mg/ml. In addition, it is established in the literature that an antimicrobial compound is considered as bactericidal/fungicidal against a microbial strain when the ratio MBC/MIC or MFC/MIC is ≤ 4 (Oussou et al., 2008; Teke et al., 2011). Based on this classification, the ethanolic extract of the plants studied in the present work can be considered as bactericidal against *S. aureus*. The ethanolic extract of *C. jacea* L. flower and HMF can be considered as bactericidal against *E. coli*. Only the ethanolic extract of HMF can be considered as fungicidal against *C. albicans*. It is important to highlight that the parts of the plants tested were not chosen arbitrarily; the unavailability of the parts not tested (*C. jacea* leaves and *C. album* flowers) during the collection period led to their exclusion from the study. Thus, these other parts could also be tested and could potentially exhibit antimicrobial activity similar to the tested parts. Moreover, we evaluated the synergy between common antibiotics and the ethanolic extracts of the plants tested in this study. The use of combination therapy has been suggested as a new approach to improve the efficacy of antimicrobial agents by screening crude extracts from medicinal plants with good indications for use in combination with antibiotics (Ngongang et

al., 2020). As observe in Table 3, no antagonism ($FIC > 4$) or indifference ($1 \leq FIC \leq 4$) was noted between the extracts and the antibiotics. However, we found an additional effect ($0.5 \leq FIC \leq 1$) in almost all the combinations antibiotics + extracts excepted for HMF + (KA or AMP) against *S. aureus* ATCC 6538 and CJF + AMP against *E. coli* where we found synergy effect (FIC) ($FIC \leq 0.5$). The lower is the FIC index, better is the synergy (Ngongang et al., 2020). The best synergies were found with HMF which well-modulated AMP (FIC = 0.250 against *S. aureus*) and KA (FIC = 0.375 against *S. aureus*). CJF also well-modulated AMP against *E. coli* with FIC = 0.375.

Finally, the larval form of *G. mellonella* was used to assess the acute toxicity of the plant extracts tested. This model has been suggested by several researchers as an ecofriendly *in vivo* approach for toxicity studies (Megaw et al., 2015; Ignasiak and Maxwell, 2017). The LD50s of HMF and leaves, CAL and CJF were 20.2, 0.58, 13.2 and 4.0 mg/ml respectively. The lower is the LD50, more toxic is the sample tested. Consequently, HML and CJF can be considered as the most toxic extracts to *G. mellonella*. In addition, according to the classification of Gosselin, Smith and Hodge (CCOHS et al., 2020), biologically active compounds can be super toxic (LD50 < 5 mg/kg bw), extremely toxic [LD50 \in (5–50 mg/kg bw)], very toxic [LD50 \in (50–500 mg/kg bw)], moderately toxic [LD50 \in (0.5–5 g/kg bw)], slightly toxic [LD50 \in (5–15 g/kg bw)], and practically non-toxic (above 15 g/kg). Therefore, the data from our study indicate that ethanolic extract of *H. mantegazzianum* leaves (0.27 g/kg bw) is very toxic, the ethanolic extract from CJF (2.28 g/kg bw) is moderately toxic, and the ethanolic extract of HMFs (11.45 g/kg bw) and CAL (6.94 g/kg bw) are slightly toxic. Plants like HMFs may be recommended for further studies in the search for new antimicrobials because their toxic doses are far greater than the doses which have shown their optimal antimicrobial activity.

Conclusion

The present study aimed to assess the antimicrobial activity of some invasive plants such as leaves and flowers of *H. mantegazzianum* Sommier and Levier, flowers of *C. jacea* L., and leaves of *C. album* L. All plants showed great activity when they were associated with antibiotics such as NIT, KA, and AMP. However, when used alone, only HMFs exhibited noteworthy antimicrobial activity against all the microorganisms tested. All other plants were only active against *S. aureus* and exhibited low antimicrobial activity against *E. coli*. It can be concluded that only HMFs have worthy antimicrobial effects, but further preclinical and clinical trials are required to evaluate the composition, the cytotoxicity and safety issues of these plant extracts and their combinations with common antibiotics before they can be recommended for antimicrobial therapy.

Author's contributions

MMJA designed the research study. MMJA, AKLD, KP, and MR performed the research. MMJA analyzed the data. SAN, MMJA, AKLD, MKM, GVS, KZS, CZA, DS, and PIV wrote the manuscript. All authors contributed to editorial changes in the manuscript. All authors read and approved the final manuscript. "

Acknowledgments

This study has been supported by the RUDN University Strategic Academic Leadership Program.

Funding

This research received no external funding.

Conflict of interest

The authors declare that there is no conflict of interest.

References

- Acosta-Gutiérrez, S., Bodrenkoc, I. and Ceccarellia, M. 2020. Permeability through bacterial porins dictates whole cell compound accumulation. ChemRxiv. Preprint. doi:10.26434/chemrxiv.11877834.v1.
- Akinsulire, O.R., Aibin, I.E., Adenipekun, T., Adelowotan, T. and Odugbemi, T. 2007. In vitro antimicrobial activity of crude extracts from plants *Bryophyllum pinnatum* and *Kalanchoe crenata*. Afr. J. Tradit. Complement Altern. Med. 4(3), 338–344.
- Amer, M.W. and Awwad, A. 2021. Green synthesis of copper nanoparticles by *Citrus limon* fruits extract, characterization, and antibacterial activity. Chem. Int. 7, 1–8.
- Andersson, D.I., Hughes, D. and Kubicek-Sutherland, J.Z. 2016. Mechanisms, and consequences of bacterial resistance to antimicrobial peptides. Drug Resist. Updat. 26, 43–57.
- Ansald, M., Boulanger, P., Brives, C., Debarbieux, L., Dufour, N., Froissart, R., Gandon, S., Hénaff, C.L., Petit, M.A., Rocha, E., and Torres-Barceló, C. 2020. Les applications antibactériennes des bactériophages. Virologie 24(1), 23–36.
- Arsene, M.M.J., Davares, A.K.L., Andreevna, S.L., Vladimirovich, E.A., Carime, B.Z., Marouf, R. and Khelifi, I. 2021a. The use of probiotics in animal feeding for safe production and as potential alternatives to antibiotics. Vet. World. 14(2), 319–328.
- Arsene, M.M.J., Podoprighora, I.V., Davares, A.K.L., Razan, M., Das, M.S. and Senyagin, A.N. 2021b. Antibacterial activity of grapefruit peel extracts and green-synthesized silver nanoparticles. Vet. World. 14(5), 1330–1341.
- Arsene, M.M.J., Jorelle, A.B., Sarra, S., Viktorovna, P.I., Davares, A.K.L., Ingrid, N.K.C., Steve, A.A.F., Andreevna, S.L., Vyacheslavovna, Y.N. and Carime, B. Z. 2022. Short review on the potential alternatives to antibiotics in the era of antibiotic resistance. J. Appl. Pharm. Sci. 12(1), 29–40.
- Arsene, M.M.J., Viktorovna, P.I., Davares, A.K.L., Esther, N. and Nikolaevich, S.A. 2021c. Urinary

- tract infections: virulence factors, resistance to antibiotics, and management of uropathogenic bacteria with medicinal plants—a review. *J. Appl. Pharm. Sci.* 11(7), 1–12.
- Arsene, M.M.J., Viktorovna, P.I. and Davares, A.K.L. 2021d. *Galleria mellonella* (greater wax moth) as an eco-friendly in vivo approach for the assessment of the acute toxicity of medicinal plants: application to some plants from Cameroon. *Open. Vet. J.* 11(4), 651–661.
- Arsene, M.M.J., Viktorovna, P.I., Grigorievna, V.E., Davares, A.K.L., Sergeevna, D.M. and Nikolaevna, S.I. 2021e. Prolonged exposure to antimicrobials induces changes in susceptibility to antibiotics, biofilm formation and pathogenicity in *Staphylococcus aureus*. *J. Pharm. Res. Int.* 33(34B), 140–151.
- Bewley, J.D. 1973. Desiccation and protein synthesis in the moss *Tortilla ruralis*. *Can. J. Bot.* 51(1), 203–206.
- CCOHS (Canadian Centre for Occupational Health and Safety Act). What is a LD50 and LC50? Available via <https://www.ccohs.ca/oshanswers/chemicals/ld50.htm> (Accessed 2 September 2021).
- Cutillo, F., D'Abrosca, B., DellaGreca, M. and Zarrelli, A. 2004. Chenoalbicin, a novel cinnamic acid amide alkaloid from *Chenopodium album*. *Chem. Biodivers.* 1(10), 1579–1583.
- Elekhrawy, E., Sonbol, F., Abdelaziz, A. and Elbanna, T. 2020. Potential impact of biocide adaptation on selection of antibiotic resistance in bacterial isolates. *Future J. Pharm. Sci.* 6(1), 1–10.
- Eloff, J.N. 2019. Avoiding pitfalls in determining antimicrobial activity of plant extracts and publishing the results. *BMC. Complement. Altern. Med.* 19(1), 1–8.
- Etame, R.M.E., Mouokeu, R.S., Poundeu, F.S.M., Voukeng, I.K., Cidjeu, C.L.P., Tiabou, A.T. Yaya A.J.G., Ngane R.A.N., Kuate J.R., and Etoa, F.X. 2019. Effect of fractioning on antibacterial activity of n-butanol fraction from *Enantia chlorantha* stem bark methanol extract. *BMC. Complement. Altern. Med.* 19(1), 1–7.
- Essiet, G.A., Anwankwo, M.U., Akuodor, G.C., Ajoku, G.A., Offor, C.C., Megwas, A.U. and Aja, D.O.J. 2019. Antibacterial and toxicological evaluation of the ethanol leaf extract of *Anthonotha macrophylla*. *J. Herb. Med. Pharmacol.* 8(3), 205–211.
- Havel, J.E., Kovalenko, K.E., Thomaz, S.M., Amalfitano, S. and Kats, L.B. 2015. Aquatic invasive species: challenges for the future. *Hydrobiologia* 750(1), 147–170.
- Ignasiak, K. and Maxwell, A. 2017. *Galleria mellonella* (greater wax moth) larvae as a model for antibiotic susceptibility testing and acute toxicity trials. *BMC. Res. Notes.* 10(1), 1–8.
- Konstantinovich, K.Y., Arsene, M.M.J., Aliya, M.V., Viktorovna, P.I., Elena, V.G., Azova, M.M., and Amira, A.A. 2022. Assessment of antimicrobial activity of ethanolic and aqueous extracts of *Aesculus hippocastanum* L. (Horse Chestnut) bark against bacteria isolated from urine of patients diagnosed positive to urinary tract infections. *Front. Biosci. (Schol Ed)*, 14(2), 11.
- Kuete, V. 2010. Potential of Cameroonian plants and derived products against microbial infections: a review. *Planta. Med.* 76, 1479–1491.
- Kuete, V. and Efferth, T. 2010. Cameroonian medicinal plants: pharmacology and derived natural products. *Front. Pharmacol.* 1, 123.
- Matoušková, M., Jurová, J., Grušová, D., Wajs-Bonikowska, A., Renčo, M., Sedlák, V., Poráčová, J., Gogařová, Z. and Kalemba, D. 2019. Phytotoxic effect of invasive *Heracleum mantegazzianum* essential oil on dicot and monocot species. *Molecules* 24(3), 425.
- Máximo, P., Ferreira, L.M., Branco, P.S. and Lourenço, A. 2020. Invasive plants: turning enemies into value. *Molecules* 25(15), 3529.
- Megaw, J., Thompson, T.P., Lafferty, R.A. and Gilmore, B.F. 2015. *Galleria mellonella* as a novel in vivo model for assessment of the toxicity of 1-alkyl-3-methylimidazolium chloride ionic liquids. *Chem* 139, 197–201.
- Mbarga, M.J.A., Desobgo, S.C.Z., Tatsadjieu, L.N., Kavhiza, N. and Kalisa, L. 2021a. Antagonistic effects of raffia sap with probiotics against pathogenic microorganisms. *Foods. Raw. Mater.* 9(1), 24–31.
- Mondal, A.H., Yadav, D., Mitra, S. and Mukhopadhyay, K. 2020. Biosynthesis of silver nanoparticles using culture supernatant of shewanella sp. Ary1 and their antibacterial activity. *Int. J. Nanomed.* 15, 8295.
- Mouafo, H.T., Tchuenchieu, A.D.K., Nguedjo, M.W., Edoun, F.L.E., Tchunte, B.R.T. and Medoua, G.N. 2021. In vitro antimicrobial activity of *Millettia laurentii* De Wild and *Lophira alata* Banks ex CF Gaertn on selected foodborne pathogens associated to gastroenteritis. *Heliyon* 7(4), e06830.
- Ngongang, F.C., Fankam, A.G., Mbaveng, A.T., Wamba, B.E., Nayim, P., Beng, V.P. and Kuete, V. 2020. Methanol extracts from *Manilkara zapota* with moderate antibacterial activity displayed strong antibiotic-modulating effects against multidrug-resistant phenotypes. *Pharmacology* 3(1), 37.
- Oren, Z., Hong, J. and Shai, Y. 1997. A repertoire of novel antibacterial diastereomeric peptides with selective cytolytic activity. *J. Biol. Chem.* 272, 14643–14649.
- Oussou, K.R., Kanko, C., Guessend, N., Koukoua, G., Dosso, M., N'Guessan, Y.T., Figueredo, G., and Chalchat, J.C. 2008. Activités antibiotiques des huiles essentielles de trois plantes aromatiques de Côte d'Ivoire. *C. R. Chim.* 7, 1081–1086.
- Papo, N. and Shai, Y. 2003. Can we predict biological activity of antimicrobial peptides from their

- interactions with model phospholipid membranes? *Peptides* 24(11), 1693–1703.
- Pimentel, D., Zuniga, R. and Morrison, D. 2005. Update on the environmental and economic costs associated with alien-invasive species in the United States. *Ecol. Econ.* 52(3), 273–288.
- Pintó-Marijuan, M. and Munné-Bosch, S. 2013. Ecophysiology of invasive plants: osmotic adjustment and antioxidants. *Trends. Plant. Sci.* 18(12), 660–666.
- Romulo, A., Zuhud, E.A.M., Rondevaldova, J. and Kokoska, L. 2018. Screening of in vitro antimicrobial activity of plants used in traditional Indonesian medicine. *Pharm. Biol.* 56(1), 287–293.
- Schaenzer, A.J. and Wright, G.D. 2020. Antibiotic resistance by enzymatic modification of antibiotic targets. *Trends. Mol. Med.* 26(8), 768–782.
- Singh, A., Gautam, P.K., Verma, A., Singh, V., Shivapriya, P.M., Shivalkar, S., Sahoo, A.K. and Samanta, S.K. 2020. Green synthesis of metallic nanoparticles as effective alternatives to treat antibiotics resistant bacterial infections: a review. *Biotechnol. Rep.* 25, e00427.
- Souadkia, S., Podoprigora, I.V., Yashina, N.V., Sarukhanova, L.E. and Kravtsov, E.G. 2020. Antibiotic-resistant uropathogenic *Escherichia coli* isolated from children with birth defects of the urinary system. *Antibiot. Chemother.* 65(7–8), 23–26.
- Su, T., Qiu, Y., Hua, X., Ye, B., Luo, H., Liu, D., Qu, P. and Qiu, Z. 2020. Novel opportunity to reverse antibiotic resistance: to explore traditional Chinese medicine with potential activity against antibiotics-resistance bacteria. *Front. Microbiol.* 11, 3372.
- Teke, G.N., Kuate, J.R., Kuete, V., Teponno, R.B., Tapondjou, L.A., Tane, P., Giacinti, G. and Vilarem, G. 2011. Bio guided isolation of potential antimicrobial and antioxidant agents from the stem bark of *Trilepisium madagascariense*. *South. Afr. J. Bot.* 77, 319–327.
- Trabelsi, A., El Kaibi, M.A., Abbassi, A., Horchani, A., Chekir-Ghedira, L. and Ghedira, K. 2020. Phytochemical study and antibacterial and antibiotic modulation activity of *Punica granatum* (pomegranate) leaves. *Scientifica* 2020, 8271203.
- Wojnicz, D., Kucharska, A.Z., Sokół-Łętowska, A., Kicia, M. and Tichaczek-Goska, D. 2012. Medicinal plants extracts affect virulence factors expression and biofilm formation by the uropathogenic *Escherichia coli*. *Urol. Res.* 40(6), 683–697.
- World Health Organization (WHO). 2019. New report calls for urgent action to avert antimicrobial resistance crisis. Geneva, Switzerland: World Health Organization. Available via <https://www.who.int/news/item/29-04-2019-new-reportcalls-for-urgent-action-to-avert-antimicrobial-resistancecrisis> (Accessed 24 May 2021).