

## Thietanes

## Consecutive and Selective Double Methylene Insertion of Lithium Carbenoids to Isothiocyanates: A Direct Assembly of Four-Membered Sulfur-Containing Cycles

Raffaele Senatore, Monika Malik, Thierry Langer, Wolfgang Holzer, and Vittorio Pace\*


**Abstract:** A formal  $\text{CH}_2\text{-CH}_2$  homologation conducted with C1 carbenoids on a carbon electrophile for the obtainment of a four-membered cycle is reported. The logic proposes the consecutive delivery of two single nucleophilic  $\text{CH}_2$  units to an isothiocyanate—as competent electrophilic partner—resulting in the assembling of a rare imino-thietane cluster. The single synthetic operation procedure documents genuine chemocontrol, as indicated by the tolerance to various reactive elements decorating the starting materials. Significantly, the double homologation protocol is accomplished directly on a carbon electrophile, thus not requiring the installation of heteroatom-centered manifolds (e.g. boron).

The progressive formal insertion of a constant methylene ( $\text{CH}_2$ ) unit into an organic array is known as homologation and, ultimately enables the modular assembly of complex skeletons,<sup>[1]</sup> thus mimicking fundamental biosynthetic pathways.<sup>[2]</sup> Regardless the essence of the adopted regime (nucleophilic, electrophilic or radical), the carbon skeleton elongation is limited to one methylene unit *per* homologation cycle, thus making multiple-homologations highly elusive processes in synthesis (Scheme 1—*path a*).<sup>[3]</sup> Although the intrinsic limited instability of reagents used for the purpose—for example, metal carbenoids<sup>[4]</sup>—may account for this reluctant behaviour, the installation of valuable linchpins on the chain results an effective solution to the problem. In this sense, the introduction by Matteson of boronic esters (**I**) as privileged electrophilic manifolds to conduct homologation with lithium carbenoids,<sup>[5]</sup> underpinned the elaboration of the elegant concept of iterative homologations illustrated more recently by Aggarwal<sup>[6]</sup> and Blakemore (Scheme 1—*path b*).<sup>[7]</sup>

How to cite: *Angew. Chem. Int. Ed.* **2021**, *60*, 24854–24858

International Edition: doi.org/10.1002/anie.202110641

German Edition: doi.org/10.1002/ange.202110641



**Scheme 1.** General context of the presented work (HMLGT=homologation).

Notably, this logic features remarkable flexibility and robust stereocontrol thus, enabling the use of (different) various C1-units as competent donors, as documented in the elaboration of acyclic complex natural products.<sup>[8]</sup> Notwithstanding, analogous elongations directly conducted on *carbon*-centered manifolds remain underdeveloped and of limited preparative significance, regardless the hybridization of the competent carbon ( $sp^3$  and  $sp^2$ ).<sup>[9]</sup> In recent years, our group developed modular homologations of *C*-electrophiles with reactive lithium carbenoids ( $\text{LiCH}_2\text{X}$ ) enabling the assembly of sophisticated structures, as a consequence of molecular rearrangements triggered by the fine tuning of reaction conditions.<sup>[3,10]</sup> In this context, we documented the halogen-driven selective homologations of trifluoroacetimidoyl chlorides (TFAIC, **I**) to *gem*-chloro-trifluoromethylaziridines (**II**) or *gem*-halomethyl-trifluoromethylaziridines (**III**)<sup>[11]</sup> and bis-trifluoromethyl- $\beta$ -diketiminates (**IV**)<sup>[12]</sup> respectively, highlighting the critical role played by the reaction stoichiometry

\*] Dr. R. Senatore, M. Sc. M. Malik, Prof. T. Langer, Prof. W. Holzer, Prof. V. Pace

University of Vienna, Department of Pharmaceutical Sciences  
Althanstrasse, 14 A-1090, Vienna (Austria)

E-mail: vittorio.pace@univie.ac.at

Homepage: <http://drugssynthesis.univie.ac.at/>

Prof. V. Pace

University of Turin, Department of Chemistry  
Via P. Giuria 7, 10125 Turin (Italy)

E-mail: vittorio.pace@unito.it

Supporting information and the ORCID identification number(s) for the author(s) of this article can be found under:  
<https://doi.org/10.1002/anie.202110641>.


© 2021 The Authors. Angewandte Chemie International Edition published by Wiley-VCH GmbH. This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

in discriminating between C1 and C1-C1 insertions (Scheme 1—*path c*). Conceptually, these precedents could be ascribed to the well-established carbonyl-oxirane (and related imine-aziridine) homologative chemistry:<sup>[13]</sup> this is, the cyclic cluster is constructed via the incorporation of a *single* C1 unit delivered by the carbenoid donor.<sup>[14]</sup> We wondered if homologation tactics with C1 carbenoids could promote the assembly of four-membered cycles<sup>[15]</sup> through the consecutive release of *two* methylene fragments on a proper electrophile. Crucial for the successful reaction design was ensuring that the homologation sequence could reach completion within the short lifetime of the carbenoid used.<sup>[16]</sup> To this end, we conceived to generate a *first* homologation reactive intermediate susceptible of rapid *second* CH<sub>2</sub>-insertion, furnishing the targeted four-membered cycle.<sup>[17]</sup> Ideally, an electrophilic platform fulfilling this criterion could be represented by an isothiocyanate (**V**): in fact, the formation of a nucleophilic thioamidate anion (**VI**)—after the first homologative event—would induce a fast cyclization to a highly reactive thiiranimine (**VII**),<sup>[18]</sup> which would then undergo the second homologation. Finally, the additional ring closure of the elongated thioamidate (**VIII**) presenting the inserted CH<sub>2</sub>-CH<sub>2</sub> motif would result in the thietane scaffold (**IX**) presenting the *exo* imine-appendix (Scheme 1—*path d*).

Collectively, our synthetic strategy would construct the rare imino-thietane skeleton through a conceptually intuitive logic starting from commercially available (or easily accessible) isothiocyanates. We anticipate the proposed direct approach conducts to the imino-thietane core presenting an integral CH<sub>2</sub>-CH<sub>2</sub> linkage. To the best of our knowledge, such a structural characteristic is unprecedented among the scattered imino-thietanes previously described (Scheme 1—*path e*). Indeed, uniformly—the recyclization of  $\beta$ -lactones (Mulzer),<sup>[19]</sup> the cycloaddition of ketenimines to thioketones (Dondoni),<sup>[20]</sup> the cycloaddition of vinyl ethers with sulfonyl isothiocyanates (Schaumann),<sup>[21]</sup> and, the photochemical isomerization of *N*-monosubstituted  $\alpha,\beta$ -unsaturated thioamides (Sakamoto)<sup>[22]</sup>—afforded imino-thietanes variously substituted on the C-C linkage.<sup>[23]</sup>

4-Ethoxyisothiocyanate **1** was selected as the model substrate for the reaction optimization (Table 1). Upon treatment with the carbenoid chloromethylithium (LiCH<sub>2</sub>Cl), only a complex mixture was obtained, even at very short reaction times (entries 1,2).<sup>[24]</sup> Because of the absence in the mixture of the product resulting from the attack of MeLi-LiBr to the electrophilic isothiocyanate (i.e. thioamide **3**), it could be assumed that the carbenoid was effectively generated. The switching to the highly unstable LiCH<sub>2</sub>F<sup>[25]</sup> enabled to assemble the targeted four-membered cycle **2**, thus validating the initial working hypothesis (entry 3). It was essential to quench the reaction after only 5 min (from the end of the addition of MeLi-LiBr) to ensure a clean crude, thus confirming the high reactivity of intermediates conceived during the process design (entries 4,5). Notably, the overall transformation is in agreement with the observation by J. Hu on the suitability of CH<sub>2</sub>-F fragments as manifolds for nucleophilic substitutions.<sup>[26]</sup> Additional improvement was achieved employing LiCH<sub>2</sub>Br, which served as a competent C1 delivery agent for constructing

**Table 1:** Reaction optimization.



EG = exchangeable group (I, Br)  
X = Cl, F, Br, I

Entry	Carbenoid precursor (equiv) <sup>[a]</sup>	Metalating agent	Reaction time [min]	Isolated yield of <b>2</b> [%]
1	ClCH <sub>2</sub> I (3.0)	MeLi-LiBr (2.8)	5	Compl. mix.
2	ClCH <sub>2</sub> I (3.0)	MeLi-LiBr (2.8)	60	Compl. mix.
3 <sup>[b]</sup>	FCH <sub>2</sub> I (3.0)	MeLi-LiBr (2.8)	5	69
4 <sup>[b]</sup>	FCH <sub>2</sub> I (3.0)	MeLi-LiBr (2.8)	15	45
5 <sup>[b]</sup>	FCH <sub>2</sub> I (3.0)	MeLi-LiBr (2.8)	60	Compl. mix.
6	BrCH <sub>2</sub> I (3.0)	MeLi-LiBr (2.8)	5	82
7	BrCH <sub>2</sub> I (3.0)	MeLi-LiBr (2.8)	15	54
8	I <sub>2</sub> CH <sub>2</sub> (3.0)	MeLi-LiBr (2.8)	5	71
9 <sup>[c]</sup>	Br <sub>2</sub> CH <sub>2</sub> (3.0)	MeLi-LiBr (2.8)	5	65
10 <sup>[d]</sup>	BrCH <sub>2</sub> I (3.0)	MeLi-LiBr (2.8)	5	Compl. mix.
11 <sup>[e]</sup>	BrCH <sub>2</sub> I (3.0)	MeLi-LiBr (2.8)	5	9
12 <sup>[f]</sup>	BrCH <sub>2</sub> I (3.0)	MeLi-LiBr (2.8)	5	Traces
13	BrCH <sub>2</sub> I (6.0)	MeLi-LiBr (5.8)	5	80
14 <sup>[g]</sup>	BrCH <sub>2</sub> I (1.3)	MeLi-LiBr (1.1)	5	27
15	BrCH <sub>2</sub> I (2.8)	<i>i</i> -PrMgCl-LiCl	5	No reaction
16 <sup>[d]</sup>	BrCH <sub>2</sub> I (2.8)	<i>i</i> -PrMgCl-LiCl	5	No reaction

[a] LiCH<sub>2</sub>X reagents were generated under Barbier conditions. [b] LiCH<sub>2</sub>F was generated in a 1:1 *v/v* mixture of THF and Et<sub>2</sub>O (ref. 25).

[c] Thioamide **3** was also formed in 21% isolated yield. [d] Reaction run at -40°C. [e] Reaction run in Et<sub>2</sub>O. [g] Reaction run in toluene. [g] No mono-homologated product (type **VII**—Scheme 1) was detected.

the thietane backbone **2** (entries 6) in an excellent 82% isolated yield. Confirming the pattern observed with LiCH<sub>2</sub>F (entries 4,5), also the reaction with the bromo carbenoid benefited from brief reaction time (entry 6 vs. 7), presumably because of avoiding the triggering of deleterious carbenoids decomposition events. Although presenting a better halogen leaving-group, the use of LiCH<sub>2</sub>I resulted in slightly dwindled yield (entry 8) presumably due to its tendency to act also as metalating agent.<sup>[12]</sup> Some additional points merit mention: a) in line with our previous investigations, the optimal LiCH<sub>2</sub>Br precursor proved to be ICH<sub>2</sub>Br which undergoes a rapid and efficient I/Li exchange,<sup>[27]</sup> thus avoiding the detection of thioamide **2a**,<sup>[28]</sup> formed at significant extent in the presence of CH<sub>2</sub>Br<sub>2</sub> (entry 9); b) conducting the process at higher temperature resulted in untractable mixtures because of compromised chemical integrity of carbenoids (entry 10); c) running the consecutive homologation in more apolar solvents (diethyl ether, toluene—entries 11,12), dramatically decreased reaction yields; d) considerable excess of nucleophile did not boost the transformation (entry 13), while using quasi-stoichiometric loading dropped the yield (entry 14); e) no reactivity was displayed by less nucleophilic magnesium carbenoids, even at higher temperatures (entries 15,16).<sup>[29]</sup>

The unambiguous assignment of the imino-thietane backbone was ascertained via X-ray crystallographic analysis of compound **2** (Figure 1). The very small value of torsion angles C7-S1-C9-C8 (0.76°), C9-C8-C7-S1 (0.91°) and C1-N1-C7-S1 (1.86°) are diagnostic for the almost complete planarity of the





Figure 1. X-ray structural analysis of compound **2**.<sup>[35]</sup>

ring, thus differentiating it from a classical non-planar thietane characterized by a puckering angle of  $26^\circ$ .<sup>[30]</sup> The internal angles C9–C8–C7 ( $84.65^\circ$ ), S1–C9–C8 ( $87.65^\circ$ ), C7–S1–C9 ( $77.01^\circ$ ) alternating from the  $sp^3$  geometry and, S1–C7–C8 ( $95.27^\circ$ ) deviating from a canonical  $C-sp^2$  confirm the formation of the thietane ring. The analysis of bond lengths indicates that the S1–C9 bond ( $1.835 \text{ \AA}$ ) is significantly longer than an average  $C(sp^3)$ –S linkage ( $1.817 \text{ \AA}$ ),<sup>[31]</sup> but slightly shorter than the C–S bond ( $1.847 \text{ \AA}$ ) of the unsubstituted thietane.<sup>[30]</sup> The constructed C8–C9 bond ( $1.541 \text{ \AA}$ ) is in the average of the analogous  $\text{CH}_2$ – $\text{CH}_2$  linkage found in the simple thietane ( $1.549 \text{ \AA}$ ).<sup>[30]</sup>

Having established the optimal conditions for the transformation, we next studied the scope of the consecutive double  $\text{CH}_2$  insertion (Scheme 2). The presence of electron-donating groups—that is, ethers—on the aromatic nucleus of the isothiocyanates is uniformly tolerated, furnishing the corresponding imino-thietanes (**2–5**) in very good yields, including when the ethereal oxygen presented phenyl- (**6**) or trifluoro- (**7**) functionalities. Running the protocol on bigger scale (20 mmol) resulted in comparable efficiency (**2**). An acetal-substituted isothiocyanate, as exemplified by the benzo[*d*][1,3]dioxole structure (**8**), is also amenable for the protocol. Chalcogens such as sulfur—embodied as a thioether (**9**) or, selenium—in the case of the selenoether (**10**), also act as productive starting materials. Notably, the adopted lithiation conditions for the carbenoid genesis do not activate a concomitant Se/Li exchange thus, leaving untouched this functionality. Aside of an unsubstituted phenyl ring (**11**), the incorporation of halogen such as fluorine (**12**, **17**, **18**), bromine (**13**), chlorine (**14–17**) at different positions of the ring are compatible with the methodology and, their presence could be advantageously exploited for further elaboration (vide infra).  $sp^3$ -Hybridized hydrocarbons do not alter the efficiency of the technique (**19–21**), including when an *i*-propyl group (**22**) or the bicyclic indanyl system (**23**) were installed on the heterocumulene starting material. Preparing imino-thietanes with the proposed strategy proceeded with remarkable chemocontrol, as documented by isothiocyanates decorated with unsaturated residues: no concomitant carbenoid-mediated cyclopropanation<sup>[32]</sup> took place on a vinyl-substituted analogue (**24**), neither deprotonation of a terminal alkyne (**25**) despite the basic conditions. Furthermore, the chemical integrity of electrophilic substituents potentially suffering the attack of nucleophilic carbenoids was preserved, as deduced by the ester (**26**) and the nitrile (**27**). The




Scheme 2. Scope of the consecutive  $\text{CH}_2$ – $\text{CH}_2$  homologation.

methodology showed that the presence of nitrogen-centered substituents—in principle able to invalidate the correct carbenoid formation—was not detrimental: accordingly, *N,N*-dimethylamino- (**28**), morpholino- (**29**), phenyldiazo- (**30**) and azido- (**31**) bearing isothiocyanates were consecutively homologated to afford the corresponding four-membered sulfur-ring. Intriguingly, 1,4-diisothiocyanatobenzene was subjected to selective assembly of the cycle on a single NCS appendix (**32**). This same pattern was confirmed also with a considerable excess of carbenoid (6.0 equiv). Finally, the flexibility to heteroaromatic systems was documented with the 3-pyridyl derivative, which furnished the desired motif in comparable good yield (**33**). The effective release of

the two carbons backbone C–C from the carbenoid source was gathered conducting the experiment—*coeteris paribus*—with di-deuterated iodomethylithium ( $\text{LiCD}_2\text{I}$ ) conveniently prepared from the commercially available  $\text{CD}_2\text{I}_2$ .<sup>[10]</sup> The evidence for the ethylene fragment in **34** featuring full deuteration—that is,  $\text{CD}_2\text{-CD}_2$ —was evidently consistent with the envisaged rationale.

Eager to understand the reactivity of the assembled system, we conducted manipulations both on the aromatic ring (proceeding from the parent isothiocyanate) and on the four-membered cycle (Scheme 3). The Buchwald-Hartwig



amination<sup>[33]</sup> with piperidine on the bromo-analogue **13** cleanly furnished the corresponding product **35** in which the thioimidate system was preserved (*path a*). Cognizant of the ring strain of four-membered rings (e.g. thietane  $80.3 \text{ kJ mol}^{-1}$ ),<sup>[34]</sup> we devised the ring-opening with a nucleophile leveraging on the electrophilicity of the  $sp^2$ -carbon (*path b*). Thus, via the addition of a reducing hydride—accomplished with  $\text{LiAlH}_4$ — $\gamma$ -amino-mercaptanes **XII** were formed in high yield, as demonstrated by the straightforward synthesis of different analogues (**36–39**). The transformation can be rationalized through the formation of a tetrahedral intermediate (**X**), whose collapse entails the breaking of the NC-S bond giving the  $\gamma$ -N-phenyliminopropyl mercapto anion (**XI**) amenable for a second hydride-reduction *en route* to the final, not previously accessible, structures (**XII**). The experiment with labelled  $\text{LiAlD}_4$  (**37**) confirmed the attack of both nucleophiles at the initially thioimidate carbon. Additional evidence for the chemical integrity of the unsubstituted iminothietanes was deduced by attempting the ring opening under different conditions, thus resulting in no modification of the nuclei (see Supporting Information).

In summary, we have documented the installation of the ethylene unit ( $-\text{CH}_2-\text{CH}_2-$ )—a formal C2-homologation—onto isothiocyanates through the consecutive delivery of two methylene fragments furnished by the carbenoid bromomethylithium ( $\text{LiCH}_2\text{Br}$ ). The tactic enables the straightforward access to four-membered cyclic imino-thietanes through

a single synthetic operation. The overall strategy relies on two sequential homologative events commenced on the electrophilic  $sp$ -carbon of a competent isothiocyanate, resulting in the formation—upon ring-closure—of a highly reactive imino-thirane which then undergoes a second homologation giving a  $\gamma$ -halothioimidate direct precursor of the final imino-thietane. A remarkable degree of chemoselectivity was uniformly observed, as evidenced by running the reaction of isothiocyanates featuring additional sensitive functionalities such as halogens, seleno-, thio-ethers, amines or electrophilic groups susceptible of the attack of the nucleophilic carbenoid. The assembled four-membered imino-thioethane—whose structure was ascertained by X-ray analysis—may serve as a versatile platform for preparing  $\gamma$ -aminopropylthiols via hydride-mediated reduction. To the best of our knowledge this concept constitutes the first example of a controlled consecutive C1-homologation on carbon electrophiles with a carbenoid without the requirement of hetero-atom linchpins (e.g. boron).

## Acknowledgements

We thank the University of Vienna and the University of Turin for generous support. ABCR-Germany chemicals is acknowledged for the kind supply of  $\text{ICH}_2\text{F}$ . We thank Ing. A. Roller and MSc. N. Lukic for X-ray analysis. Open Access Funding provided by Universita degli Studi di Torino within the CRUI-CARE Agreement.

## Conflict of Interest

The authors declare no conflict of interest.

**Keywords:** carbenoids · chemoselectivity · homologation · isothiocyanates · thietanes

- [1] For reviews, see: a) J. J. Li, *Name Reactions for Homologation*, Wiley, Hoboken, **2009**; b) N. R. Candeias, R. Paterna, P. M. P. Gois, *Chem. Rev.* **2016**, *116*, 2937–2981.
- [2] a) R. W. Hoffmann, *Angew. Chem. Int. Ed. Engl.* **1987**, *26*, 489–503; *Angew. Chem.* **1987**, *99*, 503–517; b) C. Hertweck, *Angew. Chem. Int. Ed.* **2009**, *48*, 4688–4716; *Angew. Chem.* **2009**, *121*, 4782–4811.
- [3] L. Castoldi, S. Monticelli, R. Senatore, L. Ielo, V. Pace, *Chem. Commun.* **2018**, *54*, 6692–6704.
- [4] For reviews on carbenoid chemistry, see: a) G. Boche, J. C. W. Lohrenz, *Chem. Rev.* **2001**, *101*, 697–756; b) P. R. Blakemore, R. W. Hoffmann, *Angew. Chem. Int. Ed.* **2018**, *57*, 390–407; *Angew. Chem.* **2018**, *130*, 396–413; c) V. Capriati, S. Florio, *Chem. Eur. J.* **2010**, *16*, 4152–4162; d) V. Capriati in *Contemporary Carbene Chemistry* (Eds.: R. A. Moss, M. P. Doyle), Wiley, Hoboken, **2014**, pp. 327–362; e) V. Pace, W. Holzer, N. De Kimpe, *Chem. Rec.* **2016**, *16*, 2061–2076; f) V. H. Gessner, *Chem. Commun.* **2016**, *52*, 12011–12023.
- [5] a) D. S. Matteson, *J. Org. Chem.* **2013**, *78*, 10009–10023; b) D. S. Matteson, D. Majumdar, *J. Am. Chem. Soc.* **1980**, *102*, 7588–7590. Very recently controlled mono- and double homologations of N–B bonds have been disclosed, see: c) Q. Xie, G. Dong, *J. Am. Chem. Soc.* **2021**, *143*, 14422–14427.

- [6] M. Burns, S. Essafi, J. R. Bame, S. P. Bull, M. P. Webster, S. Balieu, J. W. Dale, C. P. Butts, J. N. Harvey, V. K. Aggarwal, *Nature* **2014**, *513*, 183–188.
- [7] a) P. R. Blakemore, M. S. Burge, *J. Am. Chem. Soc.* **2007**, *129*, 3068–3069; b) C. R. Emerson, L. N. Zakharov, P. R. Blakemore, *Chem. Eur. J.* **2013**, *19*, 16342–16356.
- [8] a) C. G. Watson, A. Balanta, T. G. Elford, S. Essafi, J. N. Harvey, V. K. Aggarwal, *J. Am. Chem. Soc.* **2014**, *136*, 17370–17373; b) S. Balieu, G. E. Hallett, M. Burns, T. Bootwicha, J. Studley, V. K. Aggarwal, *J. Am. Chem. Soc.* **2015**, *137*, 4398–4403; c) G. Casoni, M. Kucukdisli, J. M. Fordham, M. Burns, E. L. Myers, V. K. Aggarwal, *J. Am. Chem. Soc.* **2017**, *139*, 11877–11886; d) J. Wu, P. Lorenzo, S. Zhong, M. Ali, C. P. Butts, E. L. Myers, V. K. Aggarwal, *Nature* **2017**, *547*, 436–440.
- [9] The homologation of R-X for preparing R-CH<sub>2</sub>-X is an uncontrolled process which may result in up to four CH<sub>2</sub>-insertions, see: a) R. Huisgen, U. Burger, *Tetrahedron Lett.* **1970**, *11*, 3053–3056; b) R. C. Hahn, J. Tompkins, *Tetrahedron Lett.* **1990**, *31*, 937–940; Knochel demonstrated that exclusively allylic iodides react with mixed Zn/Cu C1 carbenoids giving the corresponding homoallylic iodides, see: c) P. Knochel, T. S. Chou, H. G. Chen, M. C. P. Yeh, M. J. Rozema, *J. Org. Chem.* **1989**, *54*, 5202–5204; d) P. Knochel, N. Jeong, M. J. Rozema, M. C. P. Yeh, *J. Am. Chem. Soc.* **1989**, *111*, 6474–6476; e) A. Sidduri, P. Knochel, *J. Am. Chem. Soc.* **1992**, *114*, 7579–7581; f) A. Sidduri, M. J. Rozema, P. Knochel, *J. Org. Chem.* **1993**, *58*, 2694–2713; For an excellent review on mixed Zn/Cu carbenoids, see: g) I. Marek, *Tetrahedron* **2002**, *58*, 9463–9475. For selective insertions of CHXY fragments into carbonyls followed by deoxygenation, see: M. Miele, A. Citarella, T. Langer, E. Urban, M. Zehl, W. Holzer, L. Ielo, V. Pace, *Org. Lett.* **2020**, *22*, 7629–7634.
- [10] V. Pace, L. Castoldi, E. Mazzeo, M. Rui, T. Langer, W. Holzer, *Angew. Chem. Int. Ed.* **2017**, *56*, 12677–12682; *Angew. Chem.* **2017**, *129*, 12851–12856.
- [11] a) L. Ielo, S. Touqeer, A. Roller, T. Langer, W. Holzer, V. Pace, *Angew. Chem. Int. Ed.* **2019**, *58*, 2479–2484; *Angew. Chem.* **2019**, *131*, 2501–2506; b) L. Ielo, V. Pillari, M. Miele, D. Castiglione, V. Pace, *Synlett* **2021**, *32*, 551–560.
- [12] L. Ielo, L. Castoldi, S. Touqeer, J. Lombino, A. Roller, C. Prandi, W. Holzer, V. Pace, *Angew. Chem. Int. Ed.* **2020**, *59*, 20852–20857; *Angew. Chem.* **2020**, *132*, 21038–21043.
- [13] For authoritative references, see: a) A. K. Yudin, *Aziridines and Epoxides in Organic Synthesis*, Wiley-VCH, Weinheim, **2006**; b) L. Degennaro, P. Trinchera, R. Luisi, *Chem. Rev.* **2014**, *114*, 7881–7929; c) G. S. Singh, M. D'Hooghe, N. De Kimpe, *Chem. Rev.* **2007**, *107*, 2080–2135.
- [14] a) V. K. Aggarwal, C. L. Winn, *Acc. Chem. Res.* **2004**, *37*, 611–620; b) M. D. Delost, J. T. Njardarson, *Org. Lett.* **2020**, *22*, 6917–6921. Formal -CH<sub>2</sub>CH<sub>2</sub>- homologations of epoxides to tetrahydrofuranes have been conducted with sulfur ylides, see: c) E. D. Butova, A. V. Barabash, A. A. Petrova, C. M. Kleiner, P. R. Schreiner, A. A. Fokin, *J. Org. Chem.* **2010**, *75*, 6229–6235. Under analogous conditions, the bis-homologation of carbonyls to oxetanes is known. For a recent example, see: d) P. Mukherjee, M. Pettersson, J. K. Dutra, L. Xie, C. W. am Ende, *ChemMedChem* **2017**, *12*, 1574–1577.
- [15] a) J. Xu, *Beilstein J. Org. Chem.* **2020**, *16*, 1357–1410; b) M. Mato, A. Franchino, C. García-Morales, A. M. Echavarren, *Chem. Rev.* **2021**, *121*, 8613–8684; c) S. R. Roy, H. Eijsberg, J. Bruffaerts, I. Marek, *Chem. Sci.* **2017**, *8*, 334–339.
- [16] For a study on the lifetime of carbenoids, see: P. Musci, M. Colella, A. Sivo, G. Romanazzi, R. Luisi, L. Degennaro, *Org. Lett.* **2020**, *22*, 3623–3627.
- [17] For ring-expansion processes, see: a) A. K. Clarke, W. P. Unsworth, *Chem. Sci.* **2020**, *11*, 2876–2881; b) J. R. Donald, W. P. Unsworth, *Chem. Eur. J.* **2017**, *23*, 8780–8799; c) T. C. Stephens, W. P. Unsworth, *Synlett* **2020**, *31*, 133–146.
- [18] a) G. L'Abbe, J. P. Dekerk, C. Martens, S. Toppet, *J. Org. Chem.* **1980**, *45*, 4366–4371; b) J. Xu in *Synthesis of 4- to 7-membered Heterocycles by Ring Expansion: Aza-, oxa- and thiaheterocyclic small-ring systems* (Eds.: M. D'hooghe, H.-J. Ha), Springer International Publishing, Cham, **2016**, pp. 311–361; c) J. Xu, *Asian J. Org. Chem.* **2020**, *9*, 1008–1017.
- [19] J. Mulzer, T. Kerkmann, *Angew. Chem. Int. Ed. Engl.* **1980**, *19*, 466–468; *Angew. Chem.* **1980**, *92*, 470–471.
- [20] a) A. Dondoni, A. Battaglia, P. Giorgianni, *J. Org. Chem.* **1980**, *45*, 3766–3773; b) A. Dondoni, A. Battaglia, P. Giorgianni, G. Gilli, M. Sacerdoti, *J. Chem. Soc. Chem. Commun.* **1977**, 43–44.
- [21] E. Schaumann, H.-G. Bäuch, G. Adiwidjaja, *Angew. Chem. Int. Ed. Engl.* **1981**, *20*, 613–615; *Angew. Chem.* **1981**, *93*, 600–601.
- [22] M. Sakamoto, T. Ishida, T. Fujita, S. Watanabe, *J. Org. Chem.* **1992**, *57*, 2419–2422.
- [23] The imino-thietane nucleus was obtained also as side product, see: a) T. Nishio, *J. Chem. Soc. Perkin Trans. 1* **1998**, 1007–1008; b) T. Nishio, I. Iida, K. Sugiyama, *J. Chem. Soc. Perkin Trans. 1* **2000**, 3039–3046.
- [24] a) V. Pace, L. Castoldi, W. Holzer, *Chem. Commun.* **2013**, *49*, 8383–8385. The different behavior of isothiocyanates compared to isocyanates is a consequence of the higher nucleophilicity of thioimides (type VI—Scheme 1) which smoothly cyclize to the thiiranium analogues. Whereas in the case of isocyanates the oxygen-centered addition intermediate does not possess enough nucleophilicity to trigger the cyclization event, thus furnishing the mono-homologated acyclic haloacetamide. For an overview of the reactivity of iso(thio)cyanates with organometallics, see: b) V. Pace, S. Monticelli, K. de la Vega-Hernandez, L. Castoldi, *Org. Biomol. Chem.* **2016**, *14*, 7848–7854. Other heterocumulenes (carbon dioxide) react through simple insertion of CH<sub>2</sub>X fragments (interrupted homologation), see: c) S. Monticelli, E. Urban, T. Langer, W. Holzer, V. Pace, *Adv. Synth. Catal.* **2019**, *361*, 1001–1006.
- [25] G. Parisi, M. Colella, S. Monticelli, G. Romanazzi, W. Holzer, T. Langer, L. Degennaro, V. Pace, R. Luisi, *J. Am. Chem. Soc.* **2017**, *139*, 13648–13651.
- [26] C. Ni, J. Hu, *Chem. Soc. Rev.* **2016**, *45*, 5441–5454.
- [27] V. Pace, A. Pelosi, D. Antermite, O. Rosati, M. Curini, W. Holzer, *Chem. Commun.* **2016**, *52*, 2639–2642.
- [28] V. Pace, L. Castoldi, S. Monticelli, S. Safranek, A. Roller, T. Langer, W. Holzer, *Chem. Eur. J.* **2015**, *21*, 18966–18970.
- [29] T. Kimura, *Synthesis* **2017**, *49*, 5105–5119.
- [30] K. Ken-ichi, K. Kozo, *Bull. Chem. Soc. Jpn.* **1975**, *48*, 1691–1695.
- [31] F. H. Allen, O. Kennard, D. G. Watson, L. Brammer, A. G. Orpen, R. Taylor, *J. Chem. Soc. Perkin Trans. 2* **1987**, S1–S19.
- [32] M. Colella, A. Tota, A. Großjohann, C. Carlucci, A. Aramini, N. S. Sheikh, L. Degennaro, R. Luisi, *Chem. Commun.* **2019**, *55*, 8430–8433.
- [33] P. A. Forero-Cortés, A. M. Haydl, *Org. Process Res. Dev.* **2019**, *23*, 1478–1483.
- [34] E. Block in *Comprehensive Heterocyclic Chemistry* (Eds.: A. R. Katritzky, C. W. Rees), Pergamon, Oxford, **1984**, pp. 403–447.
- [35] Deposition Number 2081745 (for **2**) contains the supplementary crystallographic data for this paper. These data are provided free of charge by the joint Cambridge Crystallographic Data Centre and Fachinformationszentrum Karlsruhe Access Structures service [www.ccdc.cam.ac.uk/structures](http://www.ccdc.cam.ac.uk/structures).

Manuscript received: August 8, 2021

Accepted manuscript online: September 17, 2021

Version of record online: October 13, 2021