



Case Report

Painful torticollis due to tubercular atlantoaxial rotatory fixation: A case report

Abolfazl Rahimizadeh, Walter Williamson, Shaghayegh Rahimizadeh, Mahan Amirzadeh

Pars Advanced and Minimally Invasive Medical Manners Research Center, Pars Hospital, Iran University of Medical Sciences, Tehran, Iran.

E-mail: *Abolfazl Rahimizadeh - a_rahimizadeh@hotmail.com; Walter Williamson - wlwilliamson@gmail.com; Shaghayegh Rahimizadeh - sherryrahimi@hotmail.com; Mahan Amirzadeh - mahan.amirzade@gmail.com



*Corresponding author:

Abolfazl Rahimizadeh,
Pars Advanced and Minimally
Invasive Medical Manners
Research Center, Pars Hospital,
Iran University of Medical
Sciences, Tehran, Iran.

a_rahimizadeh@hotmail.com

Received : 20 September 2020

Accepted : 25 November 2020

Published : 16 December 2020

DOI

10.25259/SNI_663_2020

Quick Response Code:



ABSTRACT

Background: Tubercular atlantoaxial, rotary dislocation warranting fixation (AARF) is an extremely rare event.

Case Description: AARF was suspected in a 23-year-old female with painful torticollis. When diagnostic studies documented unilateral destruction of the left lateral mass of the atlas, she underwent removal of the lateral mass, reduction of the deformity, and C1-C2 fusion/reconstruction utilizing an iliac bone graft. Laboratory tests and the pathologic surveys were all consistent with the diagnosis of underlying tuberculosis.

Conclusion: We present a case of tubercular atlantoaxial, rotary dislocation (AARF) in a patient who warranted C1-C2 decompression, reduction, and fusion.

Keywords: Atlantoaxial dislocation, Atlantoaxial rotatory fixation, Atlas, Craniovertebral junction, Tuberculosis, Upper cervical

INTRODUCTION

Tuberculosis (TB) involving the atlantoaxial complex is rare, accounting for approximately 0.1% of all spinal tubercular infections.^[1,3,5,11,14] Here, we present a young woman with painful torticollis attributed to tubercular atlantoaxial rotary dislocation (AARF).

CASE REPORT

A 23-year-old female presented with a 4-week history of severe neck pain (VAS:10) and torticollis with a classic “Cock Robin” deformity. Laboratory studies demonstrated an increased erythrocyte sedimentation rate (ESR) of 42. The lateral cervical plain radiographs and computed tomography (CT) axial, 2D, and 3D reconstructed images showed C1–C2 AARF with destruction of the left lateral mass of the atlas [Figures 1-4]. The integrity of the vertebral artery at C1–C2 was also critically confirmed on the preoperative CT angiogram (CTA) [Figure 5].

Surgery

As cervical traction failed to reduce the deformity, surgical intervention was warranted. Surgery required; the initial insertion of bilateral C2 pedicle screws, isolation of the V3

This is an open-access article distributed under the terms of the Creative Commons Attribution-Non Commercial-Share Alike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as the author is credited and the new creations are licensed under the identical terms.

©2020 Published by Scientific Scholar on behalf of Surgical Neurology International

segment of the vertebral artery, removal of the destroyed left C1 lateral mass in a piecemeal fashion, and fusion (e.g., utilizing a tricortical iliac bone graft secured with left C1 laminar hook-C2 pedicle screw, and an additional right C1 lateral mass-C2 pedicle screw rod construct) [Figure 6]. Postoperatively, the patient's torticollis and intractable pain resolved. X-rays taken 1 week, and 3 months after surgery demonstrated adequate alignment of the instrumentation/construct [Figure 7]. Three years later, the patient is asymptomatic [Figure 8].



Figure 1: Lateral cervical radiograph showing C1 foramen transversarium indicating possible rotation of atlas (while arrow).




Figure 2: Axial computed tomography scan showing destruction of the left C1 lateral mass, note approximation of the odontoid to the left lateral mass.




Figure 3: Reconstructed computed tomography scan (a) coronal, (b) sagittal showing destruction of the left lateral mass of atlas.

Bacteriology and pathology

The operative specimens demonstrated: a positive polymerase chain reaction (PCR) for TB, and the pathology was compatible with a granulomatous infection.

Tubercular treatment

Four-drug therapy was warranted for TB; isoniazid (5 mg/kg), rifampicin (10 mg/kg), ethambutol (15 mg/kg), and pyrazinamide (25 mg/kg). They were administered as a first-line of treatment for 4 months. This was followed by an additional 12 months of rifampicin and isoniazid.

DISCUSSION

Spinal TB, presenting as AARF with painful torticollis, is extremely rare.

Pathogenesis

Severe painful torticollis may be the only clinical indication that AARP is present. Patients may exhibit unilateral destruction of the lateral mass of the atlas with/without infiltration/disruption of the alar ligaments. The ESRs are typically increased, and the Mantoux test is typically positive.^[1-5,9,11,14] Further, acid-fast stains of pathological material, positive cultures, and PCR obtained through CT-guided biopsy may be additional valuable adjuncts to confirm the diagnosis of TB.^[1-5,9,11,14]

Imaging

AARF is the best documented utilizing axial reconstructed 2D and 3D CT images and CTA.^[12]

The following findings are classical for TB; an osteolytic, fragmented lesion involving the C1 lateral mass, deviation of the odontoid to the affected lateral mass side, and forward displacement of the contralateral lateral mass of atlas.^[4,6,7,10,13,15] Magnetic resonance imaging also may help to establish the diagnosis of tuberculous involvement of the C1 lateral mass,




Figure 4: 3D computed tomography scan of the atlantoaxial complex (a) frontal view showing destruction of the left lateral mass and a part of anterior ring of atlas, note forward displacement of the right C1 lateral mass. (b) Occipital view demonstrates rotation of the posterior ring of atlas to the right.




Figure 5: Reconstructed coronal computed tomography angiography shows integrity of the left V3 segment of the vertebral artery passing through the destroyed lateral mass.




Figure 6: Intraoperative photograph, (a) the lesion of the left lateral mass of atlas. (b) The site after removal of the mass. (c) The lateral mass is reconstructed with tricortical iliac autogenous bone graft, note C2 pedicle and C1 hook. (d) After assembling the rod.




Figure 7: Postoperative X-ray. (a) A week after surgery showing the C2 pedicle-C1 lateral mass and hook construct used properly. (b) At 1-year follow-up.



Figure 8: Photograph of the patient, a year after surgery while holding up her cervical radiographs.

by demonstrating heterogeneous intensity on the T1, and hyperintensity on the T2-weighted and fat-suppressed images.

Differential diagnosis

The differential diagnoses for painful torticollis with unilateral involvement of the C1 lateral mass include TB tumors, rheumatoid arthritis, and other types of pyogenic spondylitis.^[4,6,7,8,10,13,15]

Management

In classic tubercular atlantoaxial dislocation, management strategies range from purely conservative treatment to radical operations.^[1,5,11,14] Surgical intervention for decompression, reduction/realignment, and instrumented fusion may also be warranted.

CONCLUSION

Here, we presented an extremely rare cause of painful torticollis due to tubercular AARF involving a unilateral C1 lateral mass requiring decompression, reduction, and fixation.

Declaration of patient consent

The authors certify that they have obtained all appropriate patient consent.

Financial support and sponsorship

Nil.

Conflicts of interest

There are no conflicts of interest.

REFERENCES

- Allali F, Benomar A, El Yahyaoui M, Chkili T, Hajjaj-Hassouni N. Atlantoaxial tuberculosis: Three cases. *Joint Bone Spine* 2000;67:481-4.
- Behari S, Nayak SR, Bhargava V, Banerji D, Chhabra DK, Jain VK. Craniocervical tuberculosis: Protocol of surgical management. *Neurosurgery* 2003;52:72-81.
- Chaudhary K, Potdar P, Bapat M, Rathod A, Laheri V. Structural odontoid lesions in craniovertebral tuberculosis: A review of 15 cases. *Spine* 2012;37:E836-43.
- Dhaon BK, Jaiswal A, Nigam V, Jain V. Atlantoaxial rotatory fixation secondary to tuberculosis of occiput: A case report. *Spine* 2003;28:E203-5.
- Gupta SK, Mohindra S, Sharma BS, Gupta R, Chhabra R, Mukherjee KK, *et al.* Tuberculosis of the craniovertebral junction: Is surgery necessary? *Neurosurgery* 2006;58:1144-50.
- Halla JT, Bliznak J, Hardin JG, Finn S. Septic arthritis of the C1-C2 lateral facet joint and torticollis: Pseudo-grisel's syndrome. *Arthritis Rheum* 1991;34:84-8.
- Halla JT, Fallahi S, Hardin JG. Nonreducible rotational head tilt and lateral mass collapse. A prospective study of frequency, radiographic findings, and clinical features in patients with rheumatoid arthritis. *Arthritis Rheum* 1982;25:1316-24.
- Jain R, Sawhney S, Berry M. Computed tomography of vertebral tuberculosis: Patterns of bone destruction. *Clin Radiol* 1993;47:196-9.
- Lal AP, Rajshekhar V, Chandy MJ. Management strategies in tuberculous atlanto-axial dislocation. *Br J Neurosurg* 1992;6:529-35.
- Ngu BB, Khanna AJ, Pak SS, McCarthy EF, Sponseller PD. Eosinophilic granuloma of the atlas presenting as torticollis in a child. *Spine* 2004;29:E98-100.
- Pandya SK. Tuberculous atlanto-axial dislocation (with remarks on the mechanism of dislocation). *Neurol India* 1971;19:116-21.
- Rahimizadeh A, Williamson W, Rahimizadeh S. Traumatic chronic irreducible atlantoaxial rotatory fixation in adults: Review of the literature, with two new examples. *Int J Spine Surg* 2019;13:350-60.
- Simsek S, Yigitkanli K, Kazanci A, Belen D, Bavbek M. Medically treated paravertebral *Brucella* abscess presenting with acute torticollis: Case report. *Surg Neurol* 2007;67:207-10.
- Sinha S, Singh AK, Gupta V, Singh D, Takayasu M, Yoshida J. Surgical management and outcome of tuberculous atlantoaxial dislocation: A 15-year experience. *Neurosurgery* 2003;52:331-9.
- White GM, Healy WL. Tumor-associated atlanto-axial rotatory fixation: A case report. *Spine* 1987;12:406-8.

How to cite this article: Rahimizadeh A, Williamson W, Rahimizadeh S, Amirzadeh M. Painful torticollis due to tubercular atlantoaxial rotatory fixation: A case report. *Surg Neurol Int* 2020;11:440.